

Assignment 1

Dishank Jain - AI20BTECH11011

Download all python codes from

[https://github.com/Dishank422/AI1103-Probability
-and-random-variables/blob/main/
Assignment_1/codes](https://github.com/Dishank422/AI1103-Probability-and-random-variables/blob/main/Assignment_1/codes)

and latex-tikz codes from

[https://github.com/Dishank422/AI1103-Probability
-and-random-variables/blob/main/
Assignment_1/main.tex](https://github.com/Dishank422/AI1103-Probability-and-random-variables/blob/main/Assignment_1/main.tex)

On substituting the values of $\Pr(A)$, $\Pr(B)$ and $\Pr(A + B)$ in (2.0.7), we get

$$\frac{3}{5} = \frac{1}{2} + p - \frac{1}{2}p \quad (2.0.8)$$

$$\Rightarrow p = \frac{1}{5} \quad (2.0.9)$$

1 PROBLEM(PROB. MISC. 6.11)

Given that the events A and B are such that $\Pr(A) = \frac{1}{2}$, $\Pr(A + B) = \frac{3}{5}$ and $\Pr(B) = p$. Find p if they are

- i) mutually exclusive
- ii) independent

2 SOLUTION

- i) Since the events are mutually exclusive, by definition

$$\Pr(AB) = 0 \quad (2.0.1)$$

$$\Rightarrow \Pr(A + B) = \Pr(A) + \Pr(B) \quad (2.0.2)$$

On substituting the values of $\Pr(A)$, $\Pr(B)$ and $\Pr(A + B)$ in (2.0.2), we get

$$\frac{3}{5} = \frac{1}{2} + p \quad (2.0.3)$$

$$\Rightarrow p = \frac{1}{10} \quad (2.0.4)$$

- ii) Since the events are independent

$$\Pr(AB) = \Pr(A) \Pr(B) \quad (2.0.5)$$

We know

$$\Pr(A + B) = \Pr(A) + \Pr(B) - \Pr(AB) \quad (2.0.6)$$

$$\Rightarrow \Pr(A + B) = \Pr(A) + \Pr(B) - \Pr(A) \Pr(B) \quad (2.0.7)$$