





All about the High-Velocity SN Ia 2022hrs

¹Divya Mishra, ¹Peter J. Brown, ¹Gabriella Cruz, and others ¹Department of Physics and Astronomy, Texas A&M University



We present the first-ever UV spectra of a HV SN Ia, SN 2022hrs, observed with HST!

Despite their optical uniformity, Type Ia supernovae (SNe Ia) exhibit considerable diversity in the ultraviolet (UV) spectrum. This variation is particularly pronounced in "High-Velocity" (HV) SNe Ia, which have been shown to possess intrinsically different colors compared to their normal counterparts [1]. Understanding the cause of this divergence is crucial for refining SNe Ia as standard candles, addressing systematic uncertainties in intrinsic color and reddening estimates.







Figure 5 – Left Panel: HST Spectra of SN 2022hrs, -3 days before B-band maximum, corrected for different R_V and E(B-V), compared to the HST spectrum of SN 2011fe at the same phase. Right Panel: Same comparison with spectra of SN 2022hrs at +6 days post B-band maximum, compared to SN 2011fe at +9 days. Each spectrum is normalized by the V band wavelength range.






Figure 6 – Left Panel: HST Spectra of SN 2022hrs (-3 days before B-band maximum) compared with SN 2017erp (-2 days before Bband maximum) [4], SN 2021fxy (-2 days before B-band maximum) [5], and SN 2011fe (-3 days before B-band maximum) [6]. Each spectrum is normalized by the V band wavelength range.




Figure 1 – Photometry of SN 2022hrs from Swift/UVOT.

intrinsic color for SN 2022hrs.




Figure 2 - Observed uvw1-v and uvm2-uvw1 color comparison for SN 2022hrs, SN 2017erp, and SN 2011by..





Figure 3 – Temporal evolution of the expansion velocities of Si II λ6355 (blue circles) and Ca II H&K (pink squares).






Figure 4 – Spectral time series of SN 2022hrs. All the spectra are at an offset from one another. The solid gray line represents the wavelengths corresponding to 0, -10,000, and -20,000 km/s around the Si II, Ca II H&K, and Ca II NIR triplet features, with 2000 km/s intervals denoted by light gray lines.




Figure 7 – Top Panel: Na I D doublet absorption feature from the Milky Way. Right Panel: Na I D doublet absorption from the host galaxy, NGC 4647. Spectra were obtained at three epochs: +3 days, +4 days, and +27 days relative to B-band maximum. The spectra are at an offset from one another.

We examined the temporal evolution of Na I D absorption features and found no significant changes. Additionally, we estimated reddening using the EW of Na I D absorption features, following the empirical relations from [2]. The MW E(B-V) was comparable to the E(B-V) values derived from [3]. However, we could not independently confirm the accuracy of the derived E(B-V) value for the host galaxy.

References

We estimated the values of R_V and E(B-V) by fitting the light

curve of SN 2022hrs using SNooPy. We performed the fitting

initially by including all available filters (B, V, I, u, g, and r), and

then excluding the bluest filters (B, u, and g). We then applied

reddening corrections for different combinations of R_V and

E(B-V). Our analysis shows that assuming a similar intrinsic

color for SN 2022hrs and SN 2011fe yields somewhat

comparable spectra in the pre-maximum phase. However, in the

post-maximum phase, better spectral agreement is achieved

when corrections are applied under the assumption of a redder

[1] Foley, R. J., & Kasen, D. (2011) (ApJ, 729, 55) [2] Poznanski, D., et al. (2012) (MNRAS, 426, 1465-1479) [3] Schlafly, E. F., & Finkbeiner, D. P. (2011) (ApJ, 737, 2, 103) [4] Brown, P. J., et al. (2019) (ApJ, 877, 2, 152) [5] Derkacy, J., et al. (2023) (MNRAS, 522, 3, 3481-3505) [6] Shappee, B. J., et al. (2017) (ApJ, 841, 1, 48)

Note: These results are preliminary. Final findings will be published in Mishra, D., et al. (2025).