

# BST 234: Lab - 1

Divy Kangeyan

February 6, 2019

# Normalized Floating Point Representation

$$x = \pm m * b^{\pm e}$$

- base  $b \in \mathbb{N}$  and  $b > 1$
- mantissa  $m = m_1 b^{-1} + \dots + m_r b^{-r} \in \mathbb{R}$
- exponent  $e = e_{s-1} b^{s-1} + \dots + e_0 b^0 \in \mathbb{N}$
- digits  $m_i, e_i \in 0, \dots, b-1$
- significant digits  $s \in \mathbb{N}$  and  $r \in \mathbb{N}$

# Practice: Normalized Floating Point Representation

Express the number  $x = 10$  in normalized floating point format for the base  $b = 2$ :

# Practice: Normalized Floating Point Representation

Express the number  $x = 10$  in normalized floating point format for the base  $b = 2$ :

- Find  $k$  such that  $b^k \leq x \leq b^{k+1}$

$$2^3 \leq 10 \leq 2^4$$

# Practice: Normalized Floating Point Representation

Express the number  $x = 10$  in normalized floating point format for the base  $b = 2$ :

- Find  $k$  such that  $b^k \leq x \leq b^{k+1}$

$$2^3 \leq 10 \leq 2^4$$

- Factor  $x$  into  $b^k, b^{k-1}, \dots$

$$10 = 2^3 \times 1 + 2^2 \times 0 + 2^1 \times 1 + 2^0 \times 0$$

# Practice: Normalized Floating Point Representation

Express the number  $x = 10$  in normalized floating point format for the base  $b = 2$ :

- Find  $k$  such that  $b^k \leq x \leq b^{k+1}$

$$2^3 \leq 10 \leq 2^4$$

- Factor  $x$  into  $b^k, b^{k-1}, \dots$

$$10 = 2^3 \times 1 + 2^2 \times 0 + 2^1 \times 1 + 2^0 \times 0$$

- Add terms and factor out  $b^{k+1}$

$$10 = 2^4(2^{-1} \times 1 + 2^{-2} \times 0 + 2^{-3} \times 1 + 2^{-4} \times 0)$$

# Practice: Normalized Floating Point Representation

Express the number  $x = 10$  in normalized floating point format for the base  $b = 2$ :

- Find  $k$  such that  $b^k \leq x \leq b^{k+1}$

$$2^3 \leq 10 \leq 2^4$$

- Factor  $x$  into  $b^k, b^{k-1}, \dots$

$$10 = 2^3 \times 1 + 2^2 \times 0 + 2^1 \times 1 + 2^0 \times 0$$

- Add terms and factor out  $b^{k+1}$

$$10 = 2^4(2^{-1} \times 1 + 2^{-2} \times 0 + 2^{-3} \times 1 + 2^{-4} \times 0)$$

- Answer:  $x = (.101)_2 * 2^4$

# Practice: Normalized Floating Point Representation

Express the number  $x = 100$  in normalized floating point format for the base  $b = 3$ :



# Practice: Normalized Floating Point Representation

Express the number  $x = 100$  in normalized floating point format for the base  $b = 3$ :

- Find  $k$  such that  $b^k \leq x \leq b^{k+1}$
- Factor  $x$  into  $b^k, b^{k-1}, \dots$
- Add terms and factor out  $b^{k+1}$

# Practice: Normalized Floating Point Representation

Express the number  $x = 100$  in normalized floating point format for the base  $b = 3$ :

- Find  $k$  such that  $b^k \leq x \leq b^{k+1}$
- Factor  $x$  into  $b^k, b^{k-1}, \dots$
- Add terms and factor out  $b^{k+1}$
- Answer:  $x = (.10201)_3 * 3^5$

# Machine Precision

Definition:

$$eps := \frac{1}{2}b^{-r+1}$$

For the IEEE-format:

$$eps_{IEEE} \leq \frac{1}{2}2^{-51}$$

- *Python demonstration*

## Machine Precision

Definition:

$$eps := \frac{1}{2}b^{-r+1}$$

For the IEEE-format:

$$eps_{IEEE} \leq \frac{1}{2}2^{-51}$$

- *Python demonstration*

$$\frac{7}{3} = 1.0011 * 2^1$$

$$\frac{4}{3} = 1.01 * 2^0$$

[illegible]

$$\frac{7}{3} - \frac{4}{3} - 1 = 2^{-52} = \epsilon$$

# Floating point arithmetic

- Since floating point arithmetic is inherently approximate and not exact following symbols are used:  $\oplus, \ominus, \otimes, \oslash$
- $(x \oplus y) \oplus z \neq x \oplus (y \oplus z)$  (Associative law doesn't hold)
- $(x \oplus y) \otimes z \neq (x \otimes z) \oplus (y \otimes z)$  (Distributive law doesn't hold)
- $x \oplus y = x$  for  $|y| \leq \frac{|x|}{b} \epsilon$
- *Python demonstration*

# Using Karatsuba algorithm for multiplication

Use Karatsuba algorithm to find the results for  $15 \times 24$  in binary