

Surgery AIIMS 2017

Question 1:

A man met with a road traffic accident, which resulted in paralysis of both his upper and lower limbs. He has not passed urine and tenderness is elicited in the cervical region. What will you advise regarding radiography?

- a) The patient should not be shifted and portable x-ray machine should be used after neck stabilization.
- b) Instruct the radiographer to take cervical and chest x-ray.
- c) Instruct the radiographer to take cervical x-ray AP and lateral view without any cervical support.
- d) Order a cervical x-ray and shift the patient from the trolley by himself.

Question 2:

A 42-year-old woman presented with a history of painless breast lump measuring 6.2 X 4.5 cm in the left upper outer quadrant and no axillary nodes. Core cut biopsy showed ductal carcinoma in situ. She underwent tumor resection with adequate margins and postoperative HPE showed DCIS with intermediate-grade necrosis and with 6-mm clearance on margins. What is the next step of treatment?

- a) No additional treatment
- b) Adjuvant radiotherapy
- c) Adjuvant chemotherapy
- d) Adjuvant chemoradiotherapy

Question 3:

Which of the following is false regarding imaging in the primary survey of a trauma patient?

- a) Hemodynamically unstable patient should not be sent for CT scan
- b) All patients should have chest X-ray PA view only
- c) X-ray of the pelvis is essential in primary survey
- d) X-rays in primary survey must also be done in pregnant women

Question 4:

A patient presents with renal calculus of > 4 cm of the staghorn type. What is the best management?

- a) Extracorporeal shockwave lithotripsy
- b) Percutaneous nephrolithotomy
- c) Ureteroscopic removal
- d) Retrorenal surgery


Question 5:

All of the following can be done to transport a polytrauma patient who is conscious except?

- a) Strap their thorax, abdomen and lower limbs on a stretcher
- b) Place in prone position to prevent aspiration
- c) Put patient on lateral position and shift him immediately
- d) Roll the patient onto a hard board

Question 6:

Identify the condition.



- a) Cholesterosis
- b) Strawberry gallbladder

- c) Gallbladder polyps
- d) Gallbladder carcinoma

Question 7:

Colour of 22G IV Cannula is?

- a) Pink
- b) Green
- c) Blue
- d) Yellow

Question 8:

Fr in size of Foley's catheter refers to:

- a) Outer Diameter
- b) Inner Diameter
- c) Inner Circumference
- d) Radius

Question 9:

A school bus full of children met with an accident. As the doctor in charge of triage, which of the following will you attend to first?

- a) Airway Obstruction
- b) Grade 4 Shock
- c) Flail Chest
- d) Severe head injury

Question 10:

Which parameter is the best guide to monitor the adequacy of fluid resuscitation in a patient?

- a) Central venous pressure
- b) Urine output

- c) Blood pressure
- d) Oxygen saturation

Question 11:

Which of the following is the most common presentation of abdominal desmoid tumor?

- a) Abdominal pain
- b) Abdominal mass
- c) Fever
- d) Rectal prolapse

Question 12:

A 10-year-old child came with complaints of pain over the right lower quadrant and lower back pain. On examination, the patient had pain over the right side of the hip on passive extension with a mass in the right lumbar region pointing towards the groin. What is the most probable diagnosis?

- a) Pyelonephritis
- b) Psoas abscess
- c) Appendicitis
- d) Torsion of right undescended testis

Question 13:

Which layer is involved in blister formation in a superficial partial thickness burn?

- a) Epidermis
- b) Subcutaneous tissue
- c) Papillary dermis
- d) Reticular dermis

Question 14:

Proper functioning of ICD tube is indicated by _____.

- a) Continuous air bubbles coming out of the underwater drain

- b) Oscillation of water column in the tube
- c) Chest X-ray
- d) Auscultation

Question 15:

In a preoperative surgical safety checklist, which of the following is not present?

- a) Signature of the doctor
- b) Consent
- c) Functioning pulse oximeter
- d) Sterility of instruments

Question 16:

A trauma patient suffers a renal laceration with urine extravasation. He is managed conservatively. However, after a few days, he presents with a urinoma. The next best management for this patient would be?

- a) Ureteral stenting
- b) Surgical repair and exploration
- c) Observation
- d) Percutaneous drainage


Question 17:

Which of the following is considered balanced resuscitation?

- a) Maintaining airway, breathing and circulation
- b) Allow permissible hypotension
- c) Maintain pH by ensuring acid-base balance
- d) Fluid resuscitation by giving colloids and crystalloids

Question 18:

The flap shown in the picture is which of the following?



- a) Serratus anterior flap
- b) Pectoral myocutaneous flap
- c) TRAM flap
- d) VRAM flap

Question 19:

A 25-year-old patient presented with neck swelling which moves with deglutition as shown below. What is your diagnosis?



- a) Branchial cyst

- b) Cystic hygroma
- c) Thyroglossal cyst
- d) Cervical lymphadenopathy

Question 20:

A 28-year-old patient was brought to the casualty ward with high fever and pain abdomen. He had a history of bloody diarrhea for several months. His past history was negative for long-term medications and previous surgeries. The abdominal X-ray image obtained is shown below. What is the most probable diagnosis?



- a) Toxic megacolon
- b) Volvulus
- c) Ogilvie syndrome
- d) Pneumatosis intestinalis

Question 21:

A USG can detect lymph node metastases in a patient with breast cancer and clinically negative lymph nodes by what percentage?

- a) 50 – 60%
- b) 10 – 20%
- c) 20 – 30%
- d) 30 – 50%

Question 22:

A young boy was brought to the casualty with a history of fever and abdominal pain. On examination, he was febrile and his pulse rate was 104 beats/min. The CT scan image is shown below. Which of the following signs is not seen in this disease?



- a) McBurney's tenderness
- b) Rovsing's sign
- c) Psoas sign
- d) Ballance sign

Question 23:

A 12-year-old boy presented to the OPD with a scald injury of his arm. The lesion was pink, oozing, and painful to moving air and touch. Which of the following is the most appropriate management?

- a) Silver sulphadiazine cream dressing
- b) Excision and grafting
- c) Paraffin gauze dressing
- d) Collagen dressing

Question 24:

A 25-year-old pregnant woman came with a history of thyroid swelling. FNAC report was suggestive of papillary carcinoma thyroid and the nodule was "cold" on scan. Which of the following will not be a treatment option for her?

- a) Total thyroidectomy
- b) Total thyroidectomy plus lymph node dissection
- c) Radioiodine ablation of thyroid
- d) Right hemithyroidectomy

Question 25:

A 41-year-old woman underwent treatment for breast cancer of 1.2 X 1 cm with positive axillary nodes. She completed radiotherapy and chemotherapy and is now receiving tamoxifen. How will you follow up the patient?

- a) Annual mammogram
- b) Regular checking of LFT once in 6 months
- c) Follow-up with tumour markers once in 6 months
- d) Yearly bone scans

Question 26:

A patient underwent laparoscopic cholecystectomy and was discharged after removing the drain tube on the first postoperative day. On day three, the patient came back to the hospital and was febrile. On ultrasound examination, there was a 5 x 5 cm collection in the subhepatic space. Next best step is _____.

- a) To give oral antibiotics since the given accumulation is normal
- b) To re-explore the wound and insert a T-tube
- c) Insert pigtail catheter for drainage
- d) ERCP

Question 27:

A 22-year-old boy was admitted to the casualty with a history of breathlessness following massage by his mother. He had a history of hairline ankle fracture and was on a below-knee cast which was removed recently. The most probable diagnosis is

- a) Pneumonitis

- b) Septicemia
- c) Acute MI
- d) Pulmonary thromboembolism

Question 28:

A 65-year-old man has the following lesion over the face. He complains of bleeding while itching. The most likely diagnosis is



- a) Dermatofibrosarcoma
- b) Marjolin's ulcer
- c) BCC
- d) Naevus

Question 29:

A 54-year-old comes to the casualty with a history of massive hemoptysis. His routine investigation and chest X-ray were normal. Which of the following you will not do to prevent hemoptysis?

- a) Bronchial artery embolization
- b) Pulmonary artery embolization
- c) Procoagulant drugs
- d) Surgical removal of lobe

Question 30:

A 30-year-old male met with an accident and came to the casualty with a contusion on the left precordium. His vitals are BP 120/80 mmHg, RR – 16/min. Normal heart sounds were heard, but decreased breath sounds on the left side, and the trachea deviated to the right. Which of the following is the first line of management?

- a) Needle thoracocentesis
- b) Pericardiocentesis
- c) Chest tube insertion
- d) Open surgery

Answer Key

Question No.	Correct Option
1	a
2	b
3	b
4	b
5	c
6	c
7	c
8	a
9	a
10	b
11	b
12	b
13	c
14	b
15	a
16	a
17	b
18	a
19	c

20	a
21	a
22	d
23	d
24	c
25	a
26	c
27	d
28	c
29	c
30	c

Detailed Explanations

Solution to Question 1:

The given clinical situation suggests a cervical spine injury due to trauma. The patient should not be shifted and a portable X-ray machine should be used for radiography after neck stabilization.

Management in a trauma patient with a suspected cervical fracture involves the following steps:

- Spinal immobilization with a rigid cervical collar and a long spine board is the immediate step in the management of blunt/penetrating cervical trauma cases.
- It is followed by a gross assessment of spinal cord function by eliciting extremity movement. Tenderness over the injured vertebrae or the presence of a deformity is indicative of an associated fracture.
- Imaging is done by X-ray or CT. Portable X-ray machines are preferred to avoid further injury while mobilizing the patient. They are useful in visualizing thoracic and lumbar vertebrae but not the cervicothoracic junction. Cervical CT has been made the modality of choice in most emergency departments owing to the high quality of images that can be obtained.

Solution to Question 2:

According to Silverstein Van Nuys prognostic index for DCIS, the total score is 9. Hence, the patient would require adjuvant radiotherapy.

Age: 42 years (score 2), size: 62mm (score 3), clearance margin: 6mm (score 2) with intermediate-grade necrotic features (score 2).

Silverstein Van Nuys prognostic index for DCIS:

Treatment:

- Scores 4-6 with ≥ 3 mm margin: Excision alone

- Score 7 with $\leq$ 3 mm margin, score 8 with $\leq$ 3mm margin, score 9 with $\leq$ 5mm margin: Excision with RT

- Score 9 with $\leq$ 5mm margin & score 10-12: Mastectomy

If the patient is ER/PR+, adjuvant hormonal therapy with tamoxifen or an aromatase inhibitor is indicated. Current data does not support the use of HER-2-directed therapy in DCIS.

Indications for adjuvant radiotherapy in breast cancer:

- Locally advanced breast cancers (T3, T4, N1, N2, N3)
- Following breast conservation surgery
- After mastectomy if tumor size $\geq$ 5 cm, skin or chest wall involvement, lymphovascular invasion and axillary node-positive for metastasis.

Post-treatment surveillance after treatment for DCIS includes history, physical examination, and routine mammography (if applicable).

Parameters	Score 1	Score 2	Score 3
Age (years)	$\geq$ 61	40-60	$<$ 40
Size (mm)	$\leq$ 15 mm	16-40 mm	$>$ 40 mm
Margin (mm)	$\geq$ 10 mm	1-9 mm	$<$ 1 mm
Class (grade and necrosis)	Grades 1 and 2 no necrosis	Grades 1 or 2 with necrosis	Grade 3 with or without necrosis

Solution to Question 3:

Chest X-ray and X-ray pelvis are taken in AP view in the adjuncts to the primary survey of a trauma patient.

Solution to Question 4:

The best management for staghorn renal calculus of $\geq$ 4 cm is percutaneous nephrolithotomy (PCNL).

The initial management of renal stones depends on the stone size and includes conservative management, extracorporeal shockwave lithotripsy (ESWL), and PCNL. PCNL is preferred for stones $\geq$ 2cm in size and for staghorn calculi, while ESWL is used to treat renal stones of size 0.6 to 1.5 cm. PCNL is indicated in

- Renal stones $\geq$ 2 cm
- Staghorn calculus
- Failed shockwave lithotripsy (SWL) or retrograde intrarenal surgery (RIRS) for renal calculi
- Lower pole renal stones with anatomy that is unfavorable for ESWL

Other options:

Option C: Ureteroscopic removal is attempted for failed SWLs or PCNLs, and is not the first choice in the surgical management of renal stones.

Option D: RIRS is a minimally invasive procedure similar to ureteroscopic removal, where the kidney is accessed using a slim and flexible ureteroscope in a retrograde manner via the ureteric orifice. It is indicated for lower pole renal stones ≤ 2 cm.

Solution to Question 5:

Patients with polytrauma are shifted into a supine or prone position (not in a lateral position) by laying them over a firm surface to restrict spinal movement.

As per ATLS guidelines, in patients with trauma, airway management is first done along with cervical spine stabilization. All trauma patients until proven otherwise are considered to have an injured vertebral column.

The patients are placed supine on a firm surface without rotating or bending the spinal column and a rigid cervical collar (philadelphia collar) is applied. Aspiration can be prevented by putting the patient in the prone position and strapping can be done to provide further stabilization.



Long spine boards with straps are helpful in the rapid transport of the patient from one place to another. The patient can be put on the long spine board by log-rolling (a technique where four people hold the patient and roll the patient like a log without allowing rotation or bending of the spine).

Conscious and alert patient indirectly means that they don't need intubation and their airway is patent. If the patient is unconscious always check the airway and if needed, endotracheal intubation must be done immediately by various methods. Preferably by direct laryngoscopy and intubation.

Solution to Question 6:

The given image shows a resected specimen of a gallbladder with a polyp on the inner surface.

Specimen of gallbladder carcinoma would include an en-bloc dissection of the gall bladder, surrounding liver tissue and regional lymph nodes (option D). In specimens of cholesterosis, also known as strawberry gallbladder, the gall bladder would appear yellow and coarsely granular due to the presence lipid nodules (options A and B).

Gall bladder polyps are usually incidental findings during abdominal imaging. Most of them are benign and comprise cholesterol polyps, adenomyomas or inflammatory polyps. Adenomatous polyps are the most common neoplastic polyps. The majority of patients with polyps are stable and smaller polyps are kept under observation. Cholecystectomy is done for symptomatic patients (biliary colic, pancreatitis), patients ≥ 50 years, sessile polyps with thickening ≥ 4 mm and polyps ≥ 10 mm.

The risk factors for malignancy in an adenomatous polyp include age ≥ 50 years, size ≥ 10 mm, sessile polyp, documented increase in size, and coexistence with gallstones.

Other options:

Option D: The gallbladder in a patient with carcinoma is removed via an en-bloc dissection with surrounding liver tissue along with regional lymph nodes. The given image below shows a cut surface of the resected gallbladder with a tumor at the neck along with liver bed involvement (marked by the red arrow).



Options A and B: Cholesterosis (strawberry gallbladder) is characterized by the accumulation of lipids (triglycerides, cholesterol precursors and cholesterol esters) in the lamina propria of the gallbladder wall. The nodules are less than 1 mm in diameter and there is mucous villous hyperplasia, which gives it a coarse, yellow and granular appearance, giving it a strawberry-like appearance. The given images below show cholesterosis.



Solution to Question 7:

The 22G IV cannula is blue in colour.

Solution to Question 8:

The French (Fr) measurement unit refers to the outer diameter.

As per the French system, the outer circumference equals 3 times the outer diameter. One French unit (1 Fr) equals 1 mm of the outer circumference and 0.3 mm of the outer diameter. Therefore, the outer diameter in mm can be determined by dividing the Fr size by 3.

Example: An 18 Fr would have a circumference of 18 mm and an outer diameter of 6 mm.

Solution to Question 9:

The patient with airway obstruction, categorized as red, should be attended to first.

Triage is used to classify the priority of treatment. It is carried out at the site of the disaster to determine the priority for immediate medical attention, transportation, and admission to the hospital. Triage decisions are made by deciding which injuries constitute the greatest immediate threat to life. In general, airway problems are more rapidly lethal than breathing problems > circulation problems > neurologic injuries.

The internationally accepted four-color coded system is commonly used. It includes:

- Red - High-priority attention or transfer
- Yellow - Medium priority

- Green - Ambulatory patients
- Black - Dead or moribund patients

Solution to Question 10:

Urine output is the best measure of organ perfusion and the best monitor for adequacy of fluid resuscitation.

Adequate end-organ perfusion is best indicated by urine output of ≥ 0.5 to 1 mL/kg/h. However, it does not give a minute-to-minute value.

The level of consciousness is an important marker for cerebral perfusion.

The only clinical indicators of perfusion of the gastrointestinal tract and muscular beds are measures of lactic acidosis (lactate and base deficit) and mixed venous oxygen saturation.

Other options:

Option A: Central venous pressure (CVP) is the mean pressure in the superior vena cava, used for estimation of preload and right atrial pressure. Normal CVP ranges from 2 to 7 mm Hg (3 to 9 cm H₂O). A sick or injured patient with a CVP ≤ 3 mm Hg is presumed to be volume depleted and may be given fluids with relative safety. However, CVP is unreliable in assessing volume status or left ventricular function with respect to resuscitation.

Options C and D: Blood pressure and oxygen saturation do not reflect body fluid levels.

Solution to Question 11:

Explanation:

The most common presentation of an abdominal desmoid tumor is an abdominal mass.

Desmoid tumors, also known as fibromatosis, aggressive fibromatosis or desmoid-type fibromatosis are locally aggressive tumors that lack metastatic potential. Desmoid tumors can be sporadic in origin or associated with familial adenomatous polyposis.

These tumors arise from fibro-aponeurotic tissue and typically are manifested as a slowly growing mass. They present as firm, non-painful, non-tender masses in the abdominal wall, shoulder, hip and limbs, mesentery, and pelvis. They are also associated with trauma, pregnancy, and oral contraceptive usage.

The investigation of choice for a desmoid tumor is MRI. It is superior to CT in defining the pattern and the extent of involvement and can determine recurrence after surgery. Diagnosis is established by a biopsy with immunohistochemistry, which stains positive for beta-catenin, actin and vimentin, and negative for desmin, cytokeratin and S-100.

Treatment:

- An initial observation is preferred for stable, asymptomatic, primary or recurrent desmoids, especially if surgery can cause high mortality.


- Surgical resection with negative margin clearance is reserved for patients with small abdominal wall tumors associated with pain or an increase in size.
- Tamoxifen and NSAIDs are used as first-line medical therapy. Other systemic treatment options are low-dose chemotherapy, tyrosinase kinase inhibitors, and full-dose chemotherapy
- Radiotherapy is preferred for candidates unfit for surgery, patients refusing surgery and when surgical resection is associated with high morbidity.


Solution to Question 12:

The given clinical history of lower back pain, mass in the right lumbar region pointing towards the groin and pain on passive extension of the ipsilateral hip is suggestive of a psoas abscess.

Psoas abscess is a retroperitoneal abscess, caused due to various etiologies. Previously, tuberculosis of the spine was the most common cause secondary to spreading infection from the digestive or urinary tract, which can cause psoas abscess. Nowadays, Crohn's disease is the most common cause. Patients present with flank pain, fever and chills, malaise and weight loss. They can have referred pain to the hip, groin or knee. When the swelling extends across the groin and below the inguinal ligament along the course of the iliopsoas, it becomes cross fluctuant across the inguinal ligament.

Imaging is done by a CT scan, which shows a low-density mass with surrounding inflammation and infiltration of surrounding fat and gas or air-fluid levels.





Treatment is done by percutaneous CT-guided drainage and appropriate antibiotics.

Other options:

Option A: Pyelonephritis is usually associated with a history of urinary tract infection and sudden onset of pain in the flank and hypochondrium, not the lumbar region.

Option C: Appendicitis (retrocaecal) can present with pain on passive extension of the hip (psoas sign), but the presence of a mass in the right lumbar region pointing towards the groin makes it an unlikely diagnosis.

Option D: Testicular torsion is a surgical emergency condition presenting with sudden onset pain in the scrotum, groin and lower abdomen, tenderness, redness and edema of the scrotal skin.

Solution to Question 13:

The papillary dermis is involved in blister formation in a superficial partial-thickness burn.

First-degree burns involve only the epidermis, with pain, redness, and mild swelling.

Second-degree (partial thickness) burns involve the epidermis and some parts of the dermis (also known as partial-thickness burns). Further divided into:

- Superficial second-degree burns - involves the upper part of the dermis (papillary) and shows pain, blisters, and severe swelling.
- Deep second-degree burns - extends up to the reticular layer of the dermis and appears more pale and mottled, does not blanch to touch, but remains painful to pinprick.

Third-degree or full-thickness burns involve the full thickness of the dermis and subcutaneous tissue.

Fourth-degree burns involve deep structures such as muscle and bone.

Solution to Question 14:

Oscillation of the water column level (moving up and down during respiration) within the tube in the water seal chamber is indicative of a functioning (patent) ICD tube.

During spontaneous respiration, the water level rises with inspiration and falls with expiration. Whereas during positive pressure ventilation, the opposite occurs. This oscillation is also known as tidaling.

Other options:

Option A: Continuous bubbling indicates an air leak in the drainage system, which should be corrected

Option B: Chest X-ray is useful for assessing the position of the ICD tube inside the chest, not its functioning.

Option C: Auscultation is not useful for assessing the functioning of an ICD tube.

Solution to Question 15:

In a preoperative patient surgical safety checklist, doctor's signature is not required.

WHO Guidelines for Surgical Safety Checklist

Surgical Safety Checklist		
World Health Organization Patient Safety		
Before induction of anaesthesia (with at least nurse and anaesthetist)	Before skin incision (with nurse, anaesthetist and surgeon)	Before patient leaves operating room (with nurse, anaesthetist and surgeon)
Has the patient confirmed his/her identity, site, procedure, and consent? ☐ Yes	☐ Confirm all team members have introduced themselves by name and role.	Nurse Verbally Confirms: ☐ The name of the procedure
Is the site marked? ☐ Yes ☐ Not applicable	☐ Confirm the patient's name, procedure, and where the incision will be made.	☐ Completion of instrument, sponge and needle counts
Is the anaesthesia machine and medication check complete? ☐ Yes	Has antibiotic prophylaxis been given within the last 60 minutes? ☐ Yes ☐ Not applicable	☐ Specimen labelling (read specimen labels aloud, including patient name)
Is the pulse oximeter on the patient and functioning? ☐ Yes	Anticipated Critical Events To Surgeon: ☐ What are the critical or non-routine steps? ☐ How long will the case take? ☐ What is the anticipated blood loss?	☐ Whether there are any equipment problems to be addressed
Does the patient have a: Known allergy? ☐ No ☐ Yes	To Anaesthetist: ☐ Are there any patient-specific concerns?	To Surgeon, Anaesthetist and Nurse: ☐ What are the key concerns for recovery and management of this patient?
Difficult airway or aspiration risk? ☐ No ☐ Yes, and equipment/assistance available	To Nursing Team: ☐ Has sterility (including indicator results) been confirmed? ☐ Are there equipment issues or any concerns?	
Risk of >500 ml blood loss (7 ml/kg in children)? ☐ No ☐ Yes, and two IV/central access and fluids planned	Is essential imaging displayed? ☐ Yes ☐ Not applicable	

FIGURE 10-4 Surgical safety checklist published by the World Health Organization.
www.who.int/patientsafety/safesurgery/checklist/en.

Solution to Question 16:

The patient with a history of grade IV renal injury (renal laceration with urine extravasation) who develops a persistent urinoma (a mass formed by encapsulated extravasated urine) should be managed with the placement of an internal ureteral stent.

In patients not cured by ureteral stenting, percutaneous nephrostomy or transcutaneous drain may be required.

Management of renal injuries

Blunt trauma is the main cause of renal injuries and is self-limiting. Blunt trauma is mostly treated conservatively. Most penetrating injuries require surgical renal exploration.

- Grading of injury through CECT
- If hemodynamically stable
- Conservative management - Bedrest & hemodynamic monitoring
- In grade IV, a repeat CT scan to ensure the extravasation of urine has resolved.
- If hemodynamically unstable
- Surgical exploration and repair
- In grade V, nephrectomy.

Solution to Question 17:

Permissive hypotension is a part of balanced resuscitation in trauma management to avoid bleeding.

Damage control resuscitation (DCR) i.e. balanced resuscitation is used in the settings of major trauma. It aims to control hemorrhage and prevent coagulopathy prior to the development of the trauma triad of death (hypothermia, acidosis, and coagulopathy). The components of DCR include:

- Permissive hypotension until definitive surgical control
- Minimize crystalloid based resuscitation
- Early use of blood products (PRBCs, FFP, platelets, cryoprecipitate) in the ratio of 1:1:1
- Use of hemostatic agents such as tranexamic acid, recombinant factor 7a, prothrombin complex concentrates

DCR has the following advantages:

- Minimizes the impact of trauma-induced coagulopathy
- Limits blood product waste
- Reduces the complications that occur with aggressive crystalloid resuscitation.

Solution to Question 18:

The given image shows a serratus anterior flap. The serratus anterior flap is a highly reliable flap with a consistently long pedicle. Its anatomy permits coverage of complex wounds. The underlying rib can be included as a myoosseous flap to expand the versatility of this flap.


Classification of muscle flaps		
Based on blood supply	Random flap Unnamed vessel supplying flap	Axial flap Named vessel supplying flap.
Method of movement	Local transfer Rotation flap Transposition flap Advancement flap	Distant transfer Pedicle flap Free flap
Based on composition	Simple Flap Cutaneous flap Musculocutaneous Fasciocutaneous	Composite flap Osseocutaneous flap Sensory flap

Solution to Question 19:

A thyroglossal cyst is the only midline neck swelling that moves with deglutition. This is because the thyroid gland is fixed to the hyoid.

Thyroglossal duct cyst:

It is a congenital midline neck swelling that typically presents during childhood often noticed after an upper respiratory infection. It represents the persistence of the track of the thyroid gland and can be found adjacent to the midline. The thyroid gland descends early in fetal life from the base of the tongue towards its position in the lower neck. At the time of descent, the hyoid bone has not been formed and the track of the thyroid gland is variable, passing in front, through, or behind the eventual position of the hyoid body.



Clinical features of thyroglossal cyst:

- Classically the cyst moves upwards on swallowing and on tongue protrusion. This indicates attachment to the hyoid bone.
- Swelling is smooth, soft, fluctuant (cystic), nontender, mobile, often transilluminant.
- Thyroid fossa is empty, if there is no thyroid in the normal location.

Complications of thyroglossal cyst:

- The most common complication is infection.
- Thyroglossal fistula is an acquired condition that either follows infection of the cyst which bursts open or due to inadequate removal of the cyst.
- 1% of thyroglossal duct cysts are found to be malignant of which the most common type of cancer is papillary (85%).

Neck swellings that move with deglutition are:

- Thyroid swellings
- Subhyoid bursitis
- Dermoid cysts attached to the hyoid bone

However, only the thyroglossal cyst moves with tongue protrusion. This is because the duct extends from the foramen caecum of the tongue to the isthmus of the gland.

Sistrunk's operation is performed to excise the cyst and prevent a recurrence. It involves the excision of the whole thyroglossal tract, involving removal of the body of the hyoid bone, and suprahyoid tract through the tongue base to the vallecula at the site of the primitive foramen caecum along with the core of tissue on either side.

Solution to Question 20:

The clinical scenario and the radiograph are suggestive of toxic megacolon.

Toxic megacolon may present with fever, abdominal pain, and abdominal distension. Abdominal radiography shows transverse or right colonic dilation of ≥ 6 cm and loss of haustrations. It is usually associated with ulcerative colitis as indicated by history of chronic bloody diarrhoea.

It can be triggered by electrolyte imbalance and narcotics which interfere with gut motility. Features of secondary bacterial infection and sepsis may be present. It can be complicated by intestinal perforation.

It usually resolves with supportive treatment (correcting electrolyte imbalance, fluids, and bowel rest) and medical therapy which includes glucocorticoids, anti-inflammatory drugs, and antibiotics. However, an urgent colectomy is indicated in patients who do not show signs of improvement.


Indications for urgent surgery include:

- Frank intraperitoneal peritonitis, secondary to perforation
- Life-threatening hemorrhage or increasing transfusion requirements
- Worsening systemic toxicity
- Worsening colonic dilatation

Other options:

Option B: Volvulus occurs when the air-filled segment of the colon twists around the mesentery. The sigmoid colon is the most commonly involved part of the large intestine (90%). The patient


can present with abdominal distension, nausea and vomiting. X-ray abdomen shows a bent inner tube or coffee bean appearance.



Option C: Ogilvie syndrome, also known as colonic pseudo-obstruction is a functional disorder with colonic dilatation without mechanical obstruction. It occurs commonly in hospitalized patients and is associated with narcotics, bed rest and comorbid disease. X-ray shows dilatation of the colon (predominantly transverse colon) with absent mechanical obstruction.



Option D: Pneumatosis intestinalis is the presence of gas within the bowel wall seen in any part of the GI tract but most commonly in the jejunum. It is a pathognomic feature of necrotizing enterocolitis. X-ray of the abdomen will show intramural gas in the bowel.



Solution to Question 21:

A USG can detect lymph node metastasis in a patient with clinically negative lymph nodes by 50-60%.

The clinical determination of axillary lymph node metastases has an accuracy of only 33%.

Ultrasound (USG) is more sensitive than physical examination alone in determining axillary lymph node involvement during the preliminary staging of breast carcinoma.

Fine-needle aspiration cytology (FNAC) or core biopsy of sonographically indeterminate or suspicious lymph nodes can provide a more definitive diagnosis than USG alone.

Ultrasonography can also be utilized to image the regional lymph nodes in patients with breast cancer. The sensitivity for the status of axillary nodes ranges from 35% to 82% and specificity ranges from 73% to 97%.

The features of a lymph node involved with cancer on USG include:

- Cortical thickening
- Change in shape of the node to a more circular appearance
- Size larger than 10 mm
- Absence of a fatty hilum
- Hypoechoic internal echoes

Solution to Question 22:

The patient's history and radiological findings are indicative of acute appendicitis. Ballance sign is not seen in acute appendicitis.

Findings on CT that are suggestive of acute appendicitis include:

- Appendiceal dilatation (>6 mm in diameter)
- Appendiceal wall thickening (>3 mm) and enhancement

Ballance sign is seen due to a ruptured spleen. The ballance sign is positive if there is a fixed dullness to percussion in the left flank and shifting dullness to percussion in the right flank. The fixed dullness is due to the presence of coagulated blood near the lacerated spleen while the shifting dullness is due to the presence of blood in the peritoneal cavity.

Option A: Deep tenderness at McBurney's point, known as McBurney's sign, is a sign of acute appendicitis. McBurney's point is a point at one-third of the distance from the anterior superior iliac spine and umbilicus.

Option B: Rovsing sign is seen in appendicitis. Deep palpation of the left iliac fossa may cause pain in the right iliac fossa.

Option C: Psoas sign is seen in appendicitis. Occasionally, an inflamed appendix lies on the psoas muscle, and the patient will lie with the right hip flexed, for pain relief.

Solution to Question 23:


Scald injury (damage from contact with hot liquids) that appears pink in colour, has oozing, and pain is suggestive of a 2nd-degree burn (i.e. partial-thickness burn). Collagen dressing is the most appropriate management for this injury.

The best dressing for burn wounds is skin; collagen is a good substitute for skin, especially in partial-thickness burns. For full-thickness burns, sequential excision and grafting should be considered.

Initial treatment for burns includes cleaning burnt areas and removing loose dead skin. Blisters are de-roofed and an appropriate dressing is applied.

Biological dressing (collagen dressings, amniotic membrane, etc.) is the preferred dressing material, especially in partial thickness burns.

If this is not available then a non-adhesive dressing like paraffin gauze impregnated with silver sulfadiazine (SSD) cream, colloidal silver, ionic silver cream, or silver nitrate cream can be applied. Since increased resistance to SSD has recently emerged, it should be judiciously used and reserved for deep dermal, full thickness, infected, and old neglected burns.



Solution to Question 24:

Radioactive ablation of the thyroid is an absolute contraindication in pregnancy as it can cause fetal goiter and hypothyroidism.

Radioactive ablation of the thyroid is done using the radioactive isotope of iodine (I-131), which has a half-life of 8 days. I-131 emits gamma rays (useful in imaging) and beta particles (useful in therapy). I-131 is concentrated within the thyroid follicles and emits radiation from within the tissue. The beta particles penetrate only 0.5-2 mm of the tissue. The thyroid follicular cells are affected from within and undergo pyknosis and necrosis followed by fibrosis without damage to neighboring tissues.

Radioactive iodine is administered as the sodium salt of I-131 dissolved in water and taken orally.

Therapeutic uses of radioactive iodine:

- Distant metastasis at diagnosis (papillary or follicular carcinoma)
- Incomplete tumor resection
- Patients at high risk for mortality or recurrence
- Graves disease
- Toxic multinodular goitre

The average therapeutic dose is 3-6 m curie, calculated based on previous tracer studies and thyroid size.

Contraindications:

- Pregnancy
- Lactation
- Young age (<25 years)

- Moderate to severe Graves ophthalmopathy

Management of differentiated thyroid cancer in pregnancy:

- Thyroidectomy after delivery for patients with no evidence of advanced disease or no rapid progression and in the second trimester for patients with rapid progression or signs of advanced disease.
- Radioactive iodine is contraindicated in pregnancy and lactation. If it is needed after delivery, breastfeeding shall be stopped at least 6 weeks before ablative therapy and mothers should be advised against nursing.
- Levothyroxine supplementation: as a suppressive treatment for those who decided to postpone surgery till the second trimester or after delivery, as a replacement therapy after thyroidectomy, and as a suppressive therapy for patients with residual disease.
- Differentiated thyroid cancer not responding to radioactive iodine can be treated with tyrosine kinase inhibitors.

Diagnostic radioiodine scan: 25–100 μ curie of I-131 is given; counting or scanning is done at intervals.

Solution to Question 25:

Follow-up of patients after treatment of breast cancer consists of a physical examination once in 3-6 months for the first 3 years and an annual mammogram.

Mammography is the only imaging modality necessary for follow-up after breast cancer surgery. There is currently no role for tumor markers, serum investigations, or any other imaging studies like bone scans in routine follow-ups.

Guidelines for follow-up after primary treatment:

- Physical examination: every 3-6 months for the first 3 years, every 6-12 months for years 4 and 5, and annually thereafter.
- Mammography: initially, 6 months after surgery, and thereafter, annually.
- Breast self-examination: Monthly
- Pelvic examination: Annually in patients taking tamoxifen

Solution to Question 26:

The given clinical situation suggests a bile duct injury identified on post-op day 3 following laparoscopic cholecystectomy. The next best step after imaging would be to control inflammation by drainage of fluid collection (pigtail insertion and drainage) and administration of parenteral broad-spectrum antibiotics.

Bile duct injury

- Usually is manifested in the postoperative period

- Clinical features include fever, increasing abdominal pain, jaundice and bile leakage from an incision.

Imaging: Ultrasonography is done to assess for fluid collection and to evaluate the biliary tree.


Treatment Strategy

- Control of infection, limiting inflammation
- Percutaneous drainage of periportal fluid collections
- Parenteral antibiotics (option A)
- Clear and thorough delineation of entire biliary anatomy
- Magnetic resonance cholangiopancreatography (MRCP) or percutaneous transhepatic cholangiography (PTC)
- Endoscopic retrograde cholangiopancreatography (ERCP), especially if a cystic duct stump leak is suspected
- Re-establishment of biliary-enteric continuity (surgery)
- Tension-free, mucosa-to-mucosa anastomosis
- Roux-en-Y hepaticojejunostomy

Other options:

Option B: Re-exploration and repair is indicated for biliary leaks identified within 48 hours post cholecystectomy.

Option D: ERCP would be indicated if drainage using a pigtail catheter fails.



Solution to Question 27:

The given clinical history of immobilization after the application of the cast is a risk factor for deep vein thrombosis (DVT). Massaging the leg may dislodge the thrombus causing pulmonary

embolism (PE). It is the commonest complication of DVT leading to death.

PE and DVT are part of the same spectrum of venous thromboembolism. PE occurs due to obstruction of the pulmonary arterial circulation by deep venous thrombi that detach from their site of formation and embolize to the vena cava, right atrium, and right ventricle. Virchow's triad of venous stasis, hypercoagulability, and endothelial injury, leading to recruitment of activated platelets, provides the ideal environment for thrombus formation.


DVT presents with pain and swelling of the calf (the most common symptom), tenderness and edema. Clinical examination is unreliable and the elicitation of signs such as Homan's or Mose's signs is not recommended owing to the risk of clot dislodgement. Patients with symptoms of PE present with fever, dyspnea, pleuritic chest pain, and hemoptysis. On examination, they can have cyanosis, jugular venous distension, fixed split-second heart sound and pleural rub, but may be subtle or absent. Some patients are asymptomatic.

Depending on the clinical presentation, patients with PE can be stratified as:

- High-risk (massive) PE is characterized by systemic hypotension. Patients can have dyspnea and syncope and may present in cardiogenic shock.
- Intermediate-risk PE is characterized by right ventricle (RV) dysfunction but normal blood pressure. RV dysfunction can be diagnosed with echocardiography and ECG.
- Low-risk PE is characterized by normal blood pressure and the absence of any RV dysfunction.

Investigations:

- Venous duplex ultrasound to look for thrombi in the leg.
- Chest x-ray in PE shows hyper lucency in an area of oligemia (Westermark sign). D-dimer levels are elevated and ECG will show sinus tachycardia (the most common ECG abnormality) and an S1Q3T3 sign (an S wave in lead I, a Q wave in lead III, and an inverted T wave in lead III).
- Definitive diagnosis of PE is by a CT-Pulmonary Angiography.



The management of PE will vary according to the risk stratification:

- A patient with high-risk PE will primarily be treated with anticoagulation and thrombolysis or catheter/surgical embolectomy.
- A patient with low-risk PE will be treated with anticoagulation alone, or an IVC filter as secondary prophylaxis.
- Patients with intermediate-risk PE require individualized treatment.

Anticoagulation can be done with the classical strategy of low-molecular-weight heparin followed by warfarin, or by low-molecular-weight heparin for 5 days, switched to novel oral anticoagulant like Dabigatran after that. Fibrinolysis, if indicated, is done with recombinant TPA administered over 2 hours. It can be done up to 14 days after the PE has occurred.

Management of DVT:

- Anticoagulation with low molecular weight heparin (LMWH) and unfractionated heparin (UFH) if renal impairment is present.
- After treatment with heparin, patients should be commenced on oral anticoagulants for at least 3 months with either factor Xa inhibitors (rivaroxaban and apixaban) or direct thrombin inhibitors (dabigatran).
- Other measures such as rest, limb elevation and compression stockings are also used.

Solution to Question 28:

The given history with the image showing an ulcer over the face with pearly, rolled-out borders points to a diagnosis of basal cell carcinoma.

Basal cell carcinoma (BCC) is an epithelial tumor arising from basal cells. It is the most common type of skin cancer most commonly seen over the face, above the line joining corner of the mouth, and lobule of the ear. Risk factors for BCC include ultraviolet radiation (strongest risk factor), age 40-80 years with excessive sun exposure, light-skinned people, and exposure to arsenic compounds, coal tar, aromatic hydrocarbons, and infrared light. It is more common in males than females. It is also associated with syndromes such as xeroderma pigmentosum and Gorlin's syndrome.

Features of BCC include:

- Pearly nodule with rolled border
- Tortuous telangiectatic vessels
- Surface which can get ulcerated with raised and beaded edge, often with a central depression or umbilication
- Bleeding on minor trauma

BCC has various subtypes such as nodular (most common type), pigmented, nodulocystic, morphoeic, infiltrative, and fibroepithelioma of Pinkus.

It is locally aggressive and can erode the bone or facial sinuses. It rarely metastasizes via lymph nodes.

The occasional presence of melanin in the nodular variant of BCC (pigmented BCC) may lead to confusion with melanoma.

Depending upon the type of the lesion, a shave, punch or excisional biopsy may be employed for diagnosis. X-ray or CT may be needed if there is a local invasion. Histology shows peripheral palisading of cells.

Treatment:

- Surgical: Wide excision (4-6 mm clearance) with a skin graft or flap or primary suturing
- Mohs' surgery for BCC close to the eyes, nose, or ear to preserve more tissue.
- Non-surgical: Radiotherapy, vismodegib and sonidegib.

Other options:

Option A: Dermatofibrosarcoma, also known as dermatofibrosarcoma protuberans (DFSP) is a rare, slow-growing, well-differentiated primary fibrosarcoma of the skin, associated with increased production of platelet-derived growth factor-beta. It presents as a protuberant nodule within a firm, indurated plaque that can ulcerate. They are locally aggressive tumors with high recurrence rates. Treatment is wide excision and Mohs microscopic surgery. Imatinib is used for unresectable, metastatic, and recurrent tumors. The given image below shows a DFSP near the umbilicus.



Option B: Marjolin's ulcer is a painless, slow-growing, locally malignant, well-differentiated squamous cell carcinoma that occurs in a chronic scar (burns scar, scar of venous ulcer). It presents as an indurated, non-tender ulcer with raised and everted edges with scarring in the surrounding area with marked fibrosis. The ulcer can spread to the lymph nodes when it reaches the normal skin. A wedge biopsy and MRI of the region involved confirm the diagnosis. Management is by wide excision, for larger ulcers, amputation is done. The given image below shows a marjolin's ulcer of the leg.



Option D: Naevus is a common benign neoplasm caused due to mutations in the components of the RAS signaling pathway. It has various types, such as congenital naevus, blue naevus, Spitz naevus, halo naevus, and dysplastic naevus. The given image below shows a patient with multiple dysplastic nevi.



Solution to Question 29:

Procoagulant drugs are not used to prevent hemoptysis. Procoagulant drugs like vasopressin are indicated only if there is an abnormal coagulation profile.

Massive hemoptysis is generally defined as the expectoration of over 600 mL of blood within a 24-h period.

Treatment priorities in the management of massive hemoptysis:

- First priorities are to secure the airway, optimize oxygenation and stabilize the hemodynamic status by administering IV fluids.
- Chest X-ray is the first investigation of choice. CT scan is done only in hemodynamically stable patients.
- If the bleeding site is known, the patient should be placed in a lateral decubitus position with the bleeding side down to prevent aspiration into the unaffected lung. Supplemental oxygen should be given.
- If large-volume bleeding continues or the airway is compromised, the patient should be intubated and undergo an emergency bronchoscopy.
- The bleeding site is localized using flexible/rigid bronchoscopy.
- Control the hemorrhage:
 - Endobronchially with Nd: YAG laser or bronchial occlusion
 - Endovascularly with bronchial and/or pulmonary artery embolization
 - Electrocauterization
 - Surgically with resection of the involved lobe

The cause of the hemoptysis must be determined to definitively prevent a recurrence.

Solution to Question 30:

The given clinical scenario of trauma, decreased breath sounds, and tracheal deviation with stable vitals are suggestive of a simple pneumothorax. Management involves inserting a chest tube.

Traumatic pneumothorax can be:

- Simple pneumothorax - with hemodynamic stability
- Tension pneumothorax - without hemodynamic stability

Initial management of simple pneumothorax with open sucking wound involves prompt closing of the defect with a sterile occlusive plastic dressing (e.g., Opsite), taped on three sides to act as a flutter-type valve followed by chest tube insertion as soon as possible in a site remote from the injury site.

Other options:

Option A: Features of tension pneumothorax include hypotension, dyspnea, distended neck veins, tracheal deviation, absent or decreased breath sounds, hyper-resonant note on percussion and normal heart sounds. The first line of management is needle thoracocentesis in the 5th intercostal space in the midaxillary line.

Definitive management - insertion of an intercostal tube in the triangle of safety. If a sucking wound is present, then it is covered with gauze pieces and taped on 3 sides.

Option B and D: Pericardial tamponade resembles tension pneumothorax but there is no tracheal deviation, normal breath sounds, normal percussion note and muffled heart sounds. Initial management is by needle pericardiocentesis followed by definitive sternotomy or lateral thoracotomy.

REMEMBER RECENT UPDATE :(Acc to ATLS 10th edition):

-
- In adults, the needle thoracotomy location changed to the 5th ICS mid-axillary line from the 2nd ICS midclavicular line.
- In the pediatric age group, the thoracotomy location remains at the 2nd ICS mid-clavicular line.

Surgery AIIMS 2018

Question 1:

All are true about intestinal transplantation except _____.

- a) Most of the grafts are multi-visceral grafts.
- b) It is not widely used due to vigorous rejection reactions.
- c) The most dangerous complication is sepsis.
- d) The most common indication is the ischemia

Question 2:

All of the following are boundaries of the triangle of doom except?

- a) Vas deferens
- b) Peritoneal Reflection
- c) Iliopubic tract
- d) Spermatic vessels

Question 3:

All of the following are true about superior mesenteric artery syndrome except?

- a) Third part of duodenum is compressed
- b) Angle between SMA and aorta is more than 70 degrees
- c) Surgical management is by duodenojejunostomy
- d) Also known as Wilkie's Syndrome

Question 4:

Which of the following type of renal calculi is resistant to the extracorporeal shockwave lithotripsy?

- a) Struvite
- b) Cystine

- c) Calcium oxalate
- d) Uric acid

Question 5:

An abdominal aortic aneurysm is operated when its size is greater than _____

- a) 5 cm
- b) 6 cm
- c) 5.5 cm
- d) 6.5 cm

Question 6:


A 3-day-old neonate presented with recurrent bilious vomiting. An X-ray erect abdomen is given below. What is the next best step in managing this condition?



- a) CT abdomen
- b) Gastrograffin follow-through
- c) USG abdomen
- d) Explorative laparotomy

Question 7:

The image shows methylene blue dye being injected in the peritumoral region. What is this procedure useful for?



- a) Localizing the tumor
- b) Identifying the margins
- c) Sentinel lymph node biopsy
- d) Identifying metastasis

Question 8:

A 3-month-old child presented with umbilical discharge. Abdomen examination was normal except for the below finding. The next step of management is?



- a) MRI Abdomen
- b) USG Abdomen
- c) CECT
- d) Exploratory laparotomy

Question 9:

Which of the following cancers rarely metastasizes to the liver?

- a) Colon
- b) Breast
- c) Prostate
- d) Melanoma

Question 10:

A 40-year-old female presents to the ER with a history of RTA. On examination, GCS is 15/15, heart rate = 110/min, BP = 90/60 mmHg, and there is elevated JVP. The right chest is hyperresonant on percussion with decreased breath sounds. The left chest appears normal. But the patient is dyspneic and ecchymosis is noted over the chest wall. The abdomen was soft. The probable diagnosis is:

- a) Cardiac tamponade
- b) Tension pneumothorax
- c) Pericardial effusion
- d) Massive hemothorax

Question 11:

A 50-year-old male presents to the emergency with a BP of 160/100 mmHg and a heart rate of 120. CECT done reveals a Stanford A aortic dissection. What is the initial management of this given condition?

- a) Beta blocker medication
- b) LMWH
- c) Surgical repair
- d) Vitamin K inhibitors


Question 12:

A 22-year-old male presented with a hard painless testicular swelling. There were no inguinal, intra-abdominal or mediastinal lymph nodes on CT Scan. The blood investigations showed AFP levels of > 3000 . USG shows uniform echotexture and small areas of necrosis. The next line of management is _____.

- a) High inguinal orchiectomy
- b) Core needle trucut biopsy
- c) FNAC
- d) Whole body PET/CT

Question 13:

A condition requires the surgery depicted in the image shown below. All the following are usually seen associated with this condition except?



- a) Absent kidney
- b) Hepatic artery anomalies
- c) Malrotation of gut
- d) Polysplenia

Question 14:

You are doing an emergency laparotomy for a case of intestinal obstruction. Which organ will you first visualize to know if this is a case of small bowel or large bowel obstruction?

- a) Ileum
- b) Caecum
- c) Jejunum
- d) Sigmoid colon

Question 15:

What is the best position for a conscious patient to be in during the insertion of a Ryle's tube?

- a) Supine with neck flexion
- b) Supine with neck extension
- c) Sitting with neck flexion
- d) Sitting with neck extension

Question 16:

Pick from the following the usual cause of thyroid storm in thyroidectomy:

- a) Rough handling of thyroid during surgery
- b) Due to parathyroid gland
- c) Inadequate patient preparation
- d) Infections

Question 17:

Which of the following colored cannula will you use to achieve maximum flow rate, in a patient who needs a large amount of fluids?

- a) Grey
- b) Green
- c) Blue
- d) Pink

Question 18:

A patient who was admitted after a road traffic accident is put on mechanical ventilation. He opens his eyes on verbal command and moves all four limbs spontaneously. Calculate his GCS.

- a) Eyes-2, Verbal-1, Motor-5
- b) Eyes-2, Verbal-NT, Motor-5
- c) Eyes-3, Verbal-1, Motor-6
- d) Eyes-3, Verbal-NT, Motor-6

Question 19:

A patient was brought to the casualty after a road traffic accident (RTA) with chest trauma. He is conscious and speaks single words. He has a respiratory rate of 44 cycles per min and a BP of 90/40 mmHg. There is a hyper-resonance on percussing the right side of his chest. What will be your next, immediate step?

- a) Intubate the patient
- b) Insert needle in appropriate intercostal space
- c) Start intravenous fluids urgently
- d) Take a chest X-ray

Question 20:

Heller's operation is done for _____.

- a) Achalasia cardia
- b) Carcinoma esophagus
- c) Esophageal stricture
- d) Hypertrophic pyloric stenosis

Answer Key

Question No.	Correct Option
1	a
2	c
3	b

4	b
5	c
6	b
7	c
8	b
9	c
10	b
11	a
12	a
13	a
14	b
15	c
16	c
17	a
18	d
19	b
20	a

Detailed Explanations

Solution to Question 1:

Multivisceral grafts are rarely used in intestinal transplantation. The most common intestinal graft is small bowel graft, with or without the colon or a combined liver and small bowel graft.

All the other statements are correct.

Small bowel transplantation is the final treatment option after the outcome of reconstruction and lengthening procedures are unsatisfactory.

The small bowel transplant is highly immunogenic since it contains a large amount of gut-associated lymphoid tissue, donor dendritic cells, and macrophages. Acute rejection has been reported in 56–75% of patients, with small bowel transplant limiting the overall survival to only one-third of recipients.

Solution to Question 2:

Iliopubic tract does not form a boundary of the triangle of doom.

The triangle of doom is bounded

- Medially by the vas deferens


- Laterally by the vessels of the spermatic cord
- Inferiorly by peritoneal reflection

Apex oriented superiorly at the internal ring

The contents of the space include

- External iliac vessels
- Deep circumflex iliac vein
- Femoral nerve
- Genital branch of the genitofemoral nerve

Note: Staples are avoided in this region during hernia repair due to the risk of injury to the vessels which might result in severe hemorrhage.



Solution to Question 3:

Option B is the false statement. In superior mesenteric artery (SMA) syndrome, the aortomesenteric angle is reduced to less than 25 degrees. The normal aortomesenteric angle is approximately between 38-65 degrees.

SMA syndrome, also called Wilkie's syndrome (option D), is characterized by an obstruction of the third part of the duodenum by the SMA (option A) due to compression against fixed retroperitoneal structures.


Symptoms include episodic epigastric pain associated with vomiting. Risk factors include rapid childhood growth without compensatory weight gain, profound weight loss, abdominal surgery, inflammatory diseases of the abdomen, and immobilization in a cast that increases lordosis and accentuates the angle of the SMA with the aorta.

Abdominal CT angiography is the investigation of choice. The CT image below shows narrowed aortomesenteric angle in SMA syndrome:



Lifestyle therapy includes frequent small feedings and patients may benefit from lying prone or on their left side postprandially.

In some cases, weight gain or removal of a body cast improves symptoms. Laparoscopic lysis of the ligament of Treitz is beneficial in few cases. The preferred surgical treatment is Strong's procedure involving duodenojejunostomy (option C).



Solution to Question 4:

Cystine stones are resistant to extracorporeal shockwave lithotripsy (ESWL) due to their hardness. Cystine, brushite, and calcium oxalate monohydrate stones are the most refractory to ESWL fragmentation, with cystine and brushite being the most resistant.

ESWL is a non-invasive method that generates shockwaves outside the body that focus on the stones. It is the most common method of treating urinary tract stones nowadays. Stones up to approximately 1.5 cm in size are suitable for this form of treatment.

Complications: Infection, haematuria, parenchymal hemorrhage, and perirenal hematoma.

Contraindications: Obese patients, pregnant patients, and patients taking oral anticoagulants.

Note: Stone fragments collecting in the distal ureter following ESWL are known as steinstrasse.

Solution to Question 5:

An asymptomatic abdominal aortic aneurysm in an otherwise fit patient should be considered for repair if ≥ 5.5 cm in diameter (measured by ultrasonography). The annual incidence of rupture rises from 1% or less in aneurysms that are ≤ 5.5 cm in diameter to a significant level of nearly 25%, in those that are 7 cm in diameter.

As open elective surgery (transabdominal) carries a 5% mortality rate, the balance is in favor of elective operation once the maximum diameter is ≥ 5.5 cm.

6 monthly follow-up ultrasonographic assessment is indicated for asymptomatic aneurysms ≤ 5.5 cm in diameter.

Solution to Question 6:

Explanation:

The given clinical history with an x-ray showing a triple bubble sign is suggestive of jejunal atresia. The next best step is to perform a contrast study to localize the site of obstruction.

Small bowel atresias present with intestinal obstruction soon after birth.

- Bilious vomiting is the dominant feature in jejunal atresia
- Abdominal distension is more prominent in ileal atresia

A small amount of pale meconium may be passed, despite the atresia.

Plain abdomen radiographs in a jejunal/ileal atresia case will show multiple air-fluid levels and dilated loops. They don't reliably distinguish small and large bowels in neonates and hence contrast studies are often required to localize the obstruction.

Water-soluble enemas (Gastrogaffin, Hypaque) are particularly useful in differentiating atresia from meconium ileus and Hirschsprung disease. Contrast studies also help to rule out distal intestinal obstruction and occasionally may act therapeutically to relieve obstructions.

In most cases of jejunal/ileal atresia, the distal end of the dilated proximal small bowel is resected and a primary end-to-end anastomosis is done. If the proximal bowel is extremely dilated, it may need to be tapered to the distal bowel before anastomosis.

Other options:

Option A: CT is not recommended for neonates owing to the harmful effects of radiation.

Option C: USG abdomen does not help to localize the site of obstruction and is more useful in the diagnosis of intussusception, where there is an abdominal mass.

Option D: Explorative laparotomy is not done for diagnosing intestinal atresias.

Solution to Question 7:

The utility of injection of methylene blue in the region around the tumor is done for sentinel lymph node biopsy (SLN) in patients without clinically positive lymph nodes.

SLN biopsy rationale:

- Afferent lymphatics from a primary tumor first drain into the sentinel node before draining into the other nodes. So, if a sentinel lymph node is positive for metastasis, it is possible that the other axillary nodes are also involved. Hence, axillary dissection would be necessary.
- If the sentinel node is tumor-free, then regional nodal metastasis is unlikely.

The nodes can be identified using fluorescein dye, methylene blue, or radioactive sulfur colloid-labeled albumin. The dye can be injected into the peritumoral tissue, subareolar, periareolar, or intradermal plane.


Contraindicated in patients with inflammatory breast cancer, stage T4 disease, a history of previous breast cancer or chest wall surgery, breast scarring, and radiotherapy.

Solution to Question 8:

The given clinical image is that of a congenital umbilical anomaly such as patent urachus, urachal cyst, and vitelline cyst (resulting from the persistence of omphalomesenteric duct). The next best step is the USG abdomen to diagnose if the anomaly is cystic, fibrous or fistula and its connections.

Omphalomesenteric duct abnormalities: Vitello intestinal duct or omphalomesenteric duct is the connection between the yolk sac and the midgut during embryonic life, which eventually gets obliterated. The persistence of the vitellointestinal duct leads to various abnormalities depending on which part of the duct remains patent.


- Persistence of the proximal intestinal end of the duct leads to Meckel's diverticula.
- Persistence of the distal end near the umbilicus leads to the umbilical polyp or sinus.
- Persistence of the distal end with mucus retention leads to the cyst, which may be associated with the intestine or the umbilicus by a fibrous band.
- A fully patent vitellointestinal duct can lead to an enterocutaneous fistula.



Patent urachus occurs due to the failure of urachus to obliterate, resulting in a connection between the bladder and the umbilicus.

Urachal cyst occurs when the urachus fails to obliterate but there is no connection between the bladder and the umbilicus.

The given image below shows a urachal cyst (UC) attached to Meckel's diverticulum (MD) on one side and urinary bladder (UB) on the other side with a patent tract- urachus (U).



Solution to Question 9:

Prostate cancers rarely metastasize to the liver.

The most common malignant tumors of the liver are metastatic lesions. The liver is a common site of metastases from gastrointestinal tumors because of dissemination through the portal

venous system. The most relevant metastatic tumor of the liver to the surgeon is colorectal cancer because of the well-documented potential for long-term survival after complete resection.

The more frequent primary sites include carcinomas of the colon, pancreas, lung, and breast in adults, and neuroblastoma, Wilms tumor, and rhabdomyosarcoma in pediatric patients. Although the origin of the tumor is often known when metastases are biopsied, carcinomas of the pancreas, stomach, and lung are particularly notorious for being occult primary neoplasms.

Solution to Question 10:

The given clinical scenario of trauma with decreased breath sounds and a hyper-resonant note on percussion indicates pneumothorax. The presence of tachycardia, hypotension and elevated JVP is suggestive of tension pneumothorax.

Tension pneumothorax develops when there is a parietal or visceral pleural tear. This tear acts as a one-way valve i.e. air enters the pleural space from the lung or chest wall during inspiration but has no way to escape during expiration, leading to the collapse of the affected lung.

It is characterized by chest pain, tachypnoea, acute dyspnea, sudden onset hemodynamic compromise (tachycardia, hypotension, low O₂ saturation, loss of consciousness), neck vein distention, tracheal deviation away from the side of injury, decreased chest movements on the affected side, hyper-resonance on percussion on the affected side and absent or decreased breath sounds on the affected side.


Tension pneumothorax is a clinical diagnosis, and treatment should never be delayed awaiting radiological confirmation.

Management includes:

- Immediate management: Needle decompression or needle thoracostomy with a wide bore needle in the 5th intercostal space anterior to the mid-axillary line. In the pediatric age group, the needle thoracostomy location is at the 2nd intercostal space in the mid-clavicular line.
- Definitive management: Tube thoracostomy or chest tube insertion in the 5th intercostal space in the triangle of safety.

The triangle of safety is bounded anteriorly by the lateral border of the pectoralis major, posteriorly by latissimus dorsi and inferiorly by a horizontal line at the level of the nipple.

The X-ray below shows a left-sided tension pneumothorax. There is a large, well-demarcated area devoid of lung markings on the left side. The yellow arrows are pointing to the tracheal deviation and the mediastinal shift to the right side.



Other options:

Options A: Pericardial tamponade can mimic tension pneumothorax, but the absence of a tympanic note on percussion rules out its possibility. A patient with pericardial tamponade can present with hypotension, raised JVP, and muffled heart sounds (Beck's triad), with normal breath sounds and no tracheal shift.

Option C: Dullness in percussion at the base of the left inferior angle of scapula in conjunction with tubular breath sounds (Ewart's sign), chest pain that improves while sitting upright and worsens when lying down and pulsus paradoxus are indicative of pericardial effusion. Pericardial effusion develops into tamponade when the effusion leads to increased pressure, impairing ventricular filling and reducing cardiac output.

Option D: Massive hemothorax is the accumulation of >1500 ml of blood or one-third or more of the patient's blood volume in the chest cavity. The patient will have collapsed neck veins, decreased breath sounds and a dull note on percussion.

Solution to Question 11:

The initial management of patients with all types of aortic dissection is beta blocker medication to control hypertension along with pain management. Patients with Stanford Type A (DeBakey Types I and II) will require surgical intervention once the patient's blood pressure is stabilized.

Aortic dissection occurs when a defect or flap occurs in the intima of the aorta.

Predisposing factors:

- Age
- Hypertension
- Marfan syndrome
- Pregnancy


- Other connective tissue disorders, for example, Ehlers–Danlos syndrome, giant cell arteritis, systemic lupus erythematosus
- Coarctation of the aorta
- Turner or Noonan syndromes
- Aortic cannulation site (iatrogenic).

The most typical presentation of aortic dissection is a tearing interscapular pain. The dissection may also result in aortic wall rupture into the pericardium (cardiac tamponade) or mediastinum (left haemothorax).

Contrast-enhanced CT is currently the most preferred imaging modality for aortic dissection. Double-lumen aorta is seen on CECT. Transesophageal echocardiography is preferred for patients who are unstable for transportation.

Acute aortic dissection can be classified by the original DeBakey system, based on the location of the proximal tear and distal extent of the dissection flap, or the Stanford system which uses the location of the proximal tear alone.

Site of dissection	DeBakey	Stanford
Both ascending and descending aorta	Type I	Type A
Only ascending aorta and arch of aorta	Type II	Type A
Descending the aorta distal to the subclavian artery	Type IIIa Limited to the thoracic aorta	Type B
Type IIIb Extends to the abdominal aorta		




Management of aortic dissection:

The initial management of patients with all types of aortic dissection is beta-blocker medication to control hypertension along with pain management. The preferred beta-blocker drug is esmolol. Hemodynamic stability is an important consideration.

- Stanford Type A (DeBakey Types I and II) requires surgical intervention once the patient's blood pressure is stabilized.
- Stanford Type B (DeBakey type III) is best managed by antihypertensive drugs. For patients with complicated type B dissection, percutaneously placed endovascular stents are currently preferred. Surgery is reserved for the rare patients not suitable for stenting.

Other options:

Options B and D: Low molecular weight heparin and vitamin K inhibitors are contraindicated in aortic dissection as they can worsen the risk of bleeding.



Solution to Question 12:

The given clinical features of painless testicular swelling with raised AFP levels without distant metastasis point towards the likely diagnosis of stage 1 non-seminomatous germ cell tumour (NSGCT) of the testis. The next step in management is a high inguinal orchidectomy.

Core needle trucut biopsy and FNAC should be never done when there is a suspicion of testicular neoplasm as it will result in the seeding of tumor cells into the skin and lymphatics, leading to upstaging of the tumor (Options B and C). Currently, there is no role for PET-CT in the routine evaluation of non-seminomatous germ cell tumor and seminoma at the time of diagnosis (Option D).

Testicular tumors are classified according to their predominant cell type:

- Germ cell tumors (90-95%): seminomatous (seminoma) or NSGCT (embryonal carcinoma, yolk sac tumor, teratoma and choriocarcinoma).
- Sex cord-stromal tumors (1-2%): Leydig cell tumors, Sertoli cell tumors

- Lymphoma (3-7%)
- Other tumors (1-2%)

The most common presentation of testicular cancer is painless testicular mass. Pain is more commonly associated with NSGCT and two-thirds of patients present with metastatic disease at the time of diagnosis. The symptoms of metastatic disease include abdominal pain, dyspnea, chest pain, cough or hemoptysis.

Management of testicular tumours:

- Ultrasound of the testis should be done in all suspected cases of a testicular tumour to confirm the diagnosis.
- Tumor markers (AFP, hCG, LDH) are measured for the staging of cancer.
- CT- chest, abdomen, and pelvis are done to detect metastasis.
- Orchidectomy via an inguinal incision (high inguinal orchidectomy) is the best line of treatment in case of a suspected testicular tumour.
- Further management is based on staging. The older staging is preferred over the TNM staging for treatment and is given below:
 - Stage I: Tumour is confined to testis and epididymis
 - Stage II: Nodal tissue is present but is confined to nodes below the diaphragm
 - Stage III: Nodal involvement above the diaphragm
 - Stage IV: Non-lymphatic metastatic spread (mostly in lungs)
- Stage I tumours:
 - Combination chemotherapy with bleomycin, etoposide, and cis-platinum (BEP chemotherapy) is given for NSGCTs, as they are radio-resistant.
- Chemotherapy with cisplatin and radiotherapy to para-aortic lymph nodes can be given in seminomas as they are radiosensitive.
- Stage II-IV tumours (seminomas and NSGCTs) are treated with BEP chemotherapy and retroperitoneal lymph node dissection if needed.

Solution to Question 13:

The given image shows a Roux-en-Y loop of the small intestine (jejunum) attached to an area of the liver capsule (Roux-en-Y hepaticojejunostomy), which is suggestive of Kasai hepatportoenterostomy procedure done for biliary atresia in infants. An absent kidney is not associated with this condition.

Extrahepatic biliary atresia (EHBA) is defined as a partial or total absence of permeable bile duct between porta hepatis and the duodenum. It is due to inflammatory fibrosis of the biliary tree. It leads to obstruction to bile flow thereby causing leaking of conjugated bilirubin into the circulation.

Associated anomalies include cardiac lesions, polysplenia, situs inversus, absent vena cava, and a preduodenal portal vein.

In the Kasai procedure, the extrahepatic biliary tree is dissected proximally till the liver capsule, the porta hepatis is transected and a jejunal loop is anastomosed in order to provide a drainage pathway for bile.

Infants with biliary atresia, who do not achieve successful drainage after Kasai's procedure, require liver transplantation within the first year of life. This is the most common reason for liver transplantation in children.

Management of extrahepatic biliary atresia		
Type I	Atresia restricted to the common bile duct	Roux-en-Y hepaticojejunostomy
Type II	Atresia of the common hepatic duct	Kasai procedure(Roux-en-Y hepatoportoenterostomy)
Type III	Atresia of the right and left hepatic duct	

Solution to Question 14:

The cecum is the first organ to be visualized to differentiate between the small and large bowel in cases of intestinal obstruction.

Distension of the cecum will confirm large bowel involvement. The identification of a collapsed distal segment of the large bowel and its sequential proximal assessment will lead to the identification of the cause.

Solution to Question 15:

The best position for a conscious patient to be in during the insertion of Ryle's tube (RT) is sitting with neck flexion. The described position helps in directing the RT towards the esophagus and also prevents it from coiling in the oropharynx or entering the trachea.

RT, also known as nasogastric (NG) tube, measures 120-125 cm. It has 4 markings at 40cm (cardia), 50cm (body), 60cm (pylorus), and 70 cm (duodenum) to indicate the approximate position after the placement.

Indications for NG tube:

- Feeding in patients with dysphagia, decreased appetite, or those who are nil per os (NPO) for medical or surgical reasons.
- Aspiration of gastric fluid
- Gastric decompression in cases of intestinal obstruction and paralytic ileus.
- Zollinger-Ellison syndrome
- Bowel irrigation

The absolute contraindications for an NG tube are severe midface trauma, recent nasal surgery, and basilar skull fractures

Steps for the correct insertion of RT in an adult:

- Examination of the patient's nostril for any anatomical abnormalities and then instillation with 10 ml of 1% lignocaine
- NEX method is used to measure the length of the RT: Place the exit port of the tube at the tip of the nose. Extend the tube to the earlobe, and then to the xiphisternum.
- The patient is asked to sit upright with partial neck flexion. The tip of the Ryle's tube is lubricated and passed along the floor of the nasal passage until a resistance is felt (corresponds to nasopharynx)
- The patient is subsequently asked to swallow water to guide the tube through the esophagus into the stomach until the marked length (based on the NEX method).
- NG tube is finally taped to secure it in place.

Confirmation of the correct position is done by:

- Auscultation in the epigastric area while injecting a bolus of air
- X-ray chest to locate the lead markers
- Aspiration of contents and subsequently confirming gastric origin (pH ≤ 6 indicates gastric placement)



Solution to Question 16:

The usual cause of thyroid storm in thyroidectomy is inadequate patient preparation before surgery. Patient preparation pre-operatively involves ensuring a euthyroid state.

Rough handling of the thyroid in thyroidectomy does not cause a thyroid storm. Though a significant amount of thyroglobulin is stored in the acinar cells, they are inactive unless converted

to thyroxine (T₃ and T₄).

Solution to Question 17:

Grey colored cannula is used to achieve maximum flow rate in a patient who needs a large amount of fluids.

Solution to Question 18:

The patients GCS is Eyes-3, Verbal-NT, Motor-6.

It is calculated as follows:

- Eye opening: Opening to verbal command - 3
- Verbal response: In the case of endotracheal intubation, the verbal response score is noted as NT (non-testable)
- Motor response: Movement of all 4 limbs spontaneously - 6

Total GCS score is not recorded since one of the components is non-testable.

Solution to Question 19:

The given clinical scenario of chest trauma with a hyper resonant note on the right side suggests pneumothorax. Hemodynamic instability in this patient suggests tension pneumothorax and this is managed by immediate decompression by insertion of a needle in the appropriate place.

Pneumothorax refers to the presence of excess air in the pleural space. Entrapment of air occurs by:

- Primary spontaneous pneumothorax - Due to the spontaneous rupture of apical pleural blebs or cystic spaces immediately below the visceral pleura.
- Secondary pneumothorax - Due to the rupture of dilated cystic spaces in lung diseases like chronic obstructive pulmonary disease (COPD)
- Tension pneumothorax - Due to a one-way valve mechanism that directs airflow into the pleural space leading to a higher than atmospheric pressure environment. This leads to the affected lung being unable to expand appropriately and leads to hemodynamic instability.

Tension pneumothorax is a surgical emergency and the excessive pressure has to be immediately relieved. A chest X-ray is not required before emergency decompression as it is a clinical diagnosis, but an X-ray should be done following chest tube insertion (option D). Tension pneumothorax is managed by immediate needle decompression in the 5th ICS (adults), followed by definitive management with a chest tube inserted in the same space.

Other options:

Option A: Intubation is performed when the patient is unresponsive post chest tube insertion.

Option C: While IV fluids are a part of the resuscitation protocol, the management of tension pneumothorax takes higher precedence as the next immediate step.

Note: According to ATLS 10th edition

- In adults, needle thoracostomy or needle decompression location is changed to the 5th ICS anterior to the midaxillary line from the 2nd ICS midclavicular line.
- In the pediatric age group, the needle thoracostomy location remains at the 2nd ICS midclavicular line.

Solution to Question 20:

The most common surgical procedure for achalasia cardia is laproscopic Heller myotomy in conjugation with partial fundoplication.

Surgery AIIMS 2019 (May & Nov)

Question 1:

A patient presents with renal calculus of > 4 cm of the staghorn type. What is the best management?

- a) Extracorporeal shockwave lithotripsy
- b) Percutaneous nephrolithotomy
- c) Ureteroscopic removal
- d) Retrorenal surgery

Question 2:

You are doing an emergency laparotomy for a case of intestinal obstruction. Which organ will you first visualize to know if this is a case of small bowel or large bowel obstruction?

- a) Ileum
- b) Caecum
- c) Jejunum
- d) Sigmoid colon

Question 3:

What is the best position for a conscious patient to be in during the insertion of a Ryle's tube?

- a) Supine with neck flexion
- b) Supine with neck extension
- c) Sitting with neck flexion
- d) Sitting with neck extension

Question 4:

Pick from the following the usual cause of thyroid storm in thyroidectomy:

- a) Rough handling of thyroid during surgery
- b) Due to parathyroid gland

- c) Inadequate patient preparation
- d) Infections

Question 5:

Which of the following colored cannula will you use to achieve maximum flow rate, in a patient who needs a large amount of fluids?

- a) Grey
- b) Green
- c) Blue
- d) Pink

Question 6:

A patient who was admitted after a road traffic accident is put on mechanical ventilation. He opens his eyes on verbal command and moves all four limbs spontaneously. Calculate his GCS.

- a) Eyes-2, Verbal-1, Motor-5
- b) Eyes-2, Verbal-NT, Motor-5
- c) Eyes-3, Verbal-1, Motor-6
- d) Eyes-3, Verbal-NT, Motor-6

Question 7:

A patient was brought to the casualty after a road traffic accident (RTA) with chest trauma. He is conscious and speaks single words. He has a respiratory rate of 44 cycles per min and a BP of 90/40 mmHg. There is a hyper-resonance on percussing the right side of his chest. What will be your next, immediate step?

- a) Intubate the patient
- b) Insert needle in appropriate intercostal space
- c) Start intravenous fluids urgently
- d) Take a chest X-ray

Question 8:

Which of the following is used for wound infection grading?

- a) ASA classification
- b) SIRS criteria
- c) Southampton grading
- d) APACHE II score

Question 9:

The most common mediastinal tumor is _____.

- a) Thymoma
- b) Neurogenic
- c) Lymphoma
- d) Bronchial cyst

Question 10:

On the 5th postoperative day post CABG, the patient develops tachypnea and hypotension. What is the most probable cause?

- a) Sepsis
- b) Acute kidney failure
- c) Acute respiratory failure
- d) Acute cardiac failure

Question 11:

The probable diagnosis based on the below clinical image of the breast is?



- a) Paget's disease
- b) Mondor's disease
- c) Zuska's disease
- d) Tietze syndrome

Question 12:

The most commonly involved artery in pseudoaneurysm formation:

- a) Aorta
- b) Femoral artery
- c) Popliteal artery
- d) None of the above

Question 13:

A chronic alcoholic patient came to the emergency with pain in the epigastrium radiating to the back and recurrent vomiting. On examination, guarding was present in the upper epigastrium. The erect chest X-ray was normal. What is the next step?

- a) Upper GI endoscopy
- b) Alcohol breath test
- c) Serum lipase
- d) CECT

Question 14:

The best way to manage the wound shown in the image is:



©MARROW

- a) Cleaning and dressing
- b) Debridement and dressing
- c) Cleaning and skin grafting
- d) Cleaning and use of allograft

Question 15:

Device shown in the image is used for which purpose?



©Marrow

- a) Prevent viral infections
- b) Prevent transfusion related reactions
- c) Prevent infections
- d) All of the above

Question 16:

In a child with a unilateral undescended testis, a diagnostic laparoscopy is performed which showed the blind-ending testicular vessels on the right side. What is the next best step?

- a) Inguinal exploration
- b) Abdominal exploration
- c) Scrotal exploration
- d) No further exploration

Question 17:

What is the surgical site infection rate expected in a patient with a clean-contaminated wound that is operated on after antibiotics prophylaxis?

- a) 1-2%
- b) 3%
- c) 13-20%
- d) 20-30%

Question 18:

Which of the following is not a component of qSOFA?

- a) Respiratory rate ≥ 22 /min
- b) Systolic BP ≤ 100 mm Hg
- c) Altered mental status
- d) Bilateral pupillary reflex

Question 19:

Which of the following does not classify as locally advanced breast cancer?

- a) Tumor more than 4 cm
- b) Inflammatory breast cancer
- c) Chest wall involvement
- d) Skin involvement

Question 20:

All of the following are features of SIRS except:

- a) Temperature > 38 degrees Celsius
- b) Respiratory Rate > 20 /minute and PaCO₂ < 35 mmHg
- c) Pulse rate > 90 /minute
- d) WBC > 12000 /mm³ or < 4000 /mm³

Question 21:

The most common neuroendocrine tumor of the pancreas is

- a) Gastrinoma
- b) Somatostatinoma
- c) Insulinoma
- d) VIPoma

Question 22:

After parotidectomy patients complains of loss of sensation while shaving. Which nerve is most likely affected?

- a) Auriculotemporal nerve
- b) Great auricular nerve
- c) Facial nerve
- d) Mandibular nerve

Question 23:

ABCDE rule applies for:

- a) Malignant melanoma
- b) BCC
- c) SCC
- d) Actinic keratosis

Question 24:

Identify the following instrument?



- a) Unipolar cautery
- b) Bipolar cautery

- c) Harmonic scalpel
- d) Ligasure

Question 25:

Radical neck dissection spares which of the following structures?

- a) Digastric muscle
- b) Sternocleidomastoid muscle
- c) Spinal accessory nerve
- d) Ipsilateral cervical lymph nodes

Question 26:

A patient comes with active variceal bleeding. Which of the following is not done?

- a) Platelet transfusion
- b) Arrange pRBC
- c) IV Octreotide
- d) Urgent Endoscopy

Question 27:

A child presents with pain and swelling in the right side of his scrotum and testicular torsion is suspected. Which is the most appropriate management?

- a) Immediate exploration on both symptomatic and asymptomatic side
- b) Immediate exploration on symptomatic side only
- c) Delayed exploration on both symptomatic and asymptomatic side
- d) Delayed exploration on symptomatic side only

Question 28:

What is the instrument shown below that is used to create a pneumoperitoneum ?



- a) Veress needle
- b) Salah's needle
- c) Hasson Trocar
- d) Von Graefe needle

Question 29:

Which of the following is the principle behind the wound-healing procedure shown in the image below?



- a) Negative pressure wound therapy

- b) Positive pressure wound therapy
- c) Foam therapy
- d) Sponge therapy

Question 30:

Stage the bed sore in the following image:



- a) 4
- b) 3
- c) 2
- d) 1

Question 31:

A patient with a road traffic accident presented to the emergency room and requires a large amount of fluids to be transfused. Which of the following IV cannula is preferred?

- a) Green
- b) Blue
- c) Pink
- d) Grey

Question 32:

A patient with a nodule of the neck presents to the surgery OPD. A junior resident performs the following method to palpate it. Identify the procedure shown in the video below.



- a) Lahey's method
- b) Crile's method
- c) Kocher's method
- d) Pemberton's method

Question 33:

The instrument given in the image is:



- a) Hemovac
- b) Jackson-Pratt drain
- c) Minivac
- d) Corrugated drain

Answer Key

Question No.	Correct Option
1	b
2	b
3	c
4	c
5	a
6	d
7	b
8	c
9	a
10	c
11	b
12	b
13	c
14	a

15	b
16	d
17	b
18	d
19	a
20	b
21	c
22	b
23	a
24	a
25	a
26	a
27	a
28	a
29	a
30	a
31	d
32	a
33	b

Detailed Explanations

Solution to Question 1:

The best management for staghorn renal calculus of ≥ 4 cm is percutaneous nephrolithotomy (PCNL).

The initial management of renal stones depends on the stone size and includes conservative management, extracorporeal shockwave lithotripsy (ESWL), and PCNL. PCNL is preferred for stones ≥ 2 cm in size and for staghorn calculi, while ESWL is used to treat renal stones of size 0.6 to 1.5 cm. PCNL is indicated in

- Renal stones ≥ 2 cm
- Staghorn calculus
- Failed shockwave lithotripsy (SWL) or retrograde intrarenal surgery (RIRS) for renal calculi
- Lower pole renal stones with anatomy that is unfavorable for ESWL

Other options:

Option C: Ureteroscopic removal is attempted for failed SWLs or PCNLs, and is not the first choice in the surgical management of renal stones.

Option D: RIRS is a minimally invasive procedure similar to ureteroscopic removal, where the kidney is accessed using a slim and flexible ureteroscope in a retrograde manner via the ureteric orifice. It is indicated for lower pole renal stones < 2 cm.

Solution to Question 2:

The cecum is the first organ to be visualized to differentiate between the small and large bowel in cases of intestinal obstruction.

Distension of the cecum will confirm large bowel involvement. The identification of a collapsed distal segment of the large bowel and its sequential proximal assessment will lead to the identification of the cause.

Solution to Question 3:

The best position for a conscious patient to be in during the insertion of Ryle's tube (RT) is sitting with neck flexion. The described position helps in directing the RT towards the esophagus and also prevents it from coiling in the oropharynx or entering the trachea.

RT, also known as nasogastric (NG) tube, measures 120-125 cm. It has 4 markings at 40cm (cardia), 50cm (body), 60cm (pylorus), and 70 cm (duodenum) to indicate the approximate position after the placement.

Indications for NG tube:

- Feeding in patients with dysphagia, decreased appetite, or those who are nil per os (NPO) for medical or surgical reasons.
- Aspiration of gastric fluid
- Gastric decompression in cases of intestinal obstruction and paralytic ileus.
- Zollinger-Ellison syndrome
- Bowel irrigation

The absolute contraindications for an NG tube are severe midface trauma, recent nasal surgery, and basilar skull fractures


Steps for the correct insertion of RT in an adult:

- Examination of the patient's nostril for any anatomical abnormalities and then instillation with 10 ml of 1% lignocaine
- NEX method is used to measure the length of the RT: Place the exit port of the tube at the tip of the nose. Extend the tube to the earlobe, and then to the xiphisternum.
- The patient is asked to sit upright with partial neck flexion. The tip of the Ryle's tube is lubricated and passed along the floor of the nasal passage until a resistance is felt (corresponds to nasopharynx)
- The patient is subsequently asked to swallow water to guide the tube through the esophagus into the stomach until the marked length (based on the NEX method).

- NG tube is finally taped to secure it in place.

Confirmation of the correct position is done by:

- Auscultation in the epigastric area while injecting a bolus of air
- X-ray chest to locate the lead markers
- Aspiration of contents and subsequently confirming gastric origin (pH ≤ 6 indicates gastric placement)



Solution to Question 4:

The usual cause of thyroid storm in thyroidectomy is inadequate patient preparation before surgery. Patient preparation pre-operatively involves ensuring a euthyroid state.

Rough handling of the thyroid in thyroidectomy does not cause a thyroid storm. Though a significant amount of thyroglobulin is stored in the acinar cells, they are inactive unless converted to thyroxine (T₃ and T₄).

Solution to Question 5:

Grey colored cannula is used to achieve maximum flow rate in a patient who needs a large amount of fluids.

Solution to Question 6:

The patient's GCS is Eyes-3, Verbal-NT, Motor-6.

It is calculated as follows:

- Eye opening: Opening to verbal command - 3
- Verbal response: In the case of endotracheal intubation, the verbal response score is noted as NT (non-testable)
- Motor response: Movement of all 4 limbs spontaneously - 6

Total GCS score is not recorded since one of the components is non-testable.

Solution to Question 7:

The given clinical scenario of chest trauma with a hyper resonant note on the right side suggests pneumothorax. Hemodynamic instability in this patient suggests tension pneumothorax and this is managed by immediate decompression by insertion of a needle in the appropriate place.

Pneumothorax refers to the presence of excess air in the pleural space. Entrapment of air occurs by:

- Primary spontaneous pneumothorax - Due to the spontaneous rupture of apical pleural blebs or cystic spaces immediately below the visceral pleura.
- Secondary pneumothorax - Due to the rupture of dilated cystic spaces in lung diseases like chronic obstructive pulmonary disease (COPD)
- Tension pneumothorax - Due to a one-way valve mechanism that directs airflow into the pleural space leading to a higher than atmospheric pressure environment. This leads to the affected lung being unable to expand appropriately and leads to hemodynamic instability.

Tension pneumothorax is a surgical emergency and the excessive pressure has to be immediately relieved. A chest X-ray is not required before emergency decompression as it is a clinical diagnosis, but an X-ray should be done following chest tube insertion (option D). Tension pneumothorax is managed by immediate needle decompression in the 5th ICS (adults), followed by definitive management with a chest tube inserted in the same space.

Other options:

Option A: Intubation is performed when the patient is unresponsive post chest tube insertion.

Option C: While IV fluids are a part of the resuscitation protocol, the management of tension pneumothorax takes higher precedence as the next immediate step.

Note: According to ATLS 10th edition

- In adults, needle thoracostomy or needle decompression location is changed to the 5th ICS anterior to the midaxillary line from the 2nd ICS midclavicular line.
- In the pediatric age group, the needle thoracostomy location remains at the 2nd ICS midclavicular line.

Solution to Question 8:

Southampton wound grading system is used to score the severity of wound site infection.

Southampton wound grading system:

- Grade 0 - Normal healing
- Grade I - Normal healing with different grades of erythema
- Grade II - Erythema with other signs of inflammation at different parts of the wound.
- Grade III - Clear or serosanguinous discharge at different sites
- Grade IV - Pus discharge

Other options:

Option A: American Society of Anesthesiologists (ASA) grading is used in pre-anesthetic checkups to assess a patient's physical health and fitness for anesthesia and surgery.

Option B: Systemic Inflammatory Response Syndrome (SIRS) is a set of clinical criteria used to identify and monitor critically ill patients with high-grade burns, severe trauma, and other major surgeries.

Option D: Acute Physiology and Chronic Health Evaluation II (APACHE II) is a scoring system used to assess the severity and predict the outcome of pancreatitis.

Solution to Question 9:

The most common mediastinal tumors in descending order of incidence are thymoma (25%), germ cell tumor (13%), lymphoma, mesenchymal, and thyroid.

The Masaoka classification and TNM staging are used in the staging of thymoma.

The most common mediastinal tumor in children is the neurogenic tumor.

Solution to Question 10:

The clinical presentation of tachypnoea and hypotension in a post-op day 5 patient who underwent coronary artery bypass grafting (CABG) points towards lung collapse (atelectasis) - type 3 respiratory failure.

Atelectasis is the most common respiratory complication in a post-operative patient.

The factors predisposing to atelectasis are general anesthesia, upper abdominal surgery, and thoracic surgery. They commonly present with tachypnoea and decreased oxygen saturation. On examination, the breath sounds are bronchial in nature and reduced in intensity.

Lung atelectasis can be reversed (within 24h-48h) with early mobilization, deep breathing (5 sequential breaths for 5s), incentive spirometry, and chest physiotherapy.

Other postoperative systemic complications include:

- Pneumonia and myocardial infarction : First few days after surgery
- Venous thromboembolism, kidney failure, and urinary tract infection: Days 4 to 7
- Superficial and deep surgical site infections: 2nd post-op week

Other options:

Option A: For a diagnosis of sepsis, SEPSIS 3 criteria needs to be satisfied:

- A documented infection
- An acute increase in ≥ 2 SOFA points

Option B: Acute kidney failure is defined as a decrease in urine output (< 0.5 mL/kg/h for 6-12h) and a rise in serum creatinine values (≥ 1.5 x baseline or ≥ 0.3 mg/dL).

Option D: Acute cardiac failure has features of hypotension and tachycardia, with peripheral signs of hypoperfusion characterized by cold clammy extremities.

Solution to Question 11:

The image shows a thrombosed subcutaneous cord attached to the skin of the breast which is the pathognomic feature of Mondor's Disease.

Mondor's disease is the thrombophlebitis of the superficial veins of the breast and anterior chest wall. It may also occur in the arm or penis. On raising the arm, the skin over the chest gets stretched and a narrow, shallow groove alongside the cord is seen. It is treated by restricting arm movements and usually subsides within a few months without recurrence, complications, or deformity.

The differential diagnosis to be considered is lymphatic permeation from an occult carcinoma of the breast.

Other options:

Option A: Paget's disease is a type of ductal carcinoma in situ arising from the nipple and commonly presents as nipple erosion.

Option B: Zuska's disease, also called recurrent periductal mastitis, is a condition of recurrent retroareolar infections and abscesses. Smoking has been implicated as a risk factor for this condition. Managed symptomatically by antibiotics coupled with incision and drainage as necessary.

Option D: Tietze's syndrome refers to the benign inflammation of one or more costal cartilages.

Solution to Question 12:


The most commonly involved artery in pseudoaneurysm is the femoral artery.

A true aneurysm involves all three vessel wall layers, while a pseudoaneurysm involves only the intimal and medial layers. Pseudoaneurysms are most commonly iatrogenic due to repeated arterial puncture for blood sampling or catheterization procedures which injure the arterial wall. The most commonly used artery in endovascular surgeries is the femoral artery, and hence, also the most commonly involved in the formation of pseudo aneurysms.

Aneurysms are usually pulsatile on palpation and a systolic bruit can be heard on auscultation.

Doppler ultrasound reveals a bidirectional, 'to-and-fro' flow of blood which is known as the yin-yang sign.

Treatment options include endovascular stent insertion, ultrasound-guided thrombin injection, and surgical repair.



Other options:

Option A: Aortic aneurysm can be types - thoracic or abdominal. They are usually seen in heritable conditions like Marfan syndrome or vascular-type Ehlers-Danlos.

Option C: The most common peripheral aneurysm is the popliteal aneurysm (80% of the cases) followed by the femoral aneurysm.

Solution to Question 13:

The clinical scenario of the patient presenting with abdominal pain, recurrent vomiting, and upper epigastric guarding in a chronic alcoholic is highly suggestive of acute pancreatitis. The normal erect chest X-ray indicates that there is no pneumoperitoneum (effectively ruling out perforation). The next step is to send blood samples for serum lipase levels.

While contrast-enhanced computed tomography (CECT) of the abdomen (option D) is the best investigation to rule out a case of acute pancreatitis, it is not the next step after an abdominal X-ray.

Acute pancreatitis is often caused by gallstones (70%) or alcoholism (25%). The cardinal symptom is acute-onset severe pain in the upper quadrant of the abdomen. The patient usually presents in shock (hypotension and tachycardia) and tachypnoea. Some may also present with signs of bleeding into the fascial planes leading to discoloration of the sites involved:

- Cullen's sign: Around the umbilicus
- Grey Turner's sign: Flanks
- Fox sign: Inguinal region

- Bryant's sign: Scrotum

The diagnosis is often clinical with a laboratory diagnosis of serum lipase preferred over serum amylase. The severity of the acute pancreatitis is assessed using the Ranson and Glasgow scoring system (≥ 3 at 48h of onset of pain). APACHE, SOFA, SAPS, and MODS can also be used to assess the severity of pancreatitis in intensive care (ICU) settings.

Pancreatitis is managed conservatively with initial nasogastric drainage followed by intravenous fluids, analgesia, and frequent monitoring of vital signs, urine output, and blood gases.

Other options:

Option A: Upper gastrointestinal (GI) endoscopy is performed by passing a flexible endoscope and is the best method for examining the upper gastrointestinal mucosa. It doesn't help us diagnose pancreatitis.

Option B: The alcohol breath test is used as a screening test to measure the concentration of alcohol in a person's breath, but doesn't help us diagnose pancreatitis.

Solution to Question 14:

The given image shows a superficial injury to the skin, an abrasion. Cleaning and dressing is the appropriate management of an abrasion.

Wound cleaning is the removal of bacteria and particles to decrease the chances of infection. It includes the "TIME" acronym:

- T - Tissue : Assessment and debridement of non-viable tissue or foreign material
- I - Inflammation/Infection : Assess the etiology of each wound
- M - Moisture Balance : Assess the etiology of wound exudate
- E - Edge of the wound : Assess the edge and surrounding skin

Wound irrigation or toilet is washing the wound with normal saline.

Wound debridement is performed using a surgical blade or fine scissors to remove any devitalized, nonviable tissue or any embedded foreign body (option B).

Skin grafting is required only in situations of wounds with larger areas soft tissue loss (options C and D).

Solution to Question 15:

The image shows a leukoreduction filter which is used for filtering out donor WBCs. This helps in reducing the incidence of transfusion-related reactions by removing WBC microaggregates. The optimum time for performing leukoreduction is before the storage of blood.

Potential benefits include reduced risk of:

- Sensitization to HLA antigens (important for transplant patients & preventing platelet refractoriness)

- Febrile non-hemolytic transfusion reactions (FNHTR)
- CMV transmission

Note: CMV also has an affinity for monocytes and macrophages (types of WBCs). Reduction in WBCs is the cause for the decrease in CMV transmission (accidental benefit), and thus, the main purpose of a leukoreduction filter is not the prevention of CMV transmission (options A and C).

Possible reduced risk of:

- Transfusion-associated graft versus host disease (TA- GVHD)
- Transfusion-related immuno-modulatory (TRIM) effects
- Transmitting variant Creutzfeldt-Jakob Disease

Other Methods of Leucodepletion are:

- Specific leucodepletion filters (Most Efficient)
- Washing
- Freezing
- Buffy coat removal
- Micro-aggregate filtration
- Low leucocyte apheresis devices

Solution to Question 16:

In non-palpable testis cases, laparoscopically encountering blind-ending vas deferens and testicular vessels prior to entering the internal ring is sufficient to diagnose a vanishing testis. In such cases, no further exploration is required.

Cryptorchidism is the failure of the descent of the intra-abdominal testes into the scrotal sac. It is the most common congenital abnormality of the genitourinary tract. The descent of the testes can arrest anywhere, and the most common site is the inguinal region.

Phases of testicular descent:

- Transabdominal phase: Testes come to lie in the lower abdomen or pelvis, and this descent is regulated by Mullerian-inhibiting substances.
- Inguinoscrotal phase: Testes descend along the inguinal canal and reach the scrotal sac, and this phase of the descent is regulated by androgen-induced calcitonin gene-related peptide from the genitofemoral nerve.

The exact pathogenesis remains unknown; however, deficiency in gonadotropins and Mullerian-inhibiting substance (MIS) and increased expression of estradiol in the placenta are thought to play a role.

Cryptorchidism is histopathologically characterized by the following:

- Testicular atrophy: thickening of the basement membrane, arrested germ cell development, marked hyalinization

- Tubules showing Sertoli cells but no spermatogenesis
- Increase in the interstitial Leydig cells

It is associated with testicular atrophy and dysfunction, which result in sterility. Cryptorchid testes carry a 3–5-fold higher risk for testicular cancer (usually seminoma).

Solution to Question 17:

The surgical site infection rate expected in a patient with a clean contaminated wound that is operated upon after antibiotics prophylaxis is 3%.

SSI rates relating to wound contamination with and without using antibiotic prophylaxis:

Type of surgery	Infection rate without prophylaxis (%)	Infection rate with prophylaxis (%)
Clean(no viscus opened)	1-2	1-2
Clean-contaminated(viscus opened, minimal spillage)	6-9	3
Contaminated(open viscus with spillage or inflammatory disease)	13-20	6
Dirty(pus or perforation, or incision through an abscess)	40	7

Solution to Question 18:

qSOFA score (quickSOFA) is a bedside prompt that may identify patients with suspected infection who are at greater risk for a poor outcome outside the intensive care unit (ICU).

It uses three criteria, assigning one point each for

- Low blood pressure (SBP $\leq$ 100 mmHg),
- High respiratory rate ($\geq$ 22 breaths per min)
- Altered mentation (Glasgow coma scale, $\leq$ 15).

The score ranges from 0 to 3 points. The presence of 2 or more qSOFA points is a simple prompt to identify infected patients outside the ICU who are likely to develop sepsis.

Solution to Question 19:

A breast tumor of size $\geq$ 5 cm (not 4 cm) is classified as locally advanced breast carcinoma (LABC).

Invasive breast carcinoma is of 3 types:

- Early breast carcinoma (EBC) - TNM combination of T1 to T2, No to N1, and Mo
- Locally advanced breast carcinoma (LABC) - T4NoMo, T3N1Mo, TanyN2-3Mo
- Advanced breast carcinoma (ABC) - TanyNanyM1

Locally advanced breast cancer (LABC) includes:

- Breast tumors more than 5 cm in diameter
- Cancers that involve the skin or the underlying muscles of the chest (T4 of TNM staging)
- Cancers that involve multiple local lymph nodes (axillary or sub clavicular)
- Inflammatory breast cancer - a rapidly growing type of cancer that makes the breast appear red and swollen

Solution to Question 20:

Respiratory Rate ≥ 20 /minute or (Not and) PaCO₂ ≤ 32 mmHg (Not ≤ 35 mmHg) would be a feature of SIRS (Systemic inflammatory response syndrome).

Systemic inflammatory response syndrome (SIRS) is an exaggerated defense response of the body to a noxious stressor (infection, trauma, surgery, acute inflammation, ischemia, or reperfusion) to localize and then eliminate the endogenous or exogenous source of the insult.

SIRS is diagnosed when any two of the following are present:

- Hyperthermia (≥ 38 Degree Celsius) or Hypothermia (≤ 36 Degree Celsius)
- Tachypnoea (≥ 20 /min) or PaCO₂ ≤ 32 mmHg
- Tachycardia (≥ 90 /min)
- White cell count $\geq 12 \times 10^9$ /liter or $\leq 4 \times 10^9$ /liter or $\geq 10\%$ Band forms

Sepsis is SIRS with a documented infection.

Solution to Question 21:

The most common neuroendocrine tumor of the pancreas is insulinoma (4 per million).

Insulinoma is an insulin-producing tumor of the pancreas, with more than 90% of them being benign. It occurs in ≥ 40 years in patients without MEN 1 syndrome and ≤ 40 years in patients with MEN 1. It presents with symptoms of hypoglycemia such as weakness, headaches, sweating, faintness, seizures, and altered behavior. These occur after fasting or exertion and improve after glucose intake.

The investigation of choice for insulinoma is the determination of 72-hour fasting glucose and serum insulin levels. C-peptide and proinsulin levels are also elevated.

Medical management includes:

- Frequent carbohydrate-rich meals
- Somatostatin analogs like octreotide are effective in controlling symptoms due to excess insulin secretion. They should be used with caution as they can suppress counterregulatory hormones, such as growth hormone (GH), glucagon, and catecholamines, and precipitously worsen hypoglycemia.
- Diazoxide can be used in the initial management of insulinoma as it inhibits insulin release. Its side effects include sodium retention and nausea.
- Everolimus has been found to be highly effective in improving glycemic control in patients with insulinoma owing to its antitumor effect and induction of insulin resistance.

Enucleation and pancreatectomy are the definitive surgical management of insulinoma.

Other options:

Option A: Gastrinoma has an incidence of 0.5 to 4 in a million.


Options B & D: Somatostatinoma and VIPoma are much rarer than gastrinoma.

Solution to Question 22:

After parotid surgery, the nerve involved in the loss of sensation while shaving is the great auricular nerve.

The great auricular nerve (GAN) originates from the second and third nerve roots of the cervical plexus (C2-C3). It supplies the skin over the lower pinna, pre-auricular region, and angle of the mandible (image below). The main trunk passes over the sternocleidomastoid and into the anterosuperior portion of the parotid glands and branches into the anterior and posterior GAN. The anterior branch of GAN is often injured during parotid surgery during the initial elevation of the skin flap.

When GAN is injured, it can cause reduced sensations in the earlobe, pre-auricular skin, and the angle of the mandible that can interfere with habits such as shaving, wearing earrings, etc.



Other options:

Option A: The auriculotemporal nerve, a branch of the mandibular nerve (V3 of trigeminal) supplies the posterior part of the temple, tragus, and adjoining auricle. It does not supply the area where a person is likely to shave. However, the injury to the auriculotemporal nerve resulting in aberrant regeneration would result in Frey's Syndrome (gustatory sweating).

Option C: Though the facial nerve is the most common nerve affected during parotidectomy, any damage would result in facial paralysis and not loss of sensation of the face.

Option D: The mandibular nerve is a branch of the trigeminal nerve (cranial nerve V) that supplies the skin over the mandible, lower lip, and cheek. However, they are not injured during parotidectomy.

Solution to Question 23:

ABCDE algorithm is used to diagnose melanoma.

It comprises of:

- Asymmetry
- Border irregularity
- Color variegation
- Diameter of more than 6 mm
- Evolution or change

Melanoma is a malignant tumor arising from melanocytes at the epidermal-dermal junction. Though it represents only 4% of skin cancers, it accounts for 80% of deaths (high mortality rate). It presents as a pigmented lesion that has enlarged, ulcerated, or bled.

Histopathological confirmation is the gold standard in the diagnosis of melanoma. It is divided into four main types: nodular melanoma (NM), superficial spreading melanoma (SSM), lentigo maligna melanoma (LMM), and acral lentiginous melanoma (ALM).

The treatment consists of wide local excision of the tumor and follow-up to detect any local recurrence around the scar and excluding nodal spread.

Solution to Question 24:

The given instrument is a unipolar cautery/monopolar cautery.

Other options:

Option B: The bipolar cautery contains two tips, unlike unipolar cautery which has only one tip. The current flows between these two tips and hence it does not need a return electrode, unlike the unipolar cautery.



©MARROW

Figure 1.10

Option C: The harmonic scalpel (shown below) is a surgical instrument used to simultaneously cut and cauterize tissue. Ultrasonic energy is used in Harmonic scalpel where ultrasonic energy is converted to mechanical energy at the active blade.



Option D: LigaSure vessel sealing system (shown below) is a bipolar apparatus for sealing vascular tissue. It applies a high coaptive pressure during the generation of tissue temperature under 1000°C , hydrogen cross-links are first ruptured and then renatured, resulting in a vascular seal that has high tensile strength.



Solution to Question 25:

Digastric muscle is spared in radical neck dissection (RND). RND is used in the treatment of malignant lesions of the neck with local spread or metastasis.

In 1905, Crile described RND where Level I-V LN, sternocleidomastoid (SCM), internal jugular vein (IJV), and XI cranial nerve (spinal accessory nerve) are removed.

In 1962, Suarez described modified RND (MRND) where Level I-V LNs are removed and the following are preserved:

- MRND I - XI cranial nerve
- MRND II - XI cranial nerve and IJV
- MRND III - XI cranial nerve, IJV, and SCM

Note: RND with an additional LN/muscle/nerve is called extended RND.

Solution to Question 26:

Platelet transfusion is not done in the case of active variceal bleeding unless coagulopathy is confirmed as the cause. Platelet transfusions are indicated only in patients with a platelet count of $<50,000/\mu\text{L}$ or in patients with evidence of microvascular bleeding.

Variceal hemorrhage is a medical emergency. Varices usually present with the acute onset large-volume haematemesis, the lower esophagus being the most common site for variceal bleeding.

The sites for portosystemic anastomosis are:

- Left gastric vein to distal oesophageal veins: oesophageal and gastric variceal bleeding


- Superior rectal veins to Middle and inferior rectal veins: hemorrhoids and rectal bleeding
- Left branch of the portal vein to Periumbilical veins: caput-medusae
- The right branch of the portal vein to retroperitoneal veins draining into azygos, hemiazygos, lumbar, intercostal, and phrenic veins
- Omental and colonic veins near hepatic and splenic anastomoses to Retroperitoneal veins near hepatic and splenic anastomoses

Medical management of variceal bleeding includes blood transfusion (option B) and drug therapy with somatostatin analogs like octreotide (option C). Short-term antibiotics like ceftriaxone are known to decrease bacterial infection and increase survival.

Surgical measures are preferred if medical management fails. They include balloon tamponade, endoscopic banding (option D), transjugular intrahepatic portosystemic shunt (TIPS), portosystemic shunts, splenectomy, and gastric revascularization

Note: Sugiura procedure is the non-shunt surgical management of refractory oesophageal variceal bleeding. It is a procedure of de-vascularizing the distal esophagus and stomach.

Management of acute variceal bleeding



©Marrow

Solution to Question 27:

In a case of suspected testicular torsion, immediate exploration needs to be done on both the symptomatic as well as the asymptomatic side. After the de-torsion, the adequacy of the blood flow has to be assessed by Doppler.

In testicular torsion, the blood supply to the testicle is compromised due to the twisting of the spermatic cord within the tunica vaginalis, resulting in ischemia of the epididymis and the testis. The patient usually presents with acute onset pain and swelling of the scrotum. On examination, the cremasteric reflex is lost, and the scrotum looks asymmetrical.

Prehn's sign is used to differentiate epididymitis from testicular torsion. Prehn's sign is said to be positive when the pain gets relieved on lifting the testis, and it is seen in epididymo-orchitis.

At the time of surgical exploration, contralateral testes also must be explored and fixed to the dartos fascia in order to prevent possible torsion occurring in this testes.

More than 80% of the testis can be salvaged if surgery is performed within 6 hours. If the testis is found to be necrosed on scrotal exploration, orchidectomy of the necrosed testis and orchidopexy of the contralateral testis is done.

Solution to Question 28:


The given image showing a long, slender laparoscopic instrument is Veress Needle. It is used in direct puncture laparoscopy to produce a pneumoperitoneum.

Direct puncture laparoscopy (closed method):

- The abdominal wall is grasped with towel clips or fingers to elevate the abdominal wall away from the underlying structures.
- A small incision is made in the umbilicus and the Veress needle is introduced into the abdomen. Two distinct pops can be felt as the needle pierces the abdominal wall fascia and the peritoneum.

The umbilicus is the preferred location for introducing a Veress needle because the abdominal wall is thin at this location.

Carbon dioxide (CO₂) gas is usually used and the maximum pressure is in the range of 14-15 mmHg. But N₂O is used if the procedure is done under local anesthesia. Insufflation with N₂O is stopped at a pressure of 10 mmHg or when 2L of gas is insufflated.



The Hasson Trocar is used to gain access to the peritoneal cavity and generate pneumoperitoneum by the open direct peritoneal access technique (option C). This is known as the Hasson technique and is done only occasionally. It is preferred for patients who have undergone previous abdominal surgeries as the small bowel may be adherent to the undersurface of the abdominal wound in such cases.



Other options:

Option B: Salah's needle is a type of reusable bone marrow biopsy needle that has a guard situated 2 cm from the tip and a side screw.



Option D: Von Graefe needle (also known as Graefe's knife) is primarily used in ophthalmic surgery.

Solution to Question 29:

The given image shows negative pressure wound therapy using a vacuum-assisted closure (VAC) device to promote healing in chronic wounds.

Principle: An intermittent negative pressure of -125 mmHg hastens debridement and formation of granulation tissue in chronic wounds.

Indications: Chronic wounds (such as diabetic foot ulcers, stage III and IV pressure ulcers), traumatic wounds, flaps and grafts, and dehiscent incisions.

Contraindications:

- Exposed vital structures (organs, blood vessels, or vascular grafts)
- Preexisting infection
- Inadequate debridement of devitalized soft tissue or bone increases the risk of infection.
- The presence of malignancy (Marjolin's ulcer).
- An allergic reaction towards the adhesive used to maintain the vacuum seal.

Though not an absolute contraindication, there is no appreciable benefit for ischemic wounds.

Solution to Question 30:

The given image shows complete skin loss that involves muscles and shows exposed bone indicating a stage 4 pressure sore.

Pressure sores are skin or soft tissue injuries that occur due to prolonged pressure over specific areas of the body. An external pressure of >30 mmHg leads to the formation of pressure sores. The common sites of pressure sores in decreasing frequency are the ischium, greater trochanter, sacrum, heel, malleolus (lateral followed by medial), and occiput.

Pressure sores can be prevented with good skincare, the use of air-fluidized beds, special pressure-dispersion cushions, or foams. Repositioning the patient every two hours can lessen the pressure on the area.

Ulcers of stages 1 and 2 often do not require operative measures. They are managed with regular wound cleaning and dressing with a good antibiotic cover.

Stage 3 and 4 ulcers are adequately debrided, and vacuum-assisted closure (VAC) is used to promote the growth of granulation tissue required for surgical closure. Large skin flaps that have muscle with sensory innervation are best for reconstruction.

Solution to Question 31:

Grey coloured IV cannula is required for this patient in order to transfuse large amount of fluids.

The ATLS 10th edition guidelines state that 2 cannulae of at least 18G should be used to gain vascular access in trauma patients. In this scenario requiring large fluid transfusions, the grey (16G) cannula would be preferable to the green (18G) cannula due to the higher flow rate.


Solution to Question 32:

The video depicts Lahey's method of palpation of the left lobe of the thyroid gland.

In Lahey's method, for palpating the left lobe, the examiner initially pushes the right lobe medially with the left hand, and then using the right hand, the border and consistency of the left lobe is palpated.

Other options:

Option B: In Crile's method the patient is asked to swallow, while the examiner palpates the gland with the thumb. This gives a clear indication regarding the slight enlargement of the thyroid gland or small nodules.



Option C: Kocher's method is done to demonstrate the pressure effect of the thyroid gland on the trachea or larynx. Lateral compression will result in stridor.

Option D: Pemberton's sign (not method), is engorgement of neck veins when the patient is asked to raise their hands above the head and the arms should touch the ears. It is seen in patients with retrosternal goiter.

Solution to Question 33:

The given image is of a Jackson-Pratt drain.

Drains are placed at surgical sites to freely drain the air or fluids that may accumulate. They promote wound healing by preventing the formation of unnecessary cavities. Drains can be:


- Open : They are based on the principle of gravity
- Semi-open : Their action is based on capillary effect
- Closed : They carry out their actions through suction

Jackson-pratt drain is a closed, air-tight drainage system that operates by self-suction. The drain is sutured in place on the skin at the site of insertion to promote stability. There are two main parts to a JP drain:

- Hollow tube

- Collection bulb

The drain should be emptied as often as possible to allow the bulb to be compressed fully in order to maintain suction. This is usually done every 4-6 hours the first few days.



Other options:

Option A: Hemovac is also a closed type of suction drain which can typically handle larger amounts of fluid than the JP drain.

Option C: Minivac is a closed wound drainage system under negative pressure used in minor surgeries.

Option D: Corrugated drain is an open drainage system to remove excess fluid from the operated site.

Surgery AIIMS 2020 (May and Nov INI-CET)

Question 1:

A 65-year-old man presents with abdominal pain for the past 6 hours, BP- 89/50mmHg, RR- 24/min, saturation 92% with 6 L of oxygen. Radiograph shows extraluminal air in the abdomen. Which of the following must be done before shifting this patient into the OT?

- a) Intubate the patient
- b) Insert central venous catheter as initial IV access of choice
- c) 1 litre of crystalloid infusion
- d) Confirm the diagnosis with CECT

Question 2:

A patient with recurrent pain in the right hypochondrium, nausea, and vomiting is diagnosed with biliary dyskinesia. Which of the following is least useful in the diagnosis of this condition?

- a) ERCP
- b) HIDA scan
- c) USG
- d) Post surgery follow-up

Question 3:

A 32-year-old woman was declared brain dead. Her family has decided to donate her organs. Who is the ideal/best recipient?

- a) 14 year old child with multi organ dysfunction syndrome
- b) 50 year alcoholic man with liver cirrhosis with 1 month abstinence
- c) 70 year old woman with end stage renal disease
- d) 32 year old woman with renal failure

Question 4:

ERCP is least useful in the diagnosis of _____

- a) Recurrent pancreatitis
- b) Unexplained jaundice
- c) Periapillary mass
- d) Acute pancreatitis without choledocholithiasis

Question 5:

Which of the following is a correct statement regarding retroperitoneal trauma?

- a) All zone 1 injury to be treated conservatively
- b) Zone 3 injury whether blunt or penetrative should always be explored
- c) Zone 2 injury with expanding hematoma should be treated surgically
- d) Mattox maneuver is when the ascending colon is mobilized to the right to visualise the aorta

Question 6:

Which of the following nerve is least injured during thyroid surgery?

- a) Recurrent laryngeal nerve
- b) Ansa cervicalis
- c) External laryngeal nerve
- d) Marginal mandibular nerve

Question 7:

eFAST is least helpful in diagnosing which of the following conditions?

- a) Pericardial Effusion
- b) Pneumothorax
- c) Retroperitoneal hematoma
- d) Renal injury

Question 8:

A 23-year-old man presented with burn injury. He is covered with burns on his head & neck, both upper limbs, front and back of the chest. Calculate the percentage of burns.

- a) 27%
- b) 36%
- c) 45%
- d) 18%

Question 9:

FAST stands for _____

- a) First assessment with sonography in trauma
- b) Focussed assessment with sonography in trauma
- c) First abdominal sonography in trauma
- d) Focussed abdominal sonography in trauma

Question 10:

A patient with unilateral breast cancer is started on tamoxifen. This will increase the risk of?

- a) Ovarian cancer
- b) Endometrial cancer
- c) Breast cancer on other side
- d) Cervical cancer

Question 11:

Which of the following is the most appropriate mesh for use in hernioplasty?

- a) Heavyweight with small pores
- b) Heavyweight mesh with large pores
- c) Lightweight mesh with small pores
- d) Lightweight mesh with large pores

Question 12:

A 30-year-old teacher underwent parotidectomy following which she developed lower lip paralysis. This probably occurred because of an injury to which of the following structures?

- a) Temporal branch of facial nerve
- b) Cervical branch of facial nerve
- c) Buccal branch of the facial nerve
- d) Mandibular branch of facial nerve

Question 13:

Which of the following is not an adverse effect seen after a massive blood transfusion?

- a) Hypocalcemia
- b) Hypercalcemia
- c) Hypokalemia
- d) Hyperkalemia

Question 14:

Which of the following gene mutations are involved in carcinoma of the colon?

- a) APC, BRCA, k-ras
- b) APC, k-ras, p53
- c) MYh, k-ras, APC
- d) MYH, p53, DCC


Question 15:

Which among the following is correct regarding adequate thyroid FNAC sampling?

- a) 6 follicular clusters with 10 cells each
- b) 3 follicular clusters with 10-15 cells each
- c) 10 follicular clusters with 6-8 cells each
- d) 12 follicular clusters with 10-12 cells each

Question 16:


Which of the following are used to stop esophageal variceal bleed?



- a) A, B, C and D
- b) A and B only
- c) A, B and C
- d) A, B and D

Question 17:

A 25-year-old male patient was brought to the ER after RTA, his RR is 25/min, BP is 80/45mmHg. On examination, hyper-resonance is heard over the left side of the chest. The chest X-ray performed is given below. What would be the appropriate management in this case?



- a) Wide bore needle insertion in the 5th intercostal space (ICS)
- b) Wide bore needle insertion in the 5th ICS followed by an intercostal drain in the 5th ICS
- c) Aspiration of the pleural fluid
- d) No intervention needed

Question 18:

A patient presents with pain in the right upper quadrant on the second postoperative day following a cholecystectomy. Evaluation reveals a collection in the subhepatic space. A bile leak was suspected. Which of the following is the best investigation for this patient?

- a) MRCP
- b) ERCP
- c) HIDA
- d) CECT

Question 19:

What is the most likely diagnosis of the lesion shown in the image below?



- a) Arterial insufficiency
- b) Neuropathic ulcer
- c) Venous ulcer
- d) Tuberculous ulcer

Question 20:

A woman was diagnosed to have metastatic liver adenocarcinoma with CK 7 (-), CK20 (+), and CDX-1 (+) IHC markers. Which of the following is the most likely primary site of origin for this metastasis?

- a) Colon carcinoma
- b) Breast Carcinoma
- c) Pancreatic carcinoma
- d) Ovarian carcinoma

Question 21:

A 35-year-old chronic smoker presents with bluish discoloration of the toe along with intermittent claudication. Which of the following will be the next appropriate investigation in this patient?

- a) CT angiography
- b) MR angiography
- c) Digital subtraction angiography

d) Duplex scan

Question 22:


A 40-year-old woman presents with complaints of swelling of the right breast over the past 15 days. On examination, there was local rise of temperature, and the swelling was mobile over the chest wall. What is the likely diagnosis?



- a)** Inflammatory carcinoma
- b)** Giant fibroadenoma
- c)** Phyllodes tumor
- d)** Galactocele

Question 23:

A middle-aged man presents with progressive difficulty in swallowing for both solids and liquids for 8 months. Barium swallow done is shown below. What other investigations need to be done in this patient?



- a) Upper GI endoscopy & Manometry
- b) Upper GI endoscopy & PET scan
- c) Upper GI endoscopy & CECT
- d) Upper GI endoscopy and 24 hour PH monitoring

Question 24:

A 55-year-old man with IHD presents with sudden, severe, generalized abdominal pain and multiple episodes of vomiting. On examination, the pulse rate is 118bpm, rhythm is irregularly irregular, BP is 90/60 mmHg. The abdomen appears soft, non-distended, and bowel sounds are absent. Which of the following could be a likely diagnosis in this case?

- a) Acute severe pancreatitis
- b) Ruptured aortic aneurysm
- c) Acute mesenteric ischemia
- d) Peptic ulcer perforation

Question 25:


A 15-year-old girl presented with a slow-growing, rubbery, non-tender swelling on the side of the neck as shown in the image. Examination shows the presence of freckles and skin patches over the face and trunk. What is the most likely diagnosis?



- a) Cystic hygroma
- b) Chemodectoma
- c) Neural tumor
- d) Lipoma

Question 26:

What is the size of the cannula shown in the below image?



- a) 16 G
- b) 18 G

- c) 20 G
- d) 22 G

Question 27:

The position of the patient in the image shown below is known as:



- a) Lloyd Davis position
- b) Jack knife position
- c) Trendelenberg position
- d) Reverse trendelenberg position

Question 28:

What will be the most likely associated condition with increased TSH and low T4 levels?

- a) Hypoparathyroidism
- b) Pituitary Adenoma
- c) Grave's disease
- d) Hashimoto's thyroiditis

Question 29:

In e-FAST which of the following body part is included that is not included in FAST?

- a) Pelvic cavity
- b) Retroperitoneum
- c) Thoracic cavity
- d) Dural and subdural spaces

Question 30:

According to Strasberg's classification of bile duct injuries, injury to the right aberrant posterior sectoral duct with bile leak is:

- a) Type A
- b) Type B
- c) Type C
- d) Type D

Question 31:

A bus conductor was diagnosed with varicose veins. On examination, skin eczema and lipodermatosclerosis were noted. According to the CEAP classification system, He would be classified as

- a) C1
- b) C2
- c) C3
- d) C4

Question 32:

Which of the following is the most commonly seen tumor in the retroperitoneal space?

- a) Liposarcoma
- b) Rhabdomyosarcoma
- c) Synovial sarcoma
- d) Neurofibroma

Question 33:

A 50-year-old female patient presents with a 4*3cm irregular mass in the upper outer quadrant of the right breast, which is not fixed to the skin or the chest wall. A lymph node of 2cm is palpated in the right axilla. No mass is palpated in the opposite breast. Which of the following investigations would you perform on this patient?

- a) 1, 2, 3 & 4
- b) 1, 2 & 3
- c) 1, 3 & 4
- d) 1 & 3

Question 34:

According to the European Hernia Society classification, M3 incisional hernia refers to:

- a) Epigastric
- b) Infraumbilical
- c) Umbilical
- d) Suprapubic

Question 35:

Which of the following is true about the Tillaux sign?

- a) Mass is felt in the periumbilical region
- b) Mass is brilliantly transilluminant
- c) Mass is mobile perpendicular to mesentery attachment line
- d) Fluctuation is positive

Question 36:

Match the correct answer related to the nerve injuries in various surgeries:

- a) A-1, B-2, C-3, D-4
- b) A-2, B-3, C-4, D-1
- c) A-3, B-4, C-1, D-2
- d) A-4, B-1, C-2, D-3

Question 37:

A child is diagnosed with congenital hypertrophic pyloric stenosis, which among the following can be seen in this child?

- a) 2, 3 and 4
- b) 3 and 4 only
- c) 1, 2 and 4
- d) 1, 2, 3 and 4

Question 38:

The components of qSOFA scoring include:

- a) 1 & 2
- b) 3 & 4
- c) 1, 2 & 3
- d) 1, 2, 3 & 4


Question 39:

Which of the following is true regarding inguinal hernia?

- a) 1 and 3
- b) 1 and 4
- c) 2 and 3
- d) 2 and 4

Question 40:

Match the following:



- a) A-3, B-4, C-2, D-1
- b) A-3, B-1, C-2, D-4
- c) A-2, B-4, C-3, D-1
- d) A-2, B-1, C-3, D-4

Question 41:

A 40-year-old man presents to the ER with severe abdominal pain. On examination, his HR is 110bpm, BP- 88/60mmHg. On per abdomen examination, guarding is present. The erect abdominal X-ray is shown below. Which of the following would you proceed with, for the management of this case?



- a) 2 & 3
- b) 2 & 4
- c) 1, 2 & 3
- d) 1, 2 & 5

Answer Key

Question No.	Correct Option
1	c
2	c
3	d
4	d
5	c
6	d
7	c
8	c
9	b
10	b
11	d
12	d
13	b
14	b
15	a
16	a
17	b
18	b
19	b
20	a
21	d
22	a
23	a
24	c
25	c
26	c
27	d

28	d
29	c
30	c
31	d
32	a
33	c
34	c
35	c
36	a
37	b
38	c
39	c
40	b
41	a

Detailed Explanations

Solution to Question 1:

Volume resuscitation with 1 liter of crystalloid infusion should be done before shifting this patient to the OT for an emergency exploratory laparotomy.

The given clinical scenario in a patient with hypotension and tachypnea with extraluminal gas in the abdomen is suggestive of perforation peritonitis with hemodynamic instability. The immediate step is volume resuscitation to maintain circulating volume and organ perfusion.

Peritonitis is the inflammation of the peritoneum and can be categorized as local or diffuse and acute or chronic.

Causes:

- Gastrointestinal perforation e.g. perforated ulcer, appendix, diverticulum
- Transmural translocation (no perforation), e.g. pancreatitis, ischaemic bowel, primary bacterial peritonitis
- Exogenous contamination, e.g. drains, open surgery, trauma, peritoneal dialysis
- Female genital tract infection, e.g. pelvic inflammatory disease
- Haematogenous spread (rare), e.g. septicaemia
- Miscellaneous- e.g. familial Mediterranean fever

Clinical features:

- Abdominal pain, worse on movement, coughing and deep respiration
- Constitutional symptoms: anorexia, malaise, fever, lassitude

- nausea +/- vomiting
- Pyrexia (may be absent)
- Tachycardia
- Tenderness +/- guarding/rigidity/rebound of the abdominal wall
- Pain/tenderness on rectal/vaginal examination (pelvic peritonitis)
- Absent or reduced bowel sounds
- Septic shock (systemic inflammatory response syndrome [SIRS] and multiorgan dysfunction syndrome [MODS]) in later stages

Management:

General care of the patient:

- Correction of fluid and electrolyte imbalance
- Insertion of a nasogastric tube and urinary catheter
- Broad-spectrum antibiotic therapy
- Appropriate analgesia

Surgical treatment: Exploratory laparotomy and repair of perforation is the definitive management. Presence of peritoneal signs and hemodynamic instability are indications for urgent surgery.

Other options

Option A: Intubation is not indicated by the patient's current vital signs.

Option B: Initial volume resuscitation can be done with the peripheral large-bore cannula. A central venous catheter may be needed if peripheral access is difficult, vasopressors are administered, patient requires TPN.

Option D: A CT scan, while helpful for locating lesion and confirming the diagnosis, is not indicated for a hemodynamically unstable patient.

Note:

According to recent ATLS guidelines, crystalloid infusion should be limited to a maximum of 1.5 litres as crystalloid resuscitation beyond 1.5 liters increases the risk of death.

Solution to Question 2:

USG is the least useful investigation in the diagnosis of biliary dyskinesia or sphincter of Oddi dysfunction since the findings can be normal or show non-specific changes such as dilatation of the common bile duct.

Sphincter of Oddi dysfunction (SOD) is caused due to a structurally or physiologically abnormal sphincter with a higher tone, which fails to relax, causing pain and recurrent episodes of pancreatitis with normal liver function tests. There are two clinical types: biliary SOD and pancreatic SOD. Each type is further classified into types 1, 2 and 3.

Other options:

Option A: The best investigation for the diagnosis of SOD is ERCP with manometry of the biliary and pancreatic sphincters.

Option B: HIDA scan will show a reduced gallbladder ejection fraction (EF). A gallbladder EF <35% is considered diagnostic of biliary dyskinesia.

Option D: Biliary-type SOD should be considered and excluded in post-cholecystectomy syndrome; hence, post-surgery follow-up is essential.

CT and MRCP can demonstrate dilatation of the biliary and pancreatic ducts.

Solution to Question 3:

Among the given options, a 32-year-old woman with renal failure is the most suitable recipient for transplantation.

This is because the prognosis for the other patients mentioned above is poor. Prioritization of organ transplant recipients should be done based on post-transplant outcomes.

Option A - 14-year-old with multi-organ dysfunction syndrome is not suitable because there is a poor prognosis for the ongoing organ dysfunction.

Option B - Alcoholic patients with just 1-month of abstinence are not considered for a transplant.

Option C - A 70-year-old woman with end-stage renal disease is not the right candidate because the chances of long-term survival and a successful outcome are very less.

Solution to Question 4:

ERCP is least useful in the diagnosis of acute pancreatitis without choledocholithiasis.

ERCP stands for Endoscopic Retrograde Cholangio-Pancreatography. It involves the use of a side-viewing endoscope which is passed through the pylorus of the stomach into the second part of the duodenum to visualise the papilla. This is then cannulated, and dye is injected into the pancreatic or bile duct to visualise the pancreatic or biliary tree. It can be therapeutic or diagnostic. It is the best diagnostic modality for viewing the distal biliary tree, but MRCP is preferred as the diagnostic modality unless a therapeutic intervention is also needed.

Indications of ERCP:

- Choledocholithiasis
- Acute cholangitis
- Biliary stricture, biliary leak
- Pancreatic duct stricture, pancreatic stones
- Extrahepatic biliary strictures related to primary sclerosing cholangitis
- Endoscopic therapy for some patients with sphincter of Oddi dysfunction

- Biliary or pancreatic ductal system tissue sampling
- Balloon dilatation of the duodenal papilla or ductal strictures
- Sphincterotomy

The most common therapeutic indication for ERCP is the relief of biliary obstruction due to gallstone disease and benign or malignant biliary strictures.

Contraindications of ERCP include patients who cannot tolerate anesthesia, patients with type 3 sphincter of Oddi dysfunction, luminal gastrointestinal obstruction (examination limited to an area proximal to the level of obstruction and untreated hemostatic disorder (high risk of bleeding).

Complications of ERCP include duodenal perforation, hemorrhage, pancreatitis and sepsis.

Other options:

Option A: ERCP can help in ruling out the etiology in cases of recurrent pancreatitis.


Option B: ERCP can act as both diagnostic and therapeutic in cases of unexplained jaundice.

Option C: ERCP can be useful in case of a periampullary mass to take a tissue biopsy and cytology.

Solution to Question 5:

A zone 2 retroperitoneal injury with expanding hematoma should be treated surgically.

Zone of retroperitoneal injury	Type of viscera	Differential management
Zone 1	Central	Always explored, after proximo-distal vascular control
Zone 2	Lateral	Only explored if the hematoma is pulsatile or expanding, else conservative/ angioembolization
Zone 3	Pelvic	Only explored if the hematoma is pulsatile or expanding, else conservative/ angioembolization



Surgical exploration is done in cases of penetrating injuries to any zone.

Other options:

Option A: Zone 1 injuries are always explored.

Option B: Depending on the situation, zone 3 injuries can be managed with surgical exploration or conservatively.

Option D: Mattox manoeuvre is left medial visceral rotation to expose the aorta, while the Cattell-Braasch manoeuvre is done by right medial visceral rotation to expose IVC.

Solution to Question 6:

The marginal mandibular nerve is least likely to be injured during thyroid surgery as it is high up (mandibular margin) than the dissection plane of the approach to thyroid surgery.

Other nerves that might be at risk of injury include:

- Recurrent laryngeal nerve may be injured if dissection is done below the cricoid cartilage or near the tracheoesophageal groove (option A).
- Ansa cervicalis may be injured if strap muscles are not cut high enough. (option B)
- External laryngeal nerve may be injured during ligation of superior thyroid vessels as it lies very close to the superior thyroid artery (option C).

Complications of thyroidectomy:

- Hemorrhage - most common complication overall; due to slipping of ligature
- Nerve injury
- External laryngeal nerve - hoarseness (more common)
- Recurrent laryngeal nerve - hoarseness if unilateral, life-threatening aspiration and aphonia if bilateral

- Respiratory obstruction
- Laryngeal edema - the most common cause of respiratory obstruction
- Bilateral recurrent laryngeal nerve palsy
- Laryngomalacia
- Tension hematoma
- Hypoparathyroidism - infarction of parathyroids
- Thyroid storm - inadequately prepared for surgery
- Hypothyroidism
- Recurrent thyrotoxicosis
- Wound infection, stitch granuloma, keloid.

Solution to Question 7:

Among the given options, eFAST (focussed assessment with sonography in trauma) is the least useful in diagnosing retroperitoneal hematoma as the sensitivity of ultrasound for the detection of retroperitoneal hematomas is extremely poor.

Computed Tomography (CT Scan) is the investigation of choice for retroperitoneal injury, but it can only be done in a hemodynamically stable patient.

FAST stands for Focussed Assessment with Sonography for Trauma. It is a rapid, non-invasive bedside test for detecting free fluid using sonography used to assess patients for potential thoracoabdominal injuries.

Technique:


- The patient is placed in a supine position and the following regions are scanned:
- Subxiphoid transverse view: assess for pericardial effusion and left lobe liver injuries
- Longitudinal view of the right upper quadrant: assess for right liver injuries, right kidney injury, and Morison pouch
- Longitudinal view of the left upper quadrant: assess for splenic injury and left kidney injury
- Transverse and longitudinal views of the suprapubic region: assess the bladder and pouch of Douglas

eFAST stands for extended FAST and incorporates left and right thoracic views to assess for pneumothorax and haemothorax.

Limitations of eFAST are:

- Operator-dependent
- Bowel gas and subcutaneous air distort images.
- Can miss diaphragm, bowel, and pancreatic injuries
- Unreliable for assessment of the retroperitoneum

- Cannot reliably exclude injury in penetrating trauma
- May need repeating or supplementing with other investigations



Solution to Question 8:

The percentage of burns calculated in the given patient is 45% according to Wallace's rule of nine.

Calculation:

- Burns on the head&neck would account for 9% of burns.
- Burns on both upper limbs (9% for each limb) would add to 18%
- Burns on both sides of the chest would account for 18%

Therefore the total percentage of burns in the given patient is $9+18+18 = 45\%$

Solution to Question 9:

FAST stands for Focussed Assessment with Sonography in Trauma.

It is a rapid, non-invasive bedside test for detecting free fluid using sonography to assess patients for potential thoracoabdominal injuries. It can be performed within 2-4 minutes.

Technique:


- The patient is placed in a supine position and the following regions are scanned:
- Subxiphoid transverse view: assess for pericardial effusion and left lobe liver injuries
- Longitudinal view of the right upper quadrant: assess for right liver injuries, right kidney injury, and Morison pouch
- Longitudinal view of the left upper quadrant: assess for splenic injury and left kidney injury

- Transverse and longitudinal views of the suprapubic region: assess the bladder and pouch of Douglas

eFAST stands for extended FAST and incorporates left and right thoracic views to assess for pneumothorax and haemothorax.

Limitations of eFAST are:

- Operator-dependent
- Bowel gas and subcutaneous air distort images.
- Can miss diaphragm, bowel, and pancreatic injuries
- Unreliable for assessment of the retroperitoneum
- Cannot reliably exclude injury in penetrating trauma
- May need repeating or supplementing with other investigations



Solution to Question 10:

Tamoxifen increases the risk of endometrial cancer by 1-2%.

The benefit of tamoxifen in breast cancer outweighs the risk of endometrial cancer.

Tamoxifen is a selective estrogen receptor modulator, approved for the treatment of estrogen receptor-positive (ER) breast cancer. It is used only in premenopausal women for 5 years in low-risk patients and for 10 years in patients with a high risk of relapse (node-positive, tumor ≥ 5 cm, lymphovascular invasion). It is also known to reduce the incidence of second primary breast cancer in the contralateral breast.

Tamoxifen increases the risk of endometrial cancer in postmenopausal women; hence aromatase inhibitors are used in postmenopausal women. Tamoxifen is also not recommended in patients who have had bilateral mastectomies with ER-positive ductal carcinoma in situ. Other side effects of tamoxifen include bone pain, hot flashes, nausea, vomiting, fluid retention and rarely

thrombosis and cataract.

Solution to Question 11:

The most appropriate mesh used in hernioplasty is a lightweight mesh with large pores.

Lightweight meshes generally contain less material and induce a less-pronounced foreign body reaction and decreased inflammatory response, which results in better tissue incorporation, increased prosthesis compliance, and decreased patient discomfort and pain.

Meshes with large pores exhibit less inflammatory infiltrate, connective tissue, and scar bridging, which allows increased soft tissue ingrowth. They have greater elasticity and flexibility and also shrink less resulting in lesser adhesions, better tissue incorporation, and decreased patient discomfort and pain. So the best mesh type among the given options is lightweight mesh with large pores.

Meshes used in hernioplasty can be synthetic or biological depending on their composition.

Synthetic meshes are made from polypropylene, polyester, or PTFE (polytetrafluoroethylene).

PTFE meshes are hydrophobic with a lower risk of adhesion.

Biological meshes are acellular matrix materials derived from humans or animals and have lesser complications. But due to poor collagen deposition and absorption, they rapidly lose mechanical strength and hence their use is restricted to infected environments.

Note: To promote better tissue integration, a composite mesh was developed which combines the attributes of polypropylene and PTFE by layering the two substances on top of one another.

POLYPROP YLENE	POLYESTE R	PTFE
monofilam ent	multifilame nt	solid
hydrophobi c	hydrophilic	hydropho bic
rigid	elastic	rigid

Solution to Question 12:

Lower lip paralysis occurs when there is an injury to the marginal mandibular branch of the facial nerve.

Branches of the facial nerve are:


- Temporal
- Zygomatic
- Buccal

- Marginal mandibular
- Cervical

The facial nerve passes through the parotid gland, hence it might be injured during parotidectomy.

In contrast to this, the upper lip is supplied by buccal branches of the facial nerve.

Complications of parotidectomy include hematoma formation, infection, deformity (unsightly scarring and retromandibular hollowing), temporary/permanent facial weakness, sialocele, facial numbness, permanent numbness of the earlobe due to great auricular nerve transection and Frey's syndrome.



Solution to Question 13:

Hypercalcemia is not an adverse effect of massive blood transfusion.

Massive blood transfusion instead leads to hypocalcemia as the anticoagulant citrate binds to calcium. The lethal triad of massive blood transfusion is acidosis, hypothermia, and coagulopathy.

Solution to Question 14:

APC, k-ras, and p53 gene mutations are implicated in carcinoma of the colon.

In the adenoma to carcinoma sequence, the loss of one normal copy of the tumor suppressor gene APC is the first hit, followed by the loss of the intact second copy of the APC gene (second hit). Other changes in the sequence include the mutation of k-ras and the inactivation of the tumor suppressor gene p53, leading to the emergence of carcinoma. The accumulation of these mutations is more important than their occurrence in a specific order. This is known as the Fearon-Vogelstein adenoma-carcinoma multistep model of colorectal neoplasia.

A number of other genes on chromosome 18q are implicated in colorectal cancer, including SMAD2, SMAD4, and DCC.

Solution to Question 15:

6 follicular clusters each with 10 cells are considered adequate thyroid FNAC sampling.

There are different criteria for the adequacy of thyroid FNAC:

- Six groups with at least 10 cells in each group, or
- Ten groups with at least 20 cells in each group, are considered adequate.

Cystic goitre may not yield thyroid follicular cells, hence, the aspiration should be repeated under USG guidance.

FNAC is the investigation of choice in discrete thyroid swellings; it has excellent patient compliance, simple and quick to perform in the outpatient department and is readily repeated.

Bethesda criteria for reporting thyroid cytopathology:


The 2023 Bethesda System for Reporting Thyroid Cytopathology

Bethesda class	Diagnostic category	Management
I	Nondiagnostic	Repeat FNA with ultrasound guidance
II	Benign	Clinical and ultrasound follow up
III	Atypia of undetermined significance Specify if AUS-nuclear atypia or AUS-other	Repeat FNA, molecular testing, diagnostic lobectomy, or surveillance
IV	Follicular neoplasm Follicular neoplasm-oncocytic follicular neoplasm*	Molecular testing ⁺ , hemithyroidectomy or lobectomy
V	Suspicious for malignancy	Molecular testing ⁺ , lobectomy or near total thyroidectomy
VI	Malignant	Lobectomy or near total thyroidectomy

* Previously called Hurthle cell neoplasm
⁺ Molecular testing can be performed to assess the type of surgery to be performed (lobectomy or total thyroidectomy)

©MARROW

Approach to thyroid nodules



Solution to Question 16:

All the given options can be used to stop esophageal variceal bleeding.

Image A shows the Sengstaken-Blakemore tube. It has 3 lumens - one aspiration port and 2 inflation balloons. They are:

- Gastric aspiration port
- Gastric balloon port
- Esophageal balloon port

The gastric balloon is inflated and tension is applied to the gastroesophageal junction. If this does not control the haemorrhage, the esophageal balloon is inflated as well, compressing the venous plexus between them. Only gastric contents can be aspirated.

Scheme of using the Sengstaken-Blakemore tube





Image B shows the Minnesota tube. It has 4 openings - 2 inflation tubes and 2 suction tubes i.e. both gastric and oesophageal aspiration ports are present, which allows the aspiration of both gastric and esophageal contents.

Images C and D both show an upper GI endoscope. They are used to identify the source of bleeding and once identified, sclerotherapy and endoscopic variceal banding are performed, which have been shown to control hemorrhage effectively.

Banding has a lower rate of complications. If bleeding does not stop by banding/sclerotherapy, balloon tamponade should be tried.

The protocol for the management of esophageal varices is given below.




Solution to Question 17:

The given clinical scenario of hemodynamic instability, hyper-resonance on the left side of the chest, and the X-ray findings are suggestive of tension pneumothorax. It is managed by inserting a wide-bore needle in the 5th intercostal space (ICS) followed by placing an intercostal drain in the 5th ICS.

Traumatic pneumothorax can be:

- Simple pneumothorax: Without hemodynamic instability
- Tension pneumothorax: With hemodynamic instability

The X-ray given below is showing a left-sided tension pneumothorax. A large, well-demarcated area is devoid of lung markings on the left side. The yellow arrows are pointing to the tracheal deviation and the mediastinal shift to the right side.



Management:

- Simple pneumothorax: Tube thoracostomy or chest tube insertion in the 5th ICS in the triangle of safety
- Tension pneumothorax:
- Immediate management: Urgent needle decompression or needle thoracostomy i.e. insertion of a wide-bore needle in the 5th ICS, anterior to the midaxillary line
- Definitive management: Tube thoracostomy or chest tube insertion in the 5th ICS in the triangle of safety

Note: According to ATLS 10th edition

- In adults, needle thoracostomy or needle decompression location is changed to 5th ICS anterior to the midaxillary line from 2nd ICS midclavicular line.
- In the pediatric age group, the needle thoracostomy location remains at the 2nd ICS midclavicular line.

Solution to Question 18:

Endoscopic retrograde cholangiopancreatography (ERCP) is the best investigation in a patient suspected of post-cholecystectomy biliary leaks.

Patients with post-cholecystectomy biliary leak usually present with fever, jaundice and abdominal pain. USG is used for an initial evaluation in such cases and can show a collection in the sub-hepatic space.


The most common bile leak following cholecystectomy is from the cystic duct. This can be treated by placing a biliary endoprosthesis (stent) in the common bile duct across the origin of the cystic duct.

MRCP shows fluid collections arising from bile leaks but cannot differentiate these from other causes of fluid collections.

ERCP is the best investigation as it has the advantage of being both diagnostic and therapeutic. ERCP is useful post-cholecystectomy as there is a high likelihood of a retained stone and stone extraction and/or temporary stenting may be required.

HIDA scan is the most sensitive investigation for biliary leakage.

CT scan can show focal fluid collections, ascites, biliary obstruction, hepatic atrophy, signs of secondary biliary cirrhosis and any vascular injury.



Solution to Question 19:

The given image which shows a punched-out ulcer with regular margins on the plantar surface of the dry, cracked foot is suggestive of a neuropathic ulcer.

Neuropathic ulcers occur as a result of repeated trauma to the insensitive part due to peripheral neuropathy. There is a lack of sensation over the pressure points, leading to unnoticed microtrauma, wounds or cuts which develop into an ulcer. Causes of neuropathic ulcers include diabetic neuropathy, B12 deficiency, alcoholism, hypothyroidism, leprosy, peripheral nerve injury, transverse myelitis and myelomeningocele.

Other options:

Option D: Tuberculous ulcers occur when a cold abscess from bone and joint tuberculosis breaks out on the surface. The most characteristic feature of this ulcer is its edge which is thin and undermined. The following is a picture of a tuberculous ulcer:

Type of Ulcer	Arterial(option A)	Venous(option C)	Neuropathic	Pressure
Underlying condition	Peripheral vascular disease, hypertension	Varicose veins, DVT, obesity, previous thrombophlebitis	Diabetes, peripheral neuropathy	Limited mobility

Type of Ulcer	Arterial(option A)	Venous(option C)	Neuropathic	Pressure
Location	Lateral malleolus, toes, and dorsum of the foot	Lower leg, above the medial malleolus	The plantar surface of the foot, toes	Bony prominences (e.g.heel)
Characteristics	Deep, punched-out, irregular margins. The wound bed has unhealthy necrotic tissue	Shallow, flat with exudate	Punched-out with regular calloused margins.	Deep, macerated
Condition of leg/foot	Cool thin shiny skin, pallor on leg elevation, absent peripheral pulses, and delayed capillary refill	Excematous and pigmented. Limb edema may be present	Dry, cracked, insensate	Atrophic skin, loss of muscle



The following is a picture of an arterial ulcer.



The following is a picture of a venous ulcer.



The following is a picture of a pressure ulcer.



Solution to Question 20:

Colorectal carcinoma is the most likely primary tumor for the given metastases. Among the given options, cytokeratin 7 is negative only in colon carcinoma.

The most common site of distant metastasis of colorectal cancer is the liver. These metastases arise from hematogenous spread via the portal venous system. The lung is the next most common site for metastasis.

Cytokeratin profiles of various primary tumors:

CK 7 positiveCK20 positive	CholangiocarcinomaOvarian mucinous carcinoma
CK 7positiveCK 20 negative	Breast carcinomaOvarian serous adenocarcinomaThyroid carcinomaPancreatic adenocarcinomaLung carcinoma
CK 7negativeCK 20positive	Colorectal adenocarcinomaMerkel cell carcinoma
CK 7 negativeCK 20 negative	Hepatocellular carcinomaRenal cell carcinomaProstatic adenocarcinomaSquamous cell carcinoma

Solution to Question 21:

Chronic smoker presenting with bluish discoloration of the toe along with intermittent claudication is suggestive of peripheral arterial disease and duplex scanning will be the next appropriate investigation.


Duplex Scanning is a major non-invasive technique that uses B-mode ultrasound to provide an image of the vessel. Most scanners now have color coding, which allows detailed visualization of blood flow in a vessel. Different colors indicate changes in the direction and velocity of flow, with areas of high flow usually indicating stenotic lesions.

Other options:

Option A: CT Angiography (CTA) is more useful in the evaluation of visceral arterial disease (eg. mesenteric ischemia). It can be beneficial in peripheral artery diseases where a duplex scan shows poor images (aortoiliac segment involvement). CTA is not routinely recommended in peripheral artery disease because of concomitant renal dysfunction in these patients which can lead to contrast-induced nephropathy.

Option B: MR Angiography is time-consuming and expensive, hence not done as the next investigation. It is contraindicated in patients with claustrophobia and metallic implants.

Option C: Digital subtraction angiography (DSA) is an invasive and expensive procedure compared to other investigations. The benefits of DSA are that it provides dynamic arterial flow information and can be combined with a definitive endovascular intervention when indicated. DSA is indicated only when there is a discrepancy in diagnosis or intervention is planned. Complications associated with DSA include bleeding, hematoma, false aneurysm formation, thrombosis, arterial dissection, distal embolisation, renal dysfunction, and allergic reaction.



Solution to Question 22:

In the given scenario, the rapid onset of the disease and the examination findings are suggestive of inflammatory carcinoma of the breast.

Inflammatory breast cancer is the most aggressive subtype of breast cancer. It mimics acute mastitis. The hallmark of inflammatory breast carcinoma is the diffuse tumor involvement of the dermal lymphatics within the breast and the overlying skin, often without a discrete underlying tumor mass (peau d' orange appearance). The patient presents with rapidly progressive swelling of the breast, warmth, erythema and edema. A breast lump may or may not be present.

A diagnosis of inflammatory breast carcinoma should be suspected when a patient with rapidly progressive swelling with inflammatory skin changes does not respond to antibiotics. The imaging features with ultrasound and a mammograph is non-specific due to the underlying erythema and edema. A biopsy should be done to confirm the diagnosis.

Treatment includes combined-modality treatment:

- Neoadjuvant chemotherapy
- Mastectomy
- Radiation therapy
- Hormonal therapy- for ER-positive tumors
- Trastuzumab for HER-2-positive tumors

Option B: Giant fibroadenomas occasionally occur during puberty. These are benign solid encapsulated slow-growing tumors over 5 cm in diameter. It presents as a painless lump in the breast. They are managed by excision via a submammary incision.

Option C: Phyllodes tumors usually occur in women over the age of 40 years but can appear in younger women. Phyllodes tumor is mostly benign even though it is also called cystosarcoma phyllodes. It presents as a large tumor with an unevenly bosselated surface. They are usually

mobile over the chest wall despite their large size. This tumor does not infiltrate the skin. Nipple retraction is not seen in this condition. Benign lesions can be treated with enucleation or wide local excision. Massive tumours, recurrent tumours and malignant lesions are treated with a mastectomy.

Option D: Galactoceles are the most common benign breast lesions in lactating women. It is a cystic enlargement of the mammary gland containing milk. They are mostly detected during lactation or in the third trimester of pregnancy but rarely, may also present after stopping breastfeeding. Clinically, the mass is usually firm and non-tender and presents as a solid tumour-like lesion on physical examination. The patient usually notes the lesion during lactation or some period after lactation. Needle aspiration, followed by excision should it recur, is the standard method of treating galactocoele.

Solution to Question 23:

The given clinical scenario and the barium swallow showing a bird beak appearance are suggestive of achalasia cardia.

Achalasia is a primary esophageal motility disorder characterized by the absence of esophageal peristalsis and impaired relaxation of the lower esophageal sphincter (LES) in response to swallowing.

Manometry is considered the most sensitive investigation to diagnose achalasia as it identifies early disease before esophageal dilatation and food retention. The diagnostic criteria for achalasia with manometry are impaired lower esophageal sphincter (LES) relaxation and absent peristalsis with an elevated LES pressure.

Endoscopy is useful for ruling out pseudo-achalasia.

Note: 24-hour pH monitoring is ideal and most accurate in patients with the features of gastroesophageal reflux disease.

Solution to Question 24:

The given clinical scenario of a patient with sudden onset severe abdominal pain which is out-of-proportion to the physical examination findings along with atrial fibrillation is suggestive of acute mesenteric ischemia.

Mesenteric ischemia may be acute or chronic. The superior mesenteric artery (SMA) is most likely affected. An embolus is the most common cause of SMA occlusion. Possible sources for SMA embolisation include:

- Left atrium in atrial fibrillation
- Previous history of ischemic heart disease/myocardial infarction
- Vegetations on mitral and aortic valves associated with endocarditis
- An atheromatous plaque from an aortic aneurysm.

Other causes of mesenteric ischemia include:

- Atherosclerosis of mesenteric vessels
- Vasculitides
- Venous thrombosis in factor V Leiden mutation
- Non-occlusive mesenteric ischemia due to alterations in splanchnic blood flow.

The most important clue to an early diagnosis of acute mesenteric ischemia is the sudden onset of severe abdominal pain in a patient with atrial fibrillation or atherosclerosis. The pain is typically in the central abdomen and out of proportion to the physical findings.

Associated clinical features include nausea, vomiting, diarrhea, abdominal distension, peritonitis, and the passage of bloody stools.

The gold standard for the diagnosis of acute mesenteric ischemia is angiography, and management is laparotomy. The goal of operative exploration is to resect the compromised bowel and restore blood supply via embolectomy.

Other options:

Option B: A pulsatile abdominal mass is seen in the case of an abdominal aortic aneurysm. Acute pain and hypotension occur with rupture of the aneurysm, which requires an emergency operation.

Options A and D: Acute severe pancreatitis and peptic ulcer perforation have no association with atrial fibrillation. They usually present with pain over the upper abdomen associated with guarding and rigidity on examination.

Solution to Question 25:

A lateral neck mass in a pediatric patient, which is non-tender and present in association with freckles, and hyperpigmented skin patches (cafe-au-lait spots) is suggestive of neural tumor i.e. neurofibroma.

Neurofibroma is a benign tumor arising from the connective tissue of the nerve containing ectodermal, neural and mesodermal connective tissue components. It can be single or multiple. It is the most common intradural extramedullary spinal tumor. Neurofibromas can be local, generalized (von Recklinghausen's disease) or plexiform types. Patients present with mild pain or a painless swelling usually in the subcutaneous or cutaneous plane with tingling, numbness and paraesthesia along the distribution of the nerve. Complications of neurofibroma include cystic degeneration, erosion into deeper planes, cord compression and paralysis (spinal neurofibroma) and sarcomatous changes.

Treatment is excision under general or regional anesthesia. Indications for surgery in neurofibroma include symptomatic patients (pain and pressure symptoms), cosmetic problems and malignant transformation (5%).

Option A: Cystic hygroma is a type of congenital lymphangioma. This swelling is soft and partially compressible, which shows positive fluctuation and translucency test.

Option B: Chemodectoma or carotid body tumor is a rare tumor, most commonly present in the fifth decade and there is an association with pheochromocytoma. These patients usually present with a slowly enlarging, painless lump at the carotid bifurcation.

Option D: A lipoma is a soft painless swelling. It shows the characteristic slip sign and is freely mobile.

Solution to Question 26:

The given cannula is 20 gauge IV cannula as its color code is pink.

Solution to Question 27:


The above image shows a reverse Trendelenburg position, where the patient's head is elevated 15 - 30 degrees higher than the feet. It is named so, as the position is reverse of the Trendelenburg position, where the head end is lower and the feet are at a higher level.

The Trendelenburg and reverse Trendelenburg positions are used in surgeries that are required to move the abdominal viscera, such as the small bowel, away from the operative field. Restraints and sideboards are placed to prevent the patient from sliding from the bed.


In laparoscopic cholecystectomy, the patient is placed in a reverse Trendelenburg position slightly rotated to the left. This exposes the fundus of the gallbladder which is retracted towards the diaphragm.

Other options:


Option A: Lloyd Davies's position is a 30-degree Trendelenburg position with the hips flexed at 15 degrees. It is used for rectal and anal canal surgeries like low anterior resection and abdominoperineal resection. It can cause compartment syndrome of the calves.



Option B: Jackknife position is a modified prone position where the patient's hips are supported on a pillow and the table is flexed at a 90-degree angle with the head and feet at a lower level. It is not used anymore due to the chest compression causing impaired ventilation leading to positional asphyxia. It was used for hemorrhoids and fissure surgery.



The image below shows commonly used patient positions in the operating room:



Solution to Question 28:

Increased TSH and low T4 levels are seen in primary hypothyroidism. Hashimoto's thyroiditis is the only cause of primary hypothyroidism in the given options.

Hashimoto's thyroiditis is an autoimmune disease that results in the destruction of the thyroid gland and gradual & progressive thyroid failure. It is the most common cause of primary hypothyroidism in iodine-sufficient areas. They also have autoantibodies against thyroid receptors, thyroid peroxidase and thyroglobulin.

Other options:

Option A: Hypoparathyroidism is generally not associated with hypothyroidism, except in certain congenital multisystem disorders such as DiGeorge's syndrome.

Option C: Grave's disease is the most common cause of hyperthyroidism. It is an autoimmune disorder caused due to autoantibodies against the TSH receptor. There will be elevated T₃ and T₄ levels with low TSH levels.

Option B: A TSH-producing pituitary adenoma can have increased TSH levels, but there will be elevated T₃, T₄ levels as well.

Solution to Question 29:

e-FAST is used to additionally assess the presence of free fluid in the thoracic cavity.

FAST stands for focused assessment with sonography for trauma. It is a rapid diagnostic examination to assess patients with potential thoracoabdominal injuries. It can be performed within 2-4 minutes.

The sequential order in which FAST is performed: pericardial sac, the right upper quadrant (RUQ), left upper quadrant (LUQ), and pelvis. 4 'Ps' are evaluated in sequence: Pericardial sac → Perihepatic region → Perisplenic region → Pelvis.

Technique:

- Following regions are scanned:
- Subxiphoid: assess for pericardial effusion and left lobe liver injuries
- Right upper quadrant: assess for right liver injuries, right kidney injury, and Morison pouch
- Left upper quadrant: assess for splenic injury and left kidney injury
- Suprapubic region: assess the bladder and pouch of Douglas

An 'extended FAST or eFAST' scan may be performed: left and right thoracic views to assess for pneumothorax and hemothorax.

Solution to Question 30:

Injury to the right aberrant posterior sectoral duct associated with bile leak is classified as Strasberg's type C injury.


Biliary injury may occur post-laparoscopic cholecystectomy. The most common site of bile leak following cholecystectomy is the cystic duct.

MRCP is the gold standard for complete morphological evaluation of the biliary tree. MRCP with magnetic resonance angiography is more useful in cases of associated vascular injury. A HIDA scan can confirm the presence of a bile leak or biliary obstruction.


The Strasberg classification of bile duct injury is widely used to anatomically define bile duct injuries.

Classification

- Type A: Bile leak from cystic duct stump or a minor biliary radical in the gallbladder fossa
- Type B: Occluded right posterior sectoral duct.
- Type C: Bile leak from the divided right posterior sectoral duct.
- Type D: Bile leak from the main bile duct without major tissue loss.
- Type E: Injury to the main hepatic duct; classified according to the level of injury
 - E1: Transected main bile duct with a stricture more than 2cm from the hilus.
 - E2: Transected main bile duct with a stricture less than 2cm from the hilus
 - E3: Stricture of the hilus with right and left ducts in communication.
 - E4: Stricture of the hilus with separation of right and left hepatic ducts.
 - E5: Stricture of the right aberrant sectoral and main bile ducts.



Management of bile duct injuries is shown below:



Solution to Question 31:

According to CEAP classification, varicose veins with skin eczema and lipodermatosclerosis come under the clinical class of C4.

The clinical class of C4 under CEAP includes changes in skin and subcutaneous tissue secondary to chronic venous disease. It is further classified as,

- C4a- pigmentation or eczema
- C4b- lipodermatosclerosis or atrophie blanche
- C4c- corona phlebectatica

Recent 2020 update on the CEAP system:

- C2r for recurrent varicose veins
- C4c for Corona phlebectatica
- C6r for recurrent active venous ulcer.

Solution to Question 32:

Liposarcoma is the most commonly seen tumor in retroperitoneal space.

Retroperitoneal tumors can be benign or malignant. Malignant retroperitoneal tumors may be mesodermal (75%), neuroectodermal (24%), or embryonic remnant (1%) in origin. Rarely it can be urologic or metastatic also. The order of frequency of primary retroperitoneal malignancy is as follows:

- Liposarcoma
- Leiomyosarcoma
- Rhabdomyosarcoma
- Fibrosarcoma
- Malignant nerve sheath tumors

Liposarcoma is composed of three histologic varieties (listed in order of decreasing frequency):

- Well-differentiated and undifferentiated liposarcoma
- Pleomorphic liposarcoma
- Myxoid/round cell liposarcoma

Among the three subtypes, pleomorphic liposarcoma is the most malignant and often metastasizes, while the well-differentiated is the least malignant. Myxoid liposarcoma has intermediate malignant behavior.

Diagnosis is confirmed by contrast-enhanced CT with CT of the chest to look for metastases.

Surgery is the main modality of treatment. Wide excision with complete en bloc dissection is done. Post-operative radiotherapy and chemotherapy are used as adjuvant methods. Radiotherapy is given at a dose of 5000 cGY, it can be external beam or intra-operative radiotherapy. Chemotherapeutic agents used are mesna, adriamycin, ifosfamide and dacarbazine (MAID regimen).

Solution to Question 33:

The given clinical scenario is suggestive of breast malignancy with locoregional spread with unlikely distant metastasis. For further evaluation of this lesion, bilateral mammography, a biopsy of the breast mass and a USG-guided biopsy of the suspicious lymph node need to be performed in order to determine staging and decide on further management.

Statement 2: PET Scan is usually performed in advanced cases of breast cancer. It is not routinely used in the assessment of locoregional disease.

Statement 1: Diagnostic mammography is used to evaluate women with abnormal findings such as a breast mass or nipple discharge. Mammography is more accurate than clinical examination for the detection of early breast cancers. Mammographic features suggestive of breast cancer include a solid mass with or without stellate features, microcalcifications, irregular soft tissue shadow, spiculations and duct distortion.

Statement 3: Pathological diagnosis of carcinoma is made by a core needle biopsy (tru-cut biopsy) or fine-needle aspiration before any type of treatment is started.

Statement 4: The presence or absence of axillary lymph node involvement in breast cancer is one of the most important prognostic factors for determining staging and treatment.

Ultrasound-guided lymph node biopsy should be done in patients with suspicious or clinically palpable lymph nodes.

Ultrasound features suggestive of breast cancer include a hypoechoic, irregular mass with indistinct, angulated or spiculated margins, acoustic shadowing, ductal extension and a branched pattern of mass.

Solution to Question 34:

According to the European Hernia Society classification, M3 incisional hernia refers to umbilical hernia.

According to the European Hernia Society classification of incisional hernias:

Midline incisional hernias are classified as-

Lateral incisional hernias are classified as-

M1	Subxiphoida l
M2	Epigastric

M1	Subxiphoidal
M3	Umbilical
M4	Infraumbilical
M5	Suprapubic

L1	Subcostal
L2	Flank
L3	Iliac
L4	Lumbar

Solution to Question 35:

Tillaux's sign is said to be positive if the mass/swelling is mobile perpendicular to the mesentery attachment line. Tillaux's sign is seen in a mesenteric cyst.

Mesenteric cysts occur either in the small intestine (60%) or the colon (40%) and can be classified as the following:

- Chylolymphatic (most common type)
- Enterogenous
- Urogenital remnant
- Dermoid
- Other causes: traumatic hematoma and cyst formation, tuberculous cold abscess of mesentery and hydatid cyst of the mesentery

They most commonly occur in adults with a mean age of 45 years. It is more common in females than males (2:1). The most common presentation is a painless abdominal swelling near the umbilicus, which is smooth and not moving with respiration with Tillaux's triad:

- Soft, smooth swelling in the umbilical region
- Freely mobile in a direction perpendicular to the mesentery (Tillaux's sign)
- Zone of resonance around the cyst

Other presentations include recurrent attacks of abdominal pain due to temporary impaction of a food bolus in a segment of the bowel narrowed by the cyst and vomiting.


Complications of mesenteric cyst include torsion, rupture, infection and hemorrhage into the cyst.

Ultrasonography and CT are used to demonstrate mesenteric cysts.

Treatment:

- Chylolymphatic cyst is managed by enucleation.
- Enterogenous cyst is managed by removal of the cyst with resection of the adjacent bowel as it arises as a diverticulum or duplication from the adjacent bowel.

In cases where complete excision is not possible due to the close proximity to vital structures, partial excision or marsupialization should be performed.



Solution to Question 36:

A-1, B-2, C-3, and D-4 is the correct option.

A: During breast surgery, removal of the axillary lymph nodes is commonly done. This carries a risk of injury to the long thoracic nerve, which originates from the C5-C7 roots of the brachial plexus. It innervates the serratus anterior muscle. Injury to this nerve during surgery results in paralysis of the serratus anterior muscle leading to winging of the scapula.

B: The external laryngeal nerve is a motor nerve that supplies the cricothyroid muscle. The close relationship of the external laryngeal nerve to the superior thyroid artery puts the nerve at potential risk when the artery is clamped during thyroid surgeries.

C: Excision of the submandibular gland results in potential risk to 3 cranial nerves:

- Marginal mandibular branch of the facial nerve
- Lingual nerve
- Hypoglossal nerve

D: The vagus nerve is at greater risk of injury during surgeries like thymectomy.

Solution to Question 37:

Patients with congenital hypertrophic pyloric stenosis (CHPS) will present with non-bilious projectile vomiting and visible peristalsis observed moving from left to right (not from right to left). Metabolic abnormalities include hypokalemia, hypochloremia and metabolic alkalosis.

CHPS occurs due to hypertrophy of the circular muscle of the pylorus, which results in constriction and obstruction of the gastric outlet.

- Patients commonly present between 2 and 8 weeks of age.
- First-born male infants are at the highest risk.
- Infants usually present with projectile, non-bilious vomiting.
- Clinically, an olive-shaped mass is palpable in the right upper quadrant or epigastrium. When the infant is given a test feed, visible peristalsis is usually observed moving from left to right (not from right to left), followed by projectile vomiting.

Metabolic abnormality:

- Infants with hypertrophic pyloric stenosis develop a characteristic metabolic abnormality i.e. hypokalemic hypochloremic metabolic alkalosis with paradoxical aciduria.
- Persistent vomiting results in loss of hydrochloric acid, leading to metabolic alkalosis (due to loss of H⁺) and hypochloremia (due to loss of Cl⁻).
- Kidneys try to compensate for alkalosis by secreting Na⁺ and HCO₃⁻. However, sodium loss stimulates aldosterone secretion through the RAAS pathway.
- Aldosterone causes active reabsorption of sodium in exchange for K⁺ and H⁺ ions leading to the development of hypokalemia and paradoxical aciduria, since the body initially loses H⁺ ions due to the loss of HCl, and instead of compensating for that loss, the body is losing more H⁺ ions through urine.

Management:

- The child should be rehydrated and the metabolic abnormality must be corrected before surgery.
- Ramstedt pyloromyotomy remains the standard surgical procedure for its treatment.

Solution to Question 38:

The quick Sequential Organ Failure Assessment (qSOFA) scoring system does not rely on the heart rate.

It is a bedside scoring system that may identify patients with suspected infection, who are at greater risk for sepsis outside the intensive care unit (ICU).

It uses three criteria, assigning one point each:

- Low blood pressure (SBP ≤ 100 mmHg)
- High respiratory rate (≥ 22 breaths per min)
- Altered mental status

The qSOFA suggests potentially life-threatening sepsis when at least two of the parameters are met.

Note :

The primary method to evaluate the risk of organ dysfunction (sepsis) in patients with suspected infection is by determining the sequential (sepsis-related) organ failure assessment (SOFA) score; which includes an assessment of:

- Circulation - Systolic BP & need for vasopressors; platelet count
- Respiration - PaO₂/FiO₂ & need for mechanical ventilation
- Mentation - Glasgow Coma Score
- Hepatic function - Bilirubin
- Renal function - Creatinine

Organ dysfunction is identified by a SOFA score of ≥ 2

The SOFA score is limited by its requirement of multiple laboratory tests which makes it complex and time-consuming. In order to simplify the scoring process, the "Sepsis -3" task force recommended the quick sepsis-related organ dysfunction assessment "qSOFA" score.

Solution to Question 39:

The medial border of the triangle of doom is bounded by vas deferens and the medial border of the myopectineal orifice is bounded by the lateral border of the rectus abdominis.

The triangle of doom is bounded


- Medially by the vas deferens
- Laterally by the vessels of the spermatic cord (gonadal vessels)
- Inferiorly by peritoneal reflection

The myopectineal orifice of Fruchaud is an osseo-myo-aponeurotic tunnel through which all groin hernias occur.

It is bounded:

- Medially by the lateral border of the rectus sheath.
- Laterally by the iliopsoas muscle.
- Above by the arched fibers of the internal oblique and transversus abdominis muscle.
- Below by the pectin pubis and fascia covering it.

Myopectineal orifice of Fruchaud



©Marrow

Solution to Question 40:

A-3, B-1, C-2, and D-4 is the correct option.

A: Uro bags are used for the collection of urine. They are usually attached to Foley's catheter, which is placed to drain the bladder (commonly in postoperative patients).

B: Nasogastric tube/Ryle's tube is used for nasogastric feeding/stomach wash/aspiration.

C: An underwater seal bag is used for the collection of drainage fluid from the thoracic cavity. It acts as a one-way valve allowing air to leave the pleural cavity, but it cannot be drawn back in by the negative pressure that is created in the intrathoracic cavity.

D: A metallic suction cannula is used to clear the operative field by suctioning secretions/blood and aspirating the fluid from various cysts and in a procedure like a hydrocele.

Solution to Question 41:

The given clinical scenario and erect X-ray abdomen showing gas under the diaphragm are suggestive of pneumoperitoneum, which is managed by initial resuscitation with IV fluids followed by exploratory laparotomy.

Pneumoperitoneum is the presence of air or gas in the peritoneal cavity. The most common cause of pneumoperitoneum is a perforation of a hollow viscus. Other causes of pneumoperitoneum in adults include peritoneal dialysis, and mechanical ventilation, after procedures like colonoscopy and endoscopy. In children, it can be caused by necrotizing enterocolitis, hirschsprung's disease, meconium ileus and mechanical ventilation.

The patient presents with abdominal pain and distension, vomiting, fever, constipation, tachycardia, tachypnea, hypotension, and reduced urine output. Per-abdomen examination can show guarding and rigidity.

An erect X-ray abdomen is highly sensitive for the detection of free intraperitoneal gas. Some of the signs seen in the X-ray include:

- Cupola sign is the presence of air underneath the diaphragm, giving a dome-like appearance.
- Rigler's sign is where both bowel surfaces are seen together as a sharp margin.
- Falciform ligament sign is the outlining of the falciform ligament with free abdominal gas.
- Liver/gallbladder edge sign is where the sharp margins of the liver/gallbladder are visible due to the collection of air.

Management includes,

- Adequate preoperative resuscitation with intravenous fluids followed by correction of electrolyte imbalance, instituting broad-spectrum intravenous antibiotics and
- Early surgical intervention/laparotomy to address the cause of pneumoperitoneum and prevent further contamination of the peritoneal cavity.

Surgery NEET 2018

Question 1:

Which of the following are indications for surgical intervention in abdominal aortic aneurysms?

- a) Size more than 5.5 cm in males
- b) Size more than 5 cm in females
- c) Expansion of >0.5 cm within 6 months
- d) All the above

Question 2:

The following image shows the retrograde ureteropyelogram of a patient with loin pain. What is the most probable clinical diagnosis?



- a) Hydronephrosis
- b) Duplex kidney
- c) Renal carcinoma
- d) Renal stone

Question 3:

Which of the following types of renal stones are produced by laxative abuse?

- a) Uric acid
- b) Ammonium urate
- c) Struvite
- d) Calcium oxalate


Question 4:

A patient is confused, opens his eyes on painful stimuli, flexes one limb on painful stimuli, and can localize the stimulus with the other limb. His Glasgow Coma Score is _____

- a) 8
- b) 9
- c) 10
- d) 11

Question 5:

A patient presents with a painless swelling of the neck. The histology of the tissue is as given in the image below. What is the diagnosis?




- a) Medullary carcinoma
- b) Non Hodgkin's lymphoma
- c) Grave's disease

d) Hashimoto's thyroiditis

Question 6:

A 45-year-old male patient who presented with a scrotal swelling underwent an orchidectomy. The image below shows the histopathology findings. What is your diagnosis?



- a) Seminoma
- b) Yolk-sac tumor
- c) Embryonal cell carcinoma
- d) Teratoma


Question 7:

A surgeon sees a necrotic bowel when operating on a child for an inguinal hernia. What is true about the resection and anastomosis of the bowel?

- a) The surgeon must use chromic catgut
- b) The surgeon must not include the submucosa in the sutures
- c) The surgeon must use the single-layer, extramucosal suture technique
- d) The surgeon must use the single-layer seromuscular suture technique

Question 8:

A lady comes with a reddish firm nodule on her chest which hasn't grown in size. Its histopathological image is shown below. What is the diagnosis?



- a) Lipoma
- b) Fibroadenoma
- c) Nevus
- d) Hemangioma

Question 9:

Which is the most common organism causing acute bacterial prostatitis?

- a) Enterococcus
- b) Proteus
- c) Streptococcus Agalactia
- d) E.coli

Question 10:

A 35-year-old man came with complaints of dark-colored stools to the OPD. After undergoing endoscopy, the patient was diagnosed with an upper GI bleed. Which of the following structures is used as a demarcation point to differentiate it from a lower GI bleed?

- a) Ampulla of Vater
- b) Ligament of Treitz

- c) Superior duodenal flexure
- d) Ileocaecal junction

Question 11:

Which is the best graft for a femoropopliteal bypass?

- a) Dacron
- b) Great saphenous vein
- c) Polytetrafluoroethylene
- d) Short saphenous vein

Question 12:

What is the most common cause of chronic pancreatitis?

- a) Gallstones
- b) Alcohol
- c) Autoimmune
- d) Drugs

Question 13:

A probe is inserted into the duct from which the nipple discharge is emerging. A tennis racquet incision is made and the flap is dissected to reach the duct. The duct is then excised. This procedure is known as?

- a) Macrodoectomy
- b) Microdoectomy
- c) Cone excision
- d) Wide excision of the lactiferous duct.

Question 14:

Which of the following is true about chromic catgut?

- a) It is made from cat gut

- b) It is made from rabbit gut
- c) It is absorbed by enzymatic degradation and macrophage phagocytosis
- d) It is tanned with nickel salts to improve handling and to resist degradation in tissue

Question 15:

Identify the instrument given in the image below:



- a) Deaver's retractor
- b) Morris retractor
- c) Czerney's retractor
- d) Double hook retractor

Question 16:

Extra mammary Paget's disease is commonly seen in the:

- a) Vulva
- b) Vagina
- c) Uterus
- d) Cervix

Question 17:

The best method to obtain tissue for the diagnosis of pancreatic carcinoma is:

- a) Transgastric FNAC guided by endoscopic ultrasound
- b) MRI-guided biopsy
- c) Laparoscopic-guided biopsy
- d) Percutaneous transperitoneal biopsy

Question 18:

An abdominal mass in a patient with congenital hypertrophic pyloric stenosis is best seen _____.

- a) During feeding
- b) Soon after birth
- c) During palpation over the left hypochondrium
- d) During palpation over the epigastrium

Question 19:

While performing a fasciotomy for compartment syndrome, the following layers are incised:

- a) Skin and Subcutaneous tissue
- b) Skin, Subcutaneous tissue, Superficial fascia
- c) Skin, Subcutaneous tissue, Superficial fascia and Deep fascia
- d) Skin, Subcutaneous tissue, Superficial fascia, Deep fascia and a few muscular fibres

Question 20:

Which of the following statements is true about Crohn's disease?

- a) For effective treatment of a Crohn's mass, only antibiotics are prescribed.
- b) Symptomatic relapse after surgery is not related to the presence of disease during resection.
- c) The proximal ileum is most commonly involved in Crohn's disease.
- d) On endoscopic examination, continuous lesions are visualised, which are diagnostic of Crohn's disease.

Question 21:

Which of the following is the earliest sign in Crohn's disease?

- a) Cobblestone appearance
- b) Deep fissured ulcers
- c) Rose-thorn appearance
- d) Aphthous ulcers

Question 22:

The Van Nuys grading system for ductal carcinoma in situ does not include:

- a) Age of the patient
- b) Size of the tumor
- c) Hormone receptor staining
- d) Presence of microcalcification

Question 23:

Which of the following statements are true with regards to a keloid scar?

- a) The extent does not cross the wound margins
- b) Wide excision of the keloid scar prevents recurrence
- c) Histopathology of a keloid scar shows reduced collagen and increased vascularity
- d) A keloid scar forms due to increase in the level of growth factors

Question 24:

Heller's operation is done for _____.

- a) Achalasia cardia
- b) Carcinoma esophagus
- c) Esophageal stricture
- d) Hypertrophic pyloric stenosis

Question 25:

Which of the following is not true about Barrett's esophagus?

- a) It is the intestinal metaplasia of esophagus
- b) Mucus secreting foveolar cells are characteristic
- c) Alician blue is used in the staining of biopsy specimen
- d) It is a reversible cell adaptation

Question 26:

Modified shock index is calculated using?

- a) HR/SBP
- b) HR/DBP
- c) HR/MAP
- d) PR/SBP

Answer Key

Question No.	Correct Option
1	d
2	a
3	b
4	d
5	d
6	a
7	c
8	d
9	d
10	b
11	b
12	b
13	b
14	c
15	b
16	a
17	a
18	a

19	c
20	b
21	d
22	c
23	d
24	a
25	b
26	c

Detailed Explanations

Solution to Question 1:

All the given are indications for surgical intervention in abdominal aortic aneurysm (AAA). The indications include:

- Size ≥ 5.5 cm in males (option A)
- Size ≥ 4.5 -5 cm in females (option B)
- Expansion of ≥ 0.5 cm within 6 months or ≥ 1 cm in one year (option C)
- Painful, tender aneurysm
- Thrombosed aneurysm, aneurysm with distal emboli

The rupture risk is high for aneurysms ≥ 5.5 cm and begins to rise exponentially with an increase in size. Women are at higher risk for rupture than men, with a lower threshold of 4.5 to 5.0 cm (depending on the reference used).


Screening is done for:

- Cardiovascular patients in men and women (60-85 years)
- Men and women ≥ 50 years with positive family history
- Annual screening in asymptomatic patients with 4-4.5 cm size, size ≥ 4.5 cm once in every 6 months.

Management of AAA can be done by open repair or endovascular aneurysm repair (EVAR). About 75% of infrarenal aneurysms are suitable for endovascular aneurysm repair (EVAR), depending on the morphology of the aneurysm assessed by a CT scan.

Aneurysms that fall below the mentioned indications may safely be followed with CT or ultrasound at 6-month intervals since long-term outcomes to routine follow-up are equivalent to earlier surgical repair of the aneurysm. Smoking cessation is very important and reduces the mortality rate.

The technique of EVAR:



Solution to Question 2:

The given image showing dilated clubbed calyces with stricture of the left ureteropelvic junction points to a diagnosis of hydronephrosis.

Hydronephrosis refers to the dilatation of the pelvicalyceal system caused by a partial or complete blockage of urinary flow.

Investigations:

- Ultrasound is the least invasive technique for diagnosing hydronephrosis.
- Retrograde pyelography is rarely indicated but will confirm the site of obstruction before corrective surgery.
- Intravenous urogram (IVU) is helpful only if the obstructed kidney is functional. In IVU, the extrarenal pelvis is dilated and the minor calyces lose their normal cupping. In early stages, there is blunting of calyces followed by its clubbing in the late stages.
- A renal isotope scan (DTPA or MAG-3) is the best modality to confirm the obstructive dilatation of the collecting system.


Solution to Question 3:

Ammonium urate stones are most likely to be seen in patients with laxative abuse.

Loss of potassium and sodium in laxative abuse causes intracellular acidosis, which increases the production of ammonia in urine. Dehydration and sodium loss further exacerbate hypokalemia and reduces the urinary volume, resulting in hyperuricemia. Highly concentrated uric acid and ammonia promote the formation of ammonium urate stones.

These are also associated with inflammatory bowel disease, chronic diarrhea, ileostomy, recurrent urinary tract infection and recurrent uric acid stone formation.

The below image shows an ammonium urate stone.



Solution to Question 4:

The GCS in this patient is 11.

- The score for verbal response: 4, because he is confused.
- The score for eye-opening: 2, because his eyes are opening to only painful stimuli.
- The score for best motor response: 5, because he can localize pain.


Therefore, the GCS score is $(4+2+5) = 11$.

Solution to Question 5:

The slide shows the histopathology of Hashimoto thyroiditis, with:

- Follicular destruction
- Reduced colloid with increased interstitial connective tissue
- Small lymphocyte and plasma cell infiltration
- A well-developed germinal centre may be seen
- Hurthle (Askanazy) cells that have abundant eosinophilic granular cytoplasm lining the follicles

Hashimoto's thyroiditis



Hashimoto's thyroiditis is the most common cause of hypothyroidism in iodine-sufficient areas. It is also known as struma lymphomatosa or lymphocytic thyroiditis. It is an autoimmune disease that results in the destruction of the thyroid gland and gradual & progressive thyroid failure.

Clinical features:

- Most prevalent between 45-65 years of age
- More common in females
- Painless enlargement of the thyroid
- Hashitoxicosis: When the auto-antibodies attack and damage the follicles, there is a sudden release of thyroid hormones from these follicles. This leads to a transient phase of thyrotoxicosis.
- Hypothyroidism: Since the follicles are damaged and cannot form the hormones anymore, these patients tend to be in a state of hypothyroidism.

Diagnosis:

- T₃ and T₄ are low
- TSH is raised
- Autoantibodies against:
 - Thyroid receptors
 - Thyroid peroxidase (TPO)
 - Thyroglobulin


These patients are managed with thyroid hormone replacement therapy.

In patients with long-standing goiter, there is a risk of compressive features and the development of thyroid lymphoma. Such cases can be treated with total thyroidectomy. It may also predispose to papillary thyroid carcinoma.

Other options:


Option A: Histological features of medullary carcinoma of the thyroid include polygonal to spindle-shaped cells in the form of nests, trabeculae or follicles. Small, round anaplastic cells may be found in some cases and amyloid deposits derived from calcitonin polypeptides are present in the stroma.

The given slide below shows medullary carcinoma of thyroid:



Option C: Grave's disease is an autoimmune disorder characterized by hyperthyroidism due to circulating autoantibodies. The thyroid-stimulating immunoglobulins bind to the TSH receptors, causing overgrowth of the thyroid gland and increased thyroid hormone synthesis. Histologically, there is acinar cell hypertrophy and hyperplasia with the absence of normal colloid in the tall columnar epithelium (normally the epithelium is flat with colloid). Lymphoid infiltrates, consisting predominantly of T cells, along with scattered B cells and mature plasma cells, are present throughout the interstitium. Germinal centers are common.

The given slide below shows Grave's disease:




Solution to Question 6:

Based on the gross and microscopic appearance of the specimen, the diagnosis is seminoma.


The gross specimen shows a well-circumscribed, pale, gray-white, fleshy, homogenous mass, with a lobulated cut surface, usually devoid of hemorrhage or necrosis. Histopathology shows uniform tumor cells with abundant clear cytoplasm, distinct cell borders, and large central nuclei. There is a sparse lymphocytic infiltrate. Classically, they have a fried egg appearance.

Other options:


Option B: Yolk-sac tumor has a yellowish-white, mucinous appearance. Histologically, they are typically composed of a lace-like (reticular) network of medium-sized cuboidal, flattened, or spindled cells, which generally show less cytologic atypia than embryonal carcinoma. In approximately 50% of tumors, structures resembling endodermal sinuses (Schiller-Duval bodies) may be seen; these consist of a mesodermal core with a central capillary and a visceral and parietal layer of cells resembling primitive glomeruli.



Option C: Embryonal cell carcinoma appears variegated with foci of hemorrhage or necrosis. Histologically, the cells grow in alveolar or tubular patterns, sometimes with papillary folds. More undifferentiated lesions may display sheets of cells with cleft-like spaces. The neoplastic cells have an epithelial appearance, are large and anaplastic, and have hyperchromatic nuclei with prominent nucleoli. The cell borders are usually indistinct, and there is considerable variation in cell and nuclear size and shape (pleomorphism). Mitotic figures and tumor giant cells are frequently seen.



Option D: Teratomas also have a variegated appearance but have cysts reflecting the multiplicity of tissue types. Histologically, collections of differentiated cells or organoid structures originating in all 3 germ layers can be seen. Elements may be mature (resembling various adult tissues) or immature (sharing histologic features with fetal or embryonic tissue).



Solution to Question 7:

The standard technique used currently for resection anastomosis of the bowel is a one-layer extra mucosal suture technique since it causes the least tissue necrosis and luminal narrowing. Interrupted and continuous single-layer suture techniques are adequate and safe.

The extra mucosal suture must also include the submucosa because this layer has a high collagen content and it is the most stable suture layer in all sections of the gastrointestinal tract.

Catgut has now been replaced with absorbable synthetic polymers. Stapling devices can be used when speed is required.

Solution to Question 8:

The given clinical scenario of a woman with reddish firm nodule on her chest and histology showing multiple dilated vascular structures with a single layer of epithelial cells is suggestive of a hemangioma.


Hemangioma is a benign tumor caused by the abnormal growth of blood vessels. Types of hemangioma are capillary hemangioma (most common), juvenile/ strawberry hemangioma, cavernous hemangioma and pyogenic hemangioma.

Indications for immediate treatment are complications such as ulceration or those pertaining to the location of the mass (e.g. visual issues due to a large hemangioma over the eyelid).

Treatment options include:

- Beta-blockers - oral propranolol
- Steroids - intralesional triamcinolone, topical clobetasol, systemic steroids
- Interferon alfa-2a and vincristine - for steroid-resistant lesions
- Local resection with cutting cautery or carbon dioxide laser

The given image shows a flat to elevated, irregular, pinkish-red mass over the skin of the abdomen suggestive of a strawberry hemangioma.



The given image shows a capillary hemangioma in a child



Solution to Question 9:

The most common organism causing acute bacterial prostatitis (ABP) is *Escherichia coli*.

It is commonly caused by an ascending urethral infection or reflux of infected urine into prostatic ducts. Sexually transmissible infections and other urinary pathogens, such as *Staphylococcus aureus* and *Streptococcus faecalis* may also be responsible.

The patient presents with fever, perineal pain, chills and rigor, and dysuria. Per rectal examination will reveal a tender, boggy prostate. Care should be taken to avoid aggressive palpation or prostatic massage as it can lead to bacterial dissemination and sepsis.

Urine examination and culture should be done. Urine examination reveals threads in urine.

Management:

Patients with systemic signs of infection should be admitted for IV antibiotics, hydration, and monitoring of laboratory parameters.

Patients who have no signs of systemic infection, who can tolerate oral intake, and who have no urinary retention can be treated with oral antibiotics. Antibiotics are given for 2-6 weeks to ensure the eradication of the infection as antibiotics don't penetrate the prostate capsule easily.


For outpatient therapy: Ciprofloxacin 500 mg oral every 12 hours or levofloxacin 500 mg OD or trimethoprim-sulfamethoxazole double-strength tablet oral every 12 hours can be used.

For hospitalized patients: Intravenous ciprofloxacin or levofloxacin with or without an aminoglycoside (gentamicin or tobramycin) or intravenous beta-lactam with or without an aminoglycoside. Gram-positive infections are treated with cephalosporins (cephalexin, cefazolin) or dicloxacillin and vancomycin.

Solution to Question 10:

The ligament of Treitz is used as the demarcation point to differentiate between upper gastrointestinal bleeding and lower gastrointestinal bleeding. It is a part of the duodenojejunal junction.

Upper GI bleeding refers to bleeding proximal to the ligament of Treitz, and lower GI bleeding refers to bleeding distal to the ligament of Treitz.



Solution to Question 11:

The great saphenous vein (GSV) is the best as it lasts longer than polytetrafluoroethylene (PTFE), dacron, and human umbilical vein as a graft in a femoropopliteal bypass.

The GSV is the conduit of choice for arterial bypass grafts below the inguinal ligament.

For bypass above the inguinal ligament, while both GSV and PTFE grafts have similar patency rates in the first 2 years, the GSV is superior thereafter and hence preferred over PTFE or any other grafts.

The ipsilateral GSV is the conduit of choice for femoropopliteal bypass. When it is not available, contralateral GSV should be used. Other veins are preferred when both ipsilateral and contralateral GSV is not available and they include a small saphenous vein, cephalic vein, and basilic vein.

Non-autogenous grafts are used for patients who lack a suitable venous conduit. The most commonly used prosthetic conduit is PTFE. Recent studies have shown that heparin-bonded PTFE is superior to compared to PTFE without heparin.

Solution to Question 12:

Alcohol consumption is the most common cause of chronic pancreatitis, accounting for 60–70% of the cases. Smoking increases the risk of alcohol-induced chronic pancreatitis.

Cystic fibrosis is the most common cause of chronic pancreatitis in the pediatric age group.

Other causes of chronic pancreatitis include:

- Pancreatic divisum
- Annular pancreas
- Hyperlipidemia
- Hypercalcemia
- Hyperparathyroidism
- Chronic duct obstruction
- Autoimmune pancreatitis
- Cystic fibrosis
- Tropical pancreatitis
- Hereditary pancreatitis

Solution to Question 13:

Microdochectomy is the excision of the entire lactiferous duct (single duct disease). It is done for duct papilloma.

Microdochectomy:

- In a duct papilloma where no palpable mass is present, the duct is expressed to observe the discharge.
- A fine probe or stiff nylon suture is gently inserted into the duct, and the probe is secured to the nipple skin using silk sutures to prevent displacement during the procedure.
- A tennis racquet shaped incision is made to elevate the skin flaps, and the duct with the probe is excised along with the skin flap for 5 cm until the papilloma is identified (The papilloma will always be located within 5 cm from the duct orifice).
- The wound is closed without a drain, and a compression bandage is applied.

It is ideal to rule out DCIS by frozen section biopsy.

Cone excision and wide excision are done under major duct excision (Adair-Haldfield operation). It is done for duct ectasias with multiple duct involvement and multiple papillomas.

Solution to Question 14:

Chronic catgut is absorbed through macrophage phagocytosis and enzymatic degradation within 90 days.

It is made from collagen derived from healthy sheep or cattle. It is then tanned with chromium salts to improve handling and resist degradation in tissue.

The tensile strength lasts for 21-28 days. It has a moderate tissue reaction as compared to plain catgut, which has a high tissue reactivity.

It is used to ligate superficial vessels, suture subcutaneous tissues, and is used for stomas and other tissues that heal rapidly. It is not used in tissues that heal slowly and require prolonged support.

Solution to Question 15:

The given image shows a Morris retractor with a single, flat transversely curved blade and a blunt projecting ridge with backward projection.

Morris retractor can be single or double-bladed. It is used to retract abdominal wall incisions, loin incisions, subcostal wounds or any wounds in depth.

It can be used to improve visibility on one side of an incision. Thus, it is valuable during the initial phase of a laparotomy.



Other options:

Option A: Deaver's retractor is a question mark-shaped stout, atraumatic retractor with a broad, gently curved long blade. It is used to retract the abdominal wall and viscera. Moist mops should be kept underneath prior to placing this retractor. It is used for major abdominal surgeries, bladder surgeries and lumbar sympathectomy.


Deaver's Retractor



©MARROW

Option C: Czerny's retractor is a superficial retractor with a thick and small blade on one side and a biflanged hook on the other side in opposite directions. It is used in hernia surgery and laparotomy, especially during closure.


Czerny's retractor



©MARROW

Option D: Double-hook retractor consists of two hooks with a handle. It is used in excision of swellings like lipoma, neurofibroma and cyst. It is also used to hold the skin flaps when raising them.

Double hook Retractor



©MARROW

Solution to Question 16:

Extramammary Paget's disease arises most frequently on the vulva. Other sites include the perineum, scrotum, perianal skin, and penis.

Extramammary Paget's disease (EMPD) can be primary or secondary.

- Primary EMPD is more common and occurs in the absence of an underlying malignancy, thought to arise from intraepidermal parts of apocrine glands or from pluripotent cells from the epidermis.
- Secondary EMPD arises from an underlying adnexal carcinoma or internal malignancy due to the epidermotropic spread of malignant cells from the underlying tumor.


Visceral malignancies associated with EMPD are carcinomas of the colon, rectum, bladder, urethra, cervix, and prostate. The patient presents with well-defined, moist, erythematous scaly plaques and severe pruritis. Lichenification and excoriations may also be seen.

Diagnosis is confirmed by punch, wedge, or excisional skin biopsy. A systemic workup should also be done to identify any underlying malignancy and they include colonoscopy, cystoscopy, colposcopy, and pelvic examination with pap smear in females.

Treatment:

- Mohs microscopic surgery with 5 cm margin clearance
- Radiotherapy
- Topical 5-fluorouracil and imiquimod
- Chemotherapy
- Photodynamic therapy

The given image below shows extramammary Paget's disease of the vulva.



Solution to Question 17:

Transgastric or transduodenal FNAC or trucut biopsy performed under endoscopic ultrasound (EUS) guidance is the best method to obtain tissue for the diagnosis of pancreatic carcinoma. This is because it avoids the spillage of tumor cells into the peritoneal cavity.

Endoscopic ultrasonography with a fine needle aspiration (EUS-FNA) is done if CT has failed to demonstrate a tumor. When a tissue diagnosis is required prior to surgery to distinguish between inflammation and neoplasia, EUS-FNA is helpful.

Diagnostic laparoscopy prior to resection can spare an unnecessary laparotomy by identifying small peritoneal and liver metastases.

Percutaneous transperitoneal biopsy of potentially resectable pancreatic tumors is avoided.

Solution to Question 18:

In a child with congenital hypertrophic pyloric stenosis (CHPS), the abdominal mass is best seen during test feed in the right upper quadrant.

Hypertrophic pyloric stenosis most presents with non-bilious projectile vomiting. It is common between 2 and 8 weeks of age. Hypertrophy of the circular muscle of the pylorus results in constriction and obstruction of the gastric outlet. Male infants are at a higher risk.

Clinically, when the infant is given a test feed, an olive-shaped mass is palpable in the right upper quadrant and visible peristalsis is usually observed moving from left to right across the abdomen.

Infants with hypertrophic pyloric stenosis develop a characteristic metabolic abnormality- hypochloraemic hypokalaemic metabolic alkalosis. Persistent vomiting results in loss of hydrochloric acid leading to the development of metabolic alkalosis (due to loss of H⁺) and hypochloremia (due to loss of Cl⁻).

All infants with congenital hypertrophic pyloric stenosis are evaluated with an ultrasound (investigation of choice). It confirms the diagnosis in the majority of cases.


Management: Rehydration and metabolic correction prior to definitive surgery - Ramstedt's pyloromyotomy.

Solution to Question 19:

The incision for a fasciotomy involves the skin, subcutaneous tissue, superficial fascia, and deep fascia.

Fasciotomy involves incising the deep muscle fascia and is best carried out via longitudinal incisions of skin, subcutaneous fat, and superficial and deep fascia. The muscle will be seen bulging out through the fasciotomy opening.

The lower limb can be decompressed via two incisions, each being lateral to the subcutaneous border of the tibia. This gives access to the two posterior compartments and the leg's peroneal and anterior compartments.



Solution to Question 20:

Symptomatic relapse of Crohn's disease after surgery is not related to the presence of the disease during resection.

The course of the disease after surgery is unpredictable since surgery does not cure Crohn's disease. Instead, the probability of relapse increases with the time since resection.

Other options:

Option A: Antibiotics alone cannot treat a Crohn's mass effectively. Abscesses can be treated with metronidazole and ciprofloxacin and should also undergo drainage (and resection if needed).

Option C: The terminal ileum is most commonly involved in Crohn's disease.

Option D: On endoscopic examination, discontinuous "skip" lesions are visualized. These are irregular areas of inflamed mucosa that are ulcerated with mucopurulent exudate separated by intervening, normal-appearing intestinal mucosa.

Solution to Question 21:

Aphthous ulcers, surrounded by a rim of erythematous mucosa, are the earliest lesions seen in a patient with Crohn's disease during endoscopy.

Crohn's colitis usually spares the rectum and is most severe in the ileocaecal region.

- In early Crohn's disease, small, punched-out aphthous ulcers are often seen (the result of submucosal lymphoid follicle expansion).
- In moderate disease, the aphthous ulcers coalesce into larger ulcers, which may take on the appearance of a star (stellate ulcers).
- As disease severity increases, submucosal edema and injury can result in cobblestoning of the mucosa (option A) and eventually the formation of serpiginous ulcers, which are typical findings in Crohn's disease.

Thorn-like extraluminal projections in barium studies are known as rose-thorn appearance, and are a typical sign of Crohn's disease. They represent deep penetrating linear ulcers or fissurings typically seen within stenosed terminal ileum with a thickened wall. (Options C and D)

Solution to Question 22:

Staining for estrogen and progesterone receptors is not part of the Van Nuys system used for grading DCIS.

The components of the Van Nuys prognostic index (VNPI) are as follows:

- Size of the tumor
- Extent of resection margin
- Histological variant of DCIS - based on nuclear grade, presence of necrosis, presence of microcalcification
- Age of the patient

Patients with a high score in this grading system benefit from radiotherapy after excision, whereas those with low-grade and completely excised tumors, need no further treatment.

A score ranging from 1 (lesions with the best prognosis) to 3 (lesions with the worst prognosis) is given for each of the four prognostic predictors. Total scores range from 4 (least likely to recur) to 12 (most likely to recur).

Note: The type of microcalcification seen on imaging, helps in determining the nuclear grade of DCIS. While the linear branching type of microcalcification is associated with high nuclear grade, fine granular type is associated with low nuclear grade.

Silverstein Van Nuys prognostic index for DCIS

Parameters	Score 1	Score 2	Score 3
Size(mm)	≤15	16-40	≥41
Margin(mm)	≥10	1-9	<1
Class	Grade 1 or 2 without necrosis	Grade 1 or 2 with necrosis	Grade 3 with or without necrosis
Age(years)	>60	40-60	<40

Scores 4-7 with >3mm margin:Excision alone
 Score 7 with < 3 mm margin, score 8 with > 3mm margin, score 9 with >5mm margin:Excision with RT
 Score 9 with < 5mm margin & score 10,11,12:Mastectomy


Solution to Question 23:

Keloid scars are associated with elevated levels of growth factor leading to increased proliferation of fibrous tissue.

A keloid scar is defined as excessive scar tissue that extends beyond the boundaries of the original incision or wound (option A) and rarely regresses on its own. It is more common in darker-pigmented ethnicities.

It can arise from surgery, burns, skin inflammation, infections, ear piercing, and even spontaneously. It occurs 3 months to years after the initial insult with even minor injuries causing keloid formation.

It most commonly occurs on the skin of the earlobe, deltoid, presternal, upper back, and rarely extends into the subcutaneous tissue.



The histology of keloid scars shows excess collagen and hypervascularity with collagen fibres arranged haphazardly in loose sheets (option C).

Treatment: Keloid scars do not improve spontaneously with time and excision alone has a high rate of recurrence (option B). The following can be done:

- Pressure - local moulds or elastic garments.
- Silicone gel sheeting.
- Intralesional triamcinolone injection.
- Excision and steroid injections.
- Excision and postoperative radiation (external beam or brachytherapy).
- Intralesional excision.
- Laser to reduce redness.
- Vitamin E and palm oil massage.

Differences between Keloids and Hypertrophic scars:

	Keloid	Hypertrophic scar
Incidence	Rare	Frequent
Ethnic groups	African, American, Asian, Hispanic	No predilection
Prior injury	Yes	Yes
Site predilection	Neck, chest, earlobes, shoulders, upper back	Anywhere
Genetics	Autosomal dominant with incomplete penetration	No
Timing	Symptom-free interval; may appear years after injury	4-6 weeks post-injury

	Keloid	Hypertrophic scar
Symptoms	Pain, pruritus, hyperesthesia, growth beyond wound margins	Raised, some pruritus, respects wound confines
Regression and contracture	No, and rare	Frequent
Histology	Hypocellular, thick, wavy collagen fibres in random orientation	Parallel orientation of collagen fibres

Solution to Question 24:

The most common surgical procedure for achalasia cardia is laproscopic Heller myotomy in conjugation with partial fundoplication.


Solution to Question 25:

Mucus-secreting goblet cells (not foveolar cells) are characteristic of Barrett's esophagus, as there is intestinal metaplasia of the esophagus. Foveolar cells are the mucus-secreting cells in the stomach.

Barrett's esophagus is a complication of chronic gastroesophageal reflux disease (GERD), and it is estimated to occur in about 10% of individuals with symptomatic GERD. Symptoms include heartburn, regurgitation and sometimes dysphagia. Obese white men over the age of 60 years represent the highest-risk group.

Intestinal metaplasia in the esophagus occurs due to chronic irritation of the esophagus in GERD (option A) since intestinal epithelium is better suited to handling gastric acid exposure. Hence, mucus-secreting goblet cells are characteristic of Barrett's esophagus. Since all metaplasia are reversible, Barrett's esophagus is also reversible (option D). Only 0.2 to 1% of Barrett's esophagus leads to dysplasia and adenocarcinoma.


Barrett's Esophagus



Alcian blue is used in the staining of the biopsy specimen, as it stains the goblet cells (option C). Mucicarmine is the specific stain.

Seattle protocol is used for the management of Barrett's esophagus.

Management of Barrett's Esophagus (Seattle Protocol)



Solution to Question 26:

- Modified shock index = HR/MAP
- Original shock index = HR/SBP


The shock index is a hemodynamic stability indicator. In patients with shock, it is a better predictor of mortality than blood pressure or heart rate alone. A shock index >0.9 is indicative of higher mortality.

The modified shock index is a more sensitive indicator of tissue perfusion than the original shock index. In critically ill patients, the DBP decreases earlier than SBP, hence, the modified shock index using MAP is better suited to assess the severity of shock.

Surgery NEET 2019

Question 1:

The type of knot seen in the image is:



- a) Half hitch knot
- b) Reef knot
- c) Surgeon's knot
- d) Crossed half hitch knot

Question 2:

Which of the following drugs causes carcinoma bladder?

- a) Cyclophosphamide
- b) Cisplatin
- c) Taxane
- d) Tamoxifen

Question 3:

The most common site of branchial cyst is _____.

- a) Junction of upper and middle third of the sternomastoid muscle
- b) Junction of middle and lower third of the sternomastoid muscle
- c) Middle of the sternomastoid muscle
- d) Between the two heads of the sternomastoid muscle

Question 4:

Dohlman's procedure is for:

- a) Meckel's diverticulum
- b) Zenker's diverticulum
- c) Bochadlek hernia
- d) Menetrier's disease

Question 5:

Parathyroid gland post surgery is commonly transferred to:

- a) Triceps
- b) Biceps
- c) Sartorius
- d) Brachioradialis

Question 6:

What is the drug used for estrogen dependant breast cancer?

- a) Estrogen
- b) Clomiphene citrate
- c) Tamoxifen
- d) Adriamycin

Question 7:

The commonest surgical approach to a retrosternal goiter is via:

- a) Neck

- b) Chest
- c) Median sternotomy
- d) None of the above

Question 8:

An intern placed an intercostal drainage tube in a patient. Which of the following steps is incorrect?

- a) Identified 5th intercostal space anterior to mid axillary line to place the tube
- b) Inserted tube along the lower border of upper rib
- c) Directed tube posteriorly to prevent injury
- d) Incised and digitally explored

Question 9:

What is the size of the given blade used for incision and drainage of a superficial abscess?



- a) 10
- b) 11
- c) 22
- d) 15

Question 10:

In a blast injury, which of the following organs is least vulnerable to the blast wave?

- a) GI tract
- b) Lungs
- c) Liver
- d) Ear drum

Question 11:

A man presents with a slowly progressive swelling over the past 1 and 1/2 years. On examination, it has a variable consistency and appears as shown in the image. What is the probable cause?



- a) Adamantinoma
- b) Dermoid cyst
- c) Pleomorphic adenoma
- d) Sebaceous cyst

Question 12:

What is shown in the given image?



- a) Dermoid cyst
- b) Sebaceous cyst
- c) Meningioma
- d) Post auricular fistula

Question 13:

A female patient presents with a breast abscess. An aspiration was attempted under antibiotic cover which failed to reveal any aspirate. What is the next step in management?

- a) Incision and Drainage
- b) Continue antibiotics only
- c) A second attempt at aspiration
- d) Stop antibiotics as no aspirate was collected

Question 14:

Which nerve is involved in Frey's syndrome?

- a) Facial Nerve
- b) Oculomotor nerve
- c) Auriculotemporal nerve
- d) Glossopharyngeal nerve

Question 15:

Which of the following statements is false regarding deep vein thrombosis (DVT)?

- a) Clinical assessment is unreliable
- b) Mostly bilateral
- c) Most common clinical presentation is pain and tenderness in calf
- d) Some cases may directly present as pulmonary thromboembolism

Question 16:

A truck driver presents with dilated, tortuous veins of 4 mm in the foot. What is the stage of clinical classification according to CEAP(Clinical-etiology-anatomy-pathophysiology for chronic venous disorders)?

- a) Co
- b) C1
- c) C2
- d) C3

Question 17:

Which of the following statements is false regarding rectal carcinoma?

- a) Hartmann's operation is done in elderly debilitated patients
- b) Adenocarcinoma is a rare variant of rectal carcinoma
- c) Early morning spurious diarrhoea and tenesmus can occur
- d) Growth confined to the rectal mucosa is stage A of modified Duke's staging

Question 18:

A gasless abdomen is seen in:

- a) Ulcerative colitis
- b) Intussusception
- c) Acute pancreatitis
- d) Crohn's disease

Question 19:

Gall stone chemistry most commonly seen amongst the following is:

- a) Crystalline cholesterol monohydrate
- b) Crystalline cholesterol dihydrate
- c) Amorphous cholesterol dihydrate
- d) Amorphous cholesterol monohydrate

Question 20:

A 56-year-old patient suffering from alcoholic cirrhosis receives a healthy liver from his biological son. What is this type of transplant known as?

- a) Autograft
- b) Allograft
- c) Xenograft
- d) Isograft

Question 21:

Esophageal endoscopy of a patient reveals the presence of serpiginous ulcers. This finding indicates:

- a) CMV esophagitis
- b) Candida esophagitis
- c) Herpetic esophagitis
- d) Radiation esophagitis

Question 22:

Which of the following is not a part of revised trauma score (RTS)?

- a) Glasgow coma scale
- b) Systolic blood pressure
- c) Pulse rate
- d) Respiratory rate

Answer Key

Question No.	Correct Option
1	b
2	a
3	a
4	b
5	d
6	c
7	a
8	b
9	b
10	c
11	c
12	b
13	c
14	c
15	b
16	c
17	b
18	c
19	a
20	b
21	a
22	c

Detailed Explanations


Solution to Question 1:

The given image shows the suture material exiting the knot is on the same side as the entry i.e no cross-over is seen. This is seen in a reef knot.

A knot is a fastening made of threads intertwined together or on themselves. Each knot consists of- 2 half-hitches placed in sequential throws. Squaring refers to tightening the knot/half-hitch in a manner that the threads lay flat due to the alternating direction of half-hitches in adjacent layers.

Types of knots:

- **Granny knot:** half-hitch over a half-hitch in the same direction. It has better-holding power than half-hitch
- **Reef knot:** half-hitch followed by a second half-hitch after squaring.
- **Triple throw knot:** Reef knot with another half-hitch knot for added security. Knots made with prolene sutures require 6-8 throws to prevent slippage.
- **Surgeon's knot:** Modification of the reef knot with two half-hitches in the first layer.
- **Slip knot:** Modification of either granny/reef knot formed when one end of the thread is kept taut while tightening the knot.



Solution to Question 2:

Cyclophosphamide (an alkylating agent) causes bladder cancer

The risk of bladder cancer with cyclophosphamide is directly proportional to the duration and intensity of cyclophosphamide treatment. Phosphoramidate mustard is the primary mutagenic metabolite that causes bladder cancer in patients exposed to cyclophosphamide.

Painless, gross hematuria is the most common symptom that is indicative of a bladder carcinoma until proven otherwise.

Depending upon the depth of invasion, bladder carcinoma can be divided into

- Flat, non-invasive carcinoma in situ (CIS)
- Non-muscle invasive bladder cancer (70% of cases) - does not invade detrusor muscle but can invade lamina propria.
- Muscle-invasive bladder cancer - invades the detrusor muscle.

Diagnosis is by cystourethroscopy (directly visualizing the growth) and biopsy. Management is as follows:

- Non-muscle invasive bladder cancer:
- Transurethral resection of bladder tumor (TURBT) followed by intravesical chemotherapy with mitomycin C or epirubicin
- Immunotherapy with regular intravesical instillations of Bacille Calmette Guerin (BCG) in high-risk tumors.
- Muscle invasive bladder cancer: Radical cystectomy along with urinary diversion procedure and lymph node dissection +/- neoadjuvant chemotherapy

Other options:

Option B: Cisplatin is an anticancer drug used to treat testicular cancer. It is associated with an increased risk for acute myeloid leukemia (AML).

Option C: Taxane is an anticancer drug and is the main component in the treatment regimens of metastatic cancers. Taxane's adverse effects are neutropenia and fluid retention (pulmonary edema).

Option D: Tamoxifen is a selective estrogen receptor modulator (SERM) and is used in the treatment of estrogen-dependent breast cancer. Due to its partial agonistic activity, it is known to increase endometrial cancer in post-menopausal women.

Solution to Question 3:

The most common site of a branchial cyst is at the junction of the upper and middle third of the sternomastoid muscle at its anterior border.

The below image shows a large right-sided branchial cyst.




Branchial cyst develops due to the persistence of the cervical sinus. The cervical sinus is formed when the 2nd arch mesoderm overgrows the succeeding arches and overhangs them as shown in the image. The space between the overhanging 2nd pharyngeal arch and the 3rd, 4th, and 6th arches is called the cervical sinus. The cavity of the cervical sinus is lined by ectoderm and

normally gets obliterated.

Branchial cysts are usually lined by squamous epithelium and contain thick, turbid fluid full of cholesterol crystals. It is fluctuant and may transilluminate when it contains clear fluid.

A branchial fistula may be acquired or congenital. The congenital fistula is seen at the junction of the middle and lower third of the sternomastoid (option B). An acquired fistula is usually seen in cases of incomplete surgery for the branchial cyst or spontaneous rupture, and it is seen at the same site as the cyst.



Solution to Question 4:

Dohlman's procedure is done for Zenker's diverticulum.

It is a transoral endoscopic approach, that ablates the septum between the diverticulum and the cervical esophagus. It can be used for diverticula between 2 and 5 cm in size.

Other options:

Option A: Meckel's diverticulum is a remnant of the vitellointestinal duct and is treated by surgical resection.

Option C: Bochadlek's hernia is a congenital posterolateral defect on the left side of the diaphragm due to defective embryological development.

Option D: Menetrier's disease is a premalignant condition with gross hypertrophy of gastric mucosal folds and is treated with gastrectomy.

Solution to Question 5:

The parathyroid is commonly transferred to the brachioradialis muscle post-surgery. This will facilitate easy removal if there is disease recurrence after surgery.

Parathyroidectomy is indicated in:

- Symptomatic primary hyperparathyroidism (HPT):
- Asymptomatic primary HPT with:
 - raise in serum calcium $>1\text{mg/dl}$ of the normal range
 - $>400\text{ mg}$ urinary calcium in 24-hour urine test
 - <-2.5 Z-score of bone mineral density
- Age $>50\text{yr}$ and failed medical treatment
- Severe cases of secondary HPT

Parathyroidectomy is performed in neck hyperextension (semi-Fowler position). The thyroid gland is mobilized, and the parathyroid gland with adenoma is identified and resected. The parathyroid tissue is transplanted to the non-dominant forearm.

Pockets are made in the belly of the brachioradialis muscle, and one to two pieces of parathyroid tissue measuring 1 mm each are placed into each pocket. A total of 12 to 14 pieces are transplanted. This is a type of heterotopic transplant.

Miami criteria (confirm the absence of another hyperactive parathyroid): A drop of parathyroid hormone intraoperatively, to more than half the maximum value at 10 minutes postresection is considered an adequate resection in a parathyroid adenoma. However, if there is no drop in PTH level, then a multi-gland disease is suspected and the surgery is converted to a bilateral neck exploration.

Complications of parathyroidectomy include:

- Persistent or recurrent hyperparathyroidism
- Hypoparathyroidism (presenting as hypocalcemia)
- Hungry bone syndrome
- Injury to the recurrent laryngeal nerve

Solution to Question 6:

Tamoxifen is a selective estrogen receptor modulator (SERM) that is mainly used for estrogen receptor-positive breast cancers in pre-menopausal women.

It exhibits selective agonistic and antagonistic activity for ER receptors ($\text{ER}\alpha$ and $\text{ER}\beta$):

- Agonist activity at the uterus, bone, liver, and pituitary
- Antagonist activity in breast and blood vessels

Uses:

- Part of the primary treatment regimen for estrogen receptor-positive (ER^+) breast cancers in pre-menopausal women
- Adjuvant therapy after breast conservative surgery to prevent a recurrence

- Tamoxifen also is effective (a 40%–50% reduction in tumor incidence) in preventing ER+ breast cancer in women at increased risk.

Adverse effects:

- Vasomotor symptoms (hot flashes) and sweating
- Endometrial cancer
- Increased risk of thromboembolic events
- Fatty liver disease

Solution to Question 7:

The commonest surgical approach to a retrosternal goiter is via the neck and the most common type of goitre approached is the secondary retrosternal goiter.

A goiter is considered to be retrosternal if at least 50% of the thyroid tissue is located intrathoracically. They can be primary or secondary.

Retrosternal goiter presents with dyspnoea, stridor, and dilated neck veins. Pemberton's sign (venous congestion of the neck on the raising of hands above the head) is seen.

Contrast-enhanced computed tomography (CECT) of the neck is performed to assess the location of the thyroid. The definitive treatment is surgical excision of the ectopic thyroid.

Other options:

Option B & C: Median sternotomy (via chest) is indicated in:

- Primary mediastinal goiter
- Malignant retrosternal goiter
- Very large retrosternal goiter
- Recurrence in the mediastinum

	Primary retrosternal goiter	Secondaryretrosternal goiter
Occurrence	1%	Majorityof retrosternal goiter s
Origin	Arise fromectopic thyroid tiss uein the chestNo connection with the thyroid tissue in the neck	Arise from thedownward extension of cervical thyroid tissue along the fascial planes of the neck
Blood supply	Intrathoracicblood vessels	Superior and inferior thyroid arteries

Solution to Question 8:


An intercostal tube (ICD) is placed along the upper border of the lower rib at the 5th intercostal space (ICS) to prevent damage to the neurovascular bundle. Hence, option B is incorrect.

Chest tube insertion (thoracostomy) is indicated in pneumothorax, hemothorax, chylothorax, and pleural effusion in order to drain air/fluid from the pleural space.

The triangle of safety, usually in the 5th intercostal space, is the safest site for chest tube insertion, it is bound by:


- anterior to the midaxillary line
- above the horizontal line marked by the level of the nipple
- below and lateral to the pectoralis major muscle

Triangle of safety



©Marrow

Insertion of chest drain



Contraindications include pulmonary adhesions, pulmonary trauma, coagulopathy, and diaphragmatic hernia.

Complications of this procedure:

- Infection at the site of insertion
- Unintentional injury to the lung - hemothorax
- Intercostal nerve/ vessel injury
- Subcutaneous hematoma/ emphysema

Solution to Question 9:

The size of the surgical blade shown is the 11-size blade (stab knife) that has a triangular shape with an oblique edge and a sharp-pointed tip.

Uses:

- Incision and drainage (I & D) of an abscess
- Insertion of drains and arteriotomy.

Other options:

Option A: The 10-size blade is used for skin incisions. It has a curved cutting edge that tapers to a pointed tip.

Option C: The 22-size blade is often used for abdominal incisions. It has a flat, unsharpened back edge and a curved front cutting edge.

Option D: The 15-size blade is used for minor surgical procedures (lipoma excision) and precise skin incisions. It is a smaller blade with a small cutting edge.



Solution to Question 10:

In a blast injury, the liver is the least vulnerable to blast wave (primary blast) injury.

Blast injury is a complex type of physical trauma resulting from direct or indirect exposure to an explosion. Primary blast injuries result from the rapid overpressure or shock waves produced by an explosion. These injuries result from the dramatic changes in barometric pressure projecting outwards from the point of detonation. They predominantly cause damage to air-filled hollow organs of the body from rapid pressure change (barotrauma). This includes the middle ear, lungs, and GIT.

The most sensitive and most frequently injured hollow organs are the tympanic membranes & lungs. Blast damage to the lungs is the most common cause of life-threatening injury following an explosion.

Solid organs are the most resistant to blast injuries as the parenchymal disruption requires very high levels of overpressure.

Classification of blast injuries:

Classification	Explanation	Injury type	Examples
Primary blast	Due to air pressure from the blast	Shockwave (overpressure)	Tympanic membrane injury, blast lung, intestinal blast injury
Secondary blast	Fragments from the blast, following a detonation	Penetrating/fragmentation	All penetrating injuries
Tertiary blast	Gross movements of persons, objects, or infrastructure by the blast wind	Blunt trauma	Blunt and crush injuries. Traumatic amputation.
Quaternary blast	Miscellaneous causes	Exposure to explosive products or toxic substances	Burns, inhalation injury
Quinary blast	The intentional addition of biological or radioactive material to the explosive device	Environmental contaminants	Radiation sickness, infection.

Solution to Question 11:

A slowly progressive swelling, with a variable consistency, lifting the ear lobe, is suggestive of pleomorphic adenoma of the parotid gland.

Pleomorphic adenoma, the most common benign salivary gland tumor, primarily affects the parotid gland, particularly its superficial lobe. It is also referred to as a mixed tumor due to its combination of epithelial and mesenchymal components. This tumor most commonly occurs in females around the age of 40.

Clinical features:

- Painless, slow-growing swelling
- Lobulated, firm, variable in consistency
- Lifts the earlobe
- No facial nerve involvement

- Curtain sign positive - cannot be moved above the zygomatic bone.
- In case of deep lobe involvement, tonsils, and faucial pillars will be pushed medially.
- Features of malignancy - a sudden increase in size, ulceration of the overlying skin, facial nerve palsy (involvement of facial nerve), painful swelling, and restriction of jaw movements.

Fine needle aspiration cytology (FNAC) is the investigation of choice in diagnosing pleomorphic adenoma. Open biopsy is contraindicated due to the chances of injury to the facial nerve.

All tumors of the superficial lobe of the parotid gland should be managed by superficial parotidectomy. Radical parotidectomy is performed for malignant pleomorphic adenoma.

Other options:

Option A: Adamantinoma is a low-grade malignant tumor that usually presents with pre-tibial pain and swelling since it most commonly affects the tibial diaphysis.

Option B: Dermoid cysts present as firm painless subcutaneous nodules and are of 4 types:

- Sequestration dermoid - sequestration of epithelium during embryological development.
- Tubulodermoid - accumulation of secretions from ectodermal tube/duct
- Implantation dermoid - implantation of the epidermis due to trauma
- Teratomatous dermoid - having all 3 germ layers (ecto, meso, and endoderm)

Option D: Sebaceous cysts are retention cysts of the sebaceous glands that present as smooth, non-tender, non-transillumination swelling with a punctum (in 70% of the patients).

Solution to Question 12:

The image shows a dark-colored summit over a swelling, usually seen in a sebaceous cyst.

Sebaceous cysts are the retention cysts of the sebaceous gland that are usually present on the face, scalp, or scrotum. They contain yellowish-white cheesy material with fat and epithelium.

Clinical features:

- The swelling is painless, soft, and freely mobile.
- Though non-transilluminating, they are Paget's test positive (fluctuant).
- Punctum is seen in 70% of the patients with sebaceous cysts. They are dark in color due to the accumulation of denuded epithelium.
- They also have an unpleasant sebum odor.
- The swelling can ulcerate and become painful (Cock's peculiar tumor).
- Some cysts over the scalp may also harden over time (Sebaceous horn).
- Many may get infected and lead to abscess formation.

Treatment:

- Excision including skin adjacent to punctum using elliptical incision-dissection method.
- Incision and avulsion of cyst wall.

- In the presence of an abscess, an initial treatment with an incision later followed by excision of the cyst

Other options:

Option A: Dermoid cysts present as firm painless subcutaneous nodules and are of 4 types:

- Sequestration dermoid - sequestration of epithelium during embryological development
- Tubulodermoid - accumulation of secretions from ectodermal tube/duct
- Implantation dermoid - implantation of the epidermis due to trauma
- Teratomatous dermoid - having all 3 germ layers (ecto, meso, and endoderm)

Option C: Meningiomas are benign tumors arising from the meningotheelial cells of the dura. They present with headaches, seizures, or focal neurological deficits.

Option D: Postauricular cutaneous mastoid fistula is a rare complication of chronic otitis media that heals very slowly and needs surgical closure.

Solution to Question 13:

In a patient with a breast abscess, the next step after a failed aspiration is to attempt aspiration once again under antibiotic cover.

Incision and drainage is not preferred, as it can potentially interfere with breastfeeding, and delicate structures such as ducts, lobules, and tubules may be injured, possibly leading to a non-healing milk fistula.

A breast abscess is a complication of acute mastitis, which is more common in new mothers at the beginning of lactation (lactational mastitis). Non-lactational mastitis is more common in diabetic, immunosuppressed people and people predisposed by duct ectasia and periareolar infections.

Clinical features of a breast abscess include:

- Fever and malaise
- Localized pain and tenderness
- Redness and swelling of the affected breast
- Positive fluctuation test.



An ultrasonogram of the affected breast is performed to confirm the abscess, and treatment proceeds as follows:

- Analgesics, antipyretics, and breast support for pain and fever relief.
- Aspiration is performed under antibiotic cover, and the pus is cultured. If aspiration proves unsuccessful, it should be reattempted after a few days.
- Incision and drainage should be considered in cases of:
 - Mastitis unresponsive to antibiotics for 48 hours.
 - Brawny induration.
 - Progressive mastitis with persistent fever

Note:

	Bacterial mastitis	Breast abscess
Condition	Localized inflammation due to ascending infection or infection due to milk stasis	Occurs as a complication of mastitis Pus formation
Symptoms	Dull pain	Throbbing pain
Signs (if any)	Tenderness Erythema	Point tenderness Erythema Induration of skin (if chronic) Lymphadenopathy Fluctuation test +ve Pus point may be present
Management	Analgesics Antibiotics (Dicloxacillin 500 mg every 6 hours orally; Erythromycin if allergic to penicillin) Adequate breast support	Analgesics Adequate breast support Aspiration of Pus under antibiotic cover; repeated aspirations if required Incision & drainage


Solution to Question 14:

The auriculotemporal nerve (V3 nerve) is involved in Frey's syndrome (gustatory sweating).

Frey's syndrome is also known as Dupuy syndrome or auriculotemporal syndrome. It commonly develops after parotid surgery. It is caused by an aberrant regeneration of post-ganglionic parasympathetic nerve fibers crossed along the sympathetic pathways to the sweat glands near the parotid region.

Clinical features are due to the result of salivary stimulation (by the smell or taste of food) leading to the sweating and flushing of the periauricular region on the involved side of the face. A starch iodine test is used for diagnosis.

Treatment ranges from antiperspirants (in mild cases) to tympanic neurectomy or botulinum toxin injection (in severe cases). It can be prevented by incorporating careful extracapsular dissection and closure of the superficial musculoaponeurotic layer during parotidectomy.



Other options:

Option A: Facial nerve is the 7th cranial nerve that supplies the muscles of the face, and results in facial paralysis if injured during parotidectomy

Option B: Oculomotor is the 3rd cranial nerve which supplies the extraocular muscles. If injured during ophthalmic surgeries, it results in ocular paralysis and the eye pointing down and out.

Option D: Glossopharyngeal nerve is the 9th cranial nerve and its injury causes loss of taste sensation from the posterior 1/3rd of the tongue.

Solution to Question 15:

Most cases of deep vein thrombosis (DVT) are unilateral. It is bilateral in only 30% of the individuals.

DVT refers to thrombosis of the deep venous system.

Caused by:

- Postoperative thrombosis (most common cause)
- Oral contraceptive pills (OCPs)
- After childbirth
- Immobility

Sites:

- Pelvic veins—common
- Leg veins—common in femoral and popliteal veins (common on the left side).

Fever is the earliest symptom, which is also most commonly accompanied by pain and tenderness in the calf and thigh (option C).

The following signs could be elicited:

- Positive Homan's sign: Resistance of calf muscles on passive forceful dorsiflexion of the foot.
- Mose's sign: Gentle squeezing of the lower part of the calf from side to side is painful.
- Neuhof's sign: Thickening and deep tenderness seen on deep palpation of the calf muscles.

Sometimes, the thrombus may get dislodged and present as pulmonary thromboembolism (option D) with features of dyspnoea, tachypnoea, and chest pain.

Clinical features are unreliable for making a definitive diagnosis (option A), and a venous Doppler is the best modality for confirming DVT. It reveals a non-compressible dilated vein. The likelihood of pulmonary embolism can be predicted using modified Wells criteria.

Patients with DVT should be immediately started with subcutaneous low molecular weight heparin (LMWH). If contraindicated due to renal insufficiency or thrombocytopenia-related conditions, they should be started on fondaparinux (indirect factor Xa inhibitor) or bivalirudin (direct thrombin inhibitor). Anticoagulants should be administered for a minimum period of 3 months. Vena cava filters are used in patients with bleeding risks (like hemophilia).

Solution to Question 16:

Dilated tortuous veins of 4 mm indicate varicose veins, so the stage is C2.

Solution to Question 17:

Adenocarcinoma is the most common variant of rectal carcinoma.

Colorectal cancer is:

- The second most common malignancy globally
- The second most common cancer in women
- The third most common cancer in men

- The fourth most common cause of death due to cancer.

Symptoms of rectal carcinoma include:


- Painless bleeding - earliest and most common
- Alteration of bowel habits - change in frequency and consistency of the stool
- Tenesmus - unable to pass a motion even when there is the sensation of needing to evacuate the bowel. It is suggestive of a tumor of the lower half of the rectum
- Early morning bloody diarrhea (option C) - passage of blood and mucus in addition to feces. Pain and weight loss are usually late symptoms

Abdominal examination and digital rectal examination are used for the primary evaluation. Rigid sigmoidoscopy is performed to obtain a biopsy. A synchronous tumor (primary tumor with a different origin than the one currently investigated) is excluded by performing a colonoscopy.

Modified Duke's classification:

A	Growth limited up to the rectal mucosa (option D)
B	B1: Tumor invading muscularis propria, no lymph node invasion. B2: Tumors invading into or through serosa, no lymph node invasion.
C	C1: The tumour did not exceed the bowel wall, lymph node metastasis. C2: Tumour exceeded the intestinal wall and lymph node metastasis.
D	Distant metastasis

Modified Dukes' staging



©Marrow

Surgery is the mainstay of management for rectal carcinoma. Non-operative management is offered to only those patients who have a complete clinical response to neoadjuvant.

The type of surgery depends on multiple factors (size, location, age of the patient, etc). However, the general consensus is:

- Early cancer (T1 and a few T2) is locally excised via the transanal approach.
- Most rectal cancer is treated by anterior resection and colorectal/coloanal anastomosis.
- Low extensive tumors are treated by laparoscopic abdominoperineal resection (APR) and permanent colostomy.
- Hartmann's operation (option A) is preferred in old and frail patients due to poor anal sphincter and postoperative incontinence.

Solution to Question 18:

A gasless abdomen is an X-ray feature that can be seen in acute pancreatitis. Patients with acute pancreatitis can have significant vomiting and diarrhea that may produce a gasless abdomen.

Common causes of gasless abdomen include:

- High obstruction: gastric outlet obstruction, congenital atresia
- Small bowel obstruction
- Bowel ischemia
- Ascites
- Acute pancreatitis
- Gastroenteritis
- Large abdominal mass - due to displacement
- Post-surgery following total colectomy, esophagogastrectomy, gastrectomy, and low anterior resection
- A paucity of small bowel gas may be a normal finding in some adults

Note: In intussusception, an X-ray shows small bowel obstruction, a mass, and a paucity of gas in the right iliac fossa.

Solution to Question 19:

The most common gallstone chemistry seen is crystalline cholesterol monohydrate.

Types of gallstones:

- Cholesterol stones (75% of gallstones) have >70% cholesterol by weight and are composed of crystalline cholesterol monohydrate. The majority of cholesterol stones have a mixed composition with bile pigments and calcium, in addition to the cholesterol. They are the most common type of gall stones worldwide and in India.

- Pigment stones have $\leq 20\%$ cholesterol and are dark due to the presence of calcium bilirubinate as the major constituent.
- Black pigment stones (20% of gallstones) - are formed due to supersaturation of unconjugated bilirubin (seen in hereditary hemolytic anemia (like sickle cell and hereditary spherocytosis))
- Brown pigment stones (5% of gallstones) - formed secondary to infections (like E. coli, and Ascaris lumbricoides)

Note: Pure cholesterol stones are uncommon and account for $\leq 10\%$ of all stones.

Solution to Question 20:


Transplant of an organ/tissue from a son to a parent is an example of an allograft.

An allograft is transplant of cells/tissues/organs from a genetically non-identical donor of the same species.

Isografts are allografts in which cells/tissues/organs are transplanted from a donor to a genetically identical recipient. (monozygotic twins)

Solution to Question 21:

The image showing serpiginous ulcers in the otherwise normal mucosa of the distal esophagus is suggestive of cytomegaloviral esophagitis.



Solution to Question 22:

Pulse rate is NOT a part of revised trauma score.

The Revised Trauma Score is made up of 3 categories: Glasgow Coma Scale, systolic blood pressure, and respiratory rate.


The score range is 0–12.

A patient with an RTS score of 12 is labelled delayed, 11 is urgent, and 3–10 is immediate. Those who have an RTS below 3 are declared dead.

Surgery NEET 2020

Question 1:

Which of the following is true about the flap given in the image.



- a) Pedicled TRAM based on superior epigastric artery
- b) Free TRAM based on superior and inferior epigastric artery
- c) DIEP based on deep inferior epigastric artery
- d) Lower Abdominal Fat based inferior epigastric vessels

Question 2:

What is the most common site of gastrinoma in MEN 1 syndrome?

- a) Jejunum
- b) Ileum
- c) Duodenum
- d) Stomach

Question 3:

A 50-year-old male who is a chronic smoker presented to the hospital with intermittent claudication pain of both thigh and buttock region on walking about 500 m. Which of the following is a likely diagnosis?

- a) Arterial disease involving the superficial femoral artery
- b) Arterial disease with aortoiliac involvement
- c) Femoral venous insufficiency
- d) Arterial disease involving the profunda femoris artery

Question 4:

Which of the following statements is true about retrosternal goitre?

- a) Operated in all patients regardless of symptoms
- b) Sternal incision is always required to operate
- c) It receives blood supply from the thoracodorsal artery
- d) Most of the retrosternal goitres can be removed by a neck incision

Question 5:

While performing surgery on a patient who presented with clinical features suggestive of peritonitis, a diverticular perforation is seen along with fecal peritonitis. What is the Hinchey's stage?

- a) Stage 1
- b) Stage 2
- c) Stage 3
- d) Stage 4

Question 6:


An 80-year-old woman presented with complaints of fatigue. On imaging, she was found to have a scleroatrophic gall bladder with impacted stones in it. A large stone was also present in the common bile duct and the biliary tree is dilated. What is the next step in management?

- a) Immediately take up patient for cholecystectomy
- b) Endoscopic retrograde cholangiopancreatography
- c) Wait and watch

d) CT scan

Question 7:

A 35-year-old man presented with recurrent episodes of fever with chills. MRCP image is shown below. What is the most likely diagnosis?



- a) Primary sclerosing cholangitis
- b) Caroli disease
- c) Oriental cholangitis
- d) Primary biliary carcinoma

Question 8:

A 5-year-old child with Wilson's disease is being considered for liver transplantation. Which of the following is not a parameter in the selection criteria for transplantation?

- a) Age < 10 years
- b) PT > 100 s
- c) Bilirubin $> 300 \mu\text{mol/L}$
- d) Jaundice less than 7 days before development of encephalopathy

Question 9:

The most common neuroendocrine tumor of the pancreas is

- a) Gastrinoma
- b) Somatostatinoma
- c) Insulinoma
- d) VIPoma

Question 10:

ASEPSIS wound scoring takes all of the following into consideration except:

- a) Wound swab from the site
- b) Serous discharge
- c) Erythema
- d) Induration

Question 11:

A patient comes with long-standing non-pitting edema in the legs. What is the most probable cause?

- a) Lymphedema
- b) Arterial Disorders
- c) Venous Disorders
- d) Coronary heart disease

Question 12:


A 41-year-old female patient presented with a tortuous swelling over the left side of the scalp and face as shown in the image given below. On local examination, the swelling was pulsatile. On auscultation, a bruit was heard. Which of the following is the most likely diagnosis?



- a) Neurofibromatosis
- b) Liposarcoma
- c) Cirroid Aneurysm
- d) Varicocele

Question 13:

What does the given barium swallow film show?



- a) Aortic aneurysm
- b) Pharyngeal pouch

- c) Carotid body tumor
- d) Esophageal carcinoma

Question 14:

True statement about intermittent claudication:

- a) Felt at rest
- b) Most commonly felt in the calf muscles
- c) Day to day distance is variable
- d) Relieved after getting out of bed

Question 15:

Which of the following is not a part of the primary survey?

- a) Measuring blood pressure
- b) Removing clothes to inspect for wounds
- c) Taking CT scan of the patient
- d) Checking airway patency

Question 16:

A 50-year-old male patient, during his routine annual checkup, is found to have an abdominal aortic aneurysm measuring 40 mm in diameter on USG. Which of the following is true about the management of this patient?

- a) Urgent endovascular aneurysm repair
- b) Surgery is indicated if aneurysmal diameter is ≥ 55 mm
- c) Urgently wheel the patient to OT for surgery
- d) Surgery is indicated only if aneurysm is ≥ 77 mm

Question 17:

Which of the following is the Miami criteria for intraoperative parathyroid hormone (PTH) measurement?

- a) Quarter fall in 5 min

- b) Quarter fall in 10 min
- c) Half fall in 5 min
- d) Half fall in 10 min

Question 18:

A 15-year-old boy undergoes bilateral adrenalectomy for pheochromocytoma. On the third postoperative day, the boy feels lethargic and tired. Blood pressure is 80/60 mm Hg, there is no sign of bleed. What could be the cause of this?

- a) Addisonian crisis
- b) SIADH
- c) Diabetes insipidus
- d) Cerebral salt wasting disease

Question 19:

Which of the following is not a component of Thoracscore?

- a) Performance status
- b) Complication of surgery
- c) ASA grading
- d) Priority of surgery

Question 20:

A middle-aged man presented to the emergency department with upper abdominal pain after having a heavy meal. On clinical examination, upper abdominal tenderness was noted. On chest radiography widening of the mediastinum with pneumomediastinum was seen. What is the probable diagnosis?

- a) Spontaneous perforation of esophagus
- b) Perforation of peptic ulcer
- c) Foreign body in esophagus
- d) Rupture of emphysematous bulla of lung

Question 21:

A 32-year-old male was diagnosed with achalasia cardia. High-resolution manometry revealed massive, simultaneous, high-pressure contractions of the distal oesophagus. According to the Chicago classification, the patient has which type of achalasia?

- a) Type II
- b) Type I
- c) Type IV
- d) Type III

Question 22:

Identify the investigation and the lesion depicted in the image given below.



- a) Retrograde cystourethrogram with membranous urethra stricture
- b) Retrograde cystourethrogram with penile urethral stricture
- c) Micturating cystourethrography with bulbar stricture
- d) Micturating cystourethrography with prostatic urethra stricture

Question 23:

According to the TNM staging, a patient with lung cancer of size 2.5 cm is staged as _____

- a) T1b
- b) T1c
- c) T2a

d) T1a

Question 24:

Which of the following should be the treatment of choice for renal cell carcinoma with tumor less than 4 cm in size?

- a) Partial nephrectomy**
- b) Partial nephrectomy + neoadjuvant chemotherapy**
- c) Radical nephrectomy**
- d) Radical nephrectomy + postoperative radiotherapy**

Question 25:

A 38-year-old man was brought to the emergency department after suffering a stab injury to the lower chest. On presentation, he had bradycardia and hypotension which did not improve despite IV fluid administration. Chest radiography showed clear lung fields. What is the next step in management?

- a) eFAST**
- b) Tube thoracostomy**
- c) CECT chest**
- d) CECT abdomen**

Question 26:

A man was brought to the emergency department after he fell into a manhole and injured his perineum. He gets an urge to micturate but is unable to urinate. On examination, blood was seen at the tip of the urethra, and swelling of the penis and scrotum was seen. What is the site of injury?

- a) Membranous urethra rupture**
- b) Bulbar urethra injury**
- c) Prostatic urethra injury**
- d) Urinary bladder rupture**

Answer Key

Question No.	Correct Option
1	c
2	c
3	b
4	d
5	d
6	b
7	a
8	d
9	c
10	d
11	a
12	c
13	b
14	b
15	c
16	b
17	d
18	a
19	b
20	a
21	d
22	b
23	b
24	a
25	a
26	b

Detailed Explanations

Solution to Question 1:

The given image showing only skin and fat moved to the chest is suggestive of a deep inferior epigastric artery perforator flap (DIEP), based on the deep inferior epigastric artery.

DIEP is a variant of TRAM flap and free flap where only the skin and fat are moved to the chest. A microvascular technique is used in which no muscle is taken and the perforating vessels are dissected out.


As the muscle is spared in DIEP, there is a reduced risk of abdominal wall weakness and its associated complications.

Other options:

Options A and B: The pedicled TRAM flap is based on the superior epigastric artery and the free TRAM flap is based on superior and inferior epigastric arteries (supercharged TRAM flap); however, the given image depicts a deep inferior epigastric artery flap. In TRAM flaps:

- The skin, fat, and muscle are moved to the chest.
- It provides good cosmesis and bulk since it has a soft tissue consistency similar to the breast.
- Disadvantages include epigastric bulge, partial flap loss, fat necrosis, abdominal wall weakness, and hernias.

The given image below shows a TRAM flap:



Solution to Question 2:

The most common site of gastrinoma in MEN 1 syndrome is the duodenum.

Gastrinoma is the most common pancreatic islet tumor in MEN 1 syndrome, arising from the G cells of the pancreas. Gastrinoma, associated with marked gastric acid production and recurrent peptic ulcers, is called Zollinger-Ellison syndrome. The majority of gastrinomas occur in the gastrinoma (Passaro's) triangle.

In MEN 1, they often occur in patients <30 years. Patients present with abdominal pain (the most common symptom), diarrhea and steatorrhea.

Management:

- Medical management for tumors ≤ 2 cm with proton-pump inhibitors.
- Surgery for tumors > 2 cm.
- Proton-pump inhibitors with somatostatin analogues for advanced locoregional disease or metastasis.

Solution to Question 3:

The given clinical scenario is suggestive of arterial disease with aortoiliac involvement. Classically, the muscle group affected by claudication is one anatomical level below the level of arterial disease.

Intermittent claudication is a feature of chronic arterial disease or occlusion of the artery. Aortoiliac disease affects the infrarenal aorta and iliac arteries.

Management involves smoking cessation, lifestyle modifications, and treatment of underlying diabetes or hyperlipidemia if present.

For claudication, cilostazol or pentoxifylline is used. Cilostazol is superior to pentoxifylline.

Surgical options include aortoiliac bypass, aortobifemoral bypass, percutaneous transluminal angioplasty (PTA), which can be with or without stent placement, and thromboendarterectomy.

Correlation of clinical findings to the Site of Disease	
Aortoiliac obstruction	Claudication in buttocks, thighs, and calves Bruit over the aortoiliac region Impotence (Leriche's syndrome)
Iliac obstruction	Claudication of thigh and calf, unilaterally Absence of femoral and distal pulses Bruit over iliac region
Femoropopliteal obstruction	Claudication in calf Femoral pulse felt but distal pulses absent
Distal obstruction	Ankle pulses absent Claudication in calf and foot Femoral and popliteal pulses palpable

Solution to Question 4:

Most of the retrosternal goiters can be removed by neck incision (transcervical approach).

Retrosternal goitre is defined as having $> 50\%$ of the goiter below the suprasternal notch. It can be either primary or secondary.

- Primary: rare and arises from ectopic thyroid tissue in the mediastinum; receives blood supply from intrathoracic blood vessels. It is not related to thyroid tissue in the neck.
- Secondary: they are the most common type; an extension of enlarged thyroid from the neck, usually arising from the lower pole of nodular goitre. It receives its blood supply from superior and inferior thyroid vessels.

The majority of patients are asymptomatic or have minimal symptoms. If the gland enlarges within the thoracic inlet, the pressure may lead to dysphagia, tracheal compression, and eventually airway symptoms. Pemberton sign can be positive (facial flushing and dilatation of cervical veins upon raising the arms above the head).

CECT neck and thorax is the investigation of choice.

Over 95% of retrosternal goiters can be removed via a transcervical incision since they are an extension from thyroid tissue in the neck.

Other options:

Option A: Surgical removal of the retrosternal goiter is indicated in patients with significant airway compression, and in young patients who have symptoms such as dyspnoea or dysphagia.

Option B: Median sternotomy approach is indicated in:

- Primary mediastinal goiter
- Malignant retrosternal goitre
- Very large retrosternal goiter
- Recurrence in the mediastinum

For a primary mediastinal goiter whose boundaries are completely within the chest, resection is often achieved through median sternotomy or thoracotomy to ensure that the vessels supplying the tumor are properly ligated. Removing the tumor through a cervical approach can result in uncontrolled bleeding and is generally avoided.

Option C: Blood supply to the retrosternal goiter is from the thyroid vessels (secondary retrosternal goiter).

Solution to Question 5:

Fecal peritonitis belongs to grade 4 in the Hinchey classification.

Diverticulitis refers to the inflammation and infection associated with the diverticula. Hinchey classification is the most commonly used classification for the severity and complication of acute diverticulitis. It is used additionally to guide the overall management of acute diverticulitis.

Hinchey classification:	
Stage 1	Mesenteric or pericolic abscess
Stage 2	Pelvic abscess
Stage 3	Purulent peritonitis

Hinchey classification:	
Stage 4	Fecal peritonitis

Solution to Question 6:

An 80-year-old woman with imaging showing a scleroatrophic gall bladder with impacted stones and a stone in the bile duct is suggestive of chronic cholelithiasis with choledocholithiasis. ERCP is the next line of management for this patient.

Choledocholithiasis is the presence of stone in the common bile duct (CBD) or biliary tree. It can be primary or secondary:

- Primary stones are formed in the CBD or biliary tree itself and are rare.
- Secondary stones are passed from the gallbladder to the CBD. They are easier to manage than primary stones.

It can be an asymptomatic and incidental finding or present with features such as jaundice (most common), abdominal pain, and fatigue. When there is inflammation of the biliary tree (cholangitis), the patient will have intermittent pain, fever, and jaundice (Charcot's triad).

Investigations:

- Liver function tests will show elevated direct bilirubin and alkaline phosphatase levels.
- Abdominal ultrasound is the first imaging modality done and may show gallstones and dilated bile duct (>8 mm).
- CT is sensitive for CBD stones. It shows stones, location, ductal stricture or block, ductal dilatation, and intrahepatic biliary changes.
- MRCP is highly sensitive and specific for CBD stones, but it is only diagnostic and intervention cannot be performed.

Management:

- ERCP with sphincterotomy is the procedure of choice, followed by the removal of stones using a Dormia basket or placement of a stent.
- Despite ERCP being an invasive procedure, it has the advantage of being both a diagnostic and therapeutic procedure and is more suited for elderly patients who cannot tolerate anesthesia and surgery.

Other options:

Option A: Cholecystectomy is not recommended for elderly patients, owing to the risk of old age and associated anesthetic and surgical complications.

Option C: Wait and watch approach can be done for asymptomatic patients, and patients with smaller stones.

Option D: CT is sensitive for CBD stones, but it is only diagnostic and intervention cannot be performed.

Solution to Question 7:

The presence of fever and chills, along with the MRCP showing multiple areas of intrahepatic beading is typical of primary sclerosing cholangitis.

Option B: Caroli's disease is a rare congenital condition characterized by irregular dilatation of the intrahepatic bile ducts with a normal extrahepatic duct system. It is common in both sexes and mostly presents before the age of 30 years with abdominal pain, cholangitis, or end-stage liver disease. MRCP shows irregular, cystic dilation of the large proximal intrahepatic bile ducts with a normal common bile duct.

Primary sclerosing cholangitis (PSC) is an autoimmune condition characterized by inflammatory strictures involving the intra and extra-hepatic biliary tree. It is a progressive condition that leads to secondary biliary cirrhosis.

MRCP or ERCP will show demonstrate multifocal, diffuse dilatation and strictures of the intrahepatic and extrahepatic biliary trees, this pattern is a beading or chain of lakes appearance. Pseudodiverticula of extrahepatic bile ducts are pathognomic of PSC. MRCP is useful in diagnosis and surveillance, while ERCP is reserved for the treatment of dominant strictures and to exclude malignancy.

The below image shows a cholangiogram of primary sclerosing cholangitis:



The common age group of primary sclerosing cholangitis is 30-45 years. Men are affected twice as commonly as women. Two-thirds of these cases are associated with ulcerative colitis. It is also associated with Riedel's thyroiditis and retroperitoneal fibrosis. 10–20% of patients develop cholangiocarcinoma.

Clinical features of a patient with PSC include pruritis, fever, chills, right upper quadrant discomfort, jaundice, and weight loss.

Liver function tests show a cholestatic pattern with an elevation of ALP and GGT and smaller rises in the aminotransferases. Bilirubin values may be variable.

Liver biopsy excludes other causes of cholestatic liver disease and defines the stage of PSC. It shows onion skin concentric periductal fibrosis. With disease progression, periportal fibrosis

occurs, progressing to bridging necrosis, and eventually biliary cirrhosis.

Currently, there is no known effective medical therapy. Surgical management involving resection of extrahepatic biliary tree and hepaticojejunostomy has shown reasonable effects. For patients with advanced liver disease, liver transplantation is the only available option.

Other options:

Option C: Oriental cholangitis, also known as recurrent pyogenic cholangitis is characterized by recurrent attacks of cholangitis due to bile stasis and stone formation. Patients between 30-40 years and both sexes are equally affected. It is caused due to infections (liver flukes, E.coli, Klebsiella, Pseudomonas, and Proteus). Patients present with Charcot's triad (intermittent fever, pain, and jaundice). MRCP shows intra or extrahepatic calculi, dilatation of ducts, and strictures.

Option D: Biliary carcinoma (cholangiocarcinoma) most commonly affects patients >60 years and presents with fever, abdominal pain, weight loss, and features of obstructive jaundice. MRCP shows irregular narrowing of the bile duct involved by the tumor, with asymmetric upstream dilatation of the intrahepatic ducts.

Solution to Question 8:

A 5-year-old child with Wilson's disease is a case of non-paracetamol-induced acute liver failure and jaundice for more than 7 days before the development of encephalopathy is one of the parameters in the King's College criteria for liver transplantation.

It includes the following parameters:

Non-paracetamol-induced acute liver failure:

- PT >100 s (INR >6.5)
- or/and
- Any three of the following:
 - Age <10 years or >40 years
 - Etiology non-A, non-B hepatitis, or idiosyncratic drug reaction
 - Jaundice more than 7 days before the development of encephalopathy i.e., not hyperacute
- PT >50 s (INR > 3.5)
- Bilirubin > 18 mg/dl (300 µmol/L)

Paracetamol-induced acute liver failure:

- pH <7.30 (irrespective of the grade of encephalopathy)
- or/and
- All three of the following:
 - Prothrombin time >100 s (INR >6.5)
 - Serum creatinine >3.4 mg/dl (300 µmol/L)
 - Grade 3 or 4 hepatic encephalopathy

Solution to Question 9:

The most common neuroendocrine tumor of the pancreas is insulinoma (4 per million).

Insulinoma is an insulin-producing tumor of the pancreas, with more than 90% of them being benign. It occurs in >40 years in patients without MEN 1 syndrome and <40 years in patients with MEN 1. It presents with symptoms of hypoglycemia such as weakness, headaches, sweating, faintness, seizures, and altered behavior. These occur after fasting or exertion and improve after glucose intake.

The investigation of choice for insulinoma is the determination of 72-hour fasting glucose and serum insulin levels. C-peptide and proinsulin levels are also elevated.

Medical management includes:

- Frequent carbohydrate-rich meals
- Somatostatin analogs like octreotide are effective in controlling symptoms due to excess insulin secretion. They should be used with caution as they can suppress counterregulatory hormones, such as growth hormone (GH), glucagon, and catecholamines, and precipitously worsen hypoglycemia.
- Diazoxide can be used in the initial management of insulinoma as it inhibits insulin release. Its side effects include sodium retention and nausea.
- Everolimus has been found to be highly effective in improving glycemic control in patients with insulinoma owing to its antitumor effect and induction of insulin resistance.

Enucleation and pancreatectomy are the definitive surgical management of insulinoma.

Other options:

Option A: Gastrinoma has an incidence of 0.5 to 4 in a million.

Options B & D: Somatostatinoma and VIPoma are much rarer than gastrinoma.

Solution to Question 10:

ASEPSIS wound score does not take induration into consideration. Instead, the I in ASEPSIS stands for the isolation of bacteria from the wound.

ASEPSIS wound score and Southampton wound score are used to determine the severity of the surgical site wound infection (SSI).

ASEPSIS score includes:

- Additional treatment
- Antibiotics for wound infection
- Drainage of pus under local anesthesia
- Debridement of the wound under general anesthesia

- Serous discharge
- Erythema
- Purulent exudate
- Separation of deep tissue
- Isolation of bacteria from the wound
- Stay as inpatient prolonged over 14 days as a result of wound infection

Solution to Question 11:

Lymphedema is the most probable cause of long-standing non-pitting edema in the limbs.

Lymphedema is the abnormal accumulation of protein-rich fluid in the interstitial spaces of the tissues.

Lymphedema can be primary or secondary.

Primary lymphedema includes:

- Lymphedema congenita (0-2 years): more common in males, involves multiple limbs, face, and genitalia. It can be familial or sporadic. The familial type is called Nonne Milroy syndrome.
- Lymphedema praecox (2-35 years): more common in females, the most common type of primary lymphedema with a peak incidence in puberty. It can be familial or sporadic. The familial type is called Meig's disease.
- Lymphedema tarda (>35 years): Rare and uncommon.

Secondary lymphedema is more common than primary and is the most common cause of lymphedema overall. Causes include lymph node dissection (axillary or inguinal), post-mastectomy, postoperative radiotherapy, obesity, varicose vein surgery, and trauma.

Complications of lymphedema include infections, skin changes, recurrent cellulitis, non-healing ulcers, and lymphangiosarcoma.

Treatment mostly includes patient education and nonsurgical interventions, which include limb elevation, and external compressive garments. When the above methods fail, surgery can be considered.

Solution to Question 12:

The presence of a tortuous, pulsatile swelling over the scalp and face is suggestive of a cirroid aneurysm.

Cirroid aneurysm is a slow-growing, tortuous swelling that develops due to abnormal arteriovenous connections among the subcutaneous blood vessels. They are usually congenital. However, in some cases, they may be traumatic. The superficial temporal artery is most commonly involved.



Patients present with a tortuous, pulsatile swelling over the scalp or face. They are usually asymptomatic and present to the clinic due to cosmetic deformity. However, some patients may have headache, tinnitus, and/or bleeding. A loud bruit is heard on auscultation.

Angiography is the investigation of choice.

Treatment includes surgical resection, endovascular occlusion or injection of percutaneous sclerosing agents.

Solution to Question 13:

The given barium swallow film shows an outpouching from the posterior wall of the pharynx suggestive of a Zenker's diverticulum (pharyngeal pouch).

Solution to Question 14:

Intermittent claudication is most commonly felt in the calf muscles because of the involvement of the superficial femoral artery in chronic vascular occlusive disease.

Intermittent claudication:

- Intermittent claudication is the debilitating cramp-like pain felt in the leg muscles due to reduced blood flow and anaerobic metabolism brought on by peripheral arterial disease.
- It usually signifies mild to moderate chronic vascular occlusive disease
- The pain is brought on by walking (option D)
- Not present on taking the first step
- Relieved by rest in both the standing and sitting positions, usually within 5 minutes (option A).

- The distance that a patient is able to walk without stopping is known as claudication distance. It varies only slightly from day to day (option C).
- The superficial femoral artery (70%) is the most commonly affected artery by claudication and manifests as pain in the calf muscles.
- Aorto-iliac occlusion may cause thigh or buttock claudication. Leriche's syndrome occurs due to aortoiliac occlusion and presents with a triad of claudication, impotence, and absent femoral pulses.

Boyd's classification of claudication:

- Grade I: Patient complains of pain after walking. If the patient continues to walk, due to increased blood flow in muscle and opening of collaterals metabolites causing pain are washed away and the pain subsides.
- Grade II: Pain still persists on continuing to walk; but can walk with effort.
- Grade III: The patient has to take rest to relieve the pain.

Solution to Question 15:

Taking a CT scan of the patient is a part of secondary survey.

Primary survey identifies and manages the most immediately life-threatening pathologies first.

The secondary survey includes investigations that help identify all other injuries, excluding the apparent ones. This includes X-rays of the spine and extremities and a CT of the abdomen, head, chest, and spine.

Solution to Question 16:

Surgery is indicated in an asymptomatic abdominal aortic aneurysm (AAA) only if its diameter is more than 55 mm in a male patient.

If the diameter is less than 55 mm, regular ultrasound monitoring is required. This is because elective open surgery has a higher rate of mortality when compared to conservative management for aneurysms of size ≤ 55 mm.

Indications for surgery in abdominal aortic aneurysm:

- Size ≥ 5.5 cm in males
- Size ≥ 5 cm in females
- Expansion of ≥ 1 cm in one year
- Painful tender aneurysm
- Thrombosed aneurysm; aneurysm with distal emboli


Management of aortic aneurysm: AAA is surgically managed by dacron graft repair. It can be either open or EVAR (endovascular aneurysm repair)

- Open surgical repair is done by exposing the aorta through a mid-sternal incision and graft replacement. To expose the aorta the descending colon should be mobilized medially and this is known as left visceral rotation or Mattox procedure.
- EVAR is a minimally invasive procedure in which an aortic stent graft is placed on the nonaneurysmal aortoiliac segment.

Complications following an AAA repair are:

- Cardiovascular (ischemia and infarction)
- Respiratory (atelectasis or consolidation).
- Colonic ischemia.
- Renal failure.
- Aortoduodenal fistula
- spinal cord ischemia

The given image below shows the technique of EVAR:



Solution to Question 17:

A drop in the level of PTH to less than half the maximum value (corresponding to the preoperative levels) at 10 min postresection is the Miami criteria.

It is used to determine the extent of resection of the parathyroid gland which is done in a case of primary hyperparathyroidism due to adenoma.

Intraoperative parathyroid hormone monitoring is a commonly used adjunct, which guides the surgeon on when to stop parathyroid exploration and helps in confirming there is no residual hyperfunctioning parathyroid tissue. The PTH levels are expected to drop after the gland is excised due to its short half-life (3-5 minutes). Blood is drawn at various time points and the hormone levels are assessed.

Based on the Miami Criteria, a more than 50% decline in PTH at 10 minutes following parathyroid gland removal compared to the highest baseline level, either the pre-incision or pre-excision, suggests a cure, and the procedure can be terminated.

Indications for parathyroidectomy:

All symptomatic patients with hyperparathyroidism should be offered surgery. Indications for surgery in asymptomatic patients include:

*BMD - bone mineral density

Serum calcium levels	0.25 mmol/L (1.0 mg/dL) above the upper limit serum calcium of normal
Skeletal	BMD by DEXA: T score $\leq$ -2.5 at the lumbar spine, total hip, femoral neck, or distal one-third of radius Vertebral fracture
Renal	Creatinine clearance $<$ 60 mL/min 24-hour urinary calcium $>$ 10 mmol/dL ($>$ 400 mg/day) Increased risk of stone formation by risk analysis
Age	$<$ 50 years

Solution to Question 18:

The given clinical scenario of hypotension with no presence of bleeding and bilateral adrenalectomy is suggestive of an Addisonian crisis. It is a life-threatening condition caused due to sudden insufficiency of adrenal hormones.

Acute adrenal insufficiency (adrenal or Addison's crisis) is a medical emergency caused due to a precipitating illness unmasking insufficiency or trauma or severe sepsis causing adrenal hemorrhage (Waterhouse-Friderichsen syndrome) or surgical removal of the glands. It presents with non-specific symptoms such as shock with fever, nausea, vomiting, abdominal pain, hypocalcemia and electrolyte imbalances. It may be often misdiagnosed as an acute abdominal emergency.

Adrenal insufficiency can be primary (due to adrenal pathology), secondary (due to pituitary pathology- lack of ACTH) or tertiary (due to hypothalamus pathology- lack of CRH).

Autoimmune adrenalitis is the most common cause of primary adrenal insufficiency worldwide.

Tuberculosis is the most common cause of primary adrenal insufficiency in India.

Adrenal crisis requires immediate intervention owing to its high mortality rate. Immediate rehydration is initiated with a saline infusion at 1 litre/hr with continuous cardiac monitoring. Glucocorticoid replacement is initiated with an initial bolus injection of hydrocortisone 100 mg followed by 200 mg over 24 hours, preferably by continuous infusion or alternatively by IV or IM

injections. Mineralocorticoid replacement is initiated once the daily dose of hydrocortisone has been reduced to <50 mg, since hydrocortisone at higher doses provides sufficient stimulation of mineralocorticoids.

Solution to Question 19:

The complication of surgery is not a component of thoracscore.

Thoracscore is used to assess the risk of operative mortality in a patient undergoing thoracic surgery. It includes 9 variables:

- Age
- Sex
- American Society of Anesthesiologists score (ASA)
- Performance status
- Dyspnoea score
- Priority of surgery
- Extent of surgery
- Malignant diagnosis
- Composite comorbidity score

Solution to Question 20:

The clinical scenario points to a diagnosis of spontaneous perforation of the esophagus.

In a perforated peptic ulcer, patients develop sudden onset severe generalized abdominal pain, tenderness, and board-like rigidity and an erect plain chest X-ray shows gas under the diaphragm (Option B). The foreign body in the esophagus can be asymptomatic or may present with perforation after ingestion. The radiopaque foreign body can be seen with a chest X-ray (Option C). Rupture of emphysematous bullae of the lung will lead to secondary pneumothorax, which manifests as chest pain and dyspnoea. A plain chest X-ray shows air in the thorax, and the mediastinum is pushed to the opposite side (Option D).

Spontaneous perforation of the esophagus needs immediate diagnosis and intervention due to the higher mortality associated with it. The most common site of perforation is the lower esophagus towards the left esophageal cavity.

Causes:

- Iatrogenic perforation (most common)
- Boerhaave's syndrome (spontaneous emetogenic perforation)
- Sharp foreign body
- Corrosive ingestion

- Esophageal carcinoma

It usually manifests as severe pain in the chest or abdomen after a meal or after a bout of drinking, commonly associated with dyspnoea. On clinical examination, there is rigidity and tenderness in the upper abdomen. The diagnosis is usually confirmed by a chest radiograph which shows pneumomediastinum, pneumoperitoneum, and pneumothorax. Subcutaneous emphysema can be seen in Boerhaave's syndrome. Patients with sepsis will have a fever, tachycardia and tachypnea.

Treatment includes adequate drainage, antibiotics, closure of the perforation site if possible and supportive measures.

Solution to Question 21:

The given features are suggestive of type III achalasia cardia.

Type III achalasia cardia is characterised by massive, simultaneous, high-pressure contractions of the distal esophagus.

Achalasia cardia occurs due to the loss of inhibitory ganglion cells in the myenteric plexus of GIT. During swallowing, the food bolus normally triggers a primary peristaltic wave, which is responsible for carrying the bolus to the stomach, accompanied by the relaxation of the lower esophageal sphincter (LES) to allow esophageal emptying into the stomach.

Achalasia is characterized by a disruption of this normal process. Failure of relaxation of the LES causes an elevated median integrated relaxation pressure and a complete failure of peristalsis. The esophagus becomes progressively more tortuous, leading to a megaesophagus.


Solution to Question 22:

The given image shows retrograde cystourethrogram with findings consistent with penile urethral stricture.

The male urethra can be visualized by retrograde urethrography (RGU) or voiding cystourethrogram (VCU), also known as micturating cystourethrogram (MCU).

In RGU, the urethra is visualized with the instillation of contrast into the anterior urethra (penile urethra and bulbar urethra). It is helpful for diagnosing anterior urethral abnormalities.

The following image shows an RGU with bulbar urethral stricture:



In VCU, after injecting contrast into the bladder using a Foleys catheter, the posterior urethra (membranous urethra and prostatic urethra) is visualized during micturition. It is helpful for diagnosing posterior urethral abnormalities and bladder abnormalities. It is the investigation of choice for vesicoureteric reflux (VUR) and posterior urethral valve.



Solution to Question 23:

According to TNM staging, a patient with lung carcinoma of size 2.5cm is staged as T1c. Stage T1-3/No-1/Mo are amenable to resection.

TNM staging system of Lung carcinoma	
T: Primary tumor	
Tx	Primary tumor cannot be assessed or tumor proven by the presence of malignant cells in sputum or bronchial washings but not visualized by imaging or bronchoscopy
To	No evidence of primary tumor
Tis	Carcinoma in situ
T1	Tumour ≤ 3 cm in the greatest dimension surrounded by lung or visceral pleura without bronchoscopic evidence of invasion more proximal than the lobar bronchus (i.e. not in the main bronchus)
T1a(mi)	Minimally invasive adenocarcinoma
T1a	Tumour ≤ 1 cm in greatest dimension
T1b	Tumor > 1 cm but ≤ 2 cm in greatest dimension
T1c	Tumor > 2 cm but ≤ 3 cm in the greatest dimension
T2	Tumor > 3 cm but ≤ 5 cm or tumor with any of the following features:– involves the main bronchus regardless of distance from the carina but without the involvement of the carina – invades visceral pleura – associated with atelectasis or obstructive pneumonitis that extends to the hilar region, involving part or all of the lung
T2a	Tumor > 3 cm but ≤ 4 cm in greatest dimension
T2b	Tumor > 4 cm but ≤ 5 cm in greatest dimension

TNM staging system of Lung carcinoma	
T3	Tumor>5 cm but ≤7 cm in greatest dimension or associated with separate tumor nodule(s) in the same lobe as the primary tumor or directly invades any of the following structures: chest wall (including the parietal pleura and superior sulcus tumors), phrenic nerve, the parietal pericardium
T4	Tumor>7 cm in greatest dimension or associated with separate tumor nodule(s) in a different ipsilateral lobe than that of the primary tumor or invades any of the following structures: diaphragm, mediastinum, heart, great vessels, trachea, recurrent laryngeal nerve, esophagus, vertebral body, and carina
N: Lymph node involvement	
Nx	Regional lymph nodes cannot be assessed
No	No regional lymph node metastasis
N1	Metastasis in ipsilateral peribronchial and/or ipsilateral hilar lymph nodes and intrapulmonary nodes, including involvement by direct extension
N2	Metastasis in ipsilateral mediastinal and/or subcarinal lymph node(s)
N3	Metastasis in contralateral mediastinal, contralateral hilar, ipsilateral or contralateral scalene, or supraclavicular lymph node(s)
M: Distant metastasis	
Mo	No distant metastasis
M1	Distant metastasis present
M1a	Separate tumor nodule(s) in a contralateral lobe; tumor with pleural or pericardial nodule(s) or malignant pleural or pericardial effusion

TNM staging system of Lung carcinoma	
M1b	Single extrathoracic metastasis
M1c	Multiple extrathoracic metastases in one or more organs

Solution to Question 24:

Partial nephrectomy (also known as nephron-sparing surgery) is the treatment of choice for tumors ≤ 4 cm.

Indications for partial nephrectomy:

- Tumors ≤ 4 cm
- Well-selected tumors between 4-7 cm
- Bilateral tumors
- Tumor in solitary kidneys
- Patients with pre-existing renal dysfunction.

Recently, partial nephrectomy is also increasingly being performed for stage T1b tumors.

Treatment of renal cell carcinoma:

RCC is a radio and chemoresistant tumor.

Radical nephrectomy consists of the removal of the kidney, perirenal fat, adrenal gland, and regional lymph nodes. The adrenal gland is no longer removed except in cases of obvious tumor involvement. Lymphadenectomy is indicated only in high-risk patients with large primary tumors and enlarged lymph nodes.

Cryoablation is indicated in the T1a stage of RCC if surgery cannot be performed. It is also indicated in advanced/metastatic tumors.

Note: Sunitinib and sorafenib are also used in GIST and hepatocellular carcinoma.

Staging	Description	Management
Stage 1	Tumor ≤ 7 cm, confined to the kidney	Partial nephrectomy
Stage 2	Tumor > 7 cm, confined to the kidney	Radical nephrectomy
Stage 3	Tumor extends into the major veins or adrenal gland with intact Gerota's fascia or regional lymph nodes involved.	Radical nephrectomy

Staging	Description	Management
Stage 4	Tumor invades beyond Gerota fascia (including contiguous extension into the ipsilateral adrenal gland)	Debulking surgery mTOR inhibitors such as everolimus, sirolimus Tyrosine kinase inhibitors: sunitinib, sorafenib

Solution to Question 25:

The given clinical scenario of a patient with a stab injury and hypotension not responding to IV fluids indicates persistent hypotension due to ongoing hemorrhage and the next step in the management is to perform an eFAST.

Cardiogenic shock or hemorrhagic shock are the most likely causes of persistent hypotension in a trauma patient. Ultrasound evaluation using eFAST to look for pericardium, pleural cavities, and abdomen in combination with plain radiographs of the chest and pelvis will usually identify the source of the bleed.

In patients with persistent hypotension due to blunt trauma, the pelvis should be wrapped in a sheet for stabilization until radiography can be done.

Evaluation of CVP or ultrasound of the IVC may further assist in distinguishing between cardiogenic and hypovolemic shock.

It is mostly fatal if not controlled, and requires immediate management. Blood and plasma transfusions are provided in a ratio of 2:1. Patients with persistent hypotension requiring surgical intervention include:

- Blunt trauma and massive hemothorax
- Penetrating trauma
- Hemoperitoneum
- Gunshot injuries to the chest or abdomen
- Penetrating weapon remaining in place.

Other options:

Options C and D: CECT can be done only when the patient is hemodynamically stable.

Option B: Tube thoracostomy has no role in this patient's management since the X-ray shows clear lung fields.

Solution to Question 26:

The given clinical scenario with a history of a blow to the perineum, blood at the tip of the meatus, swelling of the penis and scrotum is suggestive of an injury to the bulbar urethra.

Urethral injuries can be classified into:

Note: Urinary bladder rupture occurs due to an injury to the suprapubic region. Extraperitoneal bladder rupture is more common and associated with pelvic fracture. Intraperitoneal bladder rupture is caused by a blow or a fall on a distended bladder. It manifests as sudden severe pain in the hypogastrium and syncope.

Type	Proximal urethral injury	Distal urethral injury
Site of injury	Prostatic urethra Membranous urethra	Bulbar urethra Penile urethra
Mechanism	Secondary to pelvic fracture	Occurs due to a blow to the perineum by: Straddle injury Loose manhole cover Gymnasium accidents
Clinical features	Urinary retention Blood at the urethral meatus High-riding prostate on digital rectal examination (Vermooten sign) Deep perineal hematoma	Urinary retention Blood at the urethral meatus Perineal hematoma (butterfly hematoma)

Surgery INI-CET 2021

Question 1:

What is triage used for?

- a) To prepare for a disaster
- b) To assess the impact of a disaster
- c) To classify priority of treatment
- d) To rehabilitate following a disaster

Question 2:

Match the following

- a) 1-C,2-D,3-B,4-A
- b) 1-A,2-D,3-B,4-C
- c) 1-C,2-D,3-A, 4-B
- d) 1-D,2-C,3-A,4-B

Question 3:

A mother notices a swelling in the abdomen of her 3-year-old child while bathing him. He had a history of hematuria two weeks back which resolved spontaneously. On examination, a right-sided reniform ballotable mass was found. What all investigations will you do?

- a) 1 and 2
- b) 2 and 4
- c) 1, 2, and 3
- d) 1, 2, 3 and 4

Question 4:

A 26-year-old male presented with a large, round, firm, painless, right-sided testicular swelling associated with a dragging sensation. The left testis was normal. AFP was around 3000U. What is the next best step in the management?

- a) FNAC
- b) Scrotal trucut biopsy
- c) High inguinal orchiectomy
- d) Repeat tumor markers

Question 5:

A woman who had undergone cholecystectomy presented three days after the surgery with abdominal pain and tachycardia. WBC count is 11000 cells/mm³. USG showed fluid collection 5x5 cm in right sub-hepatic space. What is the next step in management?

- a) Antibiotics and send home
- b) USG guided pigtail catheter drainage
- c) Laparoscopy and re-exploration
- d) ERCP and T-tube placement

Question 6:

The instrument shown in the image is used to insufflate the abdomen for laparoscopic surgery. What is its name?



- a) Veress needle
- b) Lind's needle
- c) Jacob needle

d) Mathew needle

Question 7:

A 20-year-old was brought to the ER after falling off his bicycle. His vitals were stable, but he had a bruise over the perineum and he had not passed urine yet. Blood was seen at the meatus and the bladder was not palpable. How do you manage him?

- a) Wait and send him home**
- b) Immediate insertion of foleys catheter**
- c) Suprapubic aspiration of urine**
- d) Wait for bladder to fill and urge to urinate**

Question 8:

A patient who was admitted after a road traffic accident is put on mechanical ventilation. He opens his eyes on verbal command and moves all four limbs spontaneously. Calculate his GCS.

- a) Eyes-2, Verbal-1, Motor-5**
- b) Eyes-2, Verbal-NT, Motor-5**
- c) Eyes-3, Verbal-1, Motor-6**
- d) Eyes-3, Verbal-NT, Motor-6**

Question 9:

What is the gauge size of the cannula used here?



- a) 14
- b) 16
- c) 18
- d) 20

Question 10:

A 55-year-old female patient presented with a 4x3 cm lump in the right upper outer quadrant, with no axillary lymph node involvement. Mammography revealed BIRADS 4b staging. She underwent breast conservation surgery, and the final HPE report showed high nuclear grade DCIS with necrosis and 10 mm margin clearance. What is the further management?

- a) Adjuvant radiotherapy
- b) Adjuvant chemotherapy
- c) Follow up 6 monthly for 2 years and then yearly follow up
- d) Trastuzumab therapy

Question 11:

A lady presents with abdominal pain. USG reveals multiple gall stones with no wall thickening. CBD diameter is 12 mm and serum bilirubin was 0.8. Alkaline phosphatase is 380 and GGT is raised by 5 times. What is the next best step of management?

- a) MRCP
- b) ERCP

- c) Endoscopic ultrasound
- d) Laparoscopic cholecystectomy


Question 12:

You are suturing a laceration in the ER using the interrupted suturing technique. What is the angle of needle placement?

- a) 60 degrees
- b) 70 degrees
- c) 80 degrees
- d) 90 degrees

Question 13:

A 26-year-old male patient was brought to the emergency department with abdominal pain and obstipation for 3 days. He gives a history of bull gore to the abdomen 3 days back. His X-ray is given below. What is the probable diagnosis?



- a) Hemothorax
- b) Hollow viscus perforation
- c) Pneumothorax
- d) Intestinal obstruction


Question 14:

A man presented with a history of stab injury over the arm. On examination, there was a hematoma present, and radial and ulnar pulses were not felt. A 3.5cm longitudinal tear is noted in the brachial artery. Which among the following is the best treatment option?

- a) Primary repair with end to end anastomosis
- b) Repair with great saphenous vein graft
- c) Repair with prosthetic graft
- d) Repair with saphenous vein patch

Question 15:

A 24-year old male patient was brought to the emergency department following a road traffic accident. His heart rate was 100/min, respiratory rate was 24/min, and his chest X-ray is given below. Which of the following is contraindicated in him?



- a) Nasogastric tube insertion
- b) Chest tube insertion
- c) Log roll
- d) Epidural anaesthesia

Question 16:

A patient was received in the casualty following a road traffic accident. His BP was 90/60 mmHg and his pulse rate was 120/min. The emergency room physician asked you to replace the fluid loss in this patient. What is the maximum flow rate of fluid with the IV canula given in the image?



- a) 86ml/ min
- b) 96ml/ min
- c) 103 ml/ min
- d) 63 ml/ min

Question 17:

The most common cause of bladder outlet obstruction in a male child is

- a) Posterior urethral valve
- b) Urethral atresia
- c) Anterior urethral valve
- d) Ureterocele

Question 18:

Which of the following statements are true about colon cancer?

- a) 1, 2, and 3 only
- b) 2, 3, and 4 only

- c) 2 and 4 only
- d) 1, 2, 3 and 4

Question 19:

Frozen section was done during the surgery of carcinoma breast. This procedure will be useful for all except

- a) To know sentinel lymph node
- b) To know margin positivity
- c) To know metastasis
- d) To confirm definitive diagnosis immediately in all cases


Question 20:

A 40-year-old woman presented with jaundice, fever, and abdominal pain. She is suspected to have cholangitis. Which of the following apart from abdominal pain makes up Reynold's pentad?

- a) 1,2,3,4
- b) 2,3,4,5
- c) 1,3,4,5
- d) 1,2,4,5

Question 21:

The following drain is placed after mastectomy. Identify the type of drain.



- a) Open
- b) Closed
- c) Semi open
- d) Semi closed

Question 22:

Consider the following about hepatocellular carcinoma (HCC) and choose the correct statements.

- a) 1, 2, 3
- b) 2, 3, 4
- c) 1, 3, 4
- d) 1 and 2 only

Question 23:

A 56-year-old patient with obstructive jaundice has a distended gall bladder and a dilated common bile duct and intrahepatic biliary radicles on ultrasound. Contrast-enhanced CT confirmed the same findings with no additional findings. Which of the following will be most useful to localize the cause?

- a) Magnetic resonance imaging
- b) Endoscopic ultrasound
- c) Percutaneous transhepatic cholangiography

- d) Endoscopic retrograde cholangiopancreatography

Question 24:

All of the following are lower urinary tract symptoms except

- a) Sudden urge to urinate
- b) Incontinence while asleep
- c) Incontinence when patient lies down and coughs
- d) Incontinence during intercourse

Question 25:

Fluorescence in-situ hybridization(FISH) is required in which of the following IHC interpretations of Her2/neu?

- a) 1+
- b) 2+
- c) 3+
- d) All of the above

Question 26:

A 50-year-old male patient came to the outpatient department with complaints of hematuria. A 2x2 cm bladder mass is seen which is a low-grade transitional cell carcinoma. Which among the following is the ideal management?

- a) Transurethral resection of the tumour
- b) Resection with ileal conduit
- c) Partial cystectomy with bladder reconstruction
- d) Neo-adjuvant chemotherapy

Question 27:

Which of the following statements are true?

- a) 2, 3, 4
- b) 1, 2, 3, 4

- c) 4 only
- d) 1, 3, 4

Question 28:

Which of the following is the correct method for measurement of nasogastric tube?

- a) Tip of nose to ear lobule to xiphisternum
- b) Angle of mouth to tragus to xiphisternum
- c) Ala of nose to angle of mandible to xiphisternum
- d) Angle of mouth to ear lobule to umbilicus

Question 29:

A surgeon performs an appendicectomy with a grid iron incision. He later converts the incision to Rutherford Morrison incision. Which of the following are cut?

- a) 1, 3, 4
- b) 2, 3, 4
- c) 1, 2, 3
- d) 1, 2, 4

Answer Key

Question No.	Correct Option
1	c
2	c
3	c
4	c
5	b
6	a
7	d
8	d
9	c
10	a

11	a
12	d
13	b
14	b
15	b
16	c
17	a
18	d
19	d
20	a
21	b
22	c
23	a
24	d
25	b
26	a
27	d
28	a
29	c

Detailed Explanations

Solution to Question 1:


Triage is used to classify the priority of treatment.

Triage is carried out at the site of a disaster to determine the priority for immediate medical attention, transportation, and hospital admission. This process involves the rapid classification of the injured based on the severity of their injuries and the likelihood of survival with timely medical intervention.

Its objective is to provide maximum benefit to the largest number of injured individuals in a disaster situation. Victims whose prognosis would be favorable with simple medical care are given higher priority, while patients in critical condition, who may not benefit significantly from medical attention, are given lower priority. An internationally accepted four-colour coded is commonly used, which includes:

- Red - High priority attention or transfer
- Yellow - Medium priority
- Green - Ambulatory patients
- Black - Dead or moribund patients

Assessment of impact, rehabilitation and preparedness following a disaster are included in other phases of the disaster management cycle (options A, B and D).



Solution to Question 2:

The correct option is 1-C,2-D,3-A, 4-B


- Hirschsprung's disease - Rectal biopsy
- Posterior urethral valve - Micturating cystourethrogram
- Choledochal cyst - MRCP
- Intussusception - CT scan

Hirschsprung's disease:

- Rectal biopsy is the gold standard investigation for the diagnosis of Hirschsprung's disease. The affected part shows the absence of ganglion cells, hypertrophy of nerve trunks, and positive immunostaining for acetylcholinesterase.
- Water-soluble contrast enema: It shows the presence of a contracted aganglionic rectum, transition zone, and a dilated normally innervated colon.
- Anorectal manometry: It shows the failure of the relaxation of the internal anal sphincter.

Surgical intervention is the only definitive treatment available, and includes the Swenson, Duhamel, and Soave procedures.

Contrast enema image in Hirschsprung's disease-



Posterior urethral valve (PUV):


- Diagnosis of this condition is commonly made during the antenatal period with an ultrasound, which demonstrates bilateral hydronephrosis above a distended bladder. Pulmonary hypoplasia secondary to oligohydramnios may be detected in some complicated cases. In cases that are not detected during the antenatal period, these children usually present with urinary infection in the neonatal period or with uremia and renal failure.
- The confirmatory investigation for PUV is a micturating/voiding cystourethrogram. It demonstrates poor bladder emptying and dilated posterior urethra.
- It is treated by cystoscopic resection or ablation of the valve.

Preoperative(a) and postoperative(b) micturating cystourethrogram images in a patient with PUV is shown below. The dilated posterior urethra seen in the first image is resolved in the second image after surgery.



Choledochal cyst:

- MRI/MRCP is highly useful to access the biliary anatomy especially the relation between the lower end of the bile duct and the pancreatic duct.
 - Ultrasound is helpful to identify the presence of an abnormal cyst.
 - CT scan can be used to delineate the extent of intrahepatic or extrahepatic dilatation
- MRCP image showing multiple intrahepatic biliary duct cystic dilatations-



Intussusception:


CT scan is the investigation of choice. It may show the characteristic target sign.

USG findings in intussusception include:

- Target sign (also known as the doughnut sign)
- Pseudokidney sign
- Crescent in a doughnut sign

Barium enema shows the characteristic coiled spring appearance.

USG image showing the target sign:



Solution to Question 3:

The clinical features of hematuria and a ballotable renal mass in a 3-year-old child are suggestive of Wilms tumor (Nephroblastoma). CECT abdomen is used to identify the extent of the tumor, integrity of the other kidney, and metastasis. Wilms tumor is metabolically active and can concentrate fluorodeoxyglucose therefore PET/CT is also helpful to identify regional spread and metastatic lesions. A biopsy is discouraged as it may lead to disease upstaging. 24-hour urinary vanillyl mandelic acid is measured to exclude neuroblastoma, which is an important differential diagnosis of Wilms tumor.

Wilms tumor is the most common primary malignant renal tumor in children, typically affecting children less than 5 years. After neuroblastoma, it is the second most common malignant abdominal tumor in childhood. It most commonly presents as an asymptomatic abdominal mass. Other findings include hypertension, abdominal pain, gross painless hematuria, and constitutional symptoms such as fever and weight loss.

Imaging studies done to diagnose Wilms tumor include plain radiography, USG, and CT of the abdomen to demonstrate the renal mass. USG with doppler study of renal veins is useful in the evaluation of collecting system and to look for tumor thrombi in renal veins and inferior vena cava. A chest CT is done preoperatively to screen for pulmonary metastasis. A positron emission tomography (PET) scan is useful in identifying regional spread and metastatic lesions. A biopsy is discouraged in this condition as it results in disease upstaging, and is considered only in cases of unusual presentation or imaging.

Neuroblastoma is an important differential diagnosis to consider, as it can present similarly to Wilms tumor. It is usually seen in preschool-aged children and commonly presents as a palpable abdominal mass. The retroperitoneum is the most common site for this tumor. Therefore, it is essential to test for elevated vanillyl mandelic acid (VMA) in the urine in this patient to rule out neuroblastoma.

Children's oncology group (previously known as National Wilms Tumor study group) staging system for Wilms tumor:

Treatment: Surgery and chemotherapy, with or without radiotherapy, are the mainstays of treatment.

- Radical nephrectomy is the standard surgical procedure done. Routine sampling of the para-aortic, hilar, and paracaval lymph nodes should be done.
- Nephron-sparing nephrectomy is done for children with bilateral disease or with a solitary kidney. Preoperative chemotherapy can be used in these cases to induce tumor shrinkage.
- The classical chemotherapy includes vincristine and dactinomycin with the addition of Adriamycin(Doxorubicin).

Stage I	Tumour confined to the kidney and completely resected with negative lymph nodes.
Stage I I	Tumour extends beyond the kidney but is completely resected with negative margins and lymph nodes.
Stage I II	Residual tumour present following surgery confined to the abdomen, including gross or microscopic tumour.
Stage I V	Hematogenous metastases (lung, liver, bone, brain, etc.) or lymph node metastases outside the abdominopelvic region.
Stage V	Metastasis to opposite kidney (bilateralrenal involvement).

Solution to Question 4:

The clinical features of a male patient presenting with a large, round, firm, and painless testicular mass with elevated AFP levels are suggestive of a non-seminomatous germ cell tumor (NSGCT). Among the given options, a high inguinal orchidectomy would be the next best step in the management.

An ultrasound scan (USG) of the testis is mandatory to confirm the diagnosis of a testicular tumor and assess the contralateral testis. Since USG is not given as an option, the next best step in this scenario would be a high inguinal orchidectomy. FNAC and trucut biopsy are contraindicated due to risk of seeding of the scrotum and upstaging of the cancer.

Radical inguinal orchiectomy is the gold standard treatment for excision of the primary tumor. Sperm cryopreservation should be offered before surgery.

After removal of the affected testis, a metastatic workup is done with CT of the chest, abdomen, and pelvis to stage the disease as follows:

- Stage I: Localized tumor with no lymph node involvement or distant metastasis


- Stage II: Regional lymph node metastasis
- Stage III: Distant metastasis or nodal metastasis with elevated tumor markers

Further management is based on staging with treatment options including active surveillance, combination chemotherapy with bleomycin-etoposide-cisplatin and retroperitoneal lymph node dissection.

Solution to Question 5:

This clinical scenario involves a patient presenting with abdominal pain and tachycardia in the post-operative period following cholecystectomy. An ultrasound scan (USG) reveals a fluid collection in the right sub-hepatic space, strongly suggesting a bile duct injury following cholecystectomy. As the patient presents more than 48 hours after the surgery, the next best step in management would be to drain the fluid collection using a USG-guided pigtail catheter.


More than 80% of bile duct injuries occur during cholecystectomy. A bile duct injury can result in either bile leakage or stricture. Patients with bile leakage may present with fever, increasing abdominal pain, jaundice or bile leakage from an incision.



Management strategy:

- Imaging: Ultrasonography to assess for a fluid collection and to evaluate the biliary tree
- Control of infection, limiting inflammation
- Percutaneous drainage of periportal fluid collections
- Parenteral antibiotics
- Clear and thorough delineation of entire biliary anatomy
- Endoscopic retrograde cholangiopancreatography (ERCP), especially if cystic-duct stump leak is suspected. It is both diagnostic and therapeutic.
- Magnetic resonance cholangiopancreatography (MRCP)


- HIDA scanning is also helpful to identify and quantitate bile leaks.
- Definitive surgical correction
- Roux-en-Y choledochojejunostomy can be done for benign strictures or duct transections.
- Balloon dilatation with placement of a stent can be done for strictures of recent onset.



Solution to Question 6:

The instrument shown in the image is the Veress needle which is used to insufflate the abdomen for laparoscopic surgery.

The image below shows the Veress needle:



There are two methods for establishing abdominal access during laparoscopic surgeries:

- Direct puncture using a Veress needle
- Direct peritoneal access (Hasson's technique)

Direct puncture laparoscopy:

In this procedure, first, the abdominal wall is grasped with towel clips or fingers to elevate it away from the underlying structures. Then, a small incision is made in the umbilicus, and insufflation is performed using a Veress needle. When introducing the Veress needle, two distinct pops can be felt as the needle pierces the abdominal wall fascia first and then the peritoneum. The umbilicus is the preferred location for introducing a Veress needle because the abdominal wall is thinner at this site.




Laparoscopic surgeries can be performed under general anesthesia or local anesthesia; although, general anesthesia is preferred. Carbon dioxide gas is usually used for insufflation, and the maximum pressure is typically in the range of 14-15 mm Hg. However, if the procedure is done under local anesthesia, nitrous oxide is used. Insufflation with nitrous oxide is stopped at a pressure of 10 mmHg or when 2L of gas has been insufflated.

Direct peritoneal access technique (Hasson technique): The surgeon makes a small umbilical incision and gains access to the peritoneal cavity under direct vision by dissecting through the abdominal fascia and peritoneum. It is preferred in patients who have undergone previous abdominal operations where the small bowel may be adherent to the undersurface of the abdominal wound.

Solution to Question 7:

The clinical features of a perineal bruise and blood at the urethral meatus following a fall from a bicycle are suggestive of a bulbar urethral injury. In this case, since the bladder is not palpable, the best option would be to wait for the bladder to fill and for the urge to urinate to occur. Once the bladder is full, the urine should be drained using a suprapubic catheter.


The image given below shows the various parts of the urethra:



Urethral injuries can be classified into:

A retrograde urethrogram with water-soluble contrast is used for confirmation of the diagnosis.

Type	Proximal urethral injury	Distal urethral injury
Site of injury	Prostatic urethra Membranous urethra	Bulbar urethra Penile urethra
Mechanism	Secondary to pelvic fracture	Occurs due to a blow to the perineum by: Straddle injury Loose manhole covering Gymnasium accidents
Clinical features	Urinary retention Blood at the urethral meatus High-riding prostate on digital rectal examination (Vermooten sign) Deep perineal hematoma Injury to the membranous urethra leads to accumulation of urine in the deep perineal pouch.	Urinary retention Blood at the urethral meatus Perineal hematoma (Butterfly hematoma)



Solution to Question 8:

The patient's GCS is Eyes-3, Verbal-NT, Motor-6.

It is calculated as follows:

- Eye opening: Opening to verbal command - 3
- Verbal response: In the case of endotracheal intubation, the verbal response score is noted as NT (non-testable)
- Motor response: Movement of all 4 limbs spontaneously - 6

Total GCS score is not recorded since one of the components is non-testable.

Solution to Question 9:

The green IV cannula shown in the image corresponds to a gauge size of 18G.

Solution to Question 10:

The next step in the management in this patient with ductal carcinoma in situ (DCIS) would be adjuvant radiotherapy, according to the University of Southern California/Van Nuys Prognostic Index (USC/VNPI).

The USC/VNPI is a scoring system that evaluates the following four factors to predict recurrence in patients with DCIS who are undergoing conservative treatment:

- Tumor size

- Margin width
- Pathological classification (based on nuclear grade, and presence or absence of comedonecrosis)
- Age

This index also provides recommendations for managing patients based on the calculated score.

Scoring system for USC/VNPI:

The current recommendations for the management of DCIS according to this index are:

- Excision alone for scores 4,5, and 6
- Excision plus radiotherapy for scores 7,8, and 9
- Mastectomy for scores 10,11, and 12

In this case,

- Tumor size of 4cm (40mm) - score 2
- 10 mm margin clearance - score 1
- High nuclear grade with necrosis - score 3
- Age of 55 years - score 2

The total score is 8; hence radiotherapy is indicated after breast conservation surgery.

Parameters	Score 1	Score 2	Score 3
Size(mm)	≤15	16-40	≥41
Margin(mm)	≥10	1 - 9	<1
Class	Grade 1 or 2 without necrosis	Grade 1 or 2 with necrosis	Grade 3 with or without necrosis
Age(years)	>60	40-60	<40

Solution to Question 11:

In this clinical scenario, a patient with gallstones, a dilated common bile duct, and an elevated alkaline phosphatase level raises suspicion for common bile duct stones. The next best step would be to perform magnetic resonance cholangiopancreatography (MRCP) to confirm their presence.

A dilated common bile duct (>8 mm) detected on ultrasound in a patient with gallstones, jaundice, biliary pain and an abnormal liver function panel (elevated bilirubin, alkaline phosphatase, and transaminases; but these can be normal in some cases) strongly indicates the presence of common bile duct stones (choledocholithiasis).

If there is uncertainty regarding the presence of bile duct stones, as in this case where bilirubin levels are normal, MRCP offers precise anatomical visualization and exhibits high sensitivity and

specificity for diagnosing choledocholithiasis.

While endoscopic retrograde cholangiopancreatography (ERCP) is also highly effective in diagnosing choledocholithiasis, due to its invasive nature and the risks associated with the procedure, it is rarely used as a purely diagnostic modality. Instead, it is reserved for cases in which a therapeutic intervention, such as stone extraction or sphincterotomy, is planned.

Treatment options:

- ERCP along with sphincterotomy and stone removal.
- Laparoscopic common bile duct exploration
- Open common bile duct exploration is only done when endoscopic and laparoscopic methods are not feasible or when concomitant biliary drainage is needed.


Solution to Question 12:

In the interrupted suture technique, the needle is placed at an angle of 90 degrees (at right angles).

Interrupted sutures:

The needle should be inserted and exit the skin at right angles. It should be rotated through the tissue rather than dragged to prevent unnecessary enlargement of the needle hole. The distance from the suture entry point to the wound edge should be approximately equal to the depth of the sutured tissue. There should be twice this distance between two consecutive sutures.

The image below shows interrupted suture:




Other suturing techniques:

Continuous Suture: The first suture is inserted in a manner identical to that in an interrupted suture. The remaining sutures are inserted in a continuous manner until the far end of the wound is reached. Care should be taken to maintain the tension of the suture to avoid excessive tension


or leaving it too slack; hence, the use of an assistant is warranted. An Aberdeen knot is used to secure the end of the suture.

Continuous suture




Mattress sutures - They can be vertical or horizontal and are used to create either an eversion or inversion of the wound edges. The initial suture is inserted in the same way as an interrupted suture, but then, the needle either moves horizontally or vertically and traverses both edges of the wound once again, as seen in the image below. The main advantage of the mattress suture is that it produces an accurate approximation of wound edges, especially in settings where the edges are not easily approximated.

Mattress suture



Subcuticular sutures: It is used for a better cosmetic appearance. It is also employed in situations where the edges of the wound are easily approximated. In this technique, small bites of the subcuticular tissues are taken on alternate sites of the wound, and then, they are gently pulled together. This causes an approximation of the wound edges without the risk of the cross-hatched markings of interrupted sutures.



Solution to Question 13:


The given clinical scenario involves a patient with a history of penetrating abdominal trauma and presenting symptoms of abdominal pain and obstipation. The erect radiograph shows air under the diaphragm which strongly points to a diagnosis of hollow viscous perforation.

Perforation of the alimentary tract is a life-threatening condition. This leads to the leakage of the enteric contents, thereby leading to acute bacterial peritonitis. Peritonitis is the generalized or localized inflammation of the peritoneum either due to the free spillage of fluids into the peritoneal cavity or secondary bacterial infection.

Clinical features include abdominal pain and distension, nausea, vomiting, tachycardia, tenderness, rebound tenderness, decreased or absent bowel sounds, and guarding on palpation. Fever may be present. Features of septic shock can be seen at later stages.

Investigations include estimation of the white blood cell count, urea, serum electrolytes, and an erect abdominal radiograph showing free gas under the diaphragm. A supine radiograph may show dilated gas-filled loops of the bowel suggestive of paralytic ileus. However, CECT is the most sensitive investigation to detect minimal/ suspected pneumoperitoneum.

Hollow viscus perforation



Management of a hollow viscus perforation mainly includes resuscitation, analgesics, intravenous broad-spectrum antibiotics, and emergency exploratory laparotomy.

Other options:

Option A- Hemothorax appears as a homogenous increase in opacity of the hemithorax.

Right Hemothorax



Option C- Pneumothorax is the collection of air in the pleural space. On radiography, there may be increased lucency on the affected side, depression of the affected hemidiaphragm, and mediastinal shift to the opposite side. Associated rib fractures might also be visible.


Left Pneumothorax



Option D- Intestinal obstruction

- In small bowel obstruction, the obstructed bowel loops are centrally located and lie transversely. In cases of jejunal obstruction, valvulae conniventes, which are mucosal folds that span the entire diameter of the bowel, create a ladder effect. Additionally, a distended cecum may exhibit a gas shadow in the right iliac fossa.

Small bowel obstruction



- In large bowel obstruction, the presence of haustrations is a characteristic feature. Haustral folds are irregular and do not extend across the entire diameter of the bowel. However, it is important to note that haustral folds are absent in the caecum.

Large bowel obstruction



Solution to Question 14:

In this patient, who has sustained a long tear to the brachial artery due to a stab injury, the best treatment option is to repair it with a great saphenous vein interposition graft harvested from the patient's uninjured leg.

The most desirable graft used is the great saphenous vein . A native vein graft is preferable to a prosthetic graft because:

- Its elastic properties are comparable to that of an artery and enable pulsatile flow
- The diameter is similar to that of an extremity artery
- It is not thrombogenic and has long term patency

Other grafts that can be used are cephalic vein, lesser saphenous vein, and prosthetic conduits. Cephalic vein is not the first choice because it is less muscular than greater saphenous vein, and might also compromise the upper limb venous access in the already injured patient. Both the cephalic and the lesser saphenous veins present problems while harvesting in a trauma patient.

Other options:

Option A: Primary arterial repair with end-to-end anastomosis can be done only if the repair is tension-free. If the length of the injured area is long, vessel injuries cannot be repaired by primary end-to-end anastomoses and instead require an interposition graft.

Option C: Prosthetic grafts are only preferred when there has been a trauma to the native saphenous vein or when there is a size mis-match. Polytetrafluoroethylene (PTFE) graft is commonly used.

Option D: Saphenous vein patch is used in the closure of the arteriotomy site during carotid endarterectomies.

Solution to Question 15:

The given radiograph shows elevation of the left hemidiaphragmatic line along with the bowel loops and stomach seen in the pleural cavity. The probable cause would be a diaphragmatic hernia following blunt injury due to a road traffic accident. Chest tube insertion is contraindicated in diaphragmatic hernia due to risk of injury to bowel.

Chest tube insertion/thoracostomy is indicated cases of pneumothorax, hemothorax, chylothorax, pleural effusion, and pleural empyema. Contraindications include pulmonary adhesions, pulmonary trauma, coagulopathy, and diaphragmatic hernia.

Chest radiography performed after the placement of a nasogastric tube can be helpful in identifying any herniation of the stomach into the chest. Log roll and epidural anesthesia are not contraindicated in diaphragmatic hernia.

Blunt trauma to the diaphragm results in a linear diaphragmatic rupture, which is typically large and accompanied by the herniation of abdominal contents into the chest. Acute diaphragmatic injuries require repair through the abdominal approach to address associated visceral injuries. The diaphragmatic edges are approximated using Allis clamps, and the defect is closed with a running 1-0 polypropylene suture.

Solution to Question 16:

The green IV cannula used in this hemodynamically unstable patient is an 18G cannula, which has a maximum flow rate of 103 ml/ min.


Solution to Question 17:

The most common cause of bladder outflow obstruction in a male child is the posterior urethral valve (PUV).

A PUV is a membranous fold in the posterior urethra which results from the persistence of the urogenital membrane. It is typically located just distal to the verumontanum and is the primary cause of bladder outlet obstruction in newborn males.

Diagnosis is most commonly made antenatally with ultrasound which demonstrates bilateral hydronephrosis, a distended bladder, and a dilated posterior urethra which appears as the keyhole sign on imaging. In severe cases, there is oligohydramnios and pulmonary hypoplasia during the antenatal period.

Posterior Urethral Valve - Keyhole Sign




If the diagnosis is not made antenatally, then babies typically present with urinary tract infections in the neonatal period or with uraemia and renal failure. Clinically, a poor urinary stream, dribbling of urine, and palpable bladder (due to diverticula) can be present.

The investigation of choice is micturating cystourethrography.

The diagnosis is best made during the micturition phase of micturating cystourethrography in lateral or oblique views so that the posterior urethra can be imaged adequately. The following findings are seen:

- Dilatation and elongation of the posterior urethra - equivalent of the ultrasonographic keyhole sign
- Linear radiolucent band corresponding to the valve - seen occasionally
- Vesicoureteral reflux - seen in 50% of patients
- Bladder trabeculation/diverticula

Posterior urethral valve - Micturating cystourethrogram



Treatment is initially supportive by inserting a urinary catheter to relieve symptoms of obstruction. Definitive treatment is by urethoscopic destruction of the valves with continuing lifelong supportive treatment of the dilated urinary tract, recurrent urinary infections and uremia.

Other options:

Options B and C- Urethral atresia and anterior urethral valve are rare congenital anomalies that can also lead to urinary obstruction.

Option D- Ureterocele is the cystic enlargement of the intramural ureter. It is common in females. They can present with features of urinary tract infection, bladder obstruction, or ureteral stones. Excretion urography will show cobra head appearance.

The image shown below depicts the "cobra head" sign in a patient with bilateral ureterocele.



Solution to Question 18:

All the 4 given statements on colon cancer are true:

- Right and left colon cancers present differently. Left-sided tumors commonly cause obstruction and present with changes in bowel habits or rectal bleeding. On the other hand, right-sided tumors typically present with iron deficiency anemia or a mass.
- While the majority of colorectal cancers occur sporadically, hereditary cases are also seen in patients with a family history of colorectal cancer. Genetic testing aids in early diagnosis for individuals with a positive family history.
- Metformin, a commonly used biguanide compound for diabetes mellitus, reduces the risk of cancer, including colorectal cancer. It achieves this by disrupting cellular energy supply through the inhibition of aerobic glycolysis.
- Liver is the most frequent site of distant metastasis for colorectal cancer, with metastases occurring via the portal venous system. The risk of hepatic metastasis increases with tumor size and grade. The lung is the second most common site of distant metastasis, also due to hematogenous spread. Carcinomatosis (diffuse peritoneal metastases) results from peritoneal

seeding.

Note: Regional lymph node involvement is the most common form of spread of colorectal carcinoma and usually precedes distant metastasis.

Solution to Question 19:

Frozen section analysis performed during breast carcinoma surgery is not suitable for definitive diagnosis due to technical challenges, potential quality issues in processing and staining, and the difficulty of obtaining accurate results.

Frozen section analysis is performed when a prompt biopsy report is required, allowing the surgeon to make immediate decisions regarding the further steps of management.

- Frozen section analysis plays a valuable role in assessing involvement of sentinel lymph nodes during surgery (option A). When a positive node is identified intraoperatively, the surgeon can proceed with immediate axillary lymph node clearance. On the other hand, if the sentinel node does not show tumor involvement, further axillary dissection can be avoided.
- It also helps in diagnosing nodal metastasis in clinically identified lymph nodes (option C). However, it may not detect subcapsular micrometastases.
- While performing breast-conserving surgeries, assessment of tumor margins (option B) using frozen sections is helpful in ensuring complete tumor excision.

Solution to Question 20:

Apart from abdominal pain, 1, 2, 3, and 4 form the Reynold's pentad which is seen in severe cholangitis.

The components of Reynold's pentad are:

- Right upper quadrant abdominal pain
- Fever
- Jaundice
- Hypotension
- Altered sensorium

Cholangitis is the infection of the common bile duct (CBD) secondary to obstruction and increased intraluminal pressure due to CBD stones. Cholangitis presents with the classical triad of fever, right upper quadrant pain, and jaundice, referred to as Charcot's triad. It is a medical emergency and if left untreated, it may progress in severity and presents with Reynold's pentad - Charcot's triad with hypotension and altered mental status due to sepsis.

Abdominal ultrasound is performed to check for the presence of biliary stones and bile duct dilatations. The patient should be started immediately on a first-line broad-spectrum antibiotic and rehydration, along with biliary drainage. Biliary drainage is done using endoscopic retrograde cholangiopancreatography (ERCP). Biliary stones are the cause of cholangitis the majority of the

time, which can be removed by endoscopic sphincterotomy during ERCP.

Solution to Question 21:

The Romo Vac drain, depicted in the image, is a closed type of drain commonly used after mastectomy.

Drains are placed at surgical sites to freely collect the air or fluids that may accumulate. This enhances wound healing by preventing the formation of unnecessary cavities. Drains are prophylactic in elective surgeries and therapeutic in emergency surgeries. Drains can be -

- Open - They are based on the principle of gravity
- Semi-open - Their action is based on capillary effect
- Closed - They carry out their actions through suction

Suction drains are commonly used in chest, breast, abdominal, thyroid, and plastic surgeries. They remove intraperitoneal secretions, anastomotic leaks, and blood from hemorrhage at the surgical site. Romo Vac drain is a negative suction drain closed drain kept in the subcutaneous tissue. Romo Vac drain is indicated in modified radical mastectomy, subcutaneous swelling excision, and amputations.

Special types of drains:

- Chest drains are connected to an underwater seal to aid the exit of air from the pleural cavity. They are used in pneumothorax, pleural effusion, and hemothorax.
- After surgical exploration of the bile duct, the sphincter of Oddi goes into spasm. To facilitate drainage of bile, a T tube may be used.

Drain insertion site care: The site of insertion of the drain is kept clean by drying and patting the drain insertion site with a clean towel after bath. The dressing is changed daily with folded gauze pieces placed below and above the drain site.

Removal of drains: The catheter at the site is milked for residual collections. It is then clamped and the suture holding the catheter is cut. The tube is then gently pulled out and any clots at the perforation end are removed. The time of removal of drains for various conditions is as given below.

- Drains used in surgeries like thyroidectomy are usually removed within 24 hours. Their main purpose is to cover postoperative bleeding.
- In a mastectomy, a drain is placed for 5 days to collect serous collections.
- Drains used to cover the colorectal anastomoses are placed for 5-7 days.
- Common bile duct T-tube drains are usually placed for 10 days.
- Drains for infections are removed when the infection subsides or when the content of the drain is minimal.

After the drain is removed, a folded gauze is placed over the drain insertion area. The dressing can be removed after 24 hours. The patient may be given an antibiotic ointment to apply over the area for a week.

Solution to Question 22:

The correct statements about hepatocellular carcinoma (HCC) are 1,3 and 4. Lenvatinib is used in the advanced stages of hepatocellular carcinoma (HCC).

HCC is the most common cause of primary liver cancer. It is more commonly seen in males and the peak age group is between 65-70 years. The main risk factor for HCC is cirrhosis and associated chronic liver damage due to chronic infection by hepatitis B or C virus (HBV or HCV) infection, alcohol abuse, metabolic syndrome, or hemochromatosis.

The incidence of HCC is increasing (statement 3) worldwide. This is contributed by the increase in HCV and HBV infections, along with lifestyle disorders like obesity. Obesity is associated with non-alcoholic fatty liver disease (NAFLD). This progresses on to non-alcoholic steatohepatitis (NASH) causing cirrhosis and hepatocellular carcinoma. Thus, NASH and NAFLD are risk factors of HCC (statement 4).

Barcelona-Clinic-Liver Cancer (BCLC) Classification is the commonly used staging system. It stages the tumors as follows:

- Very early HCC - Single, well-differentiated tumor ≤ 2 cm
- Early HCC - Single tumor or a maximum of 3 nodules ≤ 3 cm in diameter
- Intermediate HCC - Any other size or number confined to the liver
- Advanced HCC - Presence of cancer-related symptoms or tumors with macrovascular invasion (of any type, including branch, hepatic, or portal vein), lymph node involvement, or extrahepatic spread
- End-stage disease - Severe liver dysfunction with impaired quality of life

Treatment

Very early HCC is treated by ablation. Early HCC with a single nodule is resected but when there are multiple nodules over 3 cm, ablation or transplantation are considered.

Intermediate stage HCC is treated by chemoembolization. Transarterial chemoembolization (TACE) is used for multinodular lesions (statement 1).


Advanced stages of the disease with macrovascular invasion are treated with systemic therapies. Lenvatinib and sorafenib are first-line drugs. Supportive care is given at the terminal stage of the disease.

Solution to Question 23:

The given clinical scenario with USG and CECT showing a distended gall bladder, dilated common bile duct, and intrahepatic biliary radicles raises suspicion of common bile duct stones (CBD stones). Among the given options, magnetic resonance imaging (MRI/MRCP) will be next most useful modality to localize the cause and level of obstruction.

Magnetic resonance cholangiopancreatography (MRCP) uses T2-weighted MRI sequences and it is highly sensitive and specific in identifying CBD stones. MRCP provides visualization of dilated bile ducts and often enables accurate assessment of the level of occlusion in the biliary tree. It is a non-invasive test and is preferred by many surgeons to assess the need for endoscopic retrograde cholangiopancreatography (ERCP). ERCP is also highly effective at diagnosing CBD stones. However, due to the risks associated with the procedure, it is rarely used as a purely diagnostic modality, rather it is now used only as a therapeutic modality such as when stone extraction or sphincterotomy is planned (option D).


Ultrasonography (USG) can detect gallstones and assess the size of the common bile duct. If a patient with gallstones, jaundice, and biliary pain shows a dilated common bile duct (>8 mm in diameter) on USG, it strongly suggests the presence of CBD stones. CT scan is also a sensitive diagnostic tool for CBD stones. USG and CT will generally be the first line modalities due to their easier availability, and lower cost and time requirements.



Other options:

Option B- Endoscopic ultrasound is used for detecting CBD stones. It is considered as good as ERCP but it is invasive and requires expertise in endoscopic procedure and hence it is not done routinely.

Option C- Percutaneous transhepatic cholangiography (PTC) is rarely performed in patients with CBD stones. In patients where endoscopic or surgical procedures are contraindicated, it may be used for both diagnostic and therapeutic purposes.



Solution to Question 24:

Incontinence during intercourse is not a lower urinary tract symptom (LUTS).

The lower urinary tract symptoms (LUTS) can be classified as:

Storage LUTS: This occurs due to the failure of the bladder to act as a urine reservoir. It is commonly seen in the overactive bladder or bladder neuropathy. Symptoms related to storage LUTS are:

- Frequency - Increased frequency of voiding
- Nocturia - a patient has to wake up at least once at night to void
- Urgency (option A) - the sudden uncontrollable desire to pass urine
- Urge incontinence - a large volume of involuntary urine leakage preceded by urgency
- Stress incontinence (option C) - dribbling of urine during coughing, laughing, sneezing, or exercising. This occurs in women who have undergone vaginal deliveries.
- Nocturnal enuresis (option B) - involuntary leakage of urine during sleep. It is commonly seen in elderly men having chronic retention of urine with overflow incontinence.
- Strangury - is a sensation of constantly needing to void. Commonly occurs due to a lower urinary tract infection.

Voiding LUTS: This is commonly seen in bladder outlet obstruction (BOO). Symptoms related to voiding LUTS are:

- Hesitancy - difficulty in initiating micturition
- Reduced stream
- Straining - muscular effort is needed to initiate or to improve the urinary stream

Post micturitional LUTS: This is also caused due to BOO. Symptoms related to post micturitional LUTS are:

- Incomplete emptying - a sensation that the bladder is full still remains after micturition
- Post micturition dribble - Involuntary passage of urine after the patient has finished passing urine.

Solution to Question 25:

Fluorescence in situ hybridization (FISH) for gene amplification should be performed in all cases of breast tumors with an immunohistochemistry (IHC) score of 2+.

HER2/neu is a protein promoting rapid growth of breast cancer cells. All invasive breast cancers show undergo HER2/neu testing on biopsy or post-surgical removal. HER2 status is crucial for guiding breast cancer treatment. Two tests, immunohistochemistry (IHC) and fluorescence in situ hybridization (FISH), determine the presence of high HER2 protein levels. As FISH is costlier and more time-consuming, IHC is often performed as the initial test.

Her-2/neu IHC interpretation:

- 0 to 1+: Negative for HER2. These cancers do not respond to drugs targetting HER2 receptors.
- 2+: Equivocal. Hence, further testing with FISH is required to clarify the HER2 status.
- 3+: Positive for HER2. These cancers respond to drugs targetting HER2 receptors.

Transtuzumab and pertuzumab are monoclonal antibodies specifically used for HER2/neu-positive tumors.

Solution to Question 26:

This patient with low-grade 2x2 cm bladder cancer is likely to have non-muscle invasive bladder carcinoma (NMIBC) should undergo transurethral resection of the bladder tumor (TURBT) for accurate staging purposes.

Bladder cancer is graded (G1/2/3) based on level of differentiation. NMIBC includes stages Tis (in-situ), Ta (non-invasive papillary carcinoma), and T1 (invades lamina propria) while muscle invasive bladder cancer (MIBC) includes stages T2 (invades muscularis propria), T3 (invades perivesicular soft tissue), and T4 (invades adjacent organs). NMIBC accounts for almost 70% of cases at presentation.

Treatment:

- NMIBC - The initial management of bladder tumors consists of transurethral resection of bladder tumor (TURBT) for accurate staging, which aids in determining the subsequent treatment plan. Intravesical therapy, involving chemotherapy (with mitomycin, doxorubicin, or epirubicin) or immunotherapy (BCG), is administered after TURBT to eradicate any remaining cancer cells. This therapy primarily targets the bladder lining and is not effective for cancers that have invaded deeper layers.

- MIBC - The standard treatment for muscle-invasive tumors is radical cystectomy and pelvic lymphadenectomy with urinary diversion (ileal conduit) (option B). Systemic adjuvant and neoadjuvant chemotherapy with M-VAC (Methotrexate, vinblastine, doxorubicin, and cisplatin) has shown to be beneficial (option D).

Option C: Partial cystectomy can be attempted in patients with focal bladder cancer, but the risk of recurrence remains high.

Solution to Question 27:

Statements 1, 3 and 4 are true.

Screening with noninvasive tests for common cancers aims to achieve early diagnosis, enable less invasive surgical treatments with reduced complications, and potentially improve surgical cure and overall survival rates.

Statement 1: Mammography as a routine screening tool has been found to lower breast cancer mortality at an acceptable cost. In the United Kingdom, an expert panel reported a 20% reduction in breast cancer mortality due to screening, despite the occurrence of overdiagnosis. The panel recommended continuing breast screening due to its significant benefits.

Statement 3: Visual inspection of the oral cavity is widely used for screening oral cancer. The target population includes individuals aged 30 and above who use tobacco and/or alcohol. Visual inspection is advantageous due to its feasibility, safety, acceptability, accuracy in detecting precancerous and cancerous lesions, and cost-effectiveness in reducing oral cancer mortality.

Statement 4: Pap smear is a screening test for cervical cancer. Cervical cancer screening should start at age 21. Women aged 21-29 should have Pap smear every 3 years, while women aged 30-65 should have both HPV and Pap co-test every 5 years.

Other statements:

Statement 2: American Cancer Society guidelines recommend colon cancer screening for average-risk individuals starting at age 50. Screening options include:

- Yearly fecal occult blood testing (FOBT) / fecal immunohistochemical testing (FIT)
- Flexible sigmoidoscopy every 5 years,
- FOBT/FIT and flexible sigmoidoscopy in combination every 5 years, or
- Colonoscopy every 10 years.

Solution to Question 28:

The correct method for measuring a nasogastric tube (Ryle's tube) is from the tip of the nose to the ear lobule to the xiphisternum.

Nasogastric (NG) tubes play a role in treating intestinal obstruction and can also be used to provide nutritional support. The commonly used method to determine the length of a nasogastric tube (NG tube) is the NEX method, which involves measuring from the tip of the nose to ear to

xiphisternum. The approximate length to be inserted is 55cms.

Ideally, Ryle's tube is inserted in the sitting position with the neck slightly flexed. Confirmation of the correct position is done by the following:

- Auscultation in the epigastric area while injecting a bolus of air
- An abdominal radiograph is the best way to confirm the location of the tube

An NG tube placed for suction should ideally be located within the stomach to avoid potential damage to the duodenum. For decreased risk of aspiration, the ideal location for an NG tube is postpyloric (55-60cm).

Indications for NG tube :

Drainage purposes:

- Conservative management of bowel obstruction and postop paralytic ileus
- Gastric decompression prior to emergency surgery
- Aspiration of gastric fluid

Feeding purposes:

- Nutritional support in patients in motor neuron disease, stroke, and following upper aerodigestive tract procedures
- Administration of medication

Contraindications:

Absolute:

- Severe midface trauma
- Recent nasal surgery

Relative:

- Coagulation abnormality
- Esophageal varices
- Recent banding of esophageal varices
- Anastomoses in the stomach and esophagus

Instruments tray used for nasogastric tube insertion:



Solution to Question 29:

The structures cut when the initial incision gridiron incision (McBurney's incision) is converted to the Rutherford Morrison incision are the external oblique, internal oblique, and transverse abdominis muscles.


When the preoperative diagnosis is reasonably certain, the widely used incision for appendectomy is known as the gridiron incision. It is made perpendicular to a line connecting the anterior superior iliac spine to the umbilicus, with its center at McBurney's point. When better access is needed, the gridiron incision can be converted to a Rutherford Morrison incision by extending the incision obliquely upwards and laterally.

Rutherford Morrison's incision is suitable for a para- or retrocecal appendix that is fixed. It is an oblique muscle-cutting incision starting from McBurney's point and extending upward and laterally as needed. All layers (skin, subcutaneous tissues, external oblique, internal oblique, transversus abdominis, and the peritoneum) are divided along the incision line.

In recent years, the transverse skin crease (Lanz) incision has gained popularity due to improved exposure and ease of extension when necessary. This incision is placed approximately 2 cm below the umbilicus along the midclavicular-midinguinal line, with its length adjusted according to the patient's size and obesity. If required, the incision can be extended medially by retracting or dividing the rectus abdominis muscle (statement 4).

When there is uncertainty in the diagnosis, especially in the presence of intestinal obstruction, a lower midline abdominal incision is preferred over a right lower paramedian incision.

Incisions used in appendicectomy



©Marrow

Surgery NEET 2021

Question 1:

A patient underwent surgery for breast carcinoma. She now presents with painless swelling involving the arm as shown in the image below. What is the most likely diagnosis?



- a) Lymphedema
- b) Lymphangiosarcoma
- c) Venous thrombosis
- d) Cellulitis

Question 2:

Which of the following statements is true regarding the condition given below?



- a) Patient must be started on anti-gas gangrene serum therapy
- b) It is a mixed flora aerobic and anaerobic infection
- c) Bilateral orchidectomy is generally performed
- d) Compulsory urinary diversion is performed

Question 3:

A baby playing unsupervised was brought with complaints of difficulty in swallowing for the last few hours. The X-ray image is given below. What is the diagnosis?



- a) Foreign body in the trachea

- b) Foreign body in the esophagus
- c) Artifact
- d) Soft tissue calcification in the neck


Question 4:

A 75-year-old man with prostate carcinoma presents to you with a PSA of 9 ng/mL with a small tumor focus. His Gleason score is 6. What will be your most likely management?

- a) Radical prostatectomy
- b) External beam radiation
- c) Brachytherapy
- d) Active surveillance

Question 5:

Identify the gross specimen obtained from a cystoprostatectomy.



©Marrow

- a) Malakoplakia
- b) Bladder carcinoma
- c) Prostate carcinoma
- d) Schistosomiasis


Question 6:

A patient after a road traffic accident presented with pain abdomen. The resident examines the child and found that vitals were stable and tenderness was present in the left lumbar region. Which is the best investigation of choice?

- a) Contrast enhanced CT scan
- b) Retrograde urethrogram
- c) Wait and watch
- d) Emergency laparotomy

Question 7:

A patient with rectal prolapse underwent surgery. The postoperative image is given below. Identify the procedure.



- a) Stapled hemorrhoidopexy
- b) Well's procedure
- c) Thiersch wiring
- d) Altemeier repair

Question 8:

A patient presents after a motorcycle road traffic accident with no reported pain or injury. During examination, blood is observed at the tip of the urethral meatus, and the bladder is

non-palpable. What would be your next step in management?

- a) Foley's catheterization
- b) Wait for bladder to fill
- c) Nephrogram
- d) Suprapubic cystostomy

Question 9:

A patient presents with a hernia medial to the inferior epigastric artery. What is the most likely diagnosis and appropriate management for the same?

- a) Direct hernia and Bassini repair
- b) Direct hernia and Lichtenstein mesh repair
- c) Indirect hernia and Bassini repair
- d) Indirect hernia and Lichtenstein mesh repair

Question 10:

A patient following a road traffic accident presented with subcutaneous emphysema. On auscultation, there was no air entry on the right side. Vitals were unstable. What will be your immediate management?

- a) Insertion of wide bore needle in the 5th intercostal space (ICS)
- b) Wide bore needle decompression and IV fluids
- c) Intubation and positive pressure ventilation
- d) eFAST

Question 11:

A 54-year-old female patient presents with abdominal pain and bilious vomiting. She had a history of abdominal hysterectomy 2 years back. On examination, abdomen auscultation revealed high-pitched bowel sounds. Investigations showed air within the biliary tree. What is the most likely diagnosis?

- a) Gallstone ileus
- b) Adhesive intestinal obstruction
- c) Ischemic enterocolitis

d) Diverticulitis

Question 12:

In a patient with abdominal pain and sterile pyuria, what is the finding shown in the image given?



- a) Putty kidney**
- b) Nephrocalcinosis**
- c) Staghorn calculus**
- d) Psoas calcification**

Question 13:

A patient was found to have a mass arising from the antrum of the stomach which involves the head of the pancreas and metastasis to the right lobe of the liver. How will you manage this patient?

- a) Palliative gastrojejunostomy followed by chemotherapy**
- b) Radical gastrectomy**
- c) Whipple's procedure**
- d) Gastrectomy with right hepatectomy**


Question 14:

A patient underwent cholecystectomy for gallstones over 2 years ago. She now presents with symptoms of obstructive jaundice. Investigation reveals a stone in the common bile duct. What is the stone most likely to be?

- a) Primary
- b) Secondary
- c) Tertiary
- d) Retained

Question 15:

A 70-year-old female patient presents with pain in the thigh during walking. Investigation reveals the following finding as given below. What will be the best surgical management?



- a) PTA with stenting
- b) Iliofemoral bypass
- c) Aortofemoral bypass
- d) Aorto-iliac bypass

Question 16:

A young air-hostess presented to the outpatient department with dilated veins on her left leg (image). According to the CEAP classification for varicose veins, what will be the C stage in this patient?



- a) C2A
- b) C2S
- c) C4a
- d) C1


Question 17:

An alcoholic patient presented with severe abdominal pain. Pancreatitis was suspected and a CT scan revealed lesions with fluid collected near the body of the pancreas. Which of the following enzymes is most likely to be elevated in this scenario?

- a) Serum GGT
- b) CEA
- c) Serum lipase
- d) Serum bilirubin

Question 18:

A 25-year-old patient presents with abdominal pain and fever for 3 days duration. The chest X-ray in the erect position is given below. What will be the most likely diagnosis?



- a) Gastric volvulus**
- b) Hollow viscus perforation**
- c) Liver abscess**
- d) Empyema thoracis**

Answer Key

Question No.	Correct Option
1	a
2	b
3	b
4	d
5	b
6	a
7	c
8	b
9	b
10	a
11	a
12	a
13	a
14	a

15	c
16	a
17	c
18	b

Detailed Explanations

Solution to Question 1:

The generalized swelling of the left arm of the patient post-surgery for breast carcinoma is suggestive of post-mastectomy lymphedema.

Lymphoedema occurs when interstitial fluid accumulates due to defective lymphatic drainage. It is a common complication of breast cancer treatment, involving axillary lymph node dissection and radiotherapy. Sentinel lymph node dissection (SLND) reduces the risk of lymphedema significantly.

Clinical signs and symptoms are observed in the ipsilateral arm. A significant increase in limb size due to swelling is seen. Other symptoms are heaviness, tightness, numbness, tingling, limb fatigue, and reduced limb mobility.

Diagnosis is predominantly based on the clinical swelling of the arm. The girth of the affected arm >2 centimetres than the normal limb, a 200-mL or more increase in limb volume, or a 5% or greater limb volume change are other diagnostic criteria.

Management:

- Limb elevation, elastic arm stockings, and pneumatic compression devices to reduce swelling.
- Using a low-pH water-based moisturizer to prevent dry skin caused by excess fluid accumulation.
- Manual lymphatic drainage (MLD) by a therapist followed by simple lymphatic drainage (SLD) by the patient daily aids interstitial fluid evacuation.
- Surgical options include lymphaticovenular anastomosis, liposuction, and debulking.

Other options:

Option B: Stewart-Treves syndrome refers to lymphangiosarcoma seen in chronic long-standing lymphedema following mastectomy (more than 10 years). They present as single/multiple bluish-red skin and subcutaneous nodules. They can ulcerate and hemorrhage.

Option C: Axillary venous thrombosis or Paget-Schrotter disease is seen in patients doing rigorous exercise and in weightlifters. The compression of the subclavian vein at the thoracic outlet causes painful swelling of the arm.

Option D: Cellulitis (lymphangitis) is a bacterial infection of the skin and subcutaneous tissue. It is characterized by a painful area of erythematous and edematous tissue.

Solution to Question 2:

The given image of extensive perineal soft tissue destruction with intact testis is suggestive of Fournier's gangrene, which is a mixed flora of aerobic and anaerobic infection.

Fournier's gangrene is a type of necrotizing fasciitis (Dartos, Scarpa, and Colle's fascia) of the perineum, external genitalia, and perianal region. Risk factors include:

- Immunocompromised states such as HIV, alcoholism, diabetes, etc.
- Minor procedures in the perineal area (opening of a periurethral abscess).

Pathogenesis is due to the obliteration of arterioles supplying the scrotal skin caused by fulminating subcutaneous polybacterial infection. After a while, the blackish discolouration of the scrotum (gangrene) with crepitus can be seen (image). The testis is usually not affected.

Clinical features of Fournier's gangrene are scrotal pain, fever, prostration, and pallor. Cellulitis spreads rapidly and may be associated with eschars, necrosis, flaking skin, and crepitus.

Diagnosis of Fournier's gangrene is majorly clinical and is a surgical emergency. It should be aggressively managed by urgent surgical debridement and skin grafting in later stages. The infection is controlled by the administration of broad-spectrum antibiotics.



Other options:

Option A: Anti-gas gangrene serum refers to refined equine antitoxin against *Clostridium* species. Hence, it has no role in Fournier's gangrene.

Option C: Bilateral orchidectomy is not performed as testes are not affected and need not be removed.

Option D: Urinary diversion may be necessary in a few cases, and is not compulsory.

Solution to Question 3:

The clinical picture of the child presenting with difficulty in swallowing and the chest X-rays (AP and lateral) showing a circular homogenous opacity in the neck region is suggestive of a foreign body (coin) lodged in the esophagus.

Foreign bodies commonly ingested by children are coins, disk batteries, and small toy items. In adults, it is usually a food bolus (mostly meat) that gets lodged.

The flat surface of a coin is seen on the anteroposterior view (AP view) and the edge is seen on the lateral view. This indicates that it is oriented in the coronal plane, which is seen when it is lodged in the esophagus (and not in the trachea).

Initial symptoms are choking, gagging, and coughing followed by excessive salivation, dysphagia, food refusal, vomiting, or pain in the neck and throat. Respiratory symptoms such as stridor, wheezing, cyanosis, or dyspnea are seen if the esophageal foreign body impinges on the larynx or tracheal wall.

Imaging with chest and abdominal X-rays should be performed in AP and lateral positions. Chest CT should always be performed if oesophageal perforation is suspected.

Endoscopic removal of the esophageal foreign body is the procedure of choice. It should be performed

- Immediately in esophageal obstruction, disk batteries, and sharp pointed objects.
- Within 12-24h in non-pointed objects, pointed objects in the stomach, and coins in the esophagus.
- Non-urgent removal in objects in stomach >2.5cm in diameter, and disk battery in the stomach (up to 48h)

Other options:

Option A: For coins lodged in the trachea, the edge is seen anteroposteriorly view and the flat side is seen in the lateral view of the X-rays (oriented in the sagittal plane).

Option C: Artifacts refer to an appearance or unintended structure that is not normally seen on the radiograph and is produced by some artificial means. It can occur due to technical or processing errors while taking a radiograph.

Option D: Soft tissue calcification in the neck is due to physiologic or pathological conditions. Commonly encountered soft tissue calcifications in the head and neck region include tonsilloliths, carotid artery calcifications, and calcified stylohyoid ligaments.

Solution to Question 4:

The given clinical vignette of a 75-year-old man, with a low PSA (<10 ng/ml), Gleason score of 6, and a small tumor focus point to a diagnosis of low-risk prostatic carcinoma. The most likely management for this patient would be active surveillance only.

Active surveillance is done when there is:

- Gleason score of 6
- Early-stage disease (T1c)

- Small volume disease on biopsy
- Low PSA (< 10 ng/ml)

Active surveillance protocols include a digital rectal examination and PSA monitoring every 3 to 6 months, with TRUS-guided prostate biopsies every 1 to 3 years.

Gleason grading system evaluates the abnormalities in the glandular architecture of the prostate. It is based on a scale of 1 to 5, with 1 being the most differentiated grade and 5 being the least differentiated. The grades are added to give the Gleason score.

The classification and management of carcinoma prostate are outlined below:

EBRT: External beam radiation therapy, TURP: Transurethral resection of the prostate

Grade	Treatment
Low Risk	Radical prostatectomy (<70 years), Active surveillance (>70 years)
Intermediate Risk	Radical prostatectomy and/or Radical radiotherapy(EBRT) (<70 years) Active surveillance +/- TURP and hormone therapy (>70 years)
High Risk	Early androgen ablation + Radiotherapy + Surgical treatment (< 70 years)
Metastatic disease	Systemic chemotherapy with palliative treatment (<70 years) Androgen ablation therapy and palliative treatment for symptomatic patients

Solution to Question 5:

The given image of the cystoprostatectomy specimen shows a large, fungating growth from the bladder wall involving the entire bladder. The two lobes of the prostate are not affected. The given specimen is that of bladder carcinoma.

Clinical features of bladder carcinoma include painless, gross hematuria (most common symptom), which may cause clots and urinary retention subsequently. Increased frequency and discomfort during urination may be seen.

Depending upon the depth of invasion, bladder carcinoma is divided into 3 groups:

- Non-muscle invasive bladder cancer - does not invade detrusor muscle but can invade lamina propria. It accounts for 70% of cases.
- Muscle-invasive bladder cancer - invades the detrusor muscle
- Flat, non-invasive carcinoma in situ (CIS)

Staging and diagnosis:

It is done by cystoscopy and resection of the tumor. A bimanual examination should be performed after resection of the tumor. Complete urinary tract imaging is required using contrast computed tomography (CT) or magnetic resonance (MR) urography.

Management:

- Non-muscle invasive bladder cancer:
- Transurethral resection of bladder tumor (TURBT) followed by intravesical chemotherapy with mitomycin C or epirubicin
- Immunotherapy with regular intravesical instillations of Bacille Calmette Guerin (BCG) in high-risk tumors.
- Muscle invasive bladder cancer: Radical cystectomy along with urinary diversion procedure and lymph node dissection +/- neoadjuvant chemotherapy

Other options:

Option A: Malakoplakia is a rare inflammatory condition that is typically seen in immunocompromised individuals. The term 'malakoplakia' means 'soft plaques'. General principles of treatment include systemic antibiotics, surgical excision, and limiting immunosuppression. Surgical excision is done for malakoplakia lesions based on the site affected, complications, and if medical treatment fails.

Option B: Prostatic carcinoma can be surgically managed by radical prostatectomy, which does not involve cystectomy. In prostatic carcinoma usually, one or more of the lobes are predominantly involved unlike in the given gross specimen.

Option D: Schistosomiasis of the bladder is a parasitic infection by *Schistosoma hematobium*. On cystoscopy, it can look like nodules, pseudo tubercles, ulceration, sandy patches, and in the end stage can cause squamous cell carcinoma of the bladder.

Solution to Question 6:

The given clinical vignette with abdominal pain and tenderness in the left lumbar region following a road traffic accident is suggestive of blunt renal trauma. The investigation of choice is CECT in a hemodynamically stable patient.

Blunt trauma is the main cause of renal injuries and is generally self-limiting. Radiological investigations are indicated when there is gross haematuria, shock in combination with microscopic haematuria, pediatric renal trauma, and penetrating injuries.

CECT is the investigation of choice. It assesses the extent of the injury, showing laceration, extravasation, surrounding hemorrhage, and vessel injury. It also shows non-renal injuries and effectively stages renal pedicle injuries. Arterial occlusion will manifest as rim enhancement of the normal renal contour.

Blunt trauma is mostly treated conservatively. Most of the penetrating injuries require surgical renal exploration.

Absolute indications for surgical intervention in renal trauma are:

- Massive hemorrhage
- Renal pedicle avulsion (grade 5 injury)
- Expanding, pulsatile, or uncontained retroperitoneal hematoma

Relative indications are:

- Large laceration of the renal pelvis or avulsion of the ureteropelvic junction
- Bowel or pancreatic injuries
- Persistent urinary leakage, postinjury urinoma, or perinephric abscess
- Complete renal artery thrombosis of both kidneys
- Renal hypertension

Other options:

Option D: Exploratory laparotomy is recommended in unstable patients and those with penetrating renal injuries.

Solution to Question 7:

The postoperative image shows a purse-string suture which is suggestive of Thiersch wiring for rectal prolapse.

In Thiersch's wiring, a steel wire or a silastic/nylon tape is placed around the anal canal. It is rarely done now due to post-operative problems like chronic perineal sepsis, anal stenosis, and obstructed defecation.

Rectal prolapse is of two types:

- Partial rectal prolapse - only the mucosal layer of the rectum is prolapsed. It is seen in children and is managed by digital repositioning. If there is a recurrence, then Thiersch's wiring or sclerotherapy is performed.
- Complete rectal prolapse - all the layers of the rectum are involved. It is seen in adults due to the weakening of the pelvic floor. It is managed by using the following approaches:
 - The perineal approach - easy to perform, and has fewer complications but a high recurrence rate. It is preferred in elderly patients with comorbidities. The surgeries done include:
 - Thiersch's wiring
 - Delorme's mucosectomy - In this procedure, the rectal mucosa is stripped circumferentially from the rectum over the length of the prolapse. The underlying muscle is plicated with a series of sutures. The excess rectal mucosa is excised and a mucosal anastomosis is performed.
 - Altemeier's recto sigmoidectomy (option D) - In this procedure, full-thickness resection of the prolapsed rectum is performed, incorporating any associated colonic prolapse. Restoration of colorectal continuity can be performed by hand-sewn/stapled anastomosis. This is the procedure of choice in patients presenting with incarcerated and strangulated prolapse.
 - The abdominal approach - It is difficult to perform and has more complications, but a lower recurrence rate. The surgeries done include:

- Ripstein's procedure - It involves the placement of a prosthetic mesh around the mobilized rectum, with the attachment of the mesh to the presacral fascia below the sacral promontory.
- Well's procedure (option B) - It involves posterior rectopexy.

Other options:


Option A: Stapled hemorrhoidopexy is used in the treatment of internal hemorrhoids, where a circular stapling device is used to excise a cylinder of mucosa and submucosa and simultaneously staple the mucosal ends together.

Solution to Question 8:

The given clinical scenario showing blood at the tip of the urethral meatus following a road traffic accident (RTA) is a clinical hallmark of a urethral injury. The next best step in this scenario is to wait and watch.

Injuries to the urethra are more common in males. They are of:

Part of the urethra	Injury type	Clinical features
Bulbarurethra(distal urethral injury)	Blunt traumato the perineu m	Butterfly hematoma(perineal bruising), urinary retention, and bleeding from the urethral tip
Bulbomembranousjunction(p roximal urethral injury)	Road traffic accidents	Blood at the tip of the meatus and urinary retentionHigh-riding prostateon digital rectal examination (Vermooten sign)



Retrograde urethrography using water-soluble contrast is used for localizing the site of urethral injury.

The patient should be managed with analgesia and antibiotics. If the bladder is full, it should be drained with a catheter placed by percutaneous suprapubic puncture using a Seldinger technique.

Other options:

Option A: Foley catheterization should not be done since it might convert a partial tear into a complete tear.

Option C: Nephrogram is not done but retrograde urethrogram is performed in urethral injury.

Option D: Suprapubic Cystostomy is done only in cases where the bladder is full or the patient is unable to urinate following a pelvic fracture/ RTA.


Solution to Question 9:

The given clinical scenario of a hernia medial to the inferior epigastric artery is suggestive of a direct hernia, for which Lichtenstein mesh repair is the appropriate surgical method for its management.

The contents of a direct hernia protrude outward, forward, and medial to the inferior epigastric artery and the deep inguinal ring. Whereas, the contents of an indirect hernia protrude lateral to the inferior epigastric artery. The sac of an indirect inguinal hernia passes from the deep inguinal ring obliquely towards the superficial inguinal ring and then to the scrotum.

A direct hernia protrudes through the Hesselbach triangle, which is a zone of weakness for direct hernias.

A Lichtenstein mesh repair is currently one of the most frequently used methods for treating an inguinal hernia. It is done for the reconstruction of the inguinal floor in adults. It is an anterior tension-free mesh repair that reduces the chances of recurrence of a hernia. A piece of prosthetic nonabsorbable mesh is used to bridge the defect in this repair.



Other surgical methods used for the repair of an inguinal hernia are:

- Bassini repair - It is done for the reconstruction of the inguinal floor in adults. In this method, a mesh is not used. It is associated with a high recurrence and is not used frequently now. It is used only in cases of hernial strangulation and peritonitis where the use of a mesh is not preferred. It is performed by suturing the transversus abdominis and internal oblique musculoaponeurotic arches/conjoined tendon to the inguinal ligament.
- Shouldice repair - It is a multilayered repair, where the posterior wall of the inguinal canal is strengthened by continuous sutures. It involves double-breasting of the transversalis fascia. It is associated with a very low recurrence rate.
- Gilbert repair (plug and patch repair) - It is an extension of Lichtenstein's original mesh repair and has now become the most commonly performed primary anterior inguinal hernia repair. In this technique, a cone-shaped plug of polypropylene mesh is used. It is deployed like an upside-down umbrella when inserted into the internal inguinal ring to occlude the hernia
- McVay repair (Cooper's ligament repair) is used for the correction of direct inguinal hernia, large indirect hernia, recurrent hernia, femoral hernias (particularly suited for strangulated femoral hernias). In this method, the transversus abdominis aponeurosis is approximated to Cooper's ligament.
- Stoppa-Rives repair - It is used for the repair of large, recurrent, or bilateral hernias. In this method, a sub-umbilical midline incision is given to place a large mesh into the preperitoneal space.
- Herniotomy - It is used to treat inguinal hernia in a child which occurs due to the presence of a patent processus vaginalis. Therefore in herniotomy, the repair of the inguinal ring is not done.

Solution to Question 10:

The given clinical scenario of a patient with h/o road traffic accident (RTA) with no air entry on the right side and hemodynamic instability suggests a diagnosis of tension pneumothorax. This is managed by immediate decompression by insertion of a needle in the appropriate place.


Pneumothorax refers to the presence of excess air in the pleural space. Entrapment of air occurs by:

- Primary spontaneous pneumothorax - due to the spontaneous rupture of apical pleural blebs or cystic spaces immediately below the visceral pleura.
- Secondary pneumothorax - due to the rupture of dilated cystic spaces in lung diseases like chronic obstructive pulmonary disease (COPD)
- Tension pneumothorax - a one-way valve mechanism that directs airflow into the pleural space leading to a higher than atmospheric pressure environment, and prevents expansion of the lung.

Tension pneumothorax is characterized by chest pain, tachypnoea, tachycardia, and hypotension. On examination, the trachea is deviated away from the side of the injury, and breath sounds are absent on the affected lung. A patient can develop subcutaneous emphysema as a complication, secondary to the pneumothorax.

Tension pneumothorax is a surgical emergency and the excessive pressure has to be immediately relieved. Tension pneumothorax is managed by immediate needle decompression in the 5th intercostal space (ICS), followed by definitive management with a chest tube inserted in the same

space.



Other options:

Option B: While IV fluids are a part of the resuscitation protocol, hypotension in a tension pneumothorax is due to an obstructive type of shock, increasing circulating volume will have minimal effect on improving blood pressure. The management of the primary cause, tension pneumothorax, takes higher precedence as the next immediate step.

Option C: Intubation is performed when the patient is unresponsive post chest tube insertion.

Option D: Extended Focused Assessment with Sonography in Trauma (eFAST) is not required before emergency decompression as tension pneumothorax is a clinical diagnosis, but it can be performed following chest tube insertion.


Solution to Question 11:

The given scenario shows abdominal pain with bilious vomiting and high-pitched bowel sounds which are suggestive of intestinal obstruction along with radiographic features of pneumobilia (air within the biliary tree) pointing to the diagnosis of gallstone ileus.

Gallstone ileus is a condition that occurs secondary to the erosion of the gallbladder wall, which results in a fistula to the duodenum. Through this, a large gallstone can pass from the gallbladder to the duodenum and usually gets impacted in the distal ileum (the most common site), where it may cause intestinal obstruction.

Rigler's triad is a characteristic radiological feature seen in gallstone ileus and consists of:

- Pneumobilia - air present in the intestine passes into the biliary tree through the fistula
- Features of small bowel obstruction (abdominal pain, bilious vomiting, hyperechoic bowel sounds)
- Ectopic gallstone - usually in the right iliac fossa



Plain X-rays show air-fluid levels consistent with a small bowel obstruction and pneumobilia can also be visualized on CT imaging.

Gallstone ileus is managed by an explorative laparotomy to relieve the bowel obstruction. The stone is milked away from the site of impacted and then crushed or removed.

Other complications of gallstones include biliary colic, acute and chronic cholecystitis, empyema of the gallbladder, mucocele, perforation, acute cholangitis, and acute pancreatitis.

Other options:

Option B: Adhesive intestinal obstruction is seen after any abdominal surgery or any cause for peritoneal irritation, the features are consistent with bowel obstruction (air-fluid levels, featureless distal ileum) but there is no pneumobilia.

Option C: Ischaemic colitis on contrast-enhanced CT of the abdomen (CECT) shows bowel wall thickening, and signs of vascular insufficiency.

Option D: Diverticulitis on CT abdomen shows bright bowel outpouching, which may be complicated by infection and abscess formation.


Solution to Question 12:

A patient with abdominal pain with sterile pyuria and the abdominal X-ray showing homogenous ground glass-like calcification is suggestive of a putty kidney.

The putty kidney is a characteristic finding of genitourinary tuberculosis (GUTB). It is caused by the dissemination of the organism through the bloodstream and is always a secondary TB (reinfection or reactivation of old TB). Genitourinary TB usually presents as sterile pyuria.

Computed tomography (CT) of the abdomen is the most sensitive modality and is the preferred means of visualizing renal calcifications. Microbiological diagnosis is confirmed by at least 3 consecutive early-morning urine samples showing acid-fast bacilli using Ziehl–Neelsen (ZN) stain and subsequently cultured (gold standard).

GUTB is treated with a short course of anti-tubercular therapy (ATT). Ureteric strictures require double J stenting or percutaneous nephrostomy. Definitive treatment is after 3-6mn of ATT, where a reconstructive procedure (open surgical repair & balloon dilatation) is performed. In case the kidney is poorly functioning, a nephrectomy is done.




Option B: Nephrocalcinosis is the deposition of calcium in the kidney parenchyma or tubules, seen in hypercalcemic conditions like hyperparathyroidism.



Option C: Staghorn calculus (a triple phosphate stone) is found in urinary tract infections (UTI) by urease-producing pathogens like proteus and is not a cause of sterile pyuria.



Option D: Psoas calcification is a muscle calcification that develops after acute renal failure and rhabdomyolysis. It is not associated with sterile pyuria.



Solution to Question 13:


The given clinical scenario of a tumour arising from the pyloric antrum of the stomach which involves the head of the pancreas with liver metastasis is suggestive of metastatic carcinoma of the stomach and is managed by palliative gastrojejunostomy followed by chemotherapy.

Gastric carcinoma has a higher incidence in Japan and affects men more than women. *H.pylori* is a significant association, and other risk factors include smoking and industrial dust ingestion. The most common sites are - the proximal stomach (Western countries) and distal stomach (Japan and the rest of the world).


In the early stages, it usually presents as dyspepsia (non-specific). Vomiting and profound weight loss is seen in advanced cases. Endoscopy and biopsy of the lesion (gold standard) are performed to diagnose gastric carcinoma. Pre-operative staging is done with contrast-enhanced computed tomography (CECT) of the abdominal/pelvis. The most important prognostic indicators in gastric cancer are - lymph node involvement and depth of tumor invasion.


Gastric carcinoma is treated with surgical resection with lymphadenectomy. Based on the tumor location, different types of gastrectomy and anastomoses can be performed.

- Billroth I gastrectomy - the lower half of the stomach is removed + cut end of the stomach is anastomosed to the first part of the duodenum.
- Billroth II gastrectomy - two-thirds of the stomach is removed, the duodenal stump is closed and the stomach is anastomosed to the jejunum.
- Roux-en Y gastrojejunostomy - a loop of jejunum is anastomosed to a proximal gastric pouch



©Marrow





Other options:

Option B: Radical gastrectomy is done in proximal gastric cancer and linitis plastica.

Option C: Whipple's procedure is done for gastric carcinoma that has spread to the head of the pancreas. Since the patient already has metastases to the liver, this will not be the preferred option.

Option D: Gastrectomy with right hepatectomy is done in gastric cancer that has hepatic metastasis.

Solution to Question 14:

This patient has a history of cholecystectomy done for gallstones 2 years ago. Thus, in this patient with choledocholithiasis, the stones found in the common bile duct would most likely be primary stones.

Choledocholithiasis or common bile duct stones are classified based on their point of origin:

- Primary stones are brown pigment stones that arise de novo in the bile duct and are composed of bile pigments and cholesterol.
- Secondary stones (option B) are usually cholesterol stones, which originate in the gall bladder and pass down into the common bile duct. Since this patient has already undergone cholecystectomy more than 2 years ago, secondary stones are unlikely.
- When stones are found in the common bile duct within 2 years of history of cholecystectomy, they are termed retained stones (option D).
- If they are found more than 2 years after cholecystectomy, they are known as recurrent stones (which are mostly primary).

Clinical features: Mostly clinically silent, but may present with biliary colic and even obstructive jaundice. When there is fever, jaundice, and associated right upper quadrant abdominal pain, it is known as Charcot's triad, seen in cholangitis.

Management of choledocholithiasis:

ERCP +/- sphincterotomy followed by laparoscopic cholecystectomy is the most preferred management of choledocholithiasis.

If stones cannot be removed through ERCP and sphincterotomy, then common bile duct exploration is carried out intraoperatively either in an open procedure or via laparoscopy. A T-tube is left in place after choledochotomy.

Solution to Question 15:

The symptom of pain in the thigh while walking and the given angiography image showing common iliac artery obstruction is suggestive of peripheral obstructive vascular disease with aortoiliac obstruction. The best surgical course, in this case, is to perform an aortofemoral bypass.

Clinical findings of aortoiliac obstruction are :

- Intermittent claudication is the debilitating cramp-like pain felt in the buttocks, leg, and thigh muscles due to reduced blood flow.
- Femoral and distal pulses are absent in both limbs.
- Arterial bruit is heard over the aortoiliac region.
- Impotence (Leriche syndrome) may occur. Buttock claudication is present.

As the disease progresses, rest pain, ulceration, gangrene, and slow capillary refilling may be seen.

CT angiography (CTA) and MR angiography (MRA) are specific investigations. Doppler ultrasound, duplex scanning, and digital subtraction percutaneous angiography (DSA) are the other available investigations.

An aortofemoral bypass is the choice of surgical intervention in aortoiliac obstruction. Dacron graft is a collagen-impregnated, knitted graft that is used to perform the proximal aortic anastomosis in aortofemoral bypass. It can also be used in aortic valve repair. In unfit patients, axillofemoral bypass may be done.

Other options:

Option A: Percutaneous Transluminal Angioplasty (PTA) is most useful in the treatment of isolated iliac stenoses (<4cm in length) than in occlusive disease.

Option B: Iliofemoral bypass is done in patients with unilateral occlusion of the distal common iliac or external iliac arteries.

Option D: Aortoiliac bypass more commonly known as the aortoiliac endarterectomy is rarely performed now a days because of greater blood loss associated with this surgery and greater sexual dysfunction caused, additionally it is difficult to perform.

Note: Aortofemoral bypass surgery has long-term patency of 70-80% at 10 years and is often done bilaterally as the occlusion is more likely to be bilateral with one side affected more severely than the other. Bilateral aortofemoral bypass does not add to physiological stress and has equal benefits comparatively.


Solution to Question 16:

The image above shows dilated, tortuous veins of the lower limb. In asymptomatic patients, they are classified as C2A according to the Clinical-Etiology-Anatomy-Pathophysiology (CEAP) classification for chronic venous disorders.

Varicose veins are characterized by the tortuous dilatation of subcutaneous veins resulting from venous insufficiency. Patients have heaviness, burning, and aching over the affected area or whole limb, which increases throughout the day or on prolonged standing. Compression and elevation can relieve the symptoms.

Diagnosed by duplex ultrasound and is treated by modalities such as compression, laser ablation, radiofrequency ablation, endovenous glue, ultrasound-guided foam sclerotherapy, and open surgery.

Image A shows tortuous dilated veins (C2), Image B is atrophie blanche (C4b), and Image C shows fan-shaped telangiectasia (C4c).



The images below show C5 and C6 - healed and active venous ulcers respectively.



Solution to Question 17:

In an alcoholic patient with acute pancreatitis, serum lipase is the enzyme most likely to be elevated.

Acute pancreatitis is often caused by gallstones (70%) or alcoholism (25%). The cardinal symptom is acute-onset severe pain in the upper quadrant of the abdomen. The patient usually presents in shock (hypotension and tachycardia) and tachypnoea. Some may also present with signs of bleeding into the fascial planes leading to discoloration of the sites involved:

- Cullen's sign: Around the umbilicus
- Grey Turner's sign: Flanks
- Fox sign: Inguinal region

- Bryant's sign: Scrotum

The diagnosis is often clinical with a laboratory diagnosis. Though serum amylase (>3 times) is indicative of pancreatitis, serum lipase is preferred over serum amylase as it is more sensitive and specific. Contrast-enhanced tomography (CECT) of the abdomen can be used to exclude pancreatitis if in doubt.

The severity of the acute pancreatitis is assessed using the Ranson and Glasgow scoring system (≥ 3 at 48h of onset of pain). APACHE, SOFA, SAPS, and MODS can also be used to assess the severity of pancreatitis in intensive care (ICU) settings. Pancreatitis is managed conservatively with initial nasogastric drainage followed by intravenous fluids, analgesia, and frequent monitoring of vital signs, urine output, and blood gases.

Other options:

Option A: Serum gamma-glutamyl transferase (GGT) can also be elevated. It is a biomarker for pancreatic disease and alcohol consumption but is not sensitive and specific to acute pancreatitis.

Option B: Carcinoembryonic antigen (CEA), a tumor marker that is elevated in the adenocarcinoma of the pancreas.

Option D: Elevated serum bilirubin is a result of (i) overproduction of bilirubin (E.g: hemolytic anemia) or (ii) impaired uptake by the liver (liver cirrhosis) or (iii) regurgitation of bilirubin (obstructive jaundice). It is not seen in acute pancreatitis.

Solution to Question 18:

The given clinical scenario of acute abdominal pain and fever with chest X-ray in an erect position showing air under the diaphragm (pneumoperitoneum) is a characteristic of hollow viscus perforation.


The most common cause of pneumoperitoneum is bowel perforation ($>90\%$ of the cases). Gastric and posterior duodenal perforations are seen in long-standing cases of peptic ulcers. Intestinal perforations are seen with ischemia, inflammation, diverticula, or obstruction (E.g:- colonic carcinoma). Other causes also include iatrogenic, penetrating injuries, prolonged mechanical ventilation, peritoneal dialysis, etc.

Perforation of a viscus is a life-threatening condition due to leakage of the enteric contents which can lead to acute bacterial peritonitis.

Clinical features of hollow viscus perforation are due to the developed peritonitis. They include abdominal pain, nausea, and vomiting. On examination, the patient has tachycardia, tenderness, rebound tenderness, decreased or absent bowel sounds, and guarding on palpation. Fever may be present.

The patient is initially investigated with an upright chest X-ray which typically shows air beneath the diaphragm. Various other signs for pneumoperitoneum in the supine abdominal X-rays are Rigler's sign, Cupola sign, falciform ligament sign, football sign, and Doge's cap sign. However, CECT is the most sensitive investigation to detect minimal/ suspected pneumoperitoneum along with the cause and site of perforation.

Hollow viscus perforation




Other options:

Option A: Gastric volvulus is a twisting or axial rotation of the stomach along its longitudinal axis (organo-axial) or vertical axis (mesentero-axial). The sudden onset of constant and severe upper abdominal pain, recurrent retching with the production of little vomitus, and the inability to pass an NG tube constitute the Borchardt triad. Plain films of the abdomen reveal a gas-filled viscus in the chest or upper abdomen.



Option C: Liver abscess will mostly have the classic presenting symptoms of fever, jaundice, and right upper quadrant pain, with a tenderness to palpation. Chest X-ray findings generally reflect subdiaphragmatic diseases, such as an elevated right hemidiaphragm, right pleural effusion, or atelectasis.

A chest radiograph shows an air-fluid level in the right upper abdomen (arrow) and multiple nodular opacities in the bilateral lungs (arrowheads)



Option D: Empyema thoracis is an infection of the pleural space. An empyema thoracis on a chest X-ray can mimic a pleural effusion or a lung abscess. The fluid is commonly drained completely with a chest tube or small-bore catheter.

The chest radiograph showing pleural empyema.



Surgery INI-CET 2022

Question 1:

Which of the following is not a perineal approach for the condition shown in the image?



- a) Ripstein's procedure
- b) Delorme's operation
- c) Thiersch operation
- d) Altemeier's procedure

Question 2:

Identify the true statement regarding split skin and full-thickness skin grafts:

- a) 2,3
- b) 2,4
- c) 1,3
- d) 1,4

Question 3:

A 37-year-old unmarried nulliparous woman, having regular intercourse is on oral contraceptive pills. Her mother was diagnosed with carcinoma breast at 60 years of age and her elder sister was diagnosed with carcinoma ovary at 40 years of age. What is the next line of management?

- a) Genetic counselling and screening for BRCA
- b) Routine mammography
- c) Stop taking oral contraceptive pills
- d) Prophylactic surgery

Question 4:

Which of the following causes maximum malnutrition?

- a) Pancreatic fistula
- b) Duodenal fistula
- c) Distal ileal fistula
- d) Colonic fistula


Question 5:

A 50-year-old alcoholic patient presents with a history of waxing and waning jaundice for the past 2 months. His CT examination reveals, dilatation of the common bile duct and pancreatic duct. What is the most probable diagnosis in this patient?

- a) Cholangiocarcinoma
- b) Periapillary carcinoma
- c) Common bile duct stone
- d) Head of pancreas carcinoma

Question 6:


Which investigation is not necessary for the following condition?



- a) Timed barium study
- b) Endoscopy
- c) Esophageal manometry
- d) 24-h pH monitoring

Question 7:

The following procedure is performed for the management of:



- a) Advanced gastric carcinoma
- b) Gallbladder carcinoma

- c) Distal cholangiocarcinoma
- d) Chronic calcific pancreatitis

Question 8:

A patient developed a loss of sensations over the root of the penis following open hernia surgery. Which of the following nerves would you implicate?

- a) Iliohypogastric nerve
- b) Ilioinguinal nerve
- c) Genitofemoral nerve
- d) Medial cutaneous nerve of thigh

Question 9:

Which of the following statements about Psoas abscess are true?

- a) 1 and 2 only
- b) 1, 3, and 4
- c) 1, 2, 3, and 4
- d) 2, 3, and 4

Question 10:

A patient who had grade 3 splenic trauma was managed conservatively in the ICU. Which of the following signs on day 2 would indicate a need for surgery?

- a) Haemoglobin fall from 12g/dL to 10g/dL
- b) Extraperitoneal bladder rupture
- c) Pneumoperitoneum
- d) Air in gallbladder

Question 11:

A 34-year-old woman was brought to the AIIMS emergency following a road traffic accident. She was hemodynamically stable and her GCS score was 3. She was intubated and sent for non-contrast CT which was found to be normal. What is the most likely diagnosis?

- a) Subdural hemorrhage
- b) Extradural hemorrhage
- c) Diffuse axonal injury
- d) Cerebral contusion

Question 12:

Which of the following findings is likely to be seen in this patient?



- a) Decreased posterior tibial & dorsalis pedis pulsations
- b) Loss of sensation over foot
- c) Positive trendelenberg sign
- d) Pain on compression of calf

Question 13:

A 30-year-old male patient presents with a pelvic fracture following blunt abdominal trauma. No blood is seen at the meatus. Catheterization does not produce significant urine output. The radiological image is given below. What is the next step in management?



- a) Suprapubic cystostomy
- b) Perurethral catheterization
- c) Laparotomy
- d) Pelvic packing along with B/L percutaneous nephrostomy

Question 14:

Which of the following would be the most commonly encountered ectopic tissue in Meckel's diverticulum?

- a) Gastric and pancreatic
- b) Colonic and pancreatic
- c) Gastric and colonic
- d) Thyroid and lung

Question 15:

What is true about gall bladder carcinoma?

- a) Palpable abdominal lump is the earliest sign
- b) Earliest sign is jaundice
- c) Clinical presentation of gall stones can be distinguished from gall bladder cancer
- d) Median survival is 6 months

Question 16:

Neurogenic shock is associated with _____.

- a) Hypertension + tachycardia
- b) Hypotension + tachycardia
- c) Hypotension + bradycardia
- d) Hypertension + bradycardia

Question 17:

An old lady sustained a fall and developed a headache and neurological symptoms a few days later. What is your diagnosis?

- a) Multi-infarct dementia
- b) Frontal lobe meningioma
- c) Korsakoff disease
- d) Subdural hematoma

Question 18:

Which of the following is not a part of damage control surgery?

- a) Control of hemorrhage
- b) Resuscitation
- c) Decontamination
- d) Vascular anastomosis

Question 19:

All the following are criteria for SIRS, except _____.

- a) Systolic BP ≤ 90 mmHg
- b) Temperature ≥ 38 degree celsius or ≤ 36 degree celsius
- c) Respiratory rate ≥ 20 bpm
- d) Heart rate ≥ 90 / min

Question 20:

Which is the correct sequence for the correction of hypospadias?

- a) 1->3->2->4
- b) 3->1->2->4
- c) 3->1->4->2
- d) 1->3->4->2

Question 21:

A female patient presents with a lump in her breast. All the following can be used to rule out malignancy, except _____.

- a) Radiological investigations
- b) PET scan
- c) Clinical examination
- d) Histology and cytology

Question 22:

An elderly woman underwent surgical resection of her bowel. After a few months, she presents with easy fatiguability and tiredness. Blood smear evaluation shows macrocytic anemia. Resection of all the following can cause this, except _____.

- a) Jejunum
- b) Duodenum
- c) Ileum
- d) Stomach

Question 23:

Match the following:

- a) A-1, B-2, C-3, D-4
- b) A-4, B-1, C-2, D-3
- c) A-3, B-2, C-1, D-4
- d) A-4, B-3, C-2, D-1

Question 24:

A female patient was diagnosed with gastric cancer. On immunohistopathology, the biopsy sample was found to be positive for CD117 and negative for LCA, cytokeratin, and S100. What is the diagnosis?

- a) GIST
- b) Adenocarcinoma
- c) Hodgkins lymphoma
- d) SCC

Question 25:

Which of these is the most life-threatening injury that can be identified by assessing the breathing component of the patient?

- a) Cervical spine injury
- b) Tension pneumothorax
- c) Blunt cardiac injury
- d) Laryngotracheal injury

Question 26:

A 32-year-old female patient with Graves' disease with eye signs and enlarged thyroid is planned for a total thyroidectomy. What can be given in the preoperative period to reduce intraoperative bleeding in the patient?

- a) Potassium iodide
- b) Propranolol
- c) Propylthiouracil
- d) Betamethasone

Question 27:

In all the following places, hand rub can be used according to standard hand hygiene protocol, except _____.

- a) If the hands are visibly soiled
- b) While moving from a contaminated site to a clean site during patient care
- c) Before donning on gloves
- d) During direct patient contact

Question 28:

Which of the following is true regarding all types of shock?

- a) Anaerobic respiration produces carbon dioxide
- b) Low pH mainly activates lysosomal enzymes
- c) Peripheral vasoconstriction
- d) Renal perfusion is maintained

Question 29:

A patient presents with multiple stab injuries to the chest. On examination, there is reduced chest movement on the right side and tracheal deviation to the left side. Additionally, the neck veins are distended. What is the next best step in the management of this patient?

- a) Resuscitative thoracotomy
- b) ICD placement
- c) Needle thoracocentesis
- d) Finger thoracotomy

Question 30:

Match the following:

- a) A-4, B-1, C-4, D-2
- b) A-3, B-1, C-4, D-2
- c) A-3, B-1, C-2, D-4
- d) A-1, B-4, C-3, D-4


Question 31:

What is the correct sequence of management in a patient who presents to the casualty with an RTA?

- a) 1, 3, 2, 4
- b) 1, 2, 3, 4
- c) 2, 1, 3, 4
- d) 2, 1, 4, 3

Question 32:

A 32-year-old male patient presents to the surgical emergency with a history of peptic ulcer disease and pain abdomen, bloating, and obstipation for 2 days. What is the next line in management?



- a) A and B
- b) A and C
- c) A and D
- d) A, B, and C

Question 33:

A female patient is planned for cholecystectomy. Which of the following is not an advantage of laparoscopic surgery?

- a) Lesser post-operative pain

- b) Cosmetically better
- c) Earlier return to normal physiological function
- d) Longer duration of stay in hospital

Question 34:

What does the magnified number marked on the Foley catheter correspond to?



- a) Inner lumen diameter
- b) Length of the tube
- c) Outer tube diameter
- d) Bulb size

Question 35:

A 49-year-old female presents with a breast lump. Which of the following findings is in accordance with basal-like breast cancer?

- a) ER +, PR +, HER2 -
- b) ER +, PR -, HER2 -
- c) ER -, PR -, HER2 -
- d) ER -, PR -, HER2 +

Question 36:

A patient presented with large intra-abdominal fluid accumulation and was found to have pancreatic necrosis. Which of the following are included in the management of the condition?

- a) 1, 2, 3, 4, and 5
- b) 2, 4, 5 and 6
- c) 1, 2, 3, and 5
- d) 1, 2, 3, 4, 5 and 6

Question 37:

A patient presents with a chest x-ray finding of pneumothorax. Which of the following is not a boundary of the triangle of safety for ICD insertion?

- a) Lateral edge of pectoralis major
- b) Mid-clavicular line
- c) Base of axilla
- d) Lateral border of latissimus dorsi

Question 38:

Which of the following cancers are correctly matched with the criteria for the minimum number of lymph nodes required for pathological staging?

- a) B, C
- b) A, B, C
- c) A, C, D
- d) A, B, C, D

Question 39:

Severe intractable pain in the leg due to sciatica in a patient with carcinoma rectum is due to:

- a) Anterior infiltration in females
- b) Anterior infiltration in males
- c) Lateral infiltration
- d) Posterior infiltration

Question 40:

Identify the correct statement regarding gastric carcinoma.

- a) 1,3,4
- b) 2, 3
- c) 2,3,4
- d) 1,4

Question 41:

All the following are true about acute cholecystitis, except _____.

- a) Gall bladder thickness ≥ 3 mm on USG
- b) Murphy's sign positive
- c) Leukocytosis
- d) Preferential visualization of gall bladder in HIDA scan

Question 42:

Lymphatic drainage from the breast is mostly drained into the axillary lymph nodes. Which of the following is considered the location of the apical lymph nodes?

- a) In the axillary pad of fat at the apex of the axilla
- b) Along the margin of the pectoralis minor
- c) Along the subscapular veins
- d) Anterior in the center

Question 43:

A 10-year-old child with hereditary spherocytosis and hypersplenism presents to the hospital. A splenectomy is planned for this child. All the following vaccines are recommended before splenectomy, except _____.

- a) Pneumococcal vaccine
- b) Typhoid vaccine
- c) Haemophilus influenzae B vaccine

- d) Meningococcal vaccine**

Question 44:

In proximal transverse colon carcinoma, which of the following arteries are resected to prevent the spread of cancer?

- a) Ileocolic, right colic and branch of middle colic**
- b) Ileocolic, left colic, and stem of middle colic**
- c) Ileocolic, right colic, and trunk of middle colic**
- d) Middle colic only**

Question 45:

Which of the following statements are true regarding central venous access and peripheral venous access?

- a) 1,3,4**
- b) 2,3,4**
- c) 1,2,3,4**
- d) 1,2,3**

Question 46:

A patient who had undergone appendicectomy 2 days ago presents with severe abdominal pain. On exploration, there is a collection of blood in the abdomen. Which of the following arteries should have been ligated during surgery?

- a) Ileocolic artery**
- b) Right colic artery**
- c) Superior rectal artery**
- d) Superior mesenteric artery**

Question 47:

A 25-year-old woman presents with a breast lump with no skin changes or induration. On histology, there was a fibroadenoma appearance. Which of the following categories does it fall

under?

- a) BIRADS 1 and 2
- b) BIRADS 2 or 4
- c) BIRADS 2, 3, 4
- d) BIRADS 2

Question 48:

Which of the following is not a contraindication for breast conservative surgery?

- a) Tumor size $\leq$ 4cm, with N1 axillary metastasis
- b) Tumor more than the size of the breast
- c) Prior chest irradiation
- d) Multifocal tumors

Answer Key

Question No.	Correct Option
1	a
2	a
3	a
4	b
5	b
6	d
7	c
8	b
9	c
10	c
11	c
12	b
13	c
14	a
15	d
16	c

17	d
18	d
19	a
20	c
21	b
22	b
23	b
24	a
25	b
26	a
27	a
28	b
29	c
30	c
31	b
32	b
33	d
34	c
35	c
36	c
37	b
38	a
39	a
40	b
41	d
42	b
43	b
44	c
45	c
46	a
47	d
48	a

Detailed Explanations

Solution to Question 1:

The condition shown in the image is a rectal prolapse and Ripstein's procedure is done using the abdominal approach.

Rectal prolapse occurs when a mucosal or full-thickness layer of rectal tissue protrudes through the anal orifice. Rectal prolapse is of two types:

- Partial rectal prolapse: Only the mucosal layer of the rectum is prolapsed. It is seen in children.
- Complete rectal prolapse: All the layers of the rectum are prolapsed. It is seen in adults due to the weakening of the pelvic floor.

Management:

- The first episode of a partial rectal prolapse is managed by digital repositioning. On recurrence, Thiersch's wiring or sclerotherapy are considered.
- Complete rectal prolapse is managed surgically, either by perineal or abdominal approaches:
 1. Perineal approach: Often preferred in patients with high operative risk and limited life expectancy, i.e. in the elderly. It is associated with less morbidity but has a higher recurrence rate.
 - Thiersch's wiring- A steel wire/nylon tape is placed around the anal canal. Largely obsolete now due to problems with chronic perineal sepsis, anal stenosis, and obstructed defecation.
 - Delorme's mucosectomy- The rectal mucosa is stripped circumferentially from the rectum over the length of the prolapse. The underlying muscle is plicated with a series of sutures. The excess rectal mucosa is excised and a mucosal anastomosis is performed.
 - Altemeier's rectosigmoidectomy- The rectum is prolapsed through the anal canal and a full-thickness resection is performed, incorporating any associated colonic prolapse. Restoration of colorectal continuity can be performed by hand-sewn/stapled anastomosis. This is the procedure of choice in patients presenting with incarcerated and strangulated prolapse.
 2. Abdominal approach: Difficult to perform, more complications but a lower recurrence rate.
 - Ripstein's procedure- involves the placement of a prosthetic mesh around the mobilized rectum, with the attachment of the mesh to the presacral fascia (anterior) below the sacral promontory.
 - Well's procedure- involves posterior rectopexy.
 - Suture rectopexy

Solution to Question 2:

Statements 2 and 3 are correct.

Split skin grafts (Thiersch grafts) are partial-thickness skin grafts consisting of the epidermis and a part of the dermis. Whereas, full-thickness grafts (Wolfe grafts) consist of the epidermis along with the entire dermis, with portions of the sweat glands, sebaceous glands, and hair follicles.

Primary contracture of the skin graft occurs due to passive recoil of the elastin fibers of the dermis. It occurs soon after the graft has been harvested.

- Since full-thickness grafts have more dermis (implying more elastin fibers), they undergo more primary contracture.

- Whereas, split skin grafts have less dermis, therefore, undergo less primary contracture.

Secondary contracture of the skin graft occurs due to the pull of myofibroblasts. It occurs over time, when the transplanted graft starts healing.

- Full-thickness grafts have more ability to resist the pull of myofibroblasts, hence they have less secondary contracture.

- Whereas, split skin grafts yield to the pull and develop more secondary contractures.

Advantages of split over full-thickness skin graft:

- Large grafts can be harvested.
- In the case of thinner split-thickness grafts, the donor site heals more rapidly and can be reused in severe cases such as burns.
- Uptake is good. Meshing can be done, i.e. small splits are done in graft, allowing exudates to drain out, thus improving graft take up.
- Meshing also allows the graft to expand.

Advantages of full-thickness over split skin graft:

- It has a lesser degree of post-graft contraction
- Best cosmetic appearance as they include the highest number of skin appendage elements.

Primary contracture	Secondary contracture
Elastic recoil	Pull of myofibroblasts
Less in split-skin grafts	More in split-skin grafts
More in full-thickness grafts	Less in full-thickness grafts

Solution to Question 3:

The given family history of two primary relatives with breast and ovarian cancer with one of the relatives with onset <50 years is suggestive of a woman with a high risk of breast cancer. The next line of management is genetic counselling and screening for BRCA.

Genetic counselling is the process of discussing various genetic tests such as BRCA testing and multiple gene panel testing.

Genetic counselling and testing for genetic mutations (BRCA 1, BRCA 2 and others such as TP53, PTEN) are recommended for high-risk women with:

- Breast cancer diagnosed before age 50
- Triple-negative breast cancer
- Bilateral breast cancer
- Breast and ovarian cancer in the same individual
- Breast cancer in men

- Family history (maternal or paternal) of two or more individuals with breast and ovarian cancer
- A close male relative with breast cancer
- A close relative with early-onset (<50 years) breast or ovarian cancer
- Known BRCA1 or BRCA2 mutation in the family

Other options:

Option B: Routine mammography is not the initial investigation of choice, but is used for surveillance after the genetic screening. Surveillance strategies for women with a family history of breast cancer or ovarian cancer include

- Self-examination of breasts starting at age 18 to 20 years - Each month
- Clinical breast examination starting at age 25 years - Every 6 months
- Mammography starting at the age of 25 years or 10 years earlier to breast cancer onset in a family member - Every year

Option C: Oral contraceptives can be continued in this patient as they don't increase the risk of breast cancer in a woman with a positive family history of breast cancer. It is also useful in reducing the risk of ovarian and endometrial cancers, which are associated with BRCA mutations.

Option D: Prophylactic surgery is advised after the identification of BRCA positivity in high-risk women. It is proven to be more effective than regular surveillance by mammography and MRI. It is usually combined with a prophylactic risk-reducing salpingo-oophorectomy that reduces the risk of ovarian and breast cancers.

Solution to Question 4:

Among the given conditions, maximum malnutrition is caused by duodenal fistulas.

Enterocutaneous fistulas are classified according to their location and volume of daily output.

- Proximal fistulas are associated with higher output, greater fluid and electrolyte loss, and greater loss of digestive capacity.
- Distal fistulas tend to have lower output, making them easier to manage and more likely to close spontaneously.

Although both gastric and duodenal fistulas are proximal fistulas, duodenal fistulas receive an additional 1 to 2 liters of enzymes, electrolytes, and water from the pancreatic and bile ducts. Hence duodenal fistulas are at greater risk for more electrolyte imbalance than gastric fistulas.

The management of enterocutaneous fistula is based on well-established principles known as SNAP which include:

- S - Sepsis elimination and skin protection
- N - A period of parenteral nutrition may well be required
- A - Anatomical assessment
- P - Definitive planned surgery

Solution to Question 5:

In the given scenario, a characteristic history of waxing and waning jaundice along with a dilated common bile duct and pancreatic duct on CT scan, points to a diagnosis of periampullary carcinoma.

Periampullary carcinoma arises from the vicinity of the ampulla of Vater. Patients with hereditary polyposis syndromes, such as familial adenomatous polyposis (FAP) and hereditary nonpolyposis colorectal cancer (HNPCC) have a higher risk of developing it.

Clinical features:

- Jaundice: It intermittently waxes and wanes because, whenever the tumor undergoes necrosis and sloughing, obstruction to bile ducts is partially relieved.
- Weight loss
- Pain: It depends on the location of the tumor and may involve back pain as well due to the involvement of retropancreatic nerves, pancreatic duct obstruction or stasis, and disruption of nerve sheath by the tumor.
- Nausea/vomiting
- Pruritus
- Fever
- Gastrointestinal bleeding

In suspected periampullary disease, a three-phase CT scan should be done routinely.

Management includes resection, along with chemotherapy/ chemoradiotherapy.

Solution to Question 6:

The given barium swallow shows a bird beak appearance and is suggestive of achalasia cardia. 24-h pH monitoring is not necessary for achalasia cardia. It is used in the diagnosis of gastroesophageal reflux disease (GERD).

Achalasia is a primary esophageal motility disorder. It is characterized by the absence of esophageal peristalsis and impaired lower esophageal sphincter (LES) relaxation in response to swallowing. This is due to the loss of ganglion cells in the myenteric (Auerbach's) plexus. These abnormalities cause a functional obstruction at the gastroesophageal junction.

Option A: On performing a timed barium study, barium will be retained for more than five minutes. The fluoroscopic examination will reveal a dilated esophagus that tapers to form a 'bird's beak' at the gastroesophageal junction as seen in the image above.

Option B: Endoscopy is an important investigation carried out to rule out cancer of the gastric cardia.

Option C: Esophageal manometry is the investigation of choice and shows:

- Incomplete LES relaxation in response to swallowing
- High resting LES pressure
- Absent esophageal peristalsis

Solution to Question 7:


The given image shows the Whipple procedure.

The Whipple procedure is a complex surgery done for carcinomas involving the head of the pancreas and distal cholangiocarcinoma. It is done only in cases when:

- The entire duodenum has to be removed e.g. in familial adenomatous polyposis (FAP)
- Tumor encroaches in the first part of the duodenum or the stomach
- A clear resection margin is not achieved by pylorus-preserving pancreatoduodenectomy

The following structures are removed during Whipple's procedure:

- Distal stomach
- Duodenum
- Head of the pancreas
- Gallbladder
- Common bile duct (CBD)
- Regional lymphatics



Solution to Question 8:

Loss of sensation in the root of the penis and adjacent part of the scrotum is suggestive of an injury to the ilioinguinal nerve. It is most commonly injured within the inguinal canal in open hernia repair due to traction or thermal injury.


Other options:

Option A: The iliohypogastric nerve carries the sensory supply from the suprapubic area. Injury to this nerve in the right iliac fossa rarely causes sensory disturbances, as the suprapubic area has a rich nerve supply from other sources too. If the nerve is severed above the anterior superior iliac spine, it can lead to the weakened posterior wall of the inguinal canal resulting in a direct inguinal hernia. The iliohypogastric nerve can also get injured following appendicectomy.

Option C: Injury to the genital branch of the genitofemoral nerve results in loss of sensation in the skin of the scrotum of males, and loss of sensation of the mons pubis and labia majora in females.

Injury to the femoral branch of the genitofemoral nerve causes hypoesthesia over the anterior thigh below the inguinal ligament.

Option D: The medial cutaneous nerve of thigh is a sensory branch of the femoral nerve. Injury to this nerve leads to loss of sensations over the areas supplied by this nerve, which include anterior-medial aspect of the thigh and knee.



Solution to Question 9:

The correct statements regarding psoas abscess are 1, 2, 3, and 4.

Psoas abscess occurs due to a collection of pus due to an infection in the iliopsoas compartment. The compartment is retroperitoneal and hence, the abscess is also called a retroperitoneal abscess. The compartment contains iliacus and iliopsoas muscles.

Psoas abscess is either primary or secondary.

- Tuberculosis of the spine leading to pott's abscess was the leading cause of psoas abscess (statement 2).

- Recently, hematogenous spread of infection from an occult primary site (gastrointestinal or urinary tract) has become common, especially in immunocompromised patients and the elderly (statement 4).


- Due to contiguous extension of infection from neighbouring structures.

Features include

- Back pain and fever.
- Soft, fluctuant swelling pointing towards the groin (hip) (statement 1).
- Pain on passive hip extension or active flexion of the psoas (statement 3).

CT scanning is useful in diagnosis. Percutaneous CT-guided drainage and antibiotic therapy are useful in the treatment of the condition.

Radiological images showing psoas abscess (mass lateral to the vertebral column):



Solution to Question 10:

The presence of pneumoperitoneum on imaging would warrant a laparotomy as it indicates visceral perforation which can lead to secondary peritonitis.

Secondary peritonitis usually requires surgery along with antibiotics.

The spleen is one of the most commonly injured abdominal organs. The severity of the injury is graded by the American association for the surgery of trauma (AAST) spleen injury scale. The management is based on the grade of the injury. Abdominal CT with intravenous contrast is the investigation of choice.

Conservative management is recommended in most cases according to current guidelines. There must be no evidence of active blood loss. Splenic angiography and embolization are also used for non-surgical management.

Surgical management is recommended when there is hemodynamic instability or failure to respond to conservative management. It can be in the form of splenic repair, partial splenectomy, or total splenectomy. Splenectomy is recommended in significant injuries to the hilum, pulverized splenic parenchyma, or any injury >grade 2 with coagulopathy.

Vaccination against encapsulated bacteria must be provided to prevent infection if a splenectomy is done.

Other options:

Option A - The initial hemoglobin level should be carefully assessed to monitor blood loss. Patients who are stabilized after crystalloid infusion can also be managed conservatively but a lower threshold for surgery must be kept. But hemoglobin fall from 12g/dL to 10g/dL is not an indication for laparotomy as the fall in hemoglobin is not marked.

Option B - Extraperitoneal bladder rupture can be managed by decompression with a urinary catheter. Later a cystography can be done to confirm healing.

Option D - Air in the gallbladder can be seen in acute cholecystitis if gas-forming organisms cause secondary bacterial infection (emphysematous cholecystitis). It is managed with antibiotics and intravenous fluids. Laparoscopic cholecystectomy is the surgery of choice.

Solution to Question 11:

The clinical scenario of a patient with a poor GCS score and a normal NCCT brain scan following a road traffic accident is suggestive of diffuse axonal injury.

Diffuse axonal injury (white matter shearing) results from shearing forces, typically from a rotational acceleration of the brain within the rigid cranium. This causes damage to microvasculature and disruption of axons in the subcortical parasagittal white matter, cerebellum, fornix, and internal and external capsule.

Most commonly it occurs in automobile accidents. Patients are usually unconscious from the moment of impact and remain the same until death.

Diffuse axonal injury is characterized by multiple focal lesions with a characteristic distribution:

- CT may be normal.

- MRI is the modality of choice for assessing suspected diffuse axonal injury, even in patients with entirely normal CT of the brain. MRI, especially SWI (susceptibility-weighted imaging) demonstrates small regions of susceptibility artifact or hemorrhage.

Other options:

Option A: Subdural hemorrhage is often due to the rupture of veins that cross the subdural space. In subdural hemorrhage, the CT shows concavo-convex opacity.

Option B: Extradural hemorrhage is mostly due to the rupture of the middle meningeal artery. Since the dura matter tends to adhere to the skull the hemorrhage is seen in CT as a biconvex dense area. They usually do not cross cranial sutures.

Option D: Cerebral contusion occurs due to traumatic brain injury, at the sites that come in direct contact with the skull. Imaging may show microhemorrhages and edema. Usually, no treatment is required.

Solution to Question 12:

In the given image, the lesion present over the plantar surface of the foot is likely to be a neuropathic ulcer. Neuropathic ulcers can lead to loss of sensation over the foot.

A neuropathic ulcer is one that occurs as a result of peripheral neuropathy. In peripheral neuropathy, lack of sensation over the pressure points leads to cumulative microtrauma, minor wounds, or cuts that go unnoticed eventually developing into ulcers. They are painless ulcers that are punched out with regular margins. They are often located on the plantar surface of the foot. Ulcer margins show the presence of calluses.

Other options:

Option A: Decreased posterior tibial and dorsalis pedis pulsations are usually seen in peripheral arterial disease, which causes arterial ulcers that are usually found on the dorsal surface of the foot.

Option C: Positive Brodie-Trendelenburg test is seen in varicose veins. This condition leads to venous ulcers which are usually seen on the medial malleolus of the foot.

Option D: Pain on compression of calf is a sign of deep venous thrombosis.

Solution to Question 13:

The given clinical scenario of blunt abdominal trauma, minimal urine output on catheterization, and the pattern of contrast extravasation in the given image indicate intraperitoneal bladder rupture. The absence of blood at the meatus rules out urethral injury. The preferred next step of management, in this case, is laparotomy.


Bladder rupture may be extraperitoneal or intraperitoneal.

- The bladder is related to the peritoneum only on its superior surface. Therefore, if the bladder rupture is towards the superior surface, the urine leaks into the peritoneal cavity, i.e. urine extravasation will be intraperitoneal.

- Whereas, bladder rupture due to injury to the anterior or lateral surfaces does not rupture the peritoneum. In such cases, the urine leaks into the perivesical space but does not enter the peritoneum, i.e. urine extravasation will be extraperitoneal.

Intraperitoneal bladder rupture is caused by a blow or fall on the distended bladder.


Intraperitoneal bladder rupture manifests as sudden severe pain in the hypogastrium and syncope. There will be an associated increase in BUN and creatinine due to systemic absorption of the extravasated urine and it may also lead to chemical peritonitis.




Investigations :


- CT scan can be done to diagnose bladder rupture.
- A plain erect X-ray may show a ground-glass appearance.
- Intravenous urography (IVU) may confirm a leak.
- Retrograde cystography will confirm the diagnosis.

Intraperitoneal bladder rupture leads to extravasation of the contrast material into the peritoneal cavity, around the bowel loops and paracolic gutters. Hence, an outline of bowel loops can be seen. The image below shows intraperitoneal bladder rupture, with arrows showing contrast material in the peritoneal cavity:



Extraperitoneal bladder rupture leads to extravasation of the contrast material into the perivesical space and the fascia planes anterior to the peritoneum. The image below shows extraperitoneal bladder rupture, with contrast material in the retro-pubic space (no contrast can be seen within the peritoneal cavity).





Solution to Question 14:

The ectopic tissue encountered in Meckel's diverticulum includes gastric and pancreatic tissue.

Meckel's diverticulum is caused by a failure of normal regression of the vitelline duct that develops during weeks 5 to 7 of gestation.

The most common presentation is painless, massive, lower GI bleeds in children younger than 5 years. A Meckel's can be the lead point for ileoileal or ileocolic intussusception. The bleeding Meckel's diverticulum may be confirmed by a 99 mTc-pertechnetate isotope scan to detect gastric mucosa and pancreatic tissue.

Surgical treatment is the definitive therapy for Meckel's diverticulum. A V-shaped diverticulectomy with transverse closure of the ileum can be done.

Solution to Question 15:

Gallbladder carcinoma is a rare malignancy that is seen in the elderly. It is associated with a poor prognosis and median survival is less than 6 months.

Gallbladder carcinoma is indistinguishable from cholecystitis or cholelithiasis (option C). It is usually diagnosed when patients undergo cholecystectomy for gallstones. Its clinical features include right upper quadrant pain, nausea, vomiting, and anorexia. Jaundice, palpable mass, and anorexia are late features (options A and B). CA 19-9 is elevated in the carcinoma gallbladder and is used for monitoring.

The preoperative diagnosis is often made on ultrasonography and followed up by a CT scan or MRI/ Magnetic resonance cholangiopancreatography.

A percutaneous biopsy under radiological guidance is done for pathological examination. PET scanning helps in detecting metastatic disease.


In select cases, a laparoscopic examination is useful in staging the disease. Laparoscopy can also detect peritoneal or liver metastases. The most important prognostic factor is the depth of invasion (T-stage).

Solution to Question 16:

Neurogenic shock is associated with Hypotension + Bradycardia.

Neurogenic shock refers to reduced tissue perfusion due to loss of vasomotor tone to peripheral arterial beds. It is a type of distributive shock.

Neurogenic shock is usually secondary to spinal cord injuries due to vertebral body fractures of the cervical or high thoracic region that disrupts the sympathetic regulation of peripheral vascular tone. Loss of vasoconstrictor impulses results in increased vascular capacitance decreased venous return and decreased cardiac output.



The classic description of neurogenic shock consists of:

- Hypotension associated with bradycardia (absence of reflexive tachycardia due to disrupted sympathetic discharge)
- Warm extremities (due to loss of peripheral vasoconstriction)
- Motor and sensory deficits indicative of a spinal cord injury
- Radiographic evidence of a vertebral column fracture

Solution to Question 17:

The given clinical scenario of an elderly suffering from headache and neurological symptoms following trauma is suggestive of subdural hematoma (SDH).

SDH (acute/chronic) is the accumulation of blood between the arachnoid membrane and the dural layers of the meninges. The source of the bleeding is usually from the rupture of bridging veins running from the cerebral cortex to the dural sinuses. Elderly, alcoholics & patients on anticoagulation are at a higher risk of developing SDH, especially after head trauma including trivial incidents.

Non-contrast CT is the investigation of choice.

Radiographic features of subdural hemorrhage are:

- Typically unilateral (85%).
- Crescent-like distribution (concavo-convex) around the periphery of the brain.
- Not limited by sutures (vs extradural hemorrhage which is limited by sutures)
- Hyperdense lesion on CT: Acute SDH
- Hypodense lesion on CT: Chronic SDH

The given image shows the hyperdense collection seen in acute SDH.



Open craniotomy for the evacuation of acute SDH is indicated for the following:

- Thickness ≥ 1 cm
- Midline shift ≥ 5 mm
- Or GCS drop by two or more points from the time of injury to hospitalization.

Other options:

Option A: Multi-infarct dementia (MID) is a type of vascular dementia occurring due to multiple small infarcts leading to brain tissue injury. It commonly causes memory loss in the elderly.

Option B: Frontal lobe meningiomas are benign brain tumors that develop from the meninges. They cause personality changes that may be mistaken for depression.

Option C: Korsakoff syndrome is a memory disorder that results from vitamin B1 deficiency and is commonly seen in alcoholics. These patients present with impairment in storing new memories, disorientation, and confabulation. Consciousness is preserved.

Solution to Question 18:

Vascular anastomosis is undertaken as a part of definitive surgery, not damage control surgery (DCS).

DCS is minimal intervention done in order to stabilize the patient and followed by definite surgery later. It aims to coordinate staged operative interventions with periods of aggressive resuscitation to salvage trauma patients sustaining major injuries.

The initial focus is hemorrhage control, resuscitation, followed by control and limitation of contamination. It includes an abbreviated laparotomy, temporary packing, and closure of the abdomen in an effort to blunt the physiologic response to prolonged shock and massive hemorrhage.

After the initial steps of DCS, definitive surgery involving vascular anastomosis may be done. This is individualized to the patient and preferred to be completed in the first 24–72 hours.

Note: Vascular anastomosis is the option least likely to be part of the initial stages of DCS. Hence, it is the best option out of the four for this testing point.

Solution to Question 19:

Systolic BP < 90 mmHg is not a criterion for SIRS (systemic inflammatory response syndrome).

SIRS is diagnosed when two of the following are present:

- Hyperthermia (> 38 degrees celsius) or hypothermia (< 36 degrees celsius)
- Tachypnoea (> 20 /min) or arterial carbon dioxide tension (PaCO_2) < 32 mmHg
- Tachycardia (> 90 /min)
- White cell count $> 12 \times 10^9$ /litre or $< 4 \times 10^9$ /litre or $> 10\%$ Band forms

Sepsis is SIRS with a documented infection.

Solution to Question 20:


The correct sequence for the correction of hypospadias is option C: 3- $>$ 1- $>$ 4- $>$ 2

The main principles of surgery in hypospadias, in sequence, include:

- Orthoplasty - Chordee correction
- Urethroplasty - To reposition the urethral opening and tubularize urethra
- Glanuloplasty - Reconstruction of glans
- Meatoplasty - Reconstruction of distal urethral meatus
- Skin cover and scrotoplasty

Hypospadias is the most common congenital abnormality of the urethra. It is characterized by the combination of a ventrally (underside of penis or perineum) placed urethral meatus, a hooded foreskin, and ventral chordee (bending of the penis).

The image given below shows the classification of hypospadias:



Perineal hypospadias is the most severe type of deformity.

Operations for hypospadias are performed for sexual, urinary stream, or cosmetic reasons. It is done between 6 and 18 months of age.

Circumcision should not be done in hypospadias as the foreskin will be used for surgical reconstruction.

Tubularized incised plate (TIP) urethroplasty is the most commonly done procedure. It is done for mid-distal hypospadias.

Solution to Question 21:

In a patient presenting with breast lump or other suspicions of breast carcinoma a triple assessment method is used to rule out malignancy. PET scan is used to assess the spread and sites of distant metastasis, it is not used to rule out malignancy. It is also not a part of the triple assessment.

The triple assessment in the examination of breast lump includes:

- Clinical assessment - Clinical history and physical examination of the lump.
- Radiological examination - USG/ mammography/ both.
- Cytological or histological examination - FNAC (Fine needle aspiration cytology)/ core biopsy of the lump.

Features of malignancy seen on clinical examination can be enlarged axillary lymph nodes, circumferential retraction of the nipple, peau'd orange appearance of the skin and ulcerations over

the breast. Benign features include non-progressive, smooth, circumscribed, soft, mobile lumps without skin changes.

Microcalcifications seen on breast radiography are also highly suggestive of breast carcinoma.

FNAC and biopsy can detect malignant and atypical cells, suggestive of malignancy.

The triple assessment method has a positive predictive value of about 99.99%. In other words, if all three modalities give negative results, breast cancer can be excluded.

Solution to Question 22:


The given clinical vignette indicates a diagnosis of macrocytic anemia due to either vitamin B12 (cyanocobalamin) and/or vitamin B9 (folate) deficiency. The resection of the duodenum is least likely to cause this condition.

The duodenum is involved in iron absorption and hence its resection causes iron deficiency anemia which shows a microcytic hypochromic picture on peripheral blood smear.

The stomach and ileum are involved in the absorption of vitamin B12, and the jejunum is mainly used in folic acid absorption. Hence, a resection of these segments will result in a deficiency of the vitamins which in turn causes megaloblastic anemia with a peripheral smear showing macro-ovalocytes and a hypersegmented neutrophil.

Vitamin B12 (cobalamin), is a water-soluble vitamin derived from animal products such as red meat, dairy, and eggs. Absorption of vitamin B12 begins when pepsin in the stomach releases vitamin B12 from the binding proteins in the food. The released B12 binds to haptocorrin (R-factor), which is a salivary protein. In the duodenum, pancreatic proteases release B12 from haptocorrin, which then binds to intrinsic factors to form the IF- B12 complex. Intrinsic factor (IF) is a glycoprotein secreted by the parietal cells of the stomach. In the terminal ileum, this complex is absorbed by the enterocytes.

Pernicious anemia is a specific form of megaloblastic anemia caused by autoimmune gastritis. It is due to impaired production of the intrinsic factor from the parietal cells. It is a common cause of vitamin B12 deficiency. The other causes of vitamin B12 deficiency are gastrectomy, gastric bypass surgery, inflammatory bowel disease, pancreatic insufficiency and so on.



Vitamin B₉ (folate) is a water-soluble vitamin, and its active form is tetrahydrofolate (THFA). They are mainly derived from liver, yeast, spinach, other greens, and nuts. Its deficiency results in megaloblastic anemia, homocysteinemia, and neural tube defects (spina bifida, Anencephaly) in the fetus.

Absorption of vitamin B₉ is mainly from the upper small intestine, specifically the proximal jejunum.

Solution to Question 23:

The correct answer is option B: A-4, B-1, C-2, D-3

Marjolin ulcer is a rare and aggressive form of squamous cell carcinoma that grows from long-standing venous ulcers, burns, and scars. A wedge biopsy is taken from the edge of the Marjolin's ulcer for diagnosis.

The image below shows a Marjolin's ulcer in a burn scar:

Marjolin ulcer over the dorsum of foot measuring 3 cm x 2 cm	Wedge biopsy
1.5 cm cervical lymph node in the neck	Excision biopsy
Soft tissue sarcoma over mid-thigh	Core needle biopsy
Solitary thyroid nodule	USG guided FNA



In a patient with lymph node pathology, an excisional biopsy is performed as it provides the entire node for histopathological examinations.

Core needle biopsy is performed for:

- Soft tissue sarcomas.
- Breast lumps (Triple assessment).

USG-guided fine-needle aspiration cytology (FNAC) for a solitary thyroid nodule helps to diagnose various benign & malignant pathologies of the thyroid gland. However, this technique cannot differentiate between follicular adenoma and follicular carcinoma of the thyroid.

Solution to Question 24:

The given clinical scenario, along with the immunohistopathology findings showing positivity for CD117 and negativity for LCA, cytokeratin, and S100, is suggestive of gastrointestinal stromal tumors (GIST).

Other IHC markers mentioned are seen in:

- S-100: In melanoma (S-100, HMB-45, Melan-A).
- Cytokeratin: In squamous cell carcinoma (SCC).
- Leukocyte common antigen: In lymphomas.

Solution to Question 25:

Tension pneumothorax should be identified as the most life-threatening injury during the assessment of the breathing component of the primary survey (ABCDE protocol).

Both breathing and ventilation are assessed. Injuries that significantly impair ventilation in the short term include tension pneumothorax, massive hemothorax, open pneumothorax, and flail chest. These injuries should be identified during the primary survey and immediate intervention must be carried out.

Other options:

Options A and D: Cervical spine injury and laryngotracheal injury are identified during the airway assessment.

Option C: Blunt cardiac injury (and dissection of the thoracic aorta) are life-threatening injuries identified during circulation assessment.

Solution to Question 26:

Potassium iodide can be given in the 7-10 days preoperative period to reduce intraoperative bleeding during total thyroidectomy.

Preoperative preparation of a hyperthyroid patient (such as in Graves disease) includes:

- Potassium iodide/ Lugol's iodine reduces the vascularity of the thyroid gland and produces transient remission in a hyperthyroid patient.
- Anti-thyroid medications such as carbimazole/propylthiouracil (Option C) are started 6-8 weeks before the surgery. They help achieve a euthyroid state by decreasing the synthesis of thyroid hormones. The last dose of anti-thyroid medication is given in the evening before the day of surgery.
- Beta blockers such as propranolol/nadolol (Option B) reduce the risk of thyroid storm by blocking the adrenergic receptors. They should be continued for 7 days post-operatively as well. The appropriate dosages are propranolol 40 mg TDS or the longer-acting nadolol 160 mg once daily. They also inhibit the peripheral conversion of T4 to T3.

Option D: Betamethasone is a glucocorticoid that reduces the peripheral conversion of T4 to T3. It is not used preoperatively in thyroidectomy.

Solution to Question 27:


According to standard hand hygiene protocol, hand rub cannot be used if the hands are visibly soiled.

Hand hygiene is the most important way of preventing infection transmission, nosocomial infections and it must be performed by all individuals who are involved in patient care.

Routinely, alcohol-based hand rubs are preferred to soap and water handwash because they are faster, more efficient, and better tolerated by the hands. But, these hand rubs cannot be used in case of visibly soiled hands.

Soap and water hand hygiene method is preferred in cases of soiled hands and while dealing with spore-forming pathogens, including outbreaks of *Clostridium difficile*.

Hand hygiene must be performed in the following cases [WHEN] : (Options B,C and D)




*NOTE: Hand hygiene must be performed in all indications described regardless of whether gloves are used or not.

The following infographic gives the steps for using hand rub [HOW] :



The following infographic gives the steps for washing hands with soap and water [HOW] :



Solution to Question 28:

The true statement among the given options is that low pH mainly activates lysosomal enzymes.

Lysosomes are membraned organelles that enclose all the hydrolyzing enzymes capable of breaking down biological polymers. The lysosomal enzymes are only active in acidic pH (low pH value) and this pH is maintained within the organelle. Hypoxic tissue injury during shock causes membrane damage, and extravasation of the enzymes from lysosomes. Thus, leading to tissue death.

Other options:

Option A: Anaerobic respiration produces lactic acid. Accumulation of lactic acid causes metabolic acidosis.

Option C: Peripheral vasoconstriction is seen in all types of shock, except septic shock.

Option D: Renal perfusion is maintained during the compensated shock stages. There is renal decompensation as shock progresses causing oliguria, and eventually anuria.

Shock is a state of circulatory failure resulting in tissue hypoperfusion and hypoxia. Prolonged shock eventually leads to irreversible tissue damage.

In cases of shock, there is inadequate oxygenation. Thus cells switch from aerobic to anaerobic respiration where lactic acid, not carbon dioxide, is produced. The accumulation of lactic acid in the blood leads to systemic acidosis.



Shock progresses through a series of well-defined stages:

- Initial nonprogressive stage: Compensatory neurohumoral mechanisms come into play to preserve tissue perfusion.


- Progressive stage: Tissue hypoperfusion leads to circulatory and metabolic disturbances. Acidosis occurs due to intracellular anaerobic glycolysis. This causes peripheral pooling, thereby leading to reduced cardiac output and disseminated intravascular coagulation (DIC).

- Irreversible stage: Tissue injury cannot be reversed even with hemodynamic correction. There is worsening myocardial contractility, renal function impairment, and bacteremia (due to the release of intestinal flora into the circulation).


The various types of shock are shown in the images below:




Hypovolemic shock



Obstructive shock



Neurogenic shock



Solution to Question 29:

Multiple stab injuries to the chest, reduced chest movement on the right side, tracheal deviation to the left side, and distension of neck veins suggest a diagnosis of tension pneumothorax on the right side. The next best step in the management would be needle thoracocentesis (emergent management).

ICD placement (Option B) is the definitive management of the tension pneumothorax, inserted after performing the initial emergency procedure.

Tension pneumothorax develops when a "one-way valve" air leak occurs causing air to get sucked in and get collected between the lung and chest wall causing the ipsilateral lung to collapse.

It is characterized by chest pain, tachypnea, tachycardia, hypotension, tracheal deviation away from the side of injury, unilateral absence of breath sound, cyanosis, and neck vein distension.

Tension pneumothorax is a clinical diagnosis and treatment is started without waiting for radiological confirmation.

The next step/ immediate management is chest decompression by rapid insertion of a large-bore cannula on the affected side (needle thoracocentesis) in the appropriate space.

If it is not successful, finger thoracostomy can be tried.

ICD placement is mandatory after needle or finger decompression of the chest and is the definitive management. This should be followed by confirmation of position by chest X-ray.

Recent update (Acc. to ATLS 10th edition):

- In adults, the needle thoracotomy location was changed from the 2nd ICS midclavicular line to the 5th intercostal space (ICS) mid-axillary line.
- In the pediatric age group, thoracotomy location remains the same, at the 2nd ICS midclavicular (MCL) line.

Solution to Question 30:

The correct option is A-3, B-1, C-2, D-4.

Nylon is a monofilament or braided multifilament type non-absorbable suture material. It is a polyamide polymer with low tissue reaction and is used in skin closure, abdominal wall mass closure, hernia repair, plastic surgery, neurosurgery, microsurgery, ophthalmic surgery etc. However, it also loses 15-20% of its tensile strength every year and degrades at the rate of 15-20% every year.

Polyglactin is a braided multifilament of type absorbable suture material. It is a copolymer of lactide and glycolide in a ratio of 90:10, coated with polyglactin and calcium stearate. It is a 'workhorse' suture used in general surgical practices gut anastomoses, vascular ligatures, subcuticular wound closures, and ophthalmic surgery.

Polypropylene is a monofilament, non-absorbable suture material. It is a propylene polymer. It is used in plastic surgery, cardiovascular surgery, hernia repair, ophthalmic surgery, and general surgical subcuticular skin closure. Polypropylene mesh is used for hernia repair.

A. Nylon 3-0	3. Midline laparotomy skin wound closure
B. Polyglactin	1. Bowel anastomosis
C. Polypropylene 1-0	2. Suturing linea alba in the midline
D. Polypropylene 2-0	4. Fixing mesh in hernia surgery

Solution to Question 31:

The correct sequence of management in a patient who presents to the casualty with an RTA is Option B: Cervical spine stabilization, Intubation, IV cannulation, CECT as per the ABCDE protocol.

Solution to Question 32:

The given clinical scenario and the chest X-ray showing air under the diaphragm is suggestive of a hollow viscus perforation occurring secondary to the peptic ulcer disease(PUD) in this patient. The next line of management in this condition is administration of IV fluids (option A) and exploratory laparotomy(option C).


Hollow viscus perforation is a life-threatening condition. In PUD, the perforation most commonly occurs as a complication of anterior duodenal ulcers and gastric ulcers.This results in leakage of gastric contents into the abdomen resulting in acute chemical peritonitis.

Clinical features of perforation include abdominal pain and distension, nausea, vomiting, tachycardia, tenderness, guarding & rigidity, rebound tenderness and decreased or absent bowel sounds. Fever may be present. Sometimes, features of septic shock can be seen at later stages.

An erect xray abdomen shows gas under the diaphragm. If the patient can't stand an x-ray can be taken in lateral decubitus position and may show dilated gas-filled loops of the bowel, suggestive of paralytic ileus.

The below image shows an x-ray radiograph of a hollow viscus perforation:

Hollow viscus perforation



CECT (option B) is performed in hemodynamically stable patients and is the most sensitive investigation to detect minimal/ suspected pneumoperitoneum.

Management of a hollow viscous perforation is mainly resuscitation, IV fluids, analgesics, IV broad-spectrum antibiotics, and emergency laparotomy.

Upper GI endoscopy is done in a patient with upper GI hemorrhage (m/c variceal veins rupture) to diagnose and manage the source of bleeding with techniques such as banding and sclerotherapy (option D).

Solution to Question 33:

Longer duration of stay in the hospital is not an advantage of laparoscopic surgeries.

Laparoscopic surgeries are performed by introducing a rigid laparoscope through a port into the peritoneal cavity. Carbon dioxide is insufflated into the abdominal cavity to create a pneumoperitoneum. The other laparoscopic instruments used for dissection are introduced via other surgically created ports.

Evidence shows that laparoscopic surgery has short-term benefits over open surgery.

Laparoscopic surgery has a better outcome with regard to blood loss, postoperative pain, length of stay, better cosmetic appearance, fewer adhesions, better anatomical perception, and earlier recovery.

The disadvantage of laparoscopic over open surgery is not knowing the extent of the thermal burns due to the relative lack of visibility of the entire instrument.

Solution to Question 34:

The number shown in the above image of the Foley catheter indicates the outer diameter.

"Fr" stands for French scale and refers to the external diameter of the catheter. 1Fr is 0.3 mm in diameter.

An 18 Fr. catheter will have a circumference of approximately 18 mm and a diameter of 6 mm.

Solution to Question 35:


Basal-like or triple-negative breast cancer on immunohistochemical staining is ER-, PR- and HER2-.

Immunohistochemistry staining:

Positive staining is indicated by the presence of brown-colored tissue.

- The estrogen receptor (ER) and progesterone receptor (PR) are both nuclear receptors, so the nucleus is stained brown if positive.
- The human epidermal growth factor receptor type 2 (HER2) is a membrane receptor, so the cell membrane is stained brown if positive.

The image below shows positive and negative staining for ER, PR, and HER2.



Molecular subtypes	ER	PR	HER2	Ki-67	Cytokeratin 5 and 6
Luminal A(Most common type and has the best prognosis)	+ve	+ve	-ve	Low	-ve
Luminal B	+ve	+ve	-ve	High	-ve
	+ve	+ve	+ve	Any	-ve
HER2 enriched	-ve	-ve	+ve	Any	-ve

Molecular subtypes	ER	PR	HER 2	Ki-67	Cytokeratin 5 and 6
Basal-like or Triple-negative breast cancer(Aggressive, has worst prognosis but shows good response to chemotherapy, high PCR rates)	-ve	-ve	-ve	Any	+ve
Unclassified, now also known asClaudin-low	-ve	-ve	-ve	Any	-ve

Solution to Question 36:

The management of acute necrotizing pancreatitis (ANC) includes administering intravenous (IV) fluids, IV antibiotics, culture, and assisted drainage.

ANC is a complication of acute pancreatitis. Necrosis is the presence of nonviable pancreatic parenchyma or peripancreatic fat and can manifest as a focal area or diffuse involvement of the gland.

Steps in the management of ANC:

- Aggressive fluid resuscitation with isotonic crystalloid solution, pain control, and early nutrition, similar to acute pancreatitis
- CECT is done to diagnose ANC. The Balthazar Score is used to grade severity based on CT findings.
- Infection screening:
 - One of the main complications of ANC is infection. If infected necrosis is suspected, fine-needle aspiration (FNA) is done, and the aspirate is subjected to Gram staining and culture sensitivity.
 - Once the infection is confirmed, IV antibiotics are started. Carbapenem is the first-choice drug. The other drugs that can be used include quinolones, metronidazole, third-generation cephalosporins, and piperacillin.
- Definitive management:
 - Traditionally, surgical debridement with necrosectomy, close continuous irrigation, or open packaging. Due to the high mortality, these surgeries are no longer performed, instead, a "step-up" approach has been adopted.
 - Step-up approach: Percutaneous drainage followed by minimally invasive video-assisted retroperitoneal debridement.

Note: Pancreatic necrosectomy via midline laparotomy is considered only if sepsis persists despite conservative measures.

Solution to Question 37:


The mid-clavicular line is not a boundary of the triangle of safety for chest tube insertion.

Chest tube insertion is carried out in the 5th intercostal space (ICS) anterior to the midaxillary line in the triangle of safety (image below).

Boundaries of Triangle of Safety:

- Anteriorly (red line) : Below and lateral to the pectoralis major muscle
- Posteriorly (yellow line) : Anterior to the mid-axillary line (some references mention the lateral border of latissimus dorsi)
- Inferiorly (blue line) : Above the horizontal level of the nipple at the level of the 5th ICS

Triangle of safety




©MARROW

Chest tube insertion/ thoracostomy is indicated in pneumothorax, hemothorax, chylothorax, pleural effusion, etc. Contraindications include pulmonary adhesions, pulmonary trauma, coagulopathy, and diaphragmatic hernia.

Possible complications of this procedure include:

- Clogging of the tube (most common)
- Incorrect placement
- Intercostal nerve/ vessel injury
- Subcutaneous hematoma/ emphysema
- Pain

The image below depicts chest tube insertion in a step-wise manner:



Solution to Question 38:

The minimum number of lymph nodes to be examined for accurate staging of colon carcinoma and gall bladder carcinoma are correctly matched.

The minimum number of lymph nodes to be examined for effective staging are:

Type of Carcinoma	Minimum no. of Lymph nodes
Stomach CA	16
Colon CA	12
Gall Bladder CA	6
Breast CA	10

Solution to Question 39:

The most appropriate and specific answer, in this case, is that sciatica in patients with rectal carcinoma is due to anterior infiltration in females.

Sciatica in rectal carcinoma occurs due to perineural invasion of the inferior hypogastric plexus, from which the sciatic nerve arises. In females, the inferior hypogastric plexus lies anterolateral to the rectum in females, and in males, it lies lateral to the rectum.

With the given options, anterior infiltration in females is the most specific.

Solution to Question 40:

The correct statements regarding gastric carcinoma are 2 and 3.

Most cases of gastric cancer are sporadic (statement 1).


Lauren's system classifies gastric cancer into Type I - intestinal and Type II - diffuse.

At the molecular level,

- the diffuse-type is characterized by decreased E-cadherin mainly due to loss-of-function mutations in the CDH1 gene (statement 2).
- the intestinal type of cancer is characterized by mutations in the APC gene or β -catenin causes increased signaling via the Wnt pathway (statement 3).

Symptoms of both types include nausea, dyspepsia, dysphagia, weight loss, anorexia, early satiety, and anemia.

The image below shows the histopathological appearances of intestinal and diffuse types of gastric carcinoma:




Endoscopy and biopsy of the lesion (gold standard) are performed to diagnose gastric carcinoma.

When possible, surgery remains the preferred treatment approach. Local invasion into the duodenum, pancreas, and retroperitoneum is common, where chemo or radiation therapy with palliative care is the treatment.

The TNM staging is most widely used. It uses depth of invasion along with nodal and distant metastases as prognostic indicators (statement 4).

**General staging and treatment strategy
for gastric adenocarcinoma**



©Marrow

Solution to Question 41:


Preferential visualization of the gall bladder in a hepatobiliary iminodiacetic acid (HIDA) scan is not a feature of acute cholecystitis. On a HIDA scan, non-visualization of the gall bladder even after 3 hours of radioisotope administration suggests the diagnosis of acute cholecystitis.

Acute cholecystitis usually occurs due to the presence of gallstones (90%) leading obstruction of the cystic duct. This leads to inflammation, distension, and edema of the gallbladder. Acalculous cholecystitis and tumor obstructing the cystic duct are rare causes.

Clinical features of acute calculous cholecystitis: Fever, pain and right upper quadrant tenderness. A positive Murphy's sign (Option B) is right upper quadrant tenderness on palpation.

Diagnosis: An ultrasound scan is sensitive in acute cholecystitis and demonstrates a distended gallbladder with a thickened wall of ≥ 3 mm (Option A), pericholecystic fluid, the presence of gallstone(s), and the sonographic Murphy's sign. CT scan findings are similar to USG. CBC will show leukocytosis (Option C).

The following image depicts the confirmatory HIDA (Technetium-99m radioisotope) scan in which the gallbladder is not visualized even after 4 hours.



Management: Patients with acute cholecystitis are kept NPO and started on parenteral antibiotics. Cholecystectomy is the treatment of choice for acute cholecystitis. Treatment including cholecystectomy done within 72 hours has better outcomes than interval surgery.

Solution to Question 42:

The apical group of the axillary lymph nodes are present partly above the superior margin of the pectoralis minor and partly posterior to the superior part of the muscle. They are present along the axillary vessels.


The central group of axillary lymph nodes are present in the axillary pad of fat at the apex of the axilla.

Axillary lymph nodes collectively drain the entire upper limb, breast, and trunk above the umbilicus and can be divided into 5 groups namely:

- Lateral/ humeral
- Anterior/ pectoral
- Posterior/ subscapular
- Central
- Apical

Given below is a diagrammatic representation and details about the individual axillary groups of lymph nodes.

Lymphatic drainage of the axilla



	Location	Afferents	Efferents
Humeral/ Lateral group	Posteromedial to axillary vein	Whole limb except vessels accompanying cephalic veins	• Central nodes • Apical nodes • Inferior deep cervical nodes
Pectoral/ Anterior group	Along lateral thoracic vessels	Skin and muscles of supra-umbilical anterolateral body wall and breast	• Central nodes • Apical nodes
Subscapular/ Posterior group	Along the subscapular vessels	Skin and muscles of the inferior posterior region of the neck, dorsal aspect of the trunk up to iliac crest	• Central nodes • Apical nodes
Central group	Embedded in the axillary fat	All preceding groups	• Apical group
Apical group	Deep to clavipectoral fascia, along the axillary vessels	Those accompanying cephalic vein	Unite as subclavian trunk, right trunk drains into Jugulo-subclavian venous junction and left trunk ends in thoracic duct

Solution to Question 43:

The typhoid vaccine is not part of recommendations for a patient undergoing splenectomy.


Splenic insufficiency following splenectomy leads to increased susceptibility to sepsis caused by encapsulated bacteria such as pneumococci, meningococci, and H. influenzae. Such patients must be vaccinated against these agents to reduce the risk of sepsis.

Solution to Question 44:

Transverse colonic carcinoma is treated with an extended right colectomy. The vessels to be ligated are ileocolic, right colic, and trunk of the middle colic vessels.

Procedures performed for tumors of the colon are summarized below:

Tumor location	Procedure of choice
Cecum/ Ascending colon	Right hemicolectomyLigation and division of the ileocolic, right colic, and right branch of the middle colic vesselsResection of cecum + approx 10 cm of the terminal ileum with ileocecal junction + ascending colon + hepatic flexure
Hepatic flexure/Transverse colon	Extended right hemicolectomyLigation and division of the ileocolic, right colic, and trunk of middle colic vesselsResection of the cecum + approx 10 cm of the terminal ileum with ileocecal junction + ascending colon + hepatic flexure + proximal transverse colon (distal transverse colon and splenic flexure are also sometimes removed)
Splenic flexure/ Descending colon	Left hemicolectomyLigation and division of the left branch of the middle colic vessels, the left colic vessel, and the first branches of the sigmoid vesselsResection of the transverse colon (left MCA territory) + Splenic flexure + Descending colon up to the sigmoid colon
Sigmoid colon	Sigmoid colectomy, or LAR (Low anterior resection)Ligation and division of the sigmoid branches of the inferior mesenteric artery. Removal of the entire sigmoid colon and rectum



Solution to Question 45:

All the given statements are true regarding central and peripheral venous catheters.

Peripheral intravenous cannulation is the insertion of a single-lumen plastic conduit across the skin into a peripheral vein. They are used in the transfusion of fluids, medications, and other therapies such as blood products. Phlebitis is the most common complication in peripheral venous access (Statement 1). Relative contraindications are coagulopathy, local infection, burns, arteriovenous fistula (Statement 3), or deep venous thrombosis on the affected limb.

Central venous catheter insertion is used in critically ill patients. It is inserted into a large, central vein. The most common veins are the internal jugular and subclavian veins. It is used in case of multiple infusions incompatible with peripheral intravenous access, and hemodynamic monitoring. Vasopressors are an integral component in the management of septic shock and are given in the central line (Statement 4).

To prevent or avoid the formation of thrombus in venous catheters, the solutions used include 0.9% sodium chloride, heparin (Statement 2), vitamin C, and sodium citrate.

Solution to Question 46:

In the given scenario, the ileocolic artery should have been ligated.

The right colic, superior rectal and superior mesenteric arteries are not ligated during an appendectomy.



When an appendectomy is performed, the inferior branch of the ileocolic artery needs to be ligated at the base of the appendix. Otherwise, postoperative hemorrhage can occur.

The appendicular artery usually arises directly from the ileocolic artery and descends posterior to the terminal ileum to enter the mesoappendix. The artery runs very close to the free margin of the

mesoappendix and terminates in multiple small branches that supply the appendix.

The terminal part of the artery lies on the wall of the appendix and may become thrombosed in appendicitis, resulting in distal gangrene or necrosis.

The blood supply of the appendix is shown in the image below.



Solution to Question 47:


Fibroadenoma is a benign tumor in a 25-year-old woman with no skin changes and no indurations and is categorized as BIRADS 2.

Fibroadenoma is a form of aberrations of normal development and involution of the breast (ANDI), and it is the most common cause of a breast lump in women aged 15–25 years.

It arises from hyperplasia of a single lobule and presents as a well-circumscribed, smooth or lobular, firm, non-tender, freely mobile mass (mouse in the breast). The histopathology shows the proliferation of intralobular stroma, and the epithelium may be surrounded by stroma (pericanalicular pattern) or compressed and distorted (intracanalicular pattern).

USG is done to confirm the diagnosis. A biopsy is needed if USG shows atypical features or the age of presentation is above 25 years.

The image below shows the histology of fibroadenoma:



Solution to Question 48:

Tumor size ≤ 4 cm, with N1 axillary metastasis is not a contraindication for breast conservative surgery.

Breast conservative therapy (BCT) involves resection of primary breast cancer with adequate normal breast tissue, adjuvant radiotherapy, and assessment of regional lymph nodes via a separate incision in the axilla. Removal of primary breast cancer is also called lumpectomy/ tylectomy/ partial mastectomy/ wide local excision.

Absolute contraindications of BCT:

- Pregnancy
- Persistently positive margin (2 microscopy shows positive margins)
- Multicentric lumps
- Diffuse microcalcification
-

Extensive ductal carcinoma in situ (DCIS)

-

Inflammatory breast cancer


Relative contraindications of BCT:

- Active SLE (systemic lupus erythematosus), scleroderma
- Large lump-to-breast ratio (>5 cm size lump)
- Known or suspected genetic predisposition to breast cancer (i.e., BRCA1, BRCA2), Li-Fraumeni syndrome (p53 mutation)
- Multifocal disease
- Prior radiation therapy to the chest wall or breast

Surgery NEET 2022

Question 1:

A 6-year-old boy came with a history of recurrent urinary tract infections. Imaging was done and is shown below. What is the diagnosis?



- a) Vesicoureteric reflux
- b) Urinary bladder diverticulum
- c) Urinary bladder hernia
- d) Vesicocolic fistula


Question 2:

A 45-year-old female patient underwent thyroidectomy. Three days after the surgery, she developed perioral numbness. Which of the following investigations need to be done for her?

- a) Free T3, T4
- b) T3, T4, Thyroid-stimulating hormone
- c) Radioiodine scan
- d) Calcium, phosphate and parathormone levels

Question 3:

A patient presented with blunt trauma to the abdomen. On evaluation, liver injury was noted for which primary repair was done. Coagulation function was monitored intraoperatively, using the method shown below. What is the method used?



©Marrow

- a) Thromboelastography
- b) Plethysmography
- c) Sonography
- d) Elastography

Question 4:

A 59-year-old lady presents with a progressive painless lump in the breast. What is the cause for the following skin change?



- a) Infiltration of subdermal lymphatics
- b) Infiltration of lactiferous duct
- c) Involvement of Cooper's ligament
- d) Spread of tumor to anterior chest wall

Question 5:

A child had a history of stab injury on the anterior abdominal wall and the image is shown below. The child is hemodynamically stable. Which will be the next course of treatment?



- a) Emergency laparotomy

- b) Observation
- c) Intravenous hydration
- d) Wait and watch


Question 6:

Which of the following is most likely to be seen due to rupture of a saccular aneurysm?

- a) Subdural hemorrhage
- b) Subarachnoid hemorrhage
- c) Intracerebral hemorrhage
- d) Hydrocephalus

Question 7:


A patient presents with sudden-onset right leg pain. An investigation was done, and the obtained image is shown below. What is the investigation?



- a) Digital subtraction angiography
- b) Ultrasound Doppler
- c) MR angiography
- d) Plethysmography

Question 8:


A male patient presented with a 0.3-cm nodule on the left nasolabial fold. It was excised and a pathological examination was done. The image is given below. What is the diagnosis?



- a) Basal cell carcinoma
- b) Melanoma
- c) Squamous cell carcinoma
- d) Nevus

Question 9:

A 45-year-old patient complained of pain in one side of the neck. She is afraid of eating food as it worsens the pain. Ultrasound imaging of the salivary glands is done and shown below. What is the most likely diagnosis?



- a) Sialolithiasis
- b) Foreign body
- c) Osteoma of floor of mouth
- d) Cervical lymphadenopathy

Question 10:

A delayed intravenous urogram of a patient is given below. What is the likely diagnosis?




- a) Pelviureteric junction obstruction
- b) Putty kidney

- c) Staghorn calculus
- d) Cystic kidney

Question 11:


A young male patient with a history of a motor vehicle accident is unable to pass urine. Blood is seen at the meatus. What is the most likely site of urethral injury?



- a) Bulbar urethra
- b) Spongy urethra
- c) Membranous urethra
- d) Penile urethra

Question 12:


A male patient presented with a midline neck swelling. He later developed cervical node enlargement. The histopathology of the lesion is shown below. Which of the following statements is false about this condition?



- a) Excellent prognosis is associated with this condition
- b) It spreads quickly via lymphatics
- c) Nuclear characteristics are used for identification
- d) Fine needle aspiration cytology (FNAC) is not diagnostic

Question 13:

A 40-year-old man who is a known case of hypertension presented with multiple episodes of hematuria and loin pain. His elder brother passed away due to stroke at the age of 40. The ultrasound abdomen is shown below. What is the probable diagnosis?




- a) Renal cell carcinoma

- b) Autosomal dominant polycystic kidney disease
- c) Tuberculosis of the kidney
- d) Autosomal recessive polycystic kidney disease

Question 14:


A patient presents with acute-onset, severe abdominal pain. He is hemodynamically stable. A chest x-ray is shown below. What is the next step in management?



- a) Gastric lavage
- b) Emergency laparotomy
- c) Ultrasound
- d) Chest tube insertion

Question 15:

Calculate the percentage of burns in a preschool child shown in the image below.



- a) 15-20%
- b) 25-30%
- c) 35-40%
- d) 10-15%

Question 16:

A 32-year-old patient who is a chronic alcoholic presents with oral ulcers and a burning sensation. A picture of the oral cavity is given below. What is the most likely diagnosis?



- a) Leukoplakia

- b) Erythroplakia
- c) Submucosal fibrosis
- d) Malakoplakia

Question 17:

A patient presents with a tender pulsatile mass as shown in the image below. What will be done next for this patient?



- a) Ultrasound doppler
- b) Needle aspiration
- c) CT angiogram with percutaneous management
- d) Intravenous antibiotics for 7 days

Question 18:

A patient presents with carcinoma of the prostate, which has now spread to the vertebra. What is the route of spread to the lumbar vertebra?

- a) Prostatic venous plexus
- b) Transcoelomic spread
- c) Inferior vesical vein
- d) Internal iliac vein

Question 19:

A male patient presents with impotence and bilateral resting pain. The following finding is seen in the legs. At which level has the pathology occurred?



- a) Bilateral popliteal arteries
- b) Bilateral internal iliac arteries
- c) Aortoiliac bifurcation
- d) Bilateral femoral arteries

Question 20:


An elderly male patient presented with a sudden onset of scrotal pain and discharge. Based on the image below, what is the likely diagnosis?



- a) Torsion of testis
- b) Acute epididymo-orchitis
- c) Scrotal carcinoma
- d) Fournier's gangrene

Question 21:

Identify the fluid shown below.




- a) Haemaccel
- b) Ringer lactate

- c) Isolyte P
- d) Isolyte M

Question 22:


The image below shows a pressure sore. Which stage does this belong to ?



- a) Stage 1
- b) Stage 2
- c) Stage 3
- d) Stage 4

Question 23:

Identify the most common site of an intra-abdominal abscess.



- a) 1
- b) 2
- c) 3
- d) 4

Question 24:

A man came with complaints of recurrent discharge and pain due to lesions around the anus for 3 years. What is the diagnosis?



- a) Pilonidal sinus

- b) Carbuncle
- c) Boil
- d) Fistula in ano

Question 25:

According to triage, which of the following categories of patients comes under green?

- a) Ambulatory patients
- b) Medium risk patients
- c) High risk patients
- d) Dead patients

Answer Key

Question No.	Correct Option
1	a
2	d
3	a
4	a
5	a
6	b
7	b
8	a
9	a
10	a
11	c
12	d
13	b
14	b
15	b
16	a
17	a
18	a
19	c

20	d
21	b
22	d
23	d
24	d
25	a

Detailed Explanations

Solution to Question 1:

In the given scenario of a 6-year-old boy with recurrent urinary tract infections and the micturating cystourethrogram showing a dilated ureter, the likely diagnosis is vesicoureteric reflux.

Vesicoureteric or vesicoureteral reflux (VUR) is a condition in which there is a retrograde urinary flow from the bladder to the ureter and kidney. It is usually congenital.

VUR can cause urinary stasis and lead to urinary tract infection (UTI). It also predisposes to pyelonephritis, which can lead to renal scarring due to parenchymal inflammation. This is known as reflux nephropathy.

Micturating cystogram is used to confirm the VUR. VUR can be graded based on the appearance of the urinary tract on the micturating cystogram. It is as follows:


Grade 1- reflux into the ureter

Grade 2- reflux into the ureter and renal pelvis

Grade 3- mild to moderate dilatation on IVU along with reflux and mild blunting of calices

Grade 4- blunting of fornices

Grade 5- absent papillary impressions



Grades 1-3 resolve spontaneously. Surgical management is done when acute pyelonephritis recurs in spite of antibiotic therapy or if severe reflux is associated with a defect that can be corrected surgically.

Other options:

Option B: Urinary bladder diverticula occurs due to pressure changes in the bladder leading to the formation of saccules that may protrude through the bladder wall. It is more common in elderly males >50 years and is usually asymptomatic or not infected. Symptoms may occur due to associated urinary obstruction, recurrent infection, or pyelonephritis. They are diagnosed incidentally on cystoscopy or ultrasonography.

Option C: Urinary bladder hernia is an uncommon condition characterized by herniation of the urinary bladder or ureter into the inguinal canal, scrotum, or femoral canal. Urinary symptoms such as dysuria, frequency, urgency, nocturia, and hematuria can be seen. Retrograde cystography is the best technique to image a bladder hernia.

Option D: Vesicocolic fistula is an abnormal communication between the bladder and the colon. As a result of this, the fecal matter may enter the bladder. Diverticulitis is the most common cause. Bubbles in the urine, pain, hematuria, and recurrent UTI can occur. Cystoscopy reveals edema and gas bubbles. The fistula can be seen in a cystogram.

Solution to Question 2:

Calcium, phosphate, and parathyroid hormone (PTH) levels should be done in this patient to evaluate hypoparathyroidism secondary to thyroidectomy.

Hypoparathyroidism is a common complication of thyroid surgery. It is temporary and usually lasts for 4-6 weeks. Hypoparathyroidism following thyroid surgery can be diagnosed when there are decreased serum calcium and/ or PTH levels, or if there is symptomatic hypocalcemia.

The features of hypocalcemia are circumoral paresthesia, and paresthesia of the neck, fingers, and toes. Muscle twitching, weakness of the forearm, hand, foot, and carpopedal spasms can be

seen. A carpal spasm can be induced by inflating a blood pressure cuff to 20 mmHg above the systolic blood pressure for 3 minutes. This is known as Trousseau's sign. Chvostek sign wherein twitching of facial muscles is demonstrated by tapping above the angle of the jaw. It is due to the stimulation of the branches of the facial nerve.


ECG shows prolonged QT interval.

Management of acute and symptomatic hypocalcemia is by giving 10 mL of intravenous 10% calcium gluconate over 5 minutes.

Solution to Question 3:

In the case of liver injury, intraoperative monitoring of coagulation function is done by thromboelastography which is shown in the given image.

Thromboelastography (TEG) is a non-invasive test that quantifies the ability of whole blood to form a clot. It is useful in intraoperative coagulation status assessment. The dynamic changes in the viscoelastic properties of a blood sample during the clotting can be detected and quantified by thromboelastography. Underlying coagulation deficiencies show different shapes in the TEG. They are shown in the image as follows



Advantages of thromboelastography:

-

Blood coagulability, including coagulation cascade, platelet function, and fibrinolysis can be assessed.

-

It is a simple bedside test

-

Can diagnose coagulopathic bleeding

-

Can be used as a guide in transfusion therapy and decrease the use of blood products

-

Can detect dynamic changes in blood coagulation during resuscitation

-

Predict the clinical efficacy of therapeutic agents affecting blood coagulability

Other options:

Option B: Plethysmography is a semiquantitative method to measure segmental blood flow and velocity in the carotid and peripheral vascular systems. It is safe, inexpensive and easy to use. It is useful in the diagnosis of vascular disease.

Option C: Sonography or ultrasound is a cheap and easily available imaging method. It is used as a first-line imaging method to assess liver, biliary, renal tract, obstetric, and gynecological disease. It is also useful in guiding needle placement during interventional procedures. Ligaments, tendon and muscle injuries can also be studied by an ultrasound.

Option D: Elastography is a newer imaging modality, which detects the changes in elasticity of tissues. It is based on the concept that the pathological process alters the elastic properties of tissue. It is useful in differentiating malignant and benign breast neoplasms, identifying traumatic changes in the muscles and tendons, helping to decide the biopsy site accurately, assess liver fibrosis and steatosis.

Solution to Question 4:

The image shows the peau d'orange appearance and it occurs due to the infiltration of the subdermal lymphatics.

Peau d'orange appearance is a skin change in which the skin of the breast appears like that of an orange. It is due to cutaneous lymphatic edema caused by dermal lymphatic infiltration by malignant cells. It is seen in several conditions such as:

- inflammatory carcinoma of the breast
- mastitis
- chronic abscess
- following breast surgery or radiation

Other options:

Option B: Infiltration of lactiferous duct leads to nipple retraction.

Option C: Involvement of Cooper's ligament leads to dimpling of the skin over the breast.

Option D: In case of spread of the tumor to the anterior chest wall, the breast will appear fixed and will not fall forwards when bending. On lifting the arm above the shoulder, the breast will not move upward.


Solution to Question 5:

The next course of treatment in the given scenario of a stab injury to the abdomen with the image showing evisceration of the omentum is emergency laparotomy.

Emergency laparotomy is performed to explore and repair abdominal injuries in case of

- Hemodynamic instability
- Peritonitis
- Evisceration
- Gunshot wounds

The algorithm for management of stab wounds of the abdomen is as follows



Solution to Question 6:

Subarachnoid hemorrhage is most likely to be seen due to the rupture of a saccular aneurysm.

Saccular aneurysms are thin-walled outpouchings measuring 2 or 3 cm in diameter with a bright, red, translucent wall. These are mainly seen at the arterial branching points along the circle of Willis.

The absence of smooth muscle and internal elastic lamina is the main pathogenesis in saccular aneurysms. Female sex, smoking, cocaine abuse, and family history are some other risk factors.


Genetic conditions predisposing to subarachnoid hemorrhage are:

- Adult polycystic kidney disease
- Fibromuscular dysplasia
- Neurofibromatosis type 1
- Ehler Danlos syndrome and Marfan's syndrome.

Clinically, subarachnoid hemorrhage presents with a sudden and severe headache commonly referred to as thunderclap headache. It may be associated with seizures, unresponsiveness, and vomiting.

CT scan is the imaging method of choice. It shows blood in basal cisterns and aneurysms appear as round enhancing lesions. Management includes craniotomy and clipping or coiling by endovascular embolization. Medical management includes fluid replacement with normal saline, oral nimodipine 60 mg every 4 hours, and supportive measures.

Hyponatremia can be a frequent complication in subarachnoid hemorrhage. The below image shows a CT of subarachnoid hemorrhage with blood in basal cisterns.



Other options:

Option A: Subdural hemorrhage or subdural hematoma occurs mainly due to rupture of the cortical bridging veins. Symptoms of raised intracranial pressure (ICP) such as headache, vomiting, blurring of vision, as well as focal deficits are seen.

Option C: Intracerebral hemorrhages may spontaneously occur in patients with hypertension, amyloid angiopathy, coagulation disorders, or as a complication of ischemic stroke. Headache, raised ICP, and focal neurological deficits can be present.

Option D: Hydrocephalus is a condition where the cerebrospinal fluid (CSF) circulation is obstructed or there is a failure to resorb thereby leading to an increase in CSF volume with enlargement of the ventricles. Symptoms of raised ICP are seen.

Solution to Question 7:

The image shows a triphasic pattern of the arterial waveform, which is seen in the ultrasound doppler.

Doppler ultrasound is used in diagnosing peripheral arterial disease. It uses a handheld doppler ultrasound probe placed at an arterial site. This transmits a continuous wave. A receiver inside the probe picks up the reflected beam. The change in frequency between the reflected beam and the

transmitted beam gets converted to a pulsatile audio signal.

A normal arterial wave has a triphasic signal. A triphasic wave has three waveforms consisting of a sharp systolic wave with a forward uprise and fall, a second wave with reverse flow occurring in diastole, and the last wave with a forward flow taking place during diastole.

Doppler ultrasound can detect moving blood. However, it does not provide information on whether the blood flow is sufficient for the maintenance of limb viability.

Other options:

Option A: Digital subtraction percutaneous angiography (DSA) is a procedure wherein a radiopaque dye is usually injected percutaneously into the femoral artery. Images obtained are digitalized and the background may be removed for a clear image. It provides information on dynamic arterial flow and can be combined with endovascular intervention if needed.

Option C: Plethysmography is a semi-quantitative method to measure segmental blood flow and velocity in the carotid and peripheral vascular systems.

Option D: Magnetic resonance angiography (MRA), or MR angiography, is an imaging modality that can be used when duplex ultrasonography cannot be done or if it cannot produce a clear image. MRA avoids ionizing radiation.

Solution to Question 8:

The given clinical scenario of a nodule on the nasolabial fold and the histopathological image showing peripheral palisading is likely to be basal cell carcinoma (BCC).

Basal cell carcinoma is a slow-growing malignant tumor, with locally invasive characteristics. It affects the pilosebaceous skin.

Ultraviolet radiation exposure is the strongest predisposing factor. Other risk factors include exposure to arsenic, coal tar, aromatic hydrocarbons, and ionizing radiation.


It mainly affects the elderly or middle-aged men who are excessively exposed to sun, common in the age group 40-80 and white-skinned people.

Macroscopically, it is categorized as localized and generalized. Nodular, nodulocystic (nodular and nodulocystic together account for 90% of BCC), pigmented, and nevoid are the localized cancers while superficial multifocal and superficial spreading, infiltrative, and cicatrizing are the generalized types. Nodular type is described as pearly waxy nodules at the margins. A central depression with umbilication or ulceration and rolled-out edges are characteristic of BCC.

The below gross image shows pearly, rolled-out borders suggestive of basal cell carcinoma.



Histopathologic features show ovoid cells in nests with a single palisading layer. The outer layer of cells is in the active division and thus incomplete excision can lead to being more aggressive. The below image shows the histopathology of BCC.




Sites near the eye, nose, and ear and size ≥ 2 cm are considered to be high risk. Direct invasion at these sites can reach the cranium.

Management is by Moh's microscopic surgery. Non-surgical measures such as topical 5-fluorouracil, and imiquimod are useful. Metastasis is extremely rare.


Other options:

Option B: Melanoma is a malignancy, commonly seen in young adults, arising from melanocytes and involves the skin, mucosa, retina, and leptomeninges. Microscopically, malignant changes are seen in the basal epidermis with breach and invasion of the dermis as shown below.



Option C: Squamous cell carcinoma is the second most common malignant tumor, seen more commonly in men. Microscopically, the proliferation of irregular masses of squamous epithelium is seen. The image is shown below.

Squamous Cell Carcinoma



Option D: Nevus are aggregates of melanocytes at the dermis or dermo-epidermal junction. It can get transformed into malignant melanoma and the features suggestive of malignant transformation are a change in size, shape, color, thickness, satellite lesions, and tingling/itching/serosanguinous discharge.

Solution to Question 9:

The given clinical scenario of pain on one side of the neck that worsens with food and the ultrasound image showing a stone within Wharton's duct is suggestive of sialolithiasis (salivary

gland stone).

The submandibular gland has highly viscous secretions and hence forms the most common site for salivary gland stone formation. The two most common sites of stone impaction are near the punctum and the hilum of the gland where the duct bends over the mylohyoid muscle.


The patient usually presents with acute painful swelling in the submandibular region, which is aggravated with food intake. Symptoms resolve 1-2 hours after the meal is completed. On examination, a tender and enlarged submandibular gland can be noted on bimanual examination. Pus may be seen draining from the sublingual papilla or can be expressed on bimanual palpation of the gland.


Most submandibular stones are radio-opaque and visible in radiographs. Stones of size ≤ 4 mm are removed via a dormia basket using endoscopic or ultrasound-guided methods under local anesthesia. Stones of size > 4 mm are broken first using intracorporeal or extracorporeal lithotripsy and then removed using a dormia basket.

Submandibular gland excision is done only when minimally invasive techniques have failed or there is evidence of a salivary gland tumor.

Nerves at risk during the removal of the submandibular gland:

- the marginal mandibular branch of the facial nerve
- the lingual nerve
- the hypoglossal nerve





Solution to Question 10:

The delayed intravenous urogram (IVU) of the patient showing hydronephrosis with an absence of contrast in the ureter makes the likely diagnosis a pelviureteric junction obstruction.

Pelviureteric junction obstruction is the commonest type of urinary obstruction in childhood, but can also be seen in young adults. It is usually caused by intrinsic stenosis. It manifests as abdominal flank or back pain, and hematuria. In newborns, a renal mass is palpable.

Antenatal sonography is the least invasive technique to diagnose PUJ obstruction in utero and reveals fetal hydronephrosis. intravenous urography is useful if the kidney is functioning. Extrarenal pelvis dilatation, clubbing of the minor calyces, and slow diffusion of contrast down the blockage can be seen. A renal isotope scan using intravenous DTPA or MAG-3 is the best modality to confirm the obstructive dilatation of the collecting system.

The treatment is surgical if there are bouts of renal pain, increasing hydronephrosis, evidence of parenchymal damage, and infection.

Other options:

Option B: Putty kidney is an end-stage manifestation of renal tuberculosis where there is amorphous dystrophic calcification involving the entire kidney. It is shown in the image below.



Option C: Staghorn calculus is seen following urinary infections with *Proteus*, *Pseudomonas*, and *Staphylococcus*. These organisms can break down urea to produce ammonia and carbon dioxide. The urine becomes alkaline and forms struvite calculi which grow into staghorn calculi. A staghorn calculus in the left kidney is shown in the image below.



Option D: Cystic kidney is seen in autosomal dominant polycystic kidney disease (ADPKD) due to mutation in genes PKD1 and PKD2. It is associated with hypertension. Multiple renal bilateral cysts will be seen, along with cysts in other organs as well such as the liver, pancreas, etc.

Solution to Question 11:

In the given scenario of a young patient with urinary retention, blood at the meatus, and a history of a motor vehicle accident, the most likely site of injury is the membranous urethra. The given

retrograde urethrogram shows contrast leakage at the membranous urethra.

Urethral injuries in men usually occur after catheterization or during cystoscopy. Injuries following external trauma are rare, but mainly affect the bulbar or the membranous urethra.

Membranous urethral rupture usually occurs following a road traffic accident, crush injury, or a fall. It is associated with pelvic fracture and extraperitoneal bladder rupture. Clinical features include urinary retention, blood at the meatus, and a high riding prostate on digital examination. Bruising is seen over the pubic area, scrotum, and penis.

Management is by inserting a suprapubic catheter. If there is an associated bladder rupture, the bladder will not be seen on the ultrasound, and management in such a case is by exploration, repair, and placement of a suprapubic catheter.

Other options:

Option A: Bulbar urethral injury occurs following a blow to the perineum with a fall astride injury during cycling or gymnasium accidents. A typical feature is a butterfly-shaped hematoma in the perineum. Other features such as urinary retention and meatal bleeding can also be present. Analgesia, antibiotics, and suprapubic catheterization are needed. The patient should be discouraged from passing urine. Definitive management is delayed urethroplasty after 8-12 weeks.

Options B and D: Spongy and penile urethral rupture can occur following a penile fracture during coitus. Hematuria and difficulty in voiding can be present in spongy urethral injury. A pop sound followed by severe penile pain and penile deformity is characteristic of penile urethral injury. Hematoma, blood at the meatus, and voiding difficulties are also seen.

Retrograde urethrogram shows loss of smooth contour of the urethra with extravasation of the dye in case of a spongy urethral rupture. Surgical repair is needed in both.

Solution to Question 12:

The given scenario of a male patient with a midline swelling, cervical lymph node enlargement, and the histopathology showing branched papillae with orphan Annie eye nuclei is suggestive of papillary thyroid carcinoma. The false statement regarding the condition is that FNAC is not diagnostic.

Papillary thyroid carcinoma is the most common thyroid malignancy. It is more commonly seen in women between the ages of 30 and 50. Radiation exposure and family history of thyroid malignancies are the common etiologies. Genetic abnormalities in either the RET gene or the BRAF gene can be seen in papillary thyroid carcinoma.

Clinical features include an asymptomatic thyroid swelling with cervical lymph node enlargement. In advanced disease, there may be features like hoarseness, dysphagia, cough, or dyspnea.

An ultrasound scan accurately assesses the size and number of nodules, as well as categorizes the lesions as benign, indeterminate, or malignant. Fine needle aspiration cytology (FNAC) is a diagnostic test, helpful in investigating an indeterminate or malignant lesion.

Histopathology shows the following features:

- Branching papillae
- Orphan Annie eye nuclei: optically clear nuclei with fine nuclear chromatin

- Pseudo inclusions: invaginations of the nuclear membrane
- Psammoma bodies: concentric calcifications in the core of the papillae.

Papillary thyroid carcinoma is known to spread via lymphatics, hence lymph node metastasis is commonly seen.

Management is by thyroidectomy with neck dissection (central or local depending upon the site involved) followed by lifelong levothyroxine supplementation. Radioactive iodine is also useful.

The disease has an excellent prognosis, with a high survival rate.

Solution to Question 13:

This clinical scenario with a family history of stroke and USG showing a cystic appearance of the kidney is suggestive of autosomal dominant polycystic kidney disease (ADPKD).

ADPKD is inherited as an autosomal dominant trait with mutations in the PKD1(chromosome 16) and PKD2 genes (chromosome 4).

The disease does not usually manifest before 30 years of age, and in some patients, it is never diagnosed. The progression of the disease is faster in males.

Hypertension is the most common presentation. Renal manifestations of ADPKD include:

- Flank/loin pain - either due to hemorrhage inside cysts or due to renal stones (most commonly uric acid stones)
- Irregular upper quadrant renal mass
- Hematuria
- Uremia
- Renal failure (seen in 10–15% of the patients who receive hemodialysis)

The most common extrarenal manifestation is hepatic cysts, which are derived from the biliary epithelium. Other extra-renal manifestations of ADPKD:

- Pancreatic cysts
- Berry aneurysms (Rupture of cerebral aneurysms is an important cause of morbidity and The prevalence of intracranial aneurysms in patients with PKD is 40%.)
- Colonic diverticula
- Abdominal hernias
- Mitral valve prolapse

Screening cerebral angiography is strongly considered in all patients with PKD and a family history of cerebral aneurysm or subarachnoid hemorrhage.

Intravenous pyelography shows characteristic spider leg sign, as a result of stretched calyces due to multiple cysts. The nephrogram shows a swiss cheese appearance. On prenatal USG, the presence of either > 3 cysts in unilateral kidneys or > 2 cysts in each kidney is suggestive of PKD.

Treatment includes

- Surgical decompression is not a routine procedure performed in these patients. It may be required only in cases of very large cysts causing persistent pain and infected cysts.
- Analgesics
- Increased fluid intake
- Prevents renal stone formation
- Reduces growth of cysts by suppressing ADH release
- ACE inhibitors and ARBs
- Blood pressure control
- Slowing of cyst growth
- Tolvaptan (V2 receptor antagonist)
- Reduce pain and slow the decline in renal function
- Long-term dialysis
- Renal transplant
- Definitive management
- Pre-transplant nephrectomy may be required in patients with very large polycystic kidneys

Autosomal recessive polycystic kidney disease (ARPKD) occurs secondary to a mutation of the PKHD1 gene. The severe form manifests in-utero or during infancy; milder cases can manifest later in childhood. 50% of affected newborns die in the first few days of life.

Renal cell carcinoma (RCC) is a relatively rare complication of ADPKD. There does not seem to be an increased frequency of RCC in these patients compared to the general population. However, when it occurs, it tends to be bilateral, multicentric, and of sarcomatoid type.

Tuberculosis kidney presents as putty kidney characterized by amorphous dystrophic calcification.


Solution to Question 14:

The given clinical scenario and the chest radiograph showing air under the diaphragm are suggestive of a hollow viscus perforation. The next best step in management among the given options is emergency laparotomy.

Perforation of a viscus is a life-threatening condition that occurs following the rupture of the alimentary tract. This leads to the leakage of the enteric contents, thereby leading to acute bacterial peritonitis. Peritonitis is the generalized or localized inflammation of the peritoneum either due to the free spillage of fluids into the peritoneal cavity or secondary bacterial infection.

Clinical features of hollow viscus perforation occur due to the developed peritonitis and the features include abdominal pain and distension, nausea, vomiting, tachycardia, tenderness, rebound tenderness, decreased or absent bowel sounds, and guarding on palpation. Fever may be present. Sometimes, features of septic shock can be seen at later stages.

Investigations include estimating the white blood cell count, urea, serum electrolytes, and an erect abdominal radiograph showing free gas under the diaphragm. A supine radiograph may show dilated gas-filled loops of the bowel, suggestive of paralytic ileus. The below image shows a hollow viscus perforation.



Management of a hollow viscous perforation is mainly resuscitation, analgesics, intravenous broad-spectrum antibiotics, and emergency laparotomy. Peritonitis is managed by nutritional support, pain relief, correction of fluid loss and circulating volume, urinary catheterization with or without gastrointestinal compression, antibiotics, analgesics, and specific surgical treatment of the cause.


Solution to Question 15:

From the given image, the percentage of burns in the child is 25-30% calculated based on Lund and Browder chart.

In the given image, the right arm, chest, abdomen, and both thighs are involved in burns. According to Lund and Browder chart, 2% is for the right arm, 13% for chest and abdomen involvement, and 4% for each thigh. Hence, the total involvement is $2+13+4+4 = 23\%$. The closest value to 23% among the given options is 25-30% (option B).

Lund and Browder chart is an accurate method to measure burns if it is used correctly. In this, different proportions of body surface areas in children are taken into consideration. This chart compensates for the changes in the shape of the body with age and can accurately assess the burns area in children.

The below image shows the Lund and Browder chart.



Note: Wallace's rule of nines is used in estimating medium to large burns in adults. It helps in the quick estimation of the burns. In this, the total burn is calculated by dividing the body into areas of 9%. It is not accurate in children.

Solution to Question 16:

The most likely diagnosis in the given scenario of a 32-year-old chronic alcoholic with oral ulcers and a burning sensation and the image showing white lesions on the oral cavity is leukoplakia.

Leukoplakia is a premalignant condition of the oral cavity characterized by whitish patches or plaques that cannot be rubbed off.

They appear as small, well-circumscribed lesions and can extensively involve a large area of the oral mucosa.

Variants of leukoplakia are

- Speckled leukoplakia is whitish lesions on an erythematous base. They have the highest risk of malignant transformation.
- Proliferative verrucous leukoplakia is rare and has a high risk for malignant transformation.

Management includes smoking cessation and reduction in alcohol consumption. A biopsy should be done and a photographic record of the lesion is used for follow-up.

Other premalignant conditions of the oral cavity are erythroplakia and oral submucosal fibrosis.

Erythroplakia (option B) refers to irregular, reddish velvety plaques on the oral cavity. The surface may be nodular and may bleed. It has a high risk of malignant transformation. Biopsy should be done followed by surgical ablation.

Oral submucosal fibrosis (option C) is characterized by the formation of fibrous bands beneath the oral cavity. The disease is progressive and causes limitation of mouth opening and restricted tongue movements. Intralesional steroids and surgical excision of fibrous bands are recommended.

Malakoplakia (Option D) is a rare inflammatory condition affecting immunocompromised individuals. It mainly affects the urinary tract but can involve any organ system. The clinical presentation includes tender ulcerations, nodules, papules, and plaques or fungating masses with perianal and genital skin involvement. Systemic antibiotics and surgical excision are recommended.

Solution to Question 17:

The given scenario of a patient with tender pulsatile mass, and the image showing an erythematous swelling at the base of the neck on the left side, is likely to be a vascular anomaly. Ultrasound Doppler should be done next in this patient.

The diagnosis of vascular anomalies is done clinically and with imaging like ultrasound Doppler. MRI plays an important role in diagnosing the type and extent of the lesion. CT angiogram can be used to find out the feeding vessel (option C).

Vascular anomalies include malformations and tumors. Vascular malformations occur as a result of developmental abnormality of an artery, vein, capillary or lymphatic system. They can be pulsatile or have a thrill when the underlying component is an artery.

Hemangiomas are the most common vascular tumors. They are seen in infants and present as bright red lesions. Common sites of a hemangioma include the head and neck. It can be of two types capillary and cavernous hemangioma. Compressibility and bluish surface are some clinical features of hemangiomas.

These lesions can be treated by complete resection. Debulking is done when complete resection is not possible. Sclerotherapy is mainly useful in venous malformation. Embolization can be done after surgical resection of the lesion.

Note: Cirroid aneurysm is a slow-growing, tortuous swelling over the scalp or face that develops due to abnormal arteriovenous connections in the scalp and extremities. They are usually congenital. However, in some cases, they may be traumatic. The superficial temporal artery is commonly involved in cirroid aneurysms of the scalp.

Solution to Question 18:

The route of the spread of prostate carcinoma to the lumbar vertebra is by prostatic venous plexus or Batson plexus.

Batson venous plexus is a valveless vertebral venous plexus communicating with prostatic venous plexus.

Carcinoma of the prostate metastasizes to the bone usually by the bloodstream through this Batson plexus. Long bones and the lower lumbar vertebrae are usually involved. Other routes of spread of prostate carcinoma are local spread to the seminal vesicles, bladder neck, trigone, and lymphatic spread.

Carcinoma of the prostate is the most common malignancy in men above 65 years. The clinical features of prostate cancer are bladder outlet obstruction, pelvic pain, hematuria, arthritis,

anemia, pancytopenia, and renal failure. Per rectal examination reveals a hard, nodular, irregularly indurated gland.

Prostate-specific antigen (PSA) is useful in assessing the follow-up. PSA >10 ng/mL is suggestive of cancer and levels >35 ng/mL are diagnostic of advanced prostate cancer. The prostatic biopsy is indicated when there is a raised PSA level or suspicion of prostate cancer.

Localized cancer is treated by radical prostatectomy and radiation therapy, while in advanced cancer palliative and hormone ablation should be carried out.

Solution to Question 19:

The given scenario of a male patient with impotence, bilateral resting pain, and the image showing a gangrenous foot with amputated toes is suggestive of Leriche's syndrome and it indicates that the level of occlusion is at the aortoiliac bifurcation.

Peripheral arterial diseases occur mainly due to atherosclerosis or thromboembolism. Leriche's syndrome is a rare condition in which aortoiliac obstruction causes impotence and claudication (crampy pain) in the thigh or buttocks. Other features of aortoiliac arterial occlusion include absent femoral and distal pulses and a bruit over the aortoiliac region.

In general, features of chronic lower limb arterial occlusion are intermittent claudication, rest pain, ulcers, gangrene, absence of arterial pulses, bruit, and delayed capillary refilling. Rest pain occurs when limb perfusion is so severely compromised that anaerobic respiration occurs even at rest. This causes pain that increases on lying down or elevation of the foot. It reduces when the foot hangs below.

Investigations that are useful include Doppler ultrasound, duplex scanning, ankle-brachial pressure index, and digital subtraction percutaneous angiography (DSA). Through a DSA, endovascular intervention can be performed along with obtaining information on the dynamic arterial flow.

Management is by exercise, smoking cessation, control of diabetes, and weight loss if obese. Drugs such as HMG CoA reductase inhibitors and 75 mg of clopidogrel or aspirin should be given. Transluminal angioplasty and stenting are useful.

Other options:

Options A and D: Femoropopliteal occlusion can lead to claudication in the calf with a palpable femoral pulse but absent distal pulses.

Option B: Iliac obstruction causes thigh and calf claudication with a bruit over the aortoiliac region. The femoral and distal pulses are absent.

Solution to Question 20:

The given scenario of an elderly male patient with sudden-onset scrotal pain and discharge, with the image showing a blackish discoloration of the scrotum, is likely to be Fournier's gangrene.

Fournier's gangrene is an uncommon polymicrobial infection involving the perineal soft tissue, external genitalia, and perianal region. There is subcutaneous tissue inflammation leading to obliterative arteritis and gangrene. It typically follows minor injuries or procedures in the perineal area, urethral dilatation, and injection for hemorrhoids are some of the etiological factors.

Clinically, sudden scrotal pain with associated prostration, pallor, and pyrexia can be seen. Other features such as crepitus and foul-smelling exudate can be present.

Emergency surgical debridement is needed after initial measures like fluid resuscitation and broad-spectrum intravenous antibiotics are undertaken. Urinary diversion and other supportive measures are useful in the management of the condition.

Other options:

Option A: Torsion of testis is a surgical emergency typically seen in the age group of 10-25 years. It is characterized by sudden groin and lower abdominal pain, nausea, and vomiting. Examination findings reveal a swollen, tender scrotum and testis with a twisted tender cord above it and loss of cremasteric reflex.

Option B: Acute epididymo-orchitis is an infection of the epididymis and testis. It is seen in sexually-active young men. In older men, it is seen following a urinary tract infection, secondary to an indwelling catheter. The features include groin ache, fever, swollen and painful epididymis and testis.

Option C: Scrotal carcinoma is a rare carcinoma that is seen in workers exposed to oil, coal products, and aromatic hydrocarbons. It starts as a wart or ulcer in the scrotum and can extend to the testis.

Solution to Question 21:

The fluid shown in the image is ringer lactate or Hartmann's solution.

Ringer lactate, or Hartmann's solution, is a crystalloid made up of sodium 131 mEq, chloride 112 mEq, potassium 5 mEq, bicarbonate (as lactate) 29 mEq, and 2 mEq of calcium. It is devoid of glucose.

It is useful in the following conditions:

- Correction of severe hypovolemia
- Metabolic acidosis
- Postoperative period, burns, diarrhea
- Correction of hypokalemia
- Diabetic ketoacidosis.

It should be avoided in severe shock, liver diseases, vomiting, nasogastric aspiration, and metabolic alkalosis. It cannot be used along with blood transfusion since calcium in the ringer lactate can bind with citrate in the transfused blood leading to clot formation.

Other options:

Option A: Haemaccel is a colloid and contains 145 mmol of sodium, 6.25 mmol of calcium, and 5.1 mmol of potassium. It is mainly used for rapid volume replacement in circulatory shock, burns, and trauma. These release histamine and hence, H₂ blocker has to be supplemented. It is also used in heart-lung machines.

Options C and D: Isolyte P and Isolyte M are crystalloids. Isolyte P is commonly used in fluid therapy for pediatrics. One liter contains 50 grams of glucose, 25 mEq of sodium, 22 mEq of potassium, 3 mEq of magnesium, and 23 mEq of acetate. Hyponatremia, renal failure, hyperkalemia, and hypovolemic shock are some of the contraindications.

Isolyte M is used for maintenance. It mainly contains 50 grams of glucose, 40 mEq sodium, 38 mEq chloride, 35 mEq potassium, 15 mEq phosphate, and 20 mEq acetate. It is useful in correcting hypokalemia in patients with good renal function, and acidosis. It is contraindicated in hyponatremia.

Solution to Question 22:

The given image shows a pressure sore with extensive tissue destruction, including damage to the muscle layer and thus it belongs to stage 4.

Pressure sores also referred known as bed sores, pressure ulcers and decubitus ulcers are ulcers that occur due to prolonged pressure on certain sites thereby leading to tissue necrosis. The commonest sites involved are ischium, greater trochanter, sacrum, heel, malleolus (lateral & medial), and occiput. Paraplegic patients, elderly patients, and severely ill patients are at a higher risk of developing pressure sores.

They are staged as:

- Stage 1 - Erythema with no blanching and no breach in the epidermis.
- Stage 2- Partial-thickness skin loss with epidermal and dermal involvement.
- Stage 3- Full-thickness skin loss with intact fascia but the involvement of the subcutaneous tissue.
- Stage 4- Full-thickness skin loss extending to the fascia with massive tissue destruction and involvement of muscle, bone, tendon, or joint.

Management is primarily proper skin care with pressure dispersion cushions, air-fluidized beds, and urinary or fecal diversion if required. Patients who are bed-bound should be turned once in 2 hours. Those in a wheelchair should lift themselves off the seat for 10 seconds every 10 minutes.

Surgery is the last resort and is done after a proper initial debridement and vacuum-assisted closure. Vacuum-assisted closure is a wound closure technique in which negative pressure is applied to the wound which leads to a decrease in edema. This hastens the debridement and formation of granulation tissue. Skin flaps with a muscle are also used for closure.

Solution to Question 23:


Among the marked parts, 4 represents the right subhepatic space. It is the most common site of an intra-abdominal abscess.

The right subhepatic space is a deep space situated in Rutherford Morison's pouch. It is located between the liver above and the transverse colon and the hepatic flexure below. The relations of the right subhepatic space are:

- the right lobe of the liver and diaphragm on the right side
- foramen of Winslow on the left
- duodenum below
- liver and the gall bladder in the front
- right kidney and the diaphragm behind.

Abscesses here usually arise from appendicitis, cholecystitis, perforated duodenal ulcer, or as sequelae of an upper abdominal surgery.

Intraperitoneal abscesses may develop following intraperitoneal sepsis in specific sites in the abdomen (intra-abdominal abscess) or pelvis (pelvic abscess). The intra-abdominal abscess may occur in subphrenic spaces, or the subhepatic spaces, on either the right or left side.



An intra-abdominal abscess is a localized collection of infected fluid within the abdominal cavity due to failure of the host's intra-abdominal defense mechanisms. Patients usually present with acute abdominal pain, fever, rigors, tachycardia, tachypnea, and leukocytosis. Other features such as nausea, anorexia, emesis, constipation, and diarrhea can occur.

CT scan with IV contrast is the imaging method of choice for an intra-abdominal abscess. Ultrasonography can also be useful.

Initial management is by early intravenous antibiotic therapy, drainage of the abscess, and fluid resuscitation with crystalloids. Vasopressors, inotropes, and packed red blood cell transfusion are given when needed.

Other measures include ultrasound or CT-guided percutaneous drainage (non-surgical) and operative drainage. Resection and debridement can also be done. It can be done through a

minimally invasive technique or laparotomy.

Solution to Question 24:


The given clinical scenario of recurrent pain and discharge due to lesions around the anus, shown in the image is likely to be a fistula-in-ano.

A fistula-in-ano is an abnormal connection between the anal canal and the perianal skin, lined with granulation tissue. It may follow the drainage of a perianal abscess or may be seen in association with diseases like tuberculosis or Crohn's disease.

Clinical features are perianal pain, swelling, and intermittent purulent discharge, which might be blood-stained. On examination, one or more external openings may be visible.

Goodsall's rule can be used as a guide in determining the location of the internal opening of the fistula. Fistulas with an external opening anteriorly connect to the internal opening by a short, radial tract. Fistulas with an external opening posteriorly track in a curvilinear fashion to the posterior midline.

However, exceptions to this rule often occur. Fistulas with an anterior external opening greater than 3 cm from the anal margin usually track to the posterior midline.



The investigations done for fistula-in-ano can include a fistulogram, endoanal ultrasound with hydrogen peroxide, MRI, and computed tomography (CT) scan. An MRI is the gold standard investigation for fistula imaging. It helps us to identify the secondary extensions of the track, which when missed may cause persistence of the fistula.

Surgical management includes fistulotomy, fistulectomy, ligation of the intersphincteric fistula tract (LIFT), setons, or advancement flaps.

Solution to Question 25:

Ambulatory patients that are stable are likely to survive even if treatment is delayed. Their medical need is not urgent and they come under green according to triage.

Triage is a method employed to identify the patients in whom treatment at the earliest produces the maximum benefit to the patient. It is used in the management of mass casualties.

Triage categories are based on the vital signs, patient's ability to walk, mental status, and presence or absence of ventilation/capillary perfusion. It is as follows:

Priority	Color	Medical need	Clinical status
First	Red	Immediate	Critical but can survive with early treatment
Second	Yellow	Urgent	Critical but can survive when treatment is given within hours
Third	Green	Non-urgent	Patients are stable and can survive with delayed treatment
Last	Black	Unsalvageable	Not breathing, no pulse, severely injured and medical care is unlikely to help.

Surgery NEET 2023

Question 1:


A 55-year-old male presented with verrucous carcinoma around the glans of the penis. Examination reveals that the inguinal lymph nodes are not enlarged. What is the appropriate management for this patient?



- a) Total penectomy.
- b) CO₂ laser excision
- c) Topical 5-fluorouracil
- d) Partial penectomy.

Question 2:

Identify the procedure being performed in the image given below.



©Marrow

- a) Intraosseous cannula for pain relief
- b) Bone marrow aspiration
- c) Intraosseous route access for giving IV fluids
- d) Intraosseous abscess drainage

Question 3:

A diabetic patient presents with sudden-onset perineal pain. On examination, foul-smelling discharge, and necrotic tissue as seen in the image is noted. Which of the following is true about the given condition?



- a) Anti-gas gangrene serum indicated for all cases

- b) Urinary diversion is the next step
- c) Bilateral orchidectomy must be done
- d) Mixed aerobic and anaerobic infection

Question 4:

Which of the following is the most common complication following ligation of the first vessel during abdominoperineal resection for rectal carcinoma?

- a) Parasympathetic - bladder dysfunction and retrograde ejaculation.
- b) Sympathetic - bladder dysfunction and impotence.
- c) Sympathetic - retrograde ejaculation and bladder dysfunction.
- d) Sympathetic - Impotence and loss of cutaneous sensation in perineal region

Question 5:

A patient presents to the clinic with the lesion given in the image. He had a traumatic injury to the chest one year ago. What is the most likely diagnosis?



- a) Hemangioma
- b) Hypertrophic scar
- c) Keloid
- d) Neurofibroma

Question 6:

A 45-year-old male underwent bilateral laparoscopic hernia repair for inguinal hernia. Postoperatively, he complained of pain in the right thigh. Which of the following nerve entrapment leads to this symptom?

- a) Ilioinguinal nerve
- b) Iliohypogastric nerve
- c) Femoral nerve
- d) Lateral cutaneous nerve of thigh

Question 7:

A patient was brought to the ER following a road traffic accident. On examination, the patient opens his eyes to a painful stimulus, speaks inappropriately, and withdraws his limbs to a painful stimulus. What is his GCS score?

- a) E2V2M3
- b) E3V3M3
- c) E2V3M4
- d) E3V2M2

Question 8:

A patient presented with fever and abdominal pain with jaundice. Investigations showed the given findings. Which of the following statements is true regarding this condition?



- a) Surgery is mandatory
- b) Fine-needle aspiration cytology is diagnostic
- c) Angioembolization is the treatment of choice
- d) Echinococcus species is involved in etiology

Question 9:

A 40-year-old male patient presents with a swelling that has been slowly growing in the past 2 years. On examination, it was variable in consistency and fully mobile. What is the most likely diagnosis?



- a) Dermoid cyst

- b) Parotid tumor
- c) Sebaceous cyst
- d) Cervical lymph node

Question 10:

A patient who is a known case of alcohol dependence syndrome presents with sudden and unintentional weight loss. What is the most likely diagnosis?

- a) Hepatic adenoma
- b) Cholangiocarcinoma
- c) Hepatocellular carcinoma
- d) Alcoholic hepatitis

Question 11:

A patient presented with features of chronic pancreatitis with recurrent attacks and has a 10 mm dilatation of the pancreatic duct with intraductal calculi present. Which of the following is the surgery of choice?

- a) Pancreaticoduodenectomy
- b) Longitudinal pancreaticojejunostomy
- c) ERCP and sphincterotomy
- d) Coring of pancreas head

Question 12:

A woman presented with a BIRADS-5 breast lesion. Which of the following is a good prognostic factor for this lesion?

- a) BRCA-1 positive
- b) p53 positive
- c) ER positive
- d) High Ki-67

Question 13:

A patient presents with sudden onset of chest pain shooting to the neck and interscapular region. X-ray shows widened mediastinum. BP is 110/90 mmHg in the right upper limb and 160/100 mmHg in the left upper limb. What is the most likely diagnosis?

- a) Acute coronary syndrome
- b) Acute pulmonary embolism
- c) Acute aortic dissection
- d) Esophageal rupture

Question 14:

A patient presented with right hypochondriac pain. He had an episode of diarrhea 1 week prior. CT scan of the abdomen reveals a liver abscess of around 25 ccs. What is the next step in management?

- a) PAIR
- b) Surgery
- c) Medical therapy
- d) Percutaneous drainage

Question 15:

A 50-year-old patient with a history of trauma one week back presents with confusion and ataxia. He is on treatment for alcohol dependence and his last intake was 2 weeks back. CT scan reveals the following finding. What is the diagnosis?



- a) Extradural hemorrhage
- b) Subdural hemorrhage
- c) Intra-parenchymal hemorrhage
- d) Subarachnoid hemorrhage

Question 16:

Which of the following is false about the given condition?



- a) Repeated radical surgery every 24 hours may be required to adequately manage this condition
- b) This is a case of Meleney's gangrene
- c) The condition is caused by beta hemolytic streptococcus and may sometimes be polymicrobial
- d) Hyperbaric oxygen has no role in the treatment of this condition

Question 17:

A patient presents with a swelling in the groin region and it is diagnosed as a direct inguinal hernia. Weakness in which of the following structures is most likely to cause this condition?

- a) Pectineal ligament
- b) Conjoint tendon
- c) Reflected part of inguinal ligament
- d) Lacunar ligament

Question 18:

A patient presents with sudden onset of hematemesis. He has been taking aspirin for his arthritis and drinks alcohol occasionally. He gives a history of occasional abdominal pain. On examination, there is no abdominal mass or tenderness. What is the likely diagnosis?

- a) Mallory-Weiss tear
- b) Esophagitis
- c) Peptic ulcer
- d) Esophageal varices

Question 19:

A construction worker came to the OPD complaining of white fingers. He has been working in the cement and concrete industry and has been working on heavy machinery drills, wood, and furniture polishing for 20 years. What is the most likely etiology for this condition?

- a) Candidal infection of the fingers due to continuous exposure to water
- b) Exposure to thinners and paints
- c) Continuous exposure to cement and concrete
- d) Continuous exposure to drills and machines

Question 20:

A 5-year-old child was admitted to the hospital for prolapsing rectal mass and painless rectal bleeding. Histopathological examination reveals enlarged and inflamed glands filled with mucin. What is the likely diagnosis?

- a) Adenoma
- b) Choristoma
- c) Hamartoma
- d) Carcinoma

Question 21:

Identify the given condition.



- a) Venous ulcer
- b) Malignant ulcer
- c) Arterial ulcer
- d) Trophic ulcer

Question 22:

A patient with varicose veins comes with complaints of eczema over the same region. What is the stage of the clinical classification according to the CEAP guidelines?



- a) C2

- b) C₃
- c) C_{4a}
- d) C_{4b}

Question 23:

A patient comes to the casualty with a severe headache. His BP was found to be 160/100 mmHg. CT scan revealed a subarachnoid hemorrhage. What is the next best step in the management of this patient?

- a) Nimodipine
- b) Angiography
- c) Surgery
- d) Fibrinolytic therapy

Question 24:

A 20-year-old female patient consumed a cleaning product containing 90% sodium hydroxide. She is having complete dysphagia. What is the best step in management?

- a) Esophagojejunostomy
- b) Stent placement
- c) Feeding jejunostomy
- d) Gastrojejunostomy

Question 25:

A 23-year-old male patient presents with midline swelling in the neck. The swelling moves with deglutition and protrusion of the tongue. What is the likely diagnosis?



- a) Brachial cyst
- b) Thyroglossal cyst
- c) Plunging ranula
- d) Dermoid cyst

Question 26:

A 54-year-old woman presents with cervical cancer stage 2A. It is decided to give neoadjuvant chemotherapy. What does neoadjuvant chemotherapy mean?

- a) Chemotherapy given along with radiation
- b) Chemotherapy given during surgery
- c) Chemotherapy given before radical surgery to reduce bulk of tumor
- d) Chemotherapy given after radical surgery for micrometastases

Question 27:

A young patient is admitted with a history of fever for 5 days. BP is 90/60 mmHg, PR-120 bpm, RR-24 breaths/min, GCS score-10; laboratory tests show leucocytosis with neutrophilia and serum creatinine level of 2.6 mg/dL. An intern doctor wants to calculate the qSOFA score to predict prognosis and hospitalization duration. Which of the following comprises the qSOFA score?

- a) BP, RR and CBC
- b) Creatinine, PR, BP

c) RR, Body temperature, PR

d) SBP, RR, GCS score

Answer Key

Question No.	Correct Option
1	d
2	c
3	d
4	c
5	c
6	d
7	c
8	d
9	b
10	c
11	b
12	c
13	c
14	c
15	b
16	d
17	b
18	c
19	d
20	c
21	d
22	c
23	b
24	c
25	b
26	c
27	d

Detailed Explanations

Solution to Question 1:

The most appropriate management for this patient with verrucous carcinoma around the glans of the penis is partial penectomy.

Carcinoma of the penis is commonly a squamous cell carcinoma. It arises from the skin of the glans penis or the prepuce. The lesion can be flat or warty in appearance.

- T1- confined to the skin
- T2- invades the corpus spongiosum or cavernosum.
- T3- invades the urethra
- T4- invades adjacent structures.


The first lymph node to be involved is the superficial inguinal lymph node and thus, the sentinel lymph node biopsy (SLNB) is helpful for evaluating the spread of penile cancer.

Tumors of low-grade and low-stage are managed by Mohs' surgery. Tumors involving the glans penis require glansectomy, and in advanced cases require partial penectomy.

Other options:

Option A: In penile carcinoma, total amputation of the penis is done if the stump is $\leq$ 2cm after resection of the tumor.

Options B, and C: Topical 5-fluorouracil cream and CO₂ laser ablation are preferred in case of penile intraepithelial neoplasia.



Solution to Question 2:

The procedure being performed in the above image is intraosseous route access for transfusing IV fluids in the anterior proximal tibia.

The intraosseous route is an alternative to the intravenous (IV) route in the transfusion of fluids in shock. The marrow of long bones has a rich network of vessels that drain into a central venous canal, emissary veins, and ultimately the central circulation. So, the bone marrow functions as a non-collapsible venous access route.


Intraosseous cannulation is recommended for patients in whom IV access proves difficult or unattainable. (e.g., in cardiopulmonary arrest). In such a case, if venous access is not available within 1 min, an intraosseous needle should be placed in the anterior proximal tibia. During a resuscitation procedure, all anesthetic drugs and fluids can be administered through intraosseous cannulae.

Insertion sites for intraosseous infusion:

- In the proximal tibia, 1-2 cm anteromedial from the tibial tuberosity
- Distal tibia at the junction of the medial malleolus and the shaft of the tibia
- Distal one-third of the femur

The response time to drug administration in intraosseous infusion is close to the IV route. Fluids should be given as boluses using a 50 mL syringe.

The image below shows the procedure of intraosseous cannulation.



Solution to Question 3:

The given condition is likely to be Fournier gangrene. It is a mixed flora aerobic and anaerobic infection.

Fournier's gangrene is a polymicrobial infection and a type of necrotizing fasciitis of the perineum, external genitalia, and perianal region. Risk factors include immunocompromised states such as HIV, alcoholism, and diabetes. The infection may be precipitated by minor

procedures in the perineal area, such as a bruise, scratch, urethral dilatation, injection of hemorrhoids, or opening of a periurethral abscess. The infection spreads along the dartos, Scarpa, and Colles fascia.

The sequence of events that lead to the development of gangrene are as follows: A mixed infection of the aerobic and anaerobic bacteria → fulminating inflammation of subcutaneous tissue → obliterative arteritis of the arterioles that supply the scrotal skin → gangrene.

Clinical features of Fournier gangrene are scrotal pain, fever, prostration, and pallor. Cellulitis spreads rapidly and may be associated with eschars, necrosis, flaking skin, and crepitus. The testis is usually not affected.

Diagnosis is majorly clinical and very rarely made by laboratory or imaging findings.

The steps in the management of this condition:

- IV fluids
- IV antibiotics
- Urgent surgical excision followed by skin grafting in later stages. Wide surgical excision is often necessary.

When the infection has been controlled, a vacuum-assisted dressing may be helpful. Skin grafting is essential for a patient surviving an acute episode. Despite the best therapy, mortality rates are around 50%.

Other options:

Option A: Anti-gas gangrene serum is a treatment for gas gangrene that is of historical importance and is no longer used. Treatment of gas gangrene is now done with antibiotics.

Option B: While urinary diversion and other supportive measures are useful in the management of the condition, it is not the next best step.

Option C: The blood supply of the testes is separate from that of the overlying skin, and hence, testes can be preserved during debridement.

Solution to Question 4:

The first vessel that is usually ligated during abdominoperineal resection (APR) is the inferior mesenteric artery (IMA). A high inferior mesenteric artery ligation during abdominoperineal resection can injure the superior hypogastric nerve, resulting in sympathetic dysfunction and leading to retrograde ejaculation and bladder dysfunction.

Abdominoperineal resection is the complete excision of the rectum, anal canal, and sphincter apparatus. This is done with the creation of a permanent colostomy. This is commonly preferred in elderly patients with distal rectal cancer with poor sphincter function. In these patients, the end colostomy offers a better quality of life.

Complications:

- Higher recurrence rates compared to low anterior resection.

- Anastomotic leaks or dehiscences
- Necrosis of the transposed colon
- Bleeding
- Strictures

The sympathetic innervation of the rectum is from L1, L2, and L3, which together form the superior hypogastric plexus. They divide into left and right hypogastric nerves at the level of the sacral promontory supplying the rectum and the genitourinary system. A high IMA ligation during surgery can result in sympathetic dysfunction causing retrograde ejaculation.

Injury to the pelvic plexus and nervi erigentes, during the division along the pelvic side wall, can cause erectile dysfunction, impotence, and atonic bladder. Injury to the periprostatic plexus can lead to sexual and bladder dysfunction.

Solution to Question 5:

The image showing a large area of scar tissue is suggestive of a keloid.

A keloid refers to excess accumulation of collagen at the scar site with growth of the scar tissue beyond the boundaries of the original wound. It does not regress. The patient's history of traumatic injury to the chest one year ago, followed by the development of an expanding scar, is suggestive of a keloid. The keloid is often glossy in appearance and rubbery in texture. They are composed of thick bundles of collagen.

Treatment of keloids usually involves intralesional injections of triamcinolone. Multiple doses are required at intervals over several weeks and may result in flattening but rarely does the wound narrow.

Option A: A hemangioma is a vascular tumor that arises from the endothelium. They typically present as red or pinkish macules or raised lesions.

Option B: Hypertrophic scars, unlike keloids, remain within the boundaries of the original wound and tend to regress over time. Since this scar has failed to regress over 1 year and has grown beyond the original wound margins, it is more likely to be a keloid.

Option D: Neurofibromas are benign tumors arising from Schwann cells. They usually present as soft, flesh-colored nodules, often in association with neurofibromatosis type 1, a genetic disorder. The patient's history of chest trauma and the appearance of the lesion make neurofibroma an unlikely diagnosis in this case.

Solution to Question 6:

Entrapment of the lateral cutaneous nerve of the thigh is the most likely cause of pain in this patient with history of laparoscopic hernia repair.

The laparoscopic approach is preferred for bilateral or recurrent inguinal hernias.

Transabdominal preperitoneal (TAPP) repair and total extraperitoneal (TEP) repair are the methods commonly used.

The nerves commonly injured in laparoscopic repair are the genitofemoral and lateral femoral cutaneous nerves of the thigh. Patients present with acute neuropathic pain along the dermatomal distribution. Injury to the lateral femoral cutaneous nerve results in meralgia paresthetica.

Treatment consists of rest, ice, NSAIDs, physical therapy, and possible local corticosteroid and anesthetic injection. Drugs like gabapentin can be used.

In laparoscopic TAPP repair, to reduce the chances of nerve injury during mesh placement, tacks are ideally placed above the iliopubic tract to avoid nerve injury.

Other options:

Option A: Ilioinguinal nerve may be commonly injured during open hernia repair. It leads to sensory disturbances and pain over the skin of the genitalia and upper medial thigh.

Option B: Iliohypogastric nerve can be injured while making an incision in the right iliac fossa during hernia surgery.

Option C: Femoral nerve can be compressed by retroperitoneal tumors or hemorrhage in patients on anticoagulants or bleeding diathesis. It presents as wasting and weakness of the quadriceps femoris muscle making the patient difficult to walk. Pain and paraesthesia are seen along the region of the distribution of the nerve.

Solution to Question 7:

The GCS score of this patient: E2V3M4

- He opens his eyes to a painful stimulus - E2
- He produces inappropriate words only - V3
- He withdraws his limbs - M4


Solution to Question 8:

The given CT image shows a cyst in the liver with serpentine linear membranes seen as wavy membranes within the cyst is suggestive of hydatid disease.


Hydatid disease or echinococcosis is a zoonotic infection caused by the Echinococcus species of flatworms. Its life cycle consists of dogs as definitive hosts and humans as abberant intermediate hosts. Primary hydatid cysts most commonly occur in the liver. Most patients are asymptomatic but when symptomatic, the patient presents with hepatomegaly, pain, and obstructive jaundice.

The first/initial imaging test is USG, and a CT scan can be used to diagnose the condition. CT scan shows the presence of a space-occupying lesion with a smooth outline and septa. CT scan has an accuracy of 98% in the detection of hydatid cysts and high sensitivity to the detection of daughter cysts and extra-hepatic cysts. On CT imaging, the serpent sign is seen as wavy membranes within the cyst.

The USG of an active hydatid cyst shows multiple septations and a rosette or honeycomb appearance:



The ball of wool sign is seen on USG when the contents settle in the dependent portion of the cyst, as seen below:



On USG, the water-lily sign is seen in hydatid infections when there is detachment of the endocyst membrane, which results in floating membranes within the pericyst that mimic the appearance of a water lily.



Other options:

Option A: While surgical treatment of hydatid cyst is useful and effective, alternatives like percutaneous aspiration, injection, re-aspiration (PAIR) are equally effective and are currently used.

Option B: IgG ELISA and indirect hemagglutinin assay are used for the diagnosis of hydatid cyst.

Option C: Angioembolization has no role in the treatment of hydatid cyst.

Solution to Question 9:

The given clinical scenario and image depict a likely case of parotid tumor, likely to be a pleomorphic adenoma.

Pleomorphic adenoma is the most common benign salivary gland tumor. The most commonly affected gland is the parotid gland. The superficial lobe is most commonly affected. It is also called as mixed tumor since it has both epithelial and mesenchymal components. It most commonly affects females around 40 years of age.

The common presenting feature is a painless swelling below, in front, or behind the ear. The involvement of the deep lobe leads to dumbbell tumor – swelling in both the neck and oral cavity.

Specific features:

- Slowly growing benign tumor
- Unicentric in origin
- Potentially malignant
- Possibility of protrusion of fringes of the tumor mass through the capsule
- Radioresistant
- Usually affects adults, females > males

- Painless swelling
- Well-defined margins
- Lobulated, firm, elastic-to-hard consistency
- Free from deeper structures
- No facial nerve involvement
- The deep lobe pushes tonsils and faucial pillars


Diagnosis is made by FNAC, which is considered the investigation of choice. Open biopsy contraindicated due to chances of injury to the facial nerve and seedling of the tumor

Surgery is the mainstay of treatment - Superficial parotidectomy (Patey's operation). Radiotherapy is considered in high-grade malignancies demonstrating extra-glandular disease, perineural invasion, direct invasion of surrounding tissue, or regional metastases.

Pathology:

Gross features:


- The tumors are rounded, well-demarcated, measuring ≤ 6 cm in greatest dimension.
- They are usually encapsulated, except in some areas such as the palate.
- The cut surface is gray-white with myxoid and blue translucent areas of the chondroid (cartilage-like) stroma.



Microscopic features:

- The epithelial elements resemble ductal cells or myoepithelial cells. They are arranged as ducts, acini, irregular tubules, strands, or sheets of cells.
- These epithelial elements are dispersed within a background of loose myxoid and hyaline tissue containing islands of cartilage and, rarely, foci of bone.

The image below shows pleomorphic adenoma of the salivary gland consisting of mixed epithelial (left) and mesenchymal cell components (right).



Dermoid cysts are slow-growing benign lesions. They are mostly seen on the scalp and skull. It is made of hair, teeth, cartilage, and skin glands. Other common sites are the forehead, postauricular, and lateral part of the eyebrow.

A sebaceous cyst is derived from hair follicle infundibulum or traumatic inclusion. The most common site of a sebaceous cyst is the postauricular sulcus or below and behind the ear lobule. Malignancy is rare.

Clinical features of a sebaceous cyst include:

- Benign swelling that is mobile, fluctuant, and fixed to the skin. It presents with a central punctum in about 70% of cases.
- It is generally an asymptomatic, non-tender swelling. It can be painful and red if inflamed.
- It may ooze out a foul-smelling, cheese-like discharge.

Solution to Question 10:

The most likely diagnosis in this patient with a history of alcohol dependence, sudden unintentional weight loss, and raised AFP is hepatocellular carcinoma (HCC).

HCC is the most common cause of primary liver cancer. It is more commonly seen in males, and the peak age group is between 65-70 years. The main risk factor for HCC is cirrhosis and associated chronic liver damage due to chronic infection by hepatitis B or C virus (HBV or HCV), alcohol abuse, metabolic syndrome, or hemochromatosis. The incidence of HCC is increasing worldwide. This is contributed by the increase in HCV and HBV infections, along with lifestyle disorders like obesity. Obesity is associated with non-alcoholic fatty liver disease (NAFLD). This progresses on to non-alcoholic steatohepatitis (NASH) causing cirrhosis and hepatocellular carcinoma. Thus, NASH and NAFLD are risk factors for HCC.

Regular surveillance, including liver imaging and monitoring of alpha-fetoprotein levels, is recommended for high-risk populations, such as those with liver cirrhosis, chronic hepatitis B or C infection, or significant alcohol consumption. Early detection of HCC can improve treatment

outcomes and overall prognosis.

Very early HCC is treated by ablation. Early HCC with a single nodule is resected, but when there are multiple nodules over 3 cm, ablation or transplantation are considered. Intermediate-stage HCC is treated by chemoembolization. Transarterial chemoembolization (TACE) is used for multinodular lesions.

Other Options:

Option A: Hepatic adenomas are benign liver tumors that are often associated with oral contraceptive use and anabolic steroid use. The patient's history of alcohol dependence and high alpha-fetoprotein levels are not consistent with hepatic adenoma. Additionally, hepatic adenomas are usually asymptomatic and do not present with sudden unintentional weight loss.

Option B: Cholangiocarcinoma is a cancer that arises from the bile ducts within the liver. While patients with alcohol dependence may have an increased risk of cholangiocarcinoma, the elevated alpha-fetoprotein levels are not typical for this type of cancer. Cholangiocarcinomas usually present with jaundice, pruritus, and abdominal pain, but sudden weight loss is less characteristic.

Option D: Alcoholic hepatitis is an inflammation of the liver caused by excessive alcohol consumption. It usually presents with jaundice, fever, and abdominal pain. While the patient's history of alcohol dependence might suggest alcoholic hepatitis, elevated alpha-fetoprotein levels and unintentional weight loss are not typical for this condition. Moreover, the AST/ALT ratio is usually greater than 2 in alcoholic hepatitis, but it is 0.5 in this case.

Solution to Question 11:

In this case of chronic pancreatitis with recurrent attacks, 10 mm dilatation of the pancreatic duct, and intraductal calculi, longitudinal pancreaticojejunostomy (also known as Puestow procedure or lateral pancreaticojejunostomy) is the surgery of choice. This procedure involves opening the pancreatic duct along its length and connecting it to a loop of the jejunum, which helps to drain the obstructed pancreatic duct and alleviate pain.

Patients with chronic pancreatitis mainly experience pain due to ductal hypertension or parenchymal inflammation. Acute exacerbations in duct pressure lead to recurrent acute attacks. The pancreatic duct dilation observed on pancreatography for chronic pancreatitis is classically described as a chain of lakes, which reflects the presence of multiple dilations and stenoses. Response to treatment has been found to be dependent upon the degree of preoperative pancreatic fibrosis.

Longitudinal pancreaticojejunostomy, also known as the Puestow procedure or lateral pancreaticojejunostomy, is a surgical treatment primarily used for chronic pancreatitis with a dilated main pancreatic duct and intraductal calculi.

In longitudinal pancreaticojejunostomy, a longitudinal incision is made along the length of the main pancreatic duct, typically from the tail to the head of the pancreas. The dilated pancreatic duct is then opened, and any intraductal calculi or obstructions are removed. The opened duct is then connected to a loop of the jejunum to create a side-to-side anastomosis. This allows for the drainage of pancreatic secretions directly into the jejunum, bypassing any narrowed or obstructed areas of the duct.

This procedure can effectively relieve pain and improve the quality of life for patients with chronic pancreatitis. However, it is important to note that it may not reverse the underlying pathology of chronic pancreatitis or prevent the progression of the disease. It is also not suitable for all patients with chronic pancreatitis, particularly those with a non-dilated pancreatic duct or isolated head involvement.

Other Options:

Pancreaticoduodenectomy: This procedure, also known as Whipple procedure, involves removing the head of the pancreas, duodenum, gallbladder, and part of the bile duct. It is typically performed for pancreatic cancer or tumors in the head of the pancreas, not for chronic pancreatitis with ductal obstruction and intraductal calculi.

ERCP and sphincterotomy: Endoscopic retrograde cholangiopancreatography (ERCP) with sphincterotomy is a procedure that involves opening the sphincter of Oddi to facilitate the drainage of bile and pancreatic secretions. Although it may provide temporary relief, it is not a definitive treatment for chronic pancreatitis with large ductal dilatation and intraductal calculi. ERCP with sphincterotomy may be useful in some cases of chronic pancreatitis for the removal of small pancreatic duct stones, but for large ductal dilatation and intraductal calculi, longitudinal pancreaticojejunostomy is more appropriate.

Coring of pancreas head: Coring of the pancreas head, also known as pancreatic head resection or Beger procedure, is a surgical technique used to treat chronic pancreatitis, particularly when the inflammation and fibrosis predominantly affect the head of the pancreas. The procedure involves removing the inflamed and fibrotic tissue from the head of the pancreas while preserving the duodenum, bile duct, and the main pancreatic duct. Longitudinal pancreaticojejunostomy is a more targeted and effective treatment for this patient's condition.

Solution to Question 12:

ER positivity is a good prognostic factor in breast cancer.

The ER and PR are two steroid hormone receptors present in breast cancer. The expression of these receptors indicates a favorable prognosis and response to hormonal therapy.

Other options:

Option A: The mutations in BRCA1 gene account for up to 40% of familial breast cancers. These tumors are high-grade, hormone receptor–negative, and aneuploid and do not have a good prognosis.

Option B: p53 is a tumor suppressor gene, which on mutation acts as an oncogene. p53 positivity is associated with highly malignant breast cancers and has poor survival.

Option D: Ki-67 is used to assess the proliferative index (activity) of the tumor. High Ki-67 indicates a poor prognosis.

Solution to Question 13:


Sudden-onset severe chest pain, with a wide discrepancy in blood pressure between the two limbs, and a widened mediastinum on x-ray point towards a diagnosis of aortic dissection.

It is initiated by an intimal tear, which allows blood to penetrate into the medial layer, producing a cleavage plane (false lumen) between the inner two-thirds and outer one-third of the media. The true lumen is separated from the false lumen by an intimomedial flap.

Classification of aortic dissection:

Acute aortic dissection can be classified by the original DeBakey system, which is based on the location of the proximal tear and distal extent of the dissection flap, or the Stanford system, which is based only on the location of the proximal tear.

	DeBakey	Stanford
Site of dissection		
Both ascending and descending aorta	Type I	Type A
Only ascending aorta and arch of the aorta	Type II	Type A
Descending the aorta distal to the subclavian artery	Type IIIa Limited to the thoracic aorta	Type B
Type IIIb Extends to the abdominal aorta		



Clinically, patients present with tearing chest/back pain associated with diaphoresis, syncope, weakness, and dyspnea or neurological manifestations.

On examination, unequal pulses or blood pressure in the extremities may be seen due to changes of flow in the true and false lumens. There can be hypertension or hypotension, loss of pulses, findings of aortic regurgitation, and pulmonary edema.


Imaging:

Transesophageal echocardiography (TEE) is the investigation of choice in a hemodynamically unstable patient. In hemodynamically stable patients, the diagnosis is established with a contrast-enhanced CT scan. The characteristic diagnostic finding is a double-lumen aorta. The image below shows aortic dissection in the thoracic aorta with the true lumen, as well as the false lumen. The true lumen is often the smaller of the two because of being compressed by the false lumen.


Acute coronary syndrome and acute pulmonary embolism may both present with sudden onset chest pain but widened mediastinum and blood pressure discrepancy make aortic dissection more likely. (Options A and B)

Both sudden onset chest pain and widened mediastinum may be seen in esophageal rupture but blood pressure discrepancy between the upper limbs makes aortic dissection more likely. (Option D)

Chest x-ray showing widened mediastinum in a patient with aortic dissection



The illustration given below shows the flowchart for the management algorithm for aortic dissection:



Solution to Question 14:

This patient with an abscess in the liver with a history of diarrhea most likely has an amebic liver abscess. The initial treatment of choice for an amebic liver abscess is medical therapy with metronidazole.

Ingestion of *Entamoeba histolytica* cysts causes intestinal diseases like diarrhea. Following this, the trophozoites reach the liver mostly through the portal venous system and cause amebic liver abscess. The common presenting features are right upper quadrant pain, tenderness, fever, hepatomegaly, and diarrhea.

CT abdomen is helpful in differentiating amebic from a pyogenic abscess. A rim enhancement is noted in case of a pyogenic abscess.

The mainstay of treatment for amebic abscesses is metronidazole, which is curative in more than 90% of patients. Luminal agents such as iodoquinol, paromomycin, or diloxanide furoate can be administered to treat the carrier state.

Aspiration of abscess is recommended in case of

- Diagnostic uncertainty
- Failure to respond to metronidazole therapy
- Abscesses larger than 5 cm in diameter and in the left liver, since they are thought to carry a high risk of rupture

Solution to Question 15:

The given clinical scenario and the plain CT scan showing a concavo-convex-shaped hematoma on the left side with a midline shift to the right is suggestive of subdural hematoma (SDH).

An acute SDH is the result of the accumulation of blood between the arachnoid membrane and the dura. It is usually due to the rupture of bridging veins running from the cerebral cortex to the dural sinuses. The level of consciousness is usually impaired at presentation and may deteriorate further as the hematoma expands. Elderly and alcoholic patients are at a higher risk of acute SDH formation after head trauma due to brain atrophy. On CT scan, acute SDH appears as a hyperdense, crescent-shaped (concavo-convex) lesion.

Open craniotomy for the evacuation of acute SDH is indicated for the following:

- Thickness ≥ 1 cm
- Midline shift ≥ 5 mm
- Glasgow Coma Scale (GCS) score drops by two or more points from the time of injury to hospitalization.

Chronic subdural hematoma results from a hematoma that expands over days or weeks by osmosis, ultimately producing symptoms of raised ICP or focal deficits. On a CT scan, diffuse hypodensity overlying the brain surface would be seen.

Solution to Question 16:

The given image is likely that of Meleney's gangrene. Hyperbaric oxygen has been shown to reduce mortality in patients with Meleney's gangrene and therefore plays a significant role in the management of this condition.

Necrotizing fasciitis of the abdominal wall is known as Meleney's gangrene. It presents with extensive necrosis of the abdominal wall, thrombosis of the microvasculature, and an acute inflammatory infiltrate. Bullae develop on the abdominal wall, which progress to overt cutaneous gangrene. The affected area is often painful, erythematous, and edematous, with possible systemic signs of infection such as fever and malaise. Early recognition and aggressive treatment are essential for a favorable outcome.

Management includes prompt surgical debridement of the necrotic tissue, broad-spectrum antibiotics to cover the polymicrobial nature of the infection, and supportive measures such as fluid resuscitation and correction of electrolyte imbalances. In some cases, hyperbaric oxygen therapy may be utilized as an adjunctive treatment to improve tissue oxygenation and enhance the healing process.

Other options:

Option A: Surgical excision of necrotic tissue is often required daily as the infection tends to spread beyond the borders of the debridement wound.

Option C: Meleney's gangrene is caused by beta-hemolytic streptococci or *Staphylococcus aureus* and may also be polymicrobial.

Solution to Question 17:

A direct inguinal hernia is caused due to weakness in the conjoint tendon.

The innermost layers of the lateral abdominal wall contain two muscles, namely internal oblique and transverse abdominis. These muscle fibers arch over the deep inguinal ring from lateral to medial, fuse, and become tendinous forming the conjoint tendon. The area below this arch lacks muscle and contains external oblique aponeurosis and fascia transversalis, making it weak.


Direct inguinal hernia is common in elderly patients. It is seen medial to the inferior epigastric vessels, in Hesselbach's triangle. This area lacks muscle and contains only fascia.

Other options:

Option A: Pectineal ligament also known as Cooper's ligament is a strong fibrous band that extends laterally along the pectineal line from the pectineal attachment of the lacunar ligament.

Option C: The inguinal ligament is external oblique aponeurosis that extends between the anterior superior iliac spine and the pubic tubercle. On the point of attachment to the pubic tubercle, some fibers pass upwards behind the superficial inguinal ring and external oblique aponeurosis to join the rectus sheath and linea alba. These form the reflected part of the inguinal ligament.

Option D: Lacunar ligament is formed from the fibers of the medial end of the inguinal ligament and the fibers from the fascia lata of the thigh. It forms the medial wall of the femoral canal.



Solution to Question 18:

This patient with a history of aspirin use, now presenting with abdominal pain and a sudden onset of hematemesis most likely has a bleeding peptic ulcer.

Peptic ulcer disease (PUD) accounts for around 2/3rd of upper gastrointestinal bleeding. The major causative factors are *Helicobacter pylori* and NSAIDs. Prostaglandins protect gastric and duodenal mucosa from acid and pepsin. They increase mucin and bicarbonate secretion and increase mucosal blood flow. They also promote epithelial cell proliferation. NSAIDs like aspirin disrupt these naturally protective mechanisms and increase the risk of peptic ulcer formation in the stomach and the duodenum.

They present with abdominal pain, bleeding, or signs of peritonitis due to perforation. Significant hemorrhage occurs when duodenal or gastric ulcers penetrate branches of the gastroduodenal or left gastric arteries, respectively.

Other options:

Option A: Mallory-Weiss tears are partial-thickness tears of the mucosa and submucosa occurring near the gastroesophageal junction. They occur in patients with a history of repeated emesis and are classical in alcoholic patients after a period of intense retching and vomiting following binge drinking.

Option B: Esophagitis commonly occurs due to repeated exposure of mucosa to the acidic gastric secretions in gastroesophageal reflux disease. It results in chronic blood loss and commonly presents as anemia or guaiac-positive stools. Ulcerations of superficial mucosa due to esophagitis generally do not bleed acutely.

Option D: Esophageal varices commonly occur due to portal hypertension in patients with cirrhosis. A history of liver disease or the stigmata of portal hypertension and cirrhosis would suggest variceal bleeding.

Solution to Question 19:

The clinical vignette suggests the diagnosis of Raynaud's syndrome due to chronic exposure to drills and machines (vibrating tools).

Raynaud's syndrome occurs due to peripheral vasospasm, and it is more common in the upper extremities. It may occur due to exposure to various stimuli, including cold temperatures, tobacco, or emotional stress, and certain occupational groups, such as those who use vibrating tools, or it can be secondary to scleroderma, systemic lupus erythematosus, Sjogren's syndrome, and antiphospholipid syndrome.

The pathophysiology behind Raynaud's syndrome is believed to be due to increased levels of α_2 -adrenergic receptors and their hypersensitivity and also due to an abnormal thermoregulatory response by the sympathetic nervous system.

A Raynaud's phenomenon attack is sudden, and it ranges from intense pallor (white attack) to cyanosis (blue attack), followed by reactive hyperemia due to subsequent relaxation of blood vessels. It typically affects fingers (distal areas), and the thumb is usually spared. If the thumb is involved, it may indicate underlying secondary etiology. Other common complaints include pin and needles sensation, numbness, or finger aches or pain. In severe attacks, digital ulcerations and gangrene may develop.

Diagnosis is primarily based on the clinical history and physical examination. Nail fold capillary microscopy can be helpful, particularly for identifying secondary Raynaud's phenomenon associated with connective tissue diseases. Angiography can be done for patients with digital ulcerations and gangrene.

Management is mainly conservative, and it includes wearing gloves, use of electric or chemically activated hand warmers, avoiding occupational exposure to vibratory tools, abstinence from tobacco, and relocating to a warmer, dryer climate.

Other management options:

- Calcium channel-blockers such as diltiazem and nifedipine are the drugs of choice.
- Fluoxetine has been shown to reduce the frequency and duration of vasospastic episodes.
- Prostaglandins can be used in patients with severe symptoms.

Complications, including digital ulcerations and gangrene, can be surgically treated by debridement and amputation, respectively.

Solution to Question 20:

The given features are suggestive of a juvenile polyp, which are a type of hamartomatous polyp. They are benign growths that are composed of a disorganized mixture of different types of tissues.

Juvenile polyps are malformations of epithelium and lamina propria commonly located in the rectum. It is seen in children <5 years of age. The typical presentation is painless rectal bleeding. Sometimes, there may be a proliferating rectal mass.


The disease may be sporadic or syndromic.

- Sporadic juvenile polyp occurs as a single mass and is called a retention polyp.
- Juvenile polyposis syndrome is characterized by multiple hamartomatous polyps, which may be premalignant.

Morphologically, both of these types are indistinguishable.

- Gross appearance: They appear as pedunculated, smooth reddish lesions, and with cystic spaces.
- Microscopic appearance: The cysts appear as dilated glands filled with mucin and inflammatory debris.

Juvenile polyp



Adenoma: Adenomas are benign tumors that arise from glandular tissue. While they may present with rectal bleeding, they are not commonly seen in young children and are not typically pedunculated like juvenile polyps.

Choristoma: Choristomas are non-neoplastic masses that arise from heterotopic tissue, i.e., tissues found in abnormal locations. They are not associated with juvenile polyps.

Solution to Question 21:

The given picture showing an ulcer with punched-out edges over a pressure point is most likely that of a trophic ulcer/neuropathic ulcer.

A neuropathic ulcer is one that occurs as a result of peripheral neuropathy. In peripheral neuropathy, lack of sensation over the pressure points leads to cumulative microtrauma, minor wounds, or cuts that remain unnoticed but eventually develop into ulcers.

Features of a neuropathic ulcer include

- Painless ulcer
- Punched out ulcer with regular margins
- Often located on the plantar surface of the foot
- Ulcer margins show the presence of calluses

	Ischemic/Arterial Ulcer	Venous Ulcer	Neuropathic Ulcer	Tuberculous Ulcer
Underlying condition	Peripheral vascular disease- mainly atherosclerosis, thromboangiitis obliterans, cervical rib, Raynaud's phenomenon	Varicose veins, previous deep vein thrombosis	Diabetes, peripheral neuropathy	Tuberculosis lymph nodes, cold abscess from bone and joint tuberculosis
Location	Toes, feet, legs, or even upper limb digits	Lower limb, commonly on the medial side of the ankle, never above the middle third of the leg	The plantar surface of the foot, toes	Neck, axilla, and groin
Characteristics	Deep, penetrates the deep fascia exposing the tendons, muscles, and underlying bone. The ulcer is painful, tender, and hyperaesthetic.	Vertically oval, painless, the floor may have pale to no granulation tissue, induration, and tenderness at the base.	Punched-out with regular calloused margins.	Edges are thin, reddish blue, undermined; pale granulation tissue, scanty discharge on the floor, and slight induration at the base
Other features	Features of ischemia in adjacent areas, like pallor, dry skin, brittle nails, loss of hair	Inguinal lymph nodes are enlarged, usually the vertical group, and the ulcer is recurrent; forms again after healing	Dry, cracked, insensate	Regional lymph nodes are enlarged, non-tender, matted

Option A) The following is an image of a venous ulcer



Option B) The following is an image of a malignant ulcer (squamous cell carcinoma)



Option C) The following is an image of an arterial ulcer



Solution to Question 22:


A patient with varicose veins and overlying eczema is classified as C4a under CEAP guidelines.

The Clinical-Etiology-Anatomy-Pathophysiology (CEAP) classification system is used to classify chronic lower extremity venous disease.

The clinical class of C4 under CEAP includes changes in skin and subcutaneous tissue secondary to chronic venous disease. It is further classified as follows:

- C4a - Pigmentation or eczema
- C4b - Lipodermatosclerosis or atrophie blanche
- C4c - Corona phlebectatica

Image A shows tortuous dilated veins - C2, image B shows atrophie blanche - C4b, and image C shows fan-shaped telangiectasia - C4c.



Solution to Question 23:

In this patient with subarachnoid hemorrhage (SAH), the next step is to do an angiography to see the vascular anatomy and identify the site of bleed.

Trauma is the commonest cause of subarachnoid hemorrhage. Spontaneous SAH usually results from the rupture of an aneurysm or an arterio-venous malformation (AVM). Patients present with a sudden and severe headache which is different from any headache they have experienced before. This is described as a thunderclap headache. Other symptoms include meningitic features like neck rigidity and photophobia, impaired consciousness, intraocular hemorrhages, with or without focal neurological deficits. Compression from a posterior communicating artery aneurysm would result in third nerve palsy.

Non-contrast CT head is the first imaging to be done in a suspected case, which would confirm bleeding in more than 98% of cases. In patients with high suspicion and a negative CT, lumbar puncture is done to detect hemoglobin breakdown products like oxyhemoglobin and bilirubin. After confirming the diagnosis, CT or MR angiography is done to locate the aneurysms or AVM. Digital subtraction angiography remains the gold standard investigation.

The definitive management of subarachnoid hemorrhage (SAH) is done surgically by craniotomy and clipping or by endovascular embolization known as coiling. This prevents rebleeding of the aneurysm.

Patients with subarachnoid hemorrhage are at risk of developing complications such as rebleeding, cerebral vasospasm, hydrocephalus, seizures, and hyponatremia. Early recognition and management of these complications are essential to improve outcomes. In addition to surgery and nimodipine, other management strategies for SAH include strict blood pressure control, analgesia for headache, antiemetics, and anticonvulsants if seizures occur. Maintaining a close monitoring of the patient's neurological status, intracranial pressure, and electrolytes is also crucial.

Other Options

Nimodipine: Although it is used to prevent cerebral vasospasm and delayed ischemic neurological deficits, it is not the immediate next step in the management of SAH. Angiography is needed first to locate the aneurysms or AVM. It is administered after the procedure.

Surgery: Although surgery is an important management option for SAH, it is not the immediate next step. Angiography must be done first to identify the site of the bleed and plan for the appropriate surgical or endovascular intervention.

Fibrinolytic therapy: This option is not appropriate for the management of SAH as it may exacerbate the bleeding. Fibrinolytic therapy is used in the treatment of ischemic stroke, not hemorrhagic stroke.

Solution to Question 24:

This patient with a corrosive esophageal injury with complete dysphagia requires a feeding jejunostomy for enteral feeding.

The injury caused by alkali ingestion is deeper due to the process of saponification of fats and dehydration. The injury caused by acids is more superficial because they cause coagulative necrosis with eschar formation, which prevents the deeper penetration of the acid.

Severe corrosive injury may lead to esophageal perforation, bleeding, stricture, and death. Patients can develop dysphagia or odynophagia, which implies esophageal damage.

Minor injuries cause mucosal edema that resolves rapidly with no late sequelae and these patients can be safely fed. In case of severe injuries, a feeding jejunostomy may be appropriate until the patient can swallow saliva satisfactorily.

Management of corrosive esophagitis:

- Hemodynamic stabilization and management of patient's airway
- Endoscopic evaluation of injuries
- Surgical management includes laparotomy for bleeding or perforation. Esophageal replacement is done for very long or multiple strictures.

Stenting is used in patients who do not respond to repeated dilatations for esophageal strictures. They are not used as a treatment in acute settings.

Solution to Question 25:

The given clinical scenario is suggestive of a thyroglossal cyst. It moves with deglutition, protrusion of the tongue, and side-to-side movement is also possible.

A thyroglossal duct cyst is a congenital midline neck swelling that typically presents during childhood often noticed after an upper respiratory infection.

A thyroglossal cyst represents the persistence of the track of the thyroid gland and can be found adjacent to the midline. The thyroid gland descends early in fetal life from the base of the tongue towards its position in the lower neck. At the time of descent, the hyoid bone has not been formed

and the track of the thyroid gland is variable, passing in front, through, or behind the eventual position of the hyoid body.

It can occur anywhere along the migratory course of the duct:

- Infrahyoid (most common site)
- Region of thyroid cartilage
- Suprahyoid
- Floor of mouth
- Beneath foramen cecum

Clinical features of thyroglossal cyst:

- Classically the cyst moves upwards on swallowing and on tongue protrusion. This indicates attachment to the hyoid bone.
- Swelling is smooth, soft, fluctuant (cystic), nontender, mobile, and shows transillumination.
- Thyroid fossa is empty, if there is no thyroid in the normal location.

Complications of thyroglossal cyst:

- The most common complication is infection.
- Thyroglossal fistula is an acquired condition that either follows infection of the cyst which bursts open or due to inadequate removal of the cyst.
- 1% of thyroglossal duct cysts are found to be malignant of which the most common type of cancer is papillary (85%).

However, only the thyroglossal cyst moves with tongue protrusion. This is because the duct extends from the foramen cecum of the tongue to the isthmus of the gland.

Sistrunk's operation is performed to excise the cyst and prevent a recurrence. It involves excision of the whole thyroglossal tract, involving removal of the body of the hyoid bone, and suprahyoid tract through the tongue base to the vallecula at the site of the primitive foramen cecum along with the core of tissue on either side.

Features of hoarseness of voice, cough, neck swelling that increases in size on coughing/Valsalva maneuver, and a gurgling sound heard on pressing the swelling are suggestive of laryngocele.

Solution to Question 26:

Neoadjuvant chemotherapy is defined as chemotherapy given before the definitive treatment (usually surgery) to reduce the bulk of the tumor.

Neoadjuvant therapy is a type of induction therapy that is given before the definitive surgery to downstage the tumor, reduce the morbidity of treatment, know the in vivo response of the tumor to therapy, and increase its chances of success. Chemotherapy, radiotherapy, or hormonal therapy can be given as neoadjuvant therapy.

Option A: Treatment that involves the administration of chemotherapy along with radiation is called chemoradiation.

Option B: Hyperthermic intraperitoneal chemotherapy (HIPEC) is used during surgery in which a hyperthermic solution of anticancer drugs is infused directly into the abdominal cavity after the definitive surgery to kill any residual tumor cells. It is often used to treat cancers that have spread to the peritoneum.

Option D: Treatment given after surgical resection of the tumor to reduce the risk of recurrence and micrometastasis is called adjuvant therapy. Adjuvant therapy can be chemotherapy, radiation therapy, hormone therapy, targeted therapy, or biological therapy.

Solution to Question 27:

qSOFA score comprises of SBP, RR, and GCS.

qSOFA score (quickSOFA) is a bedside prompt that may identify patients with suspected infection who are at greater risk for a poor outcome outside the intensive care unit (ICU).

It uses three criteria, assigning one point each for

- low blood pressure ($SBP \leq 100$ mmHg),
- high respiratory rate (≥ 22 breaths/min),
- altered mentation (GCS score, ≤ 15).

The score ranges from 0 to 3 points. The presence of 2 or more qSOFA points is a simple prompt to identify infected patients outside the ICU who are likely to develop sepsis. This is a simplified derivative of SOFA score. The primary method to evaluate the risk of organ dysfunction (sepsis) in patients with suspected infection is by determining the Sequential (sepsis-related) Organ Failure Assessment (SOFA) score, which includes assessment of:

- Circulation - Systolic BP and need for vasopressors; platelet count
- Respiration - PaO_2/FiO_2 and need for mechanical ventilation
- Mentation - Glasgow Coma Scale score
- Hepatic function - Bilirubin level
- Renal function - Creatinine level

Organ dysfunction is defined by a SOFA score of ≥ 2 .

The SOFA score is limited by its requirement of multiple laboratory tests which makes it complex and time-consuming. In order to simplify the scoring process, the "Sepsis - 3" task force recommended the quick sepsis-related organ dysfunction assessment "qSOFA" score.

SOFA vs. qSOFA:

- SOFA is a more comprehensive and accurate scoring system, which is particularly superior to qSOFA in the ICU setting.
- qSOFA is a simplified derivative of SOFA for bedside assessment of individuals at risk for sepsis, particularly useful outside the ICU setting (i.e. ward/emergency department). However, note that it was originally designed as a predictive tool that calculates the risk of death from sepsis rather than a diagnostic tool.

Thereby, the best description of sepsis based on "Sepsis-3" task force recommendations would be as follows:

- Suspected or documented infection with SOFA score of ≥ 2 or
- Suspected or documented infection with qSOFA score of ≥ 2

Another score that has been used is the National Early Warning Score (NEWS). It ranges from 0 to 20 and is based on respiratory rate, oxygen saturation, use of supplemental O₂, temperature, systolic blood pressure, pulse rate, and level of consciousness. ≥ 5 out of 20 is considered a positive score.

Other scoring systems for sepsis:

The Systemic Inflammatory Response Syndrome (SIRS) score is no longer recommended as it is not specific to infection.

The Logistic Organ Dysfunction System (LODS) score is not recommended as it is complex and has too many variables.

Surgery INI-CET 2023

Question 1:

A 25-year-old patient presents with fever and weight loss. He gives a history of blood in the stools and vomiting 2 weeks ago. On examination, there is right hypochondriac tenderness and hepatomegaly. Abdominal CT a lesion in the right lobe of the liver with ragged edges, multiple septations, and heterogenous densities. What is the most likely diagnosis?

- a) Pyogenic liver abscess
- b) Hydatid cyst
- c) Amoebic liver abscess
- d) Hemangioma

Question 2:


Which of the following tests is done before performing an A-V fistula procedure? The image of the test is given below.



- a) Allen's test
- b) Adson's test
- c) Elevated arm stress test
- d) Nicoladoni's test

Question 3:

Match the following positions of the appendix:



©Marrow

- a) 1-b, 2-a, 3-d, 4-c
- b) 1-c, 2-a, 3-b, 4-d
- c) 1-b, 2-d, 3-a, 4-c
- d) 1-c, 2-b, 3-a, 4-d

Question 4:

A patient presents with dull aching pain in both legs. He gives a history of prolonged standing and tortuous veins in the legs. Which of the following tests are done for saphenofemoral incompetence?

- a) 1, 2, 3 and 4
- b) 1, 2 and 3
- c) 1, 2
- d) 3, 4

Question 5:

Which of the following statements is true about thymoma?

- a) Most common neoplasm of thymus

- b) Most common posterior mediastinal tumor .
- c) Usually symptomatic with dyspnoea, dysphagia and SVC syndrome
- d) Chest Xray is the investigation of choice

Question 6:

In a patient who underwent laparoscopic cholecystectomy, what is the ideal urine output?

- a) 500 ml/day
- b) 0.5 ml/minute
- c) 0.5-1.5 ml/kg/hour
- d) 1 ml/kg

Question 7:

In a patient presenting with rectal prolapse, which of the following procedures is not done via the perineal approach?

- a) Ripstein operation
- b) Thiersch operation
- c) Altemeier procedure
- d) Delorme operation

Question 8:

In a patient with an inguinal hernia, the surgeon reduces the entire hernial sac into the abdominal cavity without reducing the contents. What is this known as?

- a) Maydl's hernia
- b) Reduction en masse
- c) Sliding hernia
- d) Incarcerated hernia

Question 9:

Which of the following statements about claudication pain is not true?

- a) Rest pain is seen more in the night
- b) Pain in intermittent claudication comes at first step of walking.
- c) Most common cause is atheroma
- d) The clinical stem does not reflect the site of peripheral arterial obstruction

Question 10:

A 73-year-old patient presents with painless rectal bleeding. Which of the below-mentioned investigations are done for this patient?

- a) 3 and 4
- b) 2, 3 and 4
- c) 1 and 2
- d) 1, 2 and 3


Question 11:

Smoking increases the risk of which of the following diseases?

- a) 2 & 3
- b) 1 & 4
- c) 2, 3, 4
- d) 1, 3, 4

Question 12:


Identify the following image:



- a) Hemangioma
- b) Cystic hygroma
- c) Branchial cyst
- d) Ranula

Question 13:

Feeding is done through the given method. Which of the following is not a complication of this?



- a) Refeeding syndrome

- b) Osteoporosis
- c) Pneumothorax
- d) Aspiration pneumonia

Question 14:

Identify the surgical instrument shown in the image.



- a) Adson's forceps
- b) Kocher's forceps
- c) Mixter forceps
- d) Rampley forceps

Question 15:

A 25-year-old male patient was brought to the casualty after a road traffic accident with a suspected pelvic fracture. On examination, he is found to be in hypovolemic shock. What is the earliest treatment for the control of hemorrhage in this case?

- a) Immediate internal fixation
- b) Multiple blood transfusions
- c) Use bedsheets to compress the pelvis
- d) Emergency external fixation

Question 16:

Which of the following are done in Breast Conservation Surgery?

- a) 1, 2, 3
- b) 1, 3, 4
- c) 1, 2
- d) 2, 3

Question 17:

Match the features seen due to compression of the following structures in case of thyroid swelling.

- a) 1-d, 2-b, 3-c, 4-a
- b) 1-c, 2-d, 3-b, 4-a
- c) 1-c, 2-d, 3-a, 4-b
- d) 1-a, 2-b, 3-d, 4-c

Question 18:

A 63-year-old female presented to the OPD with abdominal pain and a tense and tender swelling below and lateral to the pubic tubercle. What is the most likely diagnosis of this condition?

- a) Strangulated femoral hernia
- b) Strangulated inguinal hernia
- c) Intussusception
- d) Adhesions

Question 19:

A patient presented with a Type 1 Choledochal cyst. All of the following statements are correct except?

- a) Increased risk of cholangiocarcinoma
- b) Cyst opened and drainage is done
- c) Todani classification is used

- d) Due to anomalous pancreaticobiliary junction

Question 20:

A patient underwent laparoscopic cholecystectomy for cholelithiasis. Specimen revealed that she had gall bladder carcinoma infiltrating the muscularis layer. What is the management?

- a) Radical cholecystectomy with adjuvant chemotherapy
- b) Adjuvant chemotherapy
- c) Extended cholecystectomy
- d) Surveillance

Question 21:

A man presented with a history of stab injury over the arm. On examination, there was a hematoma present, and radial and ulnar pulses were not felt. A 3.5cm longitudinal tear is noted in the brachial artery. Which among the following is the best treatment option?

- a) Primary repair with end to end anastomosis
- b) Repair with great saphenous vein graft
- c) Repair with prosthetic graft
- d) Repair with saphenous vein patch


Question 22:

Which of the following drugs are not used in the management of a thyrotoxic crisis?

- a) Acetaminophen
- b) Aspirin
- c) Lugol's iodine
- d) Propranolol

Question 23:

Identify the incision.



- a) Battle incision
- b) Gibson incision
- c) Robson incision
- d) Lanz incision

Question 24:

Which of the following is the best method to conclusively localize a parathyroid adenoma?

- a) SPECT sestamibi
- b) Tc99 sestamibi scan
- c) USG neck
- d) MRI

Question 25:

Which classification of bile duct injury takes into account both vascular as well as the location of the injury?

- a) Strassberg
- b) Hannover
- c) Bismuth
- d) McMahon

Question 26:

If open cardiac massage were to be performed on a patient, what category would it come under?

- a) Clean
- b) Contaminated
- c) Clean-contaminated
- d) Dirty

Question 27:

A patient who was a known case of rheumatoid arthritis is brought to the ER due to dizziness. Her BP was 90/60mmHg. She was prescribed a medication by her local doctor which she stopped taking suddenly. Her blood sugar was low and serum potassium was 6mEq/L. Which of the following drugs should be given to her next?

- a) IV Saline
- b) Hydrocortisone
- c) Dopamine infusion
- d) Sodium bicarbonate

Question 28:

What is the most common cause of hypercalcemia in hospitalized patients?

- a) Hypervitaminosis D
- b) Malignancy
- c) Parathyroid adenoma
- d) Parathyroid hyperplasia

Question 29:

An elderly lady was diagnosed to have a gallbladder polyp. She was told that she has a risk for gallbladder cancer. Which of the following is not a risk factor for gallbladder cancer?

- a) Porcelain GB
- b) Primary sclerosing cholangitis

- c) A single gall bladder polyp < 1 cm
- d) Gall stone > 3cm

Question 30:

Which of the following is the most important prognostic factor of breast carcinoma?

- a) Axillary lymph node status
- b) Size
- c) Distant metastasis
- d) Hormonal status

Answer Key

Question No.	Correct Option
1	c
2	a
3	c
4	c
5	a
6	c
7	a
8	b
9	b
10	d
11	d
12	d
13	d
14	c
15	c
16	a
17	b
18	a
19	b
20	c

21	b
22	b
23	b
24	a
25	b
26	b
27	b
28	b
29	c
30	a

Detailed Explanations

Solution to Question 1:

Features of fever, hepatomegaly, and hypochondriac tenderness are suggestive of a liver abscess. A liver abscess in a patient with a history of blood in the stools and vomiting points to a diagnosis of amoebic liver abscess.


Amoebic liver abscess is more common in males in the age group of 20 - 40 years with a history of travel to endemic areas. Ingestion of *Entamoeba histolytica* cysts causes diarrhea initially. Following this, the trophozoites reach the liver through the portal venous system and cause an amoebic liver abscess. 80% of the cases involve the right lobe. The clinical features of amoebic liver abscess are right upper quadrant pain, tenderness, fever, hepatomegaly, and diarrhea.

Enzyme immunoassays detecting the presence of antibodies against the parasite have high sensitivity and specificity. Ultrasonography and CT scans can visualize the liver abscess without a peripheral rim of enhancement.



The mainstay of treatment for amebic abscesses is metronidazole, which is curative in more than 90% of patients. Luminal agents such as diloxanide furoate can be administered to treat the carrier state. Aspiration of abscess is recommended in case of a high risk of rupture. Pigtail catheter drainage is done in patients not responding to intravenous metronidazole in the first 48–72 hours to improve antibiotic penetration.

Option A: Pyogenic liver abscess is usually seen after 50 years of age and is not associated with a history of travel or diarrhea. Jaundice, bacteremia, and positive blood cultures are common in pyogenic liver abscesses compared to amoebic liver abscesses. Serology testing for *Entamoeba* is negative. A characteristic rim enhancement is noted in the case of a pyogenic abscess on a CECT abdomen and can differentiate it from an amoebic liver abscess.



Option B: Hydatid cyst a zoonotic infection caused by the *Echinococcus* species of flatworms. Primary hydatid cysts most commonly occur in the liver. Most patients are asymptomatic but when symptomatic, the patient presents with hepatomegaly, pain, and obstructive jaundice. The rupture of a hydatid cyst can result in fever, pruritus, or anaphylaxis. Imaging findings are cystic lesion with multiple daughter cysts within, detachment of the endocyst membrane (water-lily sign on USG shown below), and areas of calcification.



Option D: Hemangiomas are more common in women and are usually asymptomatic. Larger hemangiomas may present with vague upper abdominal symptoms. On contrast-enhanced CT scan, hemangiomas show asymmetrical, nodular peripheral enhancement that is isodense, with large vessels and progressive centripetal enhancement.




Solution to Question 2:

The image is suggestive of Allen's Test. It is usually done prior to radial artery cannulation and performing an A-V fistula procedure to test the adequacy of ulnar collateral blood flow to the hand.

Allen's test is performed by occluding the radial and ulnar arteries. The patient is asked to make a tight fist so as to exsanguinate the palm. Then the patient is asked to open the hand and the occlusion over the ulnar artery is released. As occlusion of the ulnar artery is released, the color of

the open palm is observed. Normally, the color will return to the palm within 5 – 6 seconds. If the palm remains pale for more than 6-10 seconds, it indicates reduced ulnar collateral flow. The modified Allen's test considers a threshold of 5-15 seconds.



In recent years, studies have shown that, although the Allen test is often used to identify patients at increased risk for ischemic complications from radial artery cannulation, the predictive value of this test is poor. It is also used for identifying patients acceptable for radial artery harvest or use for coronary angiography.


Other options:

Option B: Adson's test is a provocative maneuver for thoracic outlet syndrome. The test is performed by extending the patient's neck and turning the head towards the affected side while they breathe in deeply. This compresses the interscalene space and will result in paraesthesia and obliteration of the radial pulse.

Option C: Elevated arm stress test or Roos test is a provocative test for the diagnosis of thoracic outlet syndrome. For this test, the patient abducts the shoulders to 90 degrees and flexes the elbow to 90 degrees. The patient is made to open and close the hand slowly for three minutes. The test is positive if the patient is unable to complete the test or experiences cramping, pain, or numbness on the affected side.

Option D: The reflex bradycardia and increase in blood pressure that immediately follows the sudden occlusion of an arteriovenous (A-V) fistula is known as the Nicoladoni-Branham sign.

Modified Allen's test




Solution to Question 3:

The correctly matched positions of the appendix are:

- 1 - b: Retrocaecal - most common position, seen in 74%
- 2 - d: Paracaecal - seen in 2%
- 3 - a: Subcaecal - seen in 1.5%
- 4 - c: Inside the pelvis - seen in 21%

The appendix is a short and broad blind muscular tube that becomes tubular with the differential growth of the caecum. This continued growth of the caecum rotates the appendix into a retrocaecal but intraperitoneal position. In some cases, rotation of the appendix does not occur, resulting in a pelvic, subcaecal, or paracaecal position. During development, if the caecum does not migrate to its normal position in the right lower quadrant of the abdomen, the appendix can be found near the gallbladder. In cases of intestinal malrotation, the appendix may be found in the left iliac fossa.

Various positions of the appendix are shown in the image below.



Solution to Question 4:

Brodie Trendelenburg test and Morrissey's test are performed to test the incompetence of the saphenofemoral valve. Fegan's method is used to detect perforator incompetence while Modified Perthe's test is a test for deep vein thrombosis (DVT).

Brodie Trendelenburg test - Lay the patient down and elevate the leg. Empty the vessels and tie a tourniquet below the SFJ. Ask the patient to stand up:

- Trendelenburg's test part I - Here the tourniquet is released. A rapid filling from above shows saphenofemoral incompetence.
- Trendelenburg's test part II - Here the tourniquet is not released. A gradual filling from below to upwards indicates perforator incompetence.

Morrissey's cough impulse test - The limb is elevated to empty the varicose veins. Now it is put to bed and the patient is asked to cough forcibly. An expansile cough impulse is felt at the saphenous opening if the saphenofemoral valve is incompetent. Similarly, a bruit may be heard on auscultation. The patient is asked to cough. A cough impulse at the saphenofemoral junction indicates SFJ incompetence.

Fegan's method - Mark the places of excessive bulges within the varicosities. Ask the patient to lie down and empty the varicose vessels. Palpate along the marked area to look for gaps and pits in the deep fascia, which indicates perforator incompetence.

Modified Perthe's test - A tourniquet is tied around the upper part of the thigh to prevent reflux down the vein and the patient is asked to walk around. In the presence of normal communicating and deep veins, the varicose veins will shrink. If there is any block, the veins will get more distended. An increase in pain and prominence of dilated veins indicate DVT.

Solution to Question 5:

The true statement is that thymoma is the most common neoplasm of the thymus.

Thymoma is the most frequently encountered neoplasm of the anterior mediastinum (Option B) in adults and is rare in children. It is seen most frequently in the age group of 40 and 60 years. Most patients with thymoma are asymptomatic (Option C). Large thymic tumors may present with symptoms related to a mass effect, which may include cough, chest pain, dyspnea, or SVC syndrome. Around 10% to 50% of patients with thymoma may have symptoms suggestive of myasthenia gravis or have circulating antibodies to the acetylcholine receptor. Other paraneoplastic syndromes, including red cell aplasia, hypogammaglobulinemia, systemic lupus erythematosus, Cushing's syndrome, or SIADH, may be present.

While the mediastinal mass can be identified on screening chest X-ray, imaging with contrast-enhanced CT and MRI (Option D) have a critical role in characterizing the mediastinal masses. The diagnosis may be suspected based on history and CT imaging but imaging alone is not diagnostic. The diagnosis is usually made after surgical resection due to the relative difficulty of obtaining a needle biopsy. Biopsy should be avoided in cases where imaging is highly suggestive of thymoma. Masaoka staging system is used for thymomas and is based on the gross or microscopic invasion of the capsule and of surrounding structures, as well as on the presence of metastases.

The definitive treatment for thymoma is complete surgical removal. Thymectomy leads to improvement or resolution of symptoms of myasthenia gravis in only about 25% of patients with thymomas.

Solution to Question 6:

The ideal urine output in that patient following laparoscopic cholecystectomy is 0.5-1.5 ml/kg/hour.

Normal urine output is expected in the postoperative period of a patient who underwent laparoscopic cholecystectomy. The average normal urine output in an adult is 0.5-1.5 ml/kg/hour (Option B) or approximately 1500 mL/day (Option A). A minimum urine output of 400 mL/day is essential to excrete the end products of protein metabolism.

Option D: Normal urine output in ml/kg would be 0.5 - 1.5 ml/kg/hour multiplied by 24 which is 12-36 ml/kg per day.

Solution to Question 7:

Among the given options, the Ripstein operation is the abdominal approach for rectal prolapse.

It is managed by using the following approaches:

- The perineal approach is easy to perform with fewer complications but a high recurrence rate. The surgeries done using this approach for rectal prolapse are the following:
- Thiersch's wiring
- Delorme's mucosectomy

- Altemeier's procedure or perineal rectosigmoidectomy
- The abdominal approach is difficult to perform with more complications but a lower recurrence rate. The surgeries done using this approach are the following:
- Moschcowitz repair
- Ripstein and Wells rectopexy; ventral rectopexy or suture rectopexy
- Resection rectopexy

If the patient is fit enough for an abdominal procedure, it is preferred over the perineal procedure, since the risk of recurrence is lower. The perineal approach is preferred in elderly patients with comorbidities and in young men since the abdominal approach may injure the pelvic autonomic nerves which may result in sexual dysfunction.

Solution to Question 8:


The surgeon reducing the entire hernial sac into the abdominal cavity without reducing the contents is referred to as reduction en masse of the hernia.

Reduction en-masse of inguinal hernia refers to the reduction or migration of the hernial sac along with the incarcerated bowel into the properitoneal space. The properitoneal space lies between the parietal peritoneum and the anterior abdominal wall. This is likely to occur following forcible attempts to reduce the hernia but may occur spontaneously. Though the hernial sac no longer lies in the inguinal or femoral canal, it contains an obstructed bowel loop that causes small bowel obstruction. This is a rare condition as forcible reduction of hernias is not done and early repair is carried out.

The patient may give a history of difficult reduction and the symptoms of bowel obstruction persist even after the hernia is reduced. On a CT scan, a hernial sac in properitoneal space with a fibrous constriction at the neck may be visualized. This hernial sac will contain the bowel loop with wall enhancement and fluid that suggest strangulation. Early surgical intervention is required.

Other options:

Option A: Maydl's hernias are defined as the presence of two small bowel loops within a single hernial sac, that is, there are two efferent and two afferent loops of the bowel, forming a "W" shape.



Option B: Sliding hernia occurs when an internal organ composes a portion of the wall of the hernia sac. The most common viscus involved is the sigmoid colon and its mesentery on the left side and the caecum on the right side. It is more common on the left side. They are usually variants of indirect inguinal hernias. The failure to recognize the visceral component of the hernia sac resulting in injury to the bowel or bladder is the major concern in sliding hernia.

Option D: Hernias which cannot be reduced are referred to as incarcerated hernias while hernias with compromised blood supply to its contents are known as strangulated hernias.

Solution to Question 9:

Pain in intermittent claudication is brought on by walking and doesnot start with the first step of walking.

Peripheral artery occlusion or stenosis is predominantly caused by atherosclerosis or atheroma (Option C) and thromboembolic disease. The resulting stenosis or occlusion produces signs and symptoms related to the organ that is supplied by the affected artery. Such occlusion in the lower limbs presents with claudication pain, rest pain, and gangrene. Though the muscle groups affected by claudication are present at one anatomical level below the level of arterial disease, the clinical stem does not reflect the exact site of the peripheral arterial obstruction. (Option D)

Intermittent claudication pain is a feature of arterial stenosis or occlusion that occurs due to the anaerobic metabolism occurring in the muscle. Claudication pain may be of different types and can be differentiated as follows:

- Vascular (intermittent) claudication pain is brought on by walking and is relieved by rest within 5 minutes. (Option B)
- Claudication pain present on taking the first step of walking is classical for osteoarthritis.
- Neurogenic claudication occurs with changes in the position of the back such as bending forward and straightening. It takes more time to be relieved by rest (more than 5 minutes). It may be seen in cases of nerve compression from lumbar intervertebral disc prolapse or spinal stenosis.

Rest pain occurs when the occlusive disease progresses in severity and anaerobic respiration occurs even at rest. Though this pain is present during the day, it is exacerbated on lying down at night (Option A) due to loss of gravity dependant perfusion of the foot. This may be relieved by hanging the foot out of the bed to restore the gravitational effects.

Solution to Question 10:

Colonoscopy, CECT and CEA (Carcinoembryonic antigen) can be done in an old patient presenting with painless rectal bleeding. Serum CA 19-9 is a pancreatic cancer biomarker and has no role in this case.

Rectal bleeding occurs due to pathology of the lower gastrointestinal tract below the ligament of Treitz. The differentials that can be considered in a case of rectal bleeding are:

- Colon cancer
- Inflammatory bowel disease
- Diverticular diseases
- Hemorrhoids
- Angiodysplasia

Colonoscopy with biopsy has a key role in the diagnosis of inflammatory bowel disease, distinguishing between ulcerative colitis and Crohn's colitis and monitoring the response to treatment. Colonoscopy is also the investigation of choice if colorectal cancer is suspected, especially in old patients. Angiodysplasia is suspected in patients above the age of 50 and can be identified by a colonoscopy.

Computed tomography (CT) colonography is sensitive in picking up polyps and has the advantage of being less invasive than colonoscopy. Spiral CT can aid in the diagnosis of complicated diverticular disease.

Carcinoembryonic antigen (CEA) is a non-specific serum biomarker that is elevated in colorectal cancer and hence may be done in this case.

Solution to Question 11:

Smoking increases the risk of which of duct ectasia, Mondor's disease, and breast carcinoma.

Duct ectasia is defined as dilated major milk ducts. It is considered a disorder of involution as part of ANDI. Abnormally dilated ducts are filled with debris. This acts as an irritant and can lead to periductal inflammation and subsequent fibrosis, leading to nipple retraction. Duct ectasia appears to be associated with smoking and diabetes.

The thrombophlebitis of the superficial veins of the breast and anterior chest wall is called Mondor's disease. The pathognomonic feature is a tender thrombosed subcutaneous cord, usually attached to the skin. When the skin over the breast is stretched by raising the arm, a narrow, shallow, subcutaneous groove alongside the cord becomes apparent. Smoking was found to be a potential factor involved in the pathogenesis of Mondor's disease. Other risk factors include breast

surgery, breast biopsy, an inflammatory process, breast cancer, and trauma.

The risk factors of breast cancer can be divided into modifiable (those that can be modified by adopting a healthy diet and lifestyle) and non-modifiable risk factors. The events increasing the oestrogenic exposure of the breast are said to be risk factors for breast cancer, such as early menarche, late menopause, nulliparity, late first pregnancy, and hormone replacement with high estrogen therapy. Smoking is a modifiable risk factor with a relative risk (RR) = 1.14 for smoking 25 or more cigarettes/day and RR = 1.07 for smoking for 20 years or more.

Fibroadenoma is the most common cause of a breast lump in women aged 15–25 years. It arises from the hyperplasia of a lobule and is surrounded by a well-defined capsule. Smoking is not a risk factor for fibroadenomas.

Solution to Question 12:

The given scenario and the image of a cystic swelling on the floor of the mouth are suggestive of a ranula.

Ranula is a mucus extravasation cyst arising from the sublingual gland. It produces a characteristic translucent swelling that gives the appearance of a frog's belly. A ranula usually resolves spontaneously.

A plunging ranula is a mucus retention cyst. It has mucus collecting below the gland and perforating through the mylohyoid muscle.

Treatment should include removal of the sublingual gland as this gland has multiple ducts and ranulas can recur if the gland is left behind. Incision, drainage and marsupialisation have low success rates.

Other options:

Options A: Hemangioma is the most common vascular tumor and is characterized by increased abnormal proliferation of endothelium. Hemangiomas develop in infancy and are usually not visible at birth. They appear red or maroon if superficial, and bluish if deep.



Option B: Cystic hygroma is usually seen in neonates and early childhood. They are cystic swellings filled with lymph fluid and are characteristically brilliantly translucent. They occur in the neck and can involve the parotid, submandibular, tongue, etc. The swelling is soft and partially compressible. Definitive treatment is the excision of the swelling.



Option C: The branchial cyst develops from the vestigial remnants of the second branchial cleft. Swelling in the neck beneath the anterior border of the upper third of the sternomastoid muscle. It is smooth, soft, fluctuant, and often transilluminates. The below image shows a large right-sided branchial cyst.



Solution to Question 13:

The given image shows a central line. Pneumothorax, osteoporosis, and refeeding syndrome are likely complications in patients receiving total parenteral nutrition through a central line.

Aspiration pneumonia is more common in enteral feeding and can be prevented by placing the patient in a semi-recumbent position while feeding.

Complications of enteral nutrition are as follows:

- Tube-related (most common): Malposition, displacement, blockage, breakage/leakage, local complications (e.g. erosion of skin/mucosa)
- Gastrointestinal: Diarrhoea, bloating, nausea, vomiting, abdominal cramps, aspiration, constipation
- Metabolic/biochemical: Electrolyte disorders, vitamin, mineral, and trace element deficiencies, drug interactions
- Infective: Exogenous (handling contamination) and endogenous (patient)

Other options:

Option A: Refeeding syndrome is a significant metabolic complication of both parenteral and enteral feeding. It occurs days after feeding is commenced in patients who have been severely malnourished. The major pathological process is hypophosphatemia which causes fluid and electrolyte shifts between the intra- and extracellular compartments.

Option B: Osteoporosis and osteomalacia are complications of long-term parenteral nutrition and may lead to fractures or kidney stones. Calcium, phosphate, vitamin D supplementation, and bisphosphonates can help prevent as well as treat this complication.

Option C: Pneumothorax is the most common complication of line insertion and most commonly occurs during the insertion of subclavian lines. It is managed by insertion of a chest drain.

Solution to Question 14:

The given surgical instrument is a Mixter's right angle forceps that can be used to pass ligatures and dissect pedicles.

Other options:

Option A: Adson's forceps or Cerci-Adson-Brown forceps have a double grasping tip that ensures grasping with better stabilization. It is the most appropriate forceps for grasping cartilage in rhinoplasty. Its multi-teeth tip prevents cartilage damage or bounce.



Option B: Kocher forceps have an opposing tooth in the tip and serrations in the distal blades. It is used to hold pedicles, tough structures, and cut ends of muscles. It is also used for holding ribs during rib dissection and for holding the resected bowel.

Kocher's hemostatic forceps



©MARROW

Option D: Rampley's sponge-holding forceps have a fenestrated, serrated, and flat end. The uses are:

- to clean the operative field and swab the cavities
- to mop the oozing area
- to hold the gallbladder and cervix during surgeries
- for blunt dissections

Sponge holding forceps



©Marrow

Solution to Question 15:

The earliest treatment to control hemorrhage in a case of suspected pelvic fracture is to use bedsheets to compress the pelvis similar to a compression binder.

The most likely source of bleeding in hemodynamically unstable patients following blunt trauma is abdominal or pelvic bleeding. A pelvic binder or a sheet should be applied around the true pelvis of all hemodynamically unstable patients and not removed until after a pelvic fracture has been excluded. It must be correctly positioned at the level of the greater trochanters. This is a potentially life-saving procedure that has to be done in the emergency department. Following this, they require an immediate pelvic radiograph. Mechanical testing of the pelvis should never be performed. A pelvic fracture should always be diagnosed radiographically.

In fractures of the pelvic ring, there is a risk of injury to the major vessels (internal iliac and external iliac vessels) and to the venous plexus around the sacroiliac joint. Bleeding from the venous plexus is the major cause of pelvic hemorrhage. Pelvic fractures can have a high mortality rate of more than 40% in some open pelvic fractures. Early mortality is mostly the result of hemorrhage.

Bleeding of venous origin can be controlled by compression either by binding/ external fixator or by extraperitoneal pelvic packing to compress the pelvic veins. Persistent bleeding after packing or bleeding of arterial origin may require angioembolisation. Severe pelvic injuries require a multidisciplinary team approach. An exploratory laparotomy may be considered if

- the source of the bleeding is in doubt
- FAST/CT results are positive, showing a significant amount of blood in the peritoneal cavity
- concurrent intra-abdominal injury cannot be excluded

Solution to Question 16:

Wide local excision, axillary node dissection, and frozen section (biopsy) during surgery are done as a part of Breast Conservation Surgery (BCS). All patients with BCS must receive radiotherapy (Statement 4) and are together known as Breast Conservation Therapy (BCT).

Breast conservation surgery involves removing the tumor with a 1mm margin of normal breast tissue. Removal of primary breast cancer is also called lumpectomy/ tylectomy/ partial mastectomy/ wide local excision (Statement 1). Histological examination via intraoperative frozen section (Statement 3) is done to examine for positive margins. Patients with positive margins should have a revision of margins known as 'cavity shave'. Assessment of regional lymph nodes is done via a separate incision in the axilla (Statement 2).

Absolute contraindications of BCT:

- 1st trimester of pregnancy (may be done for some 2nd and many 3rd trimester patients who can receive radiation after delivery)
- Persistently positive margin (2 microscopy shows positive margins)
- Multicentric lumps
- Diffuse microcalcification
-

Extensive ductal carcinoma in situ (DCIS)

-

Inflammatory breast cancer

Relative contraindications of BCT:

- Active SLE (systemic lupus erythematosus), scleroderma
- Large lump-to-breast ratio (>5 cm size lump)
- Known or suspected genetic predisposition to breast cancer (i.e., BRCA1, BRCA2), Li-Fraumeni syndrome (p53 mutation)
- Multifocal disease
- Prior radiation therapy to the chest wall or breast

Solution to Question 17:

The features seen due to compression of the surrounding structures in case of thyroid swelling are as follows:

The normal thyroid gland is impalpable. Thyroid swellings may be uniform enlargement of the gland, isolated solitary nodules, or multinodular goiter. Large thyroid swelling may compress the surrounding structures and present as pressure symptoms.

- Compression of the trachea can cause respiratory symptoms like dyspnea and stridor.
- Pressure on the esophagus causes dysphagia.
- Compression of the sympathetic chain leads to Horner's syndrome.

- Unilateral paralysis of the recurrent laryngeal nerve produces hoarseness. Vocal cord paralysis from RLN compression is extremely rare and should raise concern for malignant change.

1. Trachea	c. Dyspnea
2. Esophagus	d. Dysphagia
3. Sympathetic chain	b. Horner's Syndrome
4. Recurrent Laryngeal Nerve	a. Hoarseness

Solution to Question 18:

The above clinical scenario of a tense and tender swelling present lateral and below the pubic tubercle is suggestive of a femoral hernia. The femoral hernia traverses the femoral canal, which lies medial to the femoral vein.

A femoral hernia occurs in both males and females, but it is more common in females. It will pass through the femoral ring proximally to enter the femoral canal. The femoral canal is bound by the femoral vein laterally, the inguinal ligament anteriorly, and the pelvic bone with iliopectineal ligament (Cooper's) and lacunar ligament (Gimbernat's) medially. Lacunar ligament has a sharp unyielding edge that prevents the reduction of the femoral hernia. The incidence of strangulation in femoral hernias is high because of the narrowness of the neck of the sac and the rigidity of the femoral ring.

All patients with unexplained bowel obstruction should be examined properly for a femoral hernia. Surgery is the only treatment for femoral hernia, and these patients should be treated on priority because of the high risk of strangulation. The surgical approaches which are used are:

- Low approach (Lockwood): suitable when there is no risk of strangulation; bowel can't be inspected and resected in this approach
- Inguinal approach (Lotheissen)
- High approach (McEvedy): suitable for emergency situations when the risk of bowel strangulation is high
- Laparoscopic approach: both TEP and TAPP approaches can be used; the laparoscopic approach is recommended in women.

Other Options:

Option B: Inguinal hernia is above and medial to the pubic tubercle. Another structure that is useful in differentiating between inguinal and femoral hernia is inguinal ligament.

An inguinal hernia leaves the abdominal cavity just above the inguinal ligament and a femoral hernia leaves just below the inguinal ligament.

Option C: In case of Intussusception, patients present with cramping abdominal pain, palpable sausage-shaped abdominal mass, vomiting, and passage of red currant jelly stools.

Option D: Adhesions can lead to intestinal obstruction, but there won't be any swelling below and lateral to the pubic tubercle,.

Solution to Question 19:

In the given clinical scenario, a patient presented with a Type 1 Choledochal cyst. Cyst opening and drainage will not be done in this patient. the preferred course of action involves the complete surgical excision of the cyst. Complete resection of the cyst is essential because there is a risk of malignant transformation (cholangiocarcinoma). (Option A)

Choledochal cysts are congenital cystic dilations of the intrahepatic and/or extrahepatic biliary tree. They present with a triad of jaundice, a palpable upper quadrant mass, and abdominal pain.

The clinical triad (abdominal pain, mass, and jaundice) and the MRCP (Magnetic Resonance Cholangiopancreatography) findings suggest choledochal cyst. According to the Todani classification, the type III choledochal cyst opens into the duodenum. The treatment of choice for type III choledochal cyst is Endoscopic sphincterotomy followed by surveillance.

Choledochal cysts are congenital dilations of the intra and/or extrahepatic biliary system.

Pathogenesis involves an Anomalous Pancreaticobiliary Junction (APBJ) which may allow reflux into the biliary tract, leading to inflammation, increased biliary pressure, and cyst formation. (Option D)

Clinical features of choledochal cysts are as follows

- 60% of cases are diagnosed before the age of 10 years
- more common in females than in males (4:1)
- The classic symptom triad includes abdominal pain, right upper quadrant mass, and jaundice (most consistent symptom)
- Other symptoms- nausea, weight loss, fever, pruritus, etc

Complications include

- Cholangitis (may lead to cirrhosis and portal hypertension)
- Pancreatitis
- Bile peritonitis (due to cyst rupture)
- Cholangiocarcinoma

Diagnostic investigations for choledochal cysts are as follows:

- Ultrasonography:
- Confirms the presence of an abnormal cyst
- Define the dimensions of the cyst
- Define its relationship to the vascular structures in the porta hepatis, as well as the intrahepatic ductal configuration
- Magnetic Resonance Imaging (MRI/MRCP) is the investigation of choice. It reveals the anatomy of the ducts

- CT - delineates the extent of the intra- or extrahepatic dilatation
- ERCP - defines the distal biliary tree

Todani classification (Option C) of the choledochal cysts along with their management is as shown below,

Biliary-enteric reconstruction is not required in Type III choledochal cyst as it has the lowest risk of malignancy, and it may not be associated with ABPJ (anomalous pancreaticobiliary junction). Therefore, endoscopic drainage is sufficient.

Type of choledochal cyst	Surgical management	
Type I - fusiform dilatation of the bile duct (most common - seen in 75% of patients)	complete surgical excision, cholecystectomy, and Roux-en-Y hepaticojejunostomy	
Type II - saccular diverticulum from the common bile duct	Diverticulectomy, cholecystectomy, and common bile duct repair	
Type III- diverticulum within the pancreas, also known as choledochoceles	ERCP + sphincterotomy (Only cyst in which reconstruction is not required)	
Type IV	a - multiple cysts (both intrahepatic and extrahepatic)	Liver transplant
b - multiple cysts limited to the extrahepatic biliary tree	Kasai procedure (Portoenterostomy)	
Type V - multiple cysts involving only intrahepatic ducts (Caroli's disease)	Liver transplant	

Solution to Question 20:

In the given clinical scenario, gallbladder carcinoma with T1b lesions is detected incidentally after cholecystectomy. Treatment guidelines for the same are as follows:

- Up to stage Ia and stage Ib with negative margins, a standard cholecystectomy would suffice.
- Stage Ib with perineural, lymphovascular invasion, positive cystic duct margins, or poor differentiation, and stage II, extended cholecystectomy should be performed. Extended cholecystectomy involves the removal of the gall bladder, segments IVB and V of the liver, and lymph nodes along the hepatoduodenal ligament, with or without removal of the common bile duct (CBD).

However, the likelihood of nodal disease increases significantly with T1b lesions and perineural, lymphatic, or vascular invasion, therefore, extended cholecystectomy is generally recommended for all patients who are medically fit with T1b or greater level of invasion.

If the initial procedure was performed laparoscopically, the surgeon should examine the laparoscopic port sites. The finding of disease at the port sites is a sign of generalized peritoneal disease and carries a very poor prognosis. However, resection of port sites is no longer

recommended.

Chemotherapy is not required in early gallbladder cancer. Radiation has no role.

Treatment of gallbladder carcinoma:

TNM staging	Treatment
T1a	Simple cholecystectomy
T1b, T2, T3	Radical cholecystectomy
T4	Unresectable, chemotherapy and palliative nonsurgical therapy are provided
N1	Regional lymphadenectomy
N2	Radical cholecystectomy plus hepatopancreaticoduodenectomy

Solution to Question 21:

In this patient, who has sustained a long tear to the brachial artery due to a stab injury, the best treatment option is to repair it with a great saphenous vein interposition graft harvested from the patient's uninjured leg.

The most desirable graft used is the great saphenous vein. A native vein graft is preferable to a prosthetic graft because:

- Its elastic properties are comparable to that of an artery and enable pulsatile flow
- The diameter is similar to that of an extremity artery
- It is not thrombogenic and has long term patency

Other grafts that can be used are cephalic vein, lesser saphenous vein, and prosthetic conduits. Cephalic vein is not the first choice because it is less muscular than greater saphenous vein, and might also compromise the upper limb venous access in the already injured patient. Both the cephalic and the lesser saphenous veins present problems while harvesting in a trauma patient.

Other options:

Option A: Primary arterial repair with end-to-end anastomosis can be done only if the repair is tension-free. If the length of the injured area is long, vessel injuries cannot be repaired by primary end-to-end anastomoses and instead require an interposition graft.

Option C: Prosthetic grafts are only preferred when there has been a trauma to the native saphenous vein or when there is a size mis-match. Polytetrafluoroethylene (PTFE) graft is commonly used.

Option D: Saphenous vein patch is used in the closure of the arteriotomy site during carotid endarterectomies.

Solution to Question 22:

Aspirin has no role in the management of a thyrotoxic crisis. All of the other options listed are used in managing the symptoms. Aspirin and other NSAIDs are to be avoided as they can interfere with protein binding and further exacerbate the condition.

Thyroid storm/ Thyrotoxic crisis:

It is a rare but severe life-threatening complication of hyperthyroidism induced by the release of excessive thyroid hormones.

Causes:

- Abrupt cessation of antithyroid medications
- Infection
- When a thyrotoxic patient has been inadequately prepared for thyroidectomy.
- Thyroid or non-thyroid surgery
- Trauma in patients with untreated thyrotoxicosis
- Amiodarone administration
- Exposure to iodinated contrast agents
- Following RAI (radioactive iodine) therapy

Features:

- They present 12–24 hours after surgery with hyperpyrexia and severe dehydration
- Cardiovascular: Circulatory collapse, hypotension, palpitations, tachycardia, cardiac arrhythmias, cardiac failure
- GI symptoms: vomiting, diarrhea, jaundice.
- Neurological symptoms: restlessness, irritability, delirium, tremor, convulsions, and coma

Treatment:

- Supportive measures – rehydration by proper fluid therapy, tepid sponging, cooling blankets
- Acetaminophen for fever.
- Thionamides like Propylthiouracil (PTU) or methimazole or carbimazole are used. They inhibit thyroid hormone synthesis. Propylthiouracil also inhibits the peripheral conversion of T₄ to T₃. Hence, PTU is the preferred drug in thyroid storm. However, the risk for severe liver injury and acute liver failure associated with PTU therapy makes either carbimazole or methimazole first-choice in patients with deranged liver function tests.
- Lugol's iodide is given one hour after intake of thionamides. They inhibit thyroid hormone synthesis by Wolff- Chaikoff effect. Large doses of iodide inhibit organic binding of the iodide and hence decrease thyroid hormone synthesis.

- Propranolol is used to reduce tachycardia and other adrenergic manifestations. It also reduces the peripheral conversion of T₄ to T₃.
- Glucocorticoids – reduce the peripheral conversion of T₄ to T₃ and controls the shock.
- Digitoxin, cardiac monitoring is crucial.

Solution to Question 23:

The image above shows a Gibson incision. Gibson incision is used for kidney transplantation or as an extraperitoneal approach to the distal ureter with low morbidity.

The traditional pelvic Gibson incision is an oblique or curvilinear incision that starts 3 cm above and parallel to the inguinal ligament and extends vertically 3 centimeters medial to the anterior superior iliac spine (ASIS) extending down toward the inguinal fold and terminating just lateral to the rectus muscle or continuing to above the symphysis pubis.

The Gibson incision is the traditional approach for renal transplant and transplant nephrectomy. This muscle-splitting incision provides great extraperitoneal access to pelvic vessels, the lower ureter, and the bladder for benign and malignant disease and is also utilized as a part of minimally invasive nephrectomy for specimen extraction or use of a hand port.


Other Options:

Option A: battle incision a lower right paramedian incision but placed more laterally than the standard paramedian incision. It is suitable for dealing with acute appendicitis and pathologies in the right lower quadrant of the abdomen.

Option C: Mayo Robson's incision is a right upper paramedian incision extended like a hockey stick to the midline. It is indicated in cholecystectomy.


Option D: Lanz incision, a transverse or skin crease incision for appendicitis, 2 cm below the umbilicus, centered on the midclavicular–mid-inguinal line. Due to its continuation with Langer's lines, the Lanz incision produces much more aesthetically pleasing results with reduced scarring.

An image depicting other abdominal incisions is shown below:



Solution to Question 24:

The best method to conclusively localise a parathyroid adenoma is SPECT sestamibi. Single-photon emission computed tomography (SPECT) is commonly used because of its improved spatial resolution. Sestamibi combined with CT can be used to detect smaller parathyroid tissues. It can also detect ectopic glands in the mediastinum or in the deeper cervical region.



Ultrasonography:

- It is inexpensive and noninvasive.
- Mediastinal parathyroid tissue cannot be detected because USG cannot penetrate the sternum or clavicle.

Nuclear medicine scanning with radiolabeled Sestamibi (Tc99 sestamibi scan) is also a widely used technique. Sestamibi accumulates in parathyroid adenomas. This radionuclide is concentrated in the thyroid and parathyroid tissue but usually washes out of normal thyroid tissue in under an hour. It persists in abnormal parathyroid tissue. Tc99 sestamibi scan is a 2D study, so SPECT sestamibi, which is a 3D study is preferred over Tc99 sestamibi scan. SPECT sestamibi has high specificity and sensitivity as compared to Tc99 sestamibi.

MRI is rarely used because of its low sensitivity. Occasionally, MRI may be indicated in pregnant patients in whom ultrasound is noninformative because MRI avoids radiation exposure.

Solution to Question 25:

Hannover classification of bile duct injury takes into account both vascular as well as the location of the injury.

Hannover, bile duct injuries were divided into five types from A to E. Vascular injuries are included in Type C and Type D.

- Type A is peripheral bile leakage.
- Type B is stricture of CHD or CBD without injury.
- Type C is lateral CHD or CBD injury.
- Type D is the total transection of CHD.
- Type E is bile duct stricture of the main bile duct without bile leakage at the postoperative state.

Other Options:

Option A: The Strasberg classification of bile duct injury is a widely used system to anatomically define bile duct injuries.

Option C: The Bismuth–Corlette classification system provides an anatomic description of the tumor location and longitudinal extension in the biliary tree.

Option D: The McMahon classification separated the injuries by the width of bile duct injury

Solution to Question 26:

Open cardiac massage is classified as a contaminated wound.

Contaminated wounds are class III surgical wounds. They include fresh open accidental wounds, surgeries with a major break in sterile technique, gross spillage of the gastrointestinal tract, or incision through an inflamed nonpurulent tissue. Examples include:

- Open cardiac massage
- Penetrating abdominal trauma

Surgical wounds are classified based on the expected amount of bacterial load at the time of surgery.

Open cardiac massage is performed when there is inadequate cardiac output, after emergency thoracotomy for penetrating chest trauma, during pericardial tamponade.

Class I surgical wounds are clean wounds (option A). They are noninfected surgical wounds without any inflammation. Examples include:

- Hernia repair
- Breast biopsy specimen

Class II surgical wounds are clean/contaminated wounds (option C). They are surgical wounds that include an entry hollow viscus such as the respiratory tract or gastrointestinal tract under controlled conditions without significant spillage. Examples include:

- Cholecystectomy
- Elective gastrointestinal surgery

Class IV surgical wounds are dirty wounds (option D). They are old traumatic wounds with the presence of necrotic tissue and perforated viscus with a heavy load of contamination. Examples

include:

- Perforated diverticulitis
- Necrotizing soft tissue infection

Solution to Question 27:

The given clinical scenario is suggestive of an Addisonian crisis. It is a life-threatening condition caused due to sudden insufficiency of adrenal hormones. If a patient displays features of adrenal insufficiency, treatment must immediately be commenced, before awaiting the biochemical diagnosis. Intravenous administration of hydrocortisone 100 mg every 6 hours, & 2 litres of saline is given in 6 hours under careful cardiovascular monitoring.

The symptoms of adrenal insufficiency are as follows:

- Glucocorticoid deficiency leads to:
 - Fatigue, lack of energy
 - Weight loss, anorexia
 - Myalgia
 - Hypoglycemia
 - Low blood pressure, postural hypotension
- Mineralocorticoid deficiency leads to:
 - Abdominal pain, nausea, vomiting
 - Dizziness, postural hypotension
 - Low blood pressure, postural hypotension
 - Hyponatremia
 - Hyperkalemia
- Adrenal androgen deficiency leads to dry and itchy skin, loss of libido
- The hyperpigmentation of sun-exposed areas results from excess ACTH (adrenocorticotrophic hormone) secretion which occurs in response to decreased adrenal cortical hormones.

Primary causes of the Addisonian crisis include:

- Adrenal haemorrhage
- Acute infection

Secondary causes of the Addisonian crisis include:

- Surgery
- Cessation of steroid therapy
- Postpartum pituitary necrosis.

Solution to Question 28:

Cancer is the most common cause of hypercalcemia in hospitalized patients, whereas hyperparathyroidism is the most common cause of hypercalcemia detected incidentally in outpatients. The causes of hypercalcemia in cancer :

- Cancers affecting bone like multiple myeloma, leukemia, bony metastasis - cause the bone to break down which releases excess calcium into the blood
- Pseudohyperparathyroidism - tumours producing parathyroid hormone-related peptide (PTHrP) cause the bone to release calcium into the blood. The most common carcinoma associated is squamous cell carcinoma of lung.
- Cancers affecting kidneys - failure to remove excess calcium from the blood
- Dehydration caused by nausea and vomiting - difficult for the kidneys to remove calcium from the blood properly
- Lack of physical activity - causes bone breakdown, releasing calcium into the blood

Solution to Question 29:

A single gallbladder polyp ≤ 1 cm is not a risk factor for carcinoma gallbladder. Gallbladder polyps ≥ 10 mm have malignant potential.

Risk factors for carcinoma gallbladder:

- Gallstones of size ≥ 3 cm (most important risk factor)
- Porcelain gallbladder
- Anomalous pancreato-biliary junction
- Choledochal cyst
- Adenomatous polyps
- Primary sclerosing cholangitis
- Obesity
- Salmonella typhi infection.

Gallbladder carcinoma is a rare malignancy that is seen in the elderly. It is associated with a poor prognosis and is indistinguishable from cholecystitis or cholelithiasis. It is usually diagnosed when patients undergo cholecystectomy for gallstones.

Its clinical features include right upper quadrant pain, nausea, vomiting, and anorexia. Jaundice is a late feature. CA 19-9 is elevated in the carcinoma gallbladder and is used for monitoring.

The preoperative diagnosis is often made on ultrasonography and followed up by a CT scan or MRI/Magnetic resonance cholangiopancreatography.

A percutaneous biopsy under radiological guidance is done for pathological examination. PET scanning helps in detecting metastatic disease.

In select cases, a laparoscopic examination is useful in staging the disease. Laparoscopy can also detect peritoneal or liver metastases. The most important prognostic factor is the depth of invasion (T-stage).

Solution to Question 30:

Axillary lymph node status is the most important prognostic factor in patients with breast cancer (non-metastatic). Women with node-negative disease have less than a 30% risk of recurrence, compared with as much as a 75% risk for women with node-positive disease. The presence of lymph node involvement at initial diagnosis also serves as a prognostic factor in recurrent breast cancer.

Prognostic factors in metastatic breast cancer include:

- Relapse-free intervals
- Site of metastasis
- Estrogen receptor (ER) or progesterone receptor (PR) status

Hormone receptor status (estrogen receptor [ER]/progesterone receptor [PR]) is the most important prognostic factor in patients with metastatic breast cancer. Long relapse-free intervals, absence of brain or visceral metastases, and ER/PR positivity are associated with better prognosis in metastatic breast cancer.

Surgery INI-CET 2024


Question 1:

Which of the following are extra-esophageal complications associated with GERD?

- a) 1,2
- b) 2,3,4
- c) 1,2,3,4
- d) 3,4

Question 2:


A 25 year old male presented with difficulty in swallowing food and weight loss for past few months. He also complains of regurgitation of undigested food and halitosis. Barium swallow was performed. What is the diagnosis?



- a) Achalasia cardia
- b) Esophageal carcinoma
- c) Barrett's esophagus
- d) Esophageal stricture

Question 3:

The instruments shown below are used for which of the following procedures?



- a) Orogastric tube insertion
- b) Chest tube drainage
- c) Percutaneous endoscopic gastrostomy
- d) Flatus tube insertion


Question 4:

Patients diagnosed with breast cancer are usually advised to undergo ER screening test. What is its significance?

- a) Diagnostic marker
- b) Prognostic marker
- c) Monitoring biomarker
- d) Surrogate molecular marker

Question 5:

A 45-year-old female presents with recurrent episodes of right flank pain, hematuria, and a history of urinary tract infections. Urine analysis shows alkaline urine with a pH of 8.0. Which of the following is the most likely underlying cause of this patient's condition?



- a) Triple phosphate stones
- b) Calcium oxalate
- c) Uric acid stones
- d) Ammonium urate stones

Question 6:

All of the following statements are true regarding Wilm's tumour except?

- a) It presents most commonly as abdominal mass
- b) Most common age of presentation is within 5 years
- c) 95% of patients live long term with radiotherapy and chemotherapy
- d) Metastasis to lungs is rare and occurs late

Question 7:

What is the colour code in trauma triage for patients with life threatening conditions?

- a) Red
- b) Yellow
- c) Green
- d) Black

Question 8:

Which of the following conditions is not a risk factor for the development of cholangiocarcinoma?

- a) Hepatitis B
- b) Caroli disease
- c) Primary sclerosing cholangitis
- d) Infection with *Opisthorchis viverrini*


Question 9:

A woman presented with ulcerative lesion over the breast associated with erythema of surrounding skin and history of retraction of nipple. Biopsy revealed invasive ductal carcinoma. According to 8th edition of AJCC classification, the patient can be classified under which stage?

- a) Stage cannot be M1
- b) Stage IIIb
- c) Stage 4, if N1-N3
- d) T4b

Question 10:

A 35-year-old male had an episode of loss of consciousness subsequent to a head injury from a bike accident. Upon awakening, he began experiencing severe headache and nausea. He appears drowsy with a Glasgow Coma Scale (GCS) score of 10. One of his pupils is dilated and unresponsive. CT head given. What is the management?



- a) Anticoagulants to prevent further clotting of blood
- b) Observe and serial monitoring in the ICU
- c) Osmotic diuresis to decrease intracranial pressure
- d) Urgent surgical evacuation

Question 11:

A patient with papillary carcinoma thyroid underwent thyroidectomy. She developed hoarseness of voice after surgery. Which of the following steps in surgery could not have resulted in this complication?

- a) Division of strap muscles
- b) Dissection close to Zuckerkandl tubercle
- c) Dissection along triangle of Beahr
- d) Ligation of superior thyroid vessels

Question 12:

Which is the most common hernia in females?

- a) Indirect inguinal
- b) Direct inguinal
- c) Femoral
- d) Obturator

Question 13:

Which of the following is not done during the primary survey in trauma management?

- a) 1, 3 and 4
- b) 3 and 4
- c) 1 only
- d) 1 and 3

Question 14:

A 45-year-old male arrived at the ER with altered mental status and hematemesis that began a day ago. He has a 20-year history of alcoholism, with his most recent binge occurring a week prior. An ultrasound revealed altered echo texture of liver and mild portal vein dilation. CT scan brain was normal. He was started on intravenous pantoprazole and antiemetics. What should be the next step in his management?

- a) Start a course of broad spectrum antibiotics
- b) Order a liver biopsy to assess for fibrosis
- c) Perform an endoscopy to look for varices
- d) Administer intravenous fluids and electrolyte replacement

Question 15:

A 23-year-old female presented with complaints of neck swelling, difficulty swallowing, and hoarseness of voice. Radiological evaluation revealed a mass compressing the trachea. Biopsy showed cells arranged in nests with amyloid deposits. What is the origin of these cells?

- a) Follicular cells
- b) Parafollicular cells
- c) Hurthle cells
- d) Ashkanazy cells

Question 16:

A 34-year-old male, on investigation serum calcium level of 12.8 mg/dl (N- 8.5-10.25mg/dl) with raised PTH. On ultrasonography, a 1.5x1.5 mass was detected on the right lower lobe of

the thyroid gland. Which of the following is true regarding this condition?

- a) A normal DEXA scan rules out the need for surgery
- b) MIBI followed by focussed parathyroidectomy to be done
- c) No surgery is required, only surveillance
- d) Creatinine clearance $>80\text{ml/min}$ mandates surgery


Question 17:

A middle-aged woman presents with flank pain and painful micturition. The pain was relieved by IV analgesics given at a nearby primary healthcare centre. The patient is diagnosed with ureteric colic due to ureteric calculi. What is the best diagnostic modality in this case?

- a) NCCT KUB
- b) CECT
- c) USG KUB
- d) Ureteroscopy

Question 18:

Identify the marked structure.



- a) Corpora amylacea
- b) Corpora arenacea

- c) Lewy body
- d) Psammoma bodies

Question 19:

A 25-year-old patient sustains a fall from the 2nd floor of a building and is brought to the emergency department. Which of the following is not an adjunct to the primary survey?

- a) X-ray pelvis AP
- b) X-ray chest AP
- c) Extended FAST
- d) CT abdomen and pelvis

Question 20:

A 20-year-old male presents with acute epigastric pain and fever. He has a past history of cholelithiasis. On examination, abdominal guarding and rigidity present. His serum amylase, lipase are raised and patient has leucocytosis. What is true regarding his condition?

- a) Serum amylase remains elevated even after 30 days of resolution of acute pancreatitis
- b) Serum lipase is more specific than serum amylase
- c) > 75 percent patients present with hypercalcemia
- d) Increase in the level of serum lipase is directly proportional to the severity

Question 21:

A 34-year-old female presents with a lump in her left breast for 2 years. On examination, a 6 x 4 cm mass is palpated. Dimpling of the skin and bloody discharge from the nipple is present. The lesion is invading the pectoralis major muscle. Ipsilateral Level I and Level II matted axillary lymph nodes present. There was no distant metastasis. What is the TNM staging (AJCC 8th)?

- a) T4bN1Mo
- b) T3N2Mo
- c) T4aN1Mo
- d) T2N1M1

Question 22:

An 84-year-old man with a history of stroke 2 years ago, who is on prolonged bed rest, develops a pressure ulcer with partial thickness skin loss and a clean base. What is the grade of his pressure sore?

- a) Grade 1
- b) Grade 4
- c) Grade 3
- d) Grade 2

Question 23:

During ward rounds, you see a patient who is admitted with a ca esophagus. What is the most likely presentation in this patient?

- a) Dysphagia of solids followed by liquids
- b) Dysphagia of liquids followed by solids
- c) Chest pain
- d) Neck mass

Question 24:

What is the T stage of this oral cavity tumour?



- a) T4a
- b) T4b
- c) T3
- d) T2

Question 25:

A 30-year-old patient with peptic ulcer disease presents with abdominal pain that is not responding despite treatment in multiple hospitals. He was found to have ulcers in the jejunum and ileum. Gastric pH ≤ 2 , and gastrin levels after adequate cessation of PPIs are 10 times the normal value. Identify the lesion present and the boundaries of the triangle containing it.

- a) Gastrinoma; triangle formed by common bile duct, junction of D2-D3, neck and body of pancreas
- b) Insulinoma; triangle formed by cystic duct, junction of D2-D3, neck and body of pancreas
- c) Gastrinoma; triangle formed by cystic duct, junction of D2-D3, neck and body of pancreas
- d) Insulinoma; triangle formed by common bile duct, junction of D2-D3, neck and body of pancreas

Question 26:

An intern is assisting with laparoscopic surgery, and the resident is orienting the intern to the instruments and the operating ports. Your surgeon asks you what the angle between the camera port and the working port is.

- a) Azimuth's angle
- b) Manipulation angle
- c) Crossing angle
- d) Elevation angle

Question 27:

A 35-year-old male underwent surgery for an ileal perforation. Due to heavy contamination, only the sheath was closed, and the wound was left open. Daily wound washing was performed, and the wound was closed on postoperative day 5. By which mechanism does wound healing occur in this case?

- a) Primary intention

- b) Secondary intention
- c) Tertiary intention
- d) Quarternary intention

Question 28:

A 40-year-old patient who is a known case of Hodgkin's lymphoma with a previous history of irradiation presents with right-sided thyroid swelling. FNAC reveals papillary carcinoma of the thyroid and there is no lymph node involvement. What is the next step in management?

- a) Hemi-thyroidectomy
- b) Hemi-thyroidectomy with modified radical neck dissection
- c) Total thyroidectomy
- d) Total thyroidectomy with modified radical neck dissection

Question 29:

One of your friends is teaching you about enhanced recovery after surgery. Which of the following he states is incorrect?

- a) Prolonged fasting is necessary
- b) Epidural anaesthesia is recommended in laparoscopic surgery
- c) Reduction of the hospital stay by 2-3 weeks
- d) Beta-blockers are given after major surgery done under general anaesthesia to improve outcome

Question 30:

A chronic smoker comes with complaints of pain in his legs after walking 100 meters. On examination, he has an absent pulse in the bilateral dorsalis pedis, anterior tibial, and posterior tibial arteries, but no gangrene is present. Intermittent claudication is implicated. Which of the following statements is true regarding this condition?

- a) Pain is more at night and sleep
- b) Development of pain on the 1st step of walking
- c) Pain aggravated by standing and relieved by bending forward
- d) Pain develops after walking some distance and is relieved by rest

Question 31:

A 25-year-old male with a history of a road traffic accident is brought to the emergency department. His GCS is 8/15, oxygen saturation is 85%, BP is 100/ 60 mmHg, and heart rate is 60 beats per minute. On auscultation his right side breath sounds were absent. What is the next step in management?

- a) Secure airway, give oxygen, insert ICD, secure IV line
- b) Administer IV dextrose followed by NCCT head
- c) Start IV fluids, give atropine and secure airway
- d) Evaluate for head injury and do CT-head

Question 32:

An elderly male patient comes with foul-smelling breath and dysphagia. A barium swallow X-ray is attached. What is the most likely diagnosis?



- a) Achalasia cardia
- b) Zenker's diverticulum
- c) Tracheoesophageal fistula
- d) Carcinoma esophagus

Question 33:

Truelove and Witts classification is used for?

- a)** Acute encephalitis
- b)** Acute pancreatitis
- c)** ARDS
- d)** Acute severe ulcerative colitis

Answer Key

Question No.	Correct Option
1	c
2	a
3	c
4	b
5	a
6	c
7	a
8	a
9	b
10	d
11	a
12	a
13	c
14	d
15	b
16	b
17	a
18	a
19	d
20	b
21	b
22	d
23	a
24	a
25	c
26	a
27	c

28	c
29	a
30	d
31	a
32	b
33	d

Detailed Explanations

Solution to Question 1:

Asthma, laryngitis, pulmonary fibrosis, and cough are all extra esophageal complications in gastroesophageal reflux disease (GERD)

Extraesophageal symptoms of GERD are:

- Chronic cough
- Laryngitis,
- Asthma
- Dental erosions (especially on the lingual and palatal tooth surfaces).
- Pulmonary fibrosis
- Sinusitis
- Pharyngitis
- Recurrent otitis media

Extraesophageal symptoms of GERD originate from the respiratory tract and include both laryngeal and pulmonary issues. These symptoms can arise through two mechanisms. The first and most common mechanism involves proximal esophageal reflux and microaspiration of gastroduodenal contents, which cause direct injury to the larynx and lower respiratory tract. The second mechanism involves distal esophageal acid exposure triggering a vagal nerve reflex, leading to bronchospasm and cough, due to the shared vagal innervation of the trachea and esophagus.

The incidence of Barrett's esophagus is also more in patients with long term GERD, which in turn increases the risk of adenocarcinoma of esophagus.

Solution to Question 2:

The given complaints of difficulty in swallowing, weight loss, regurgitation of undigested food and halitosis along with the barium swallow showing bird beak appearance is suggestive of achalasia

cardia.

Achalasia is a primary esophageal motility disorder caused due to the degeneration of neurons in the myenteric plexus of the esophagus. It is characterized by the absence of esophageal peristalsis and impaired relaxation of the lower esophageal sphincter (LES), thereby causing a functional obstruction at the gastroesophageal junction.

The patients present with a triad of dysphagia, regurgitation, and weight loss. The dysphagia initially begins with liquid and then progresses to solid foods, halitosis is present due to the accumulation of food. Patients may develop pneumonia, lung abscess, etc. at a later stage.

The initial diagnosis is made by barium swallow or endoscopy, which shows the presence of dilated esophagus due to the constriction of the lower esophageal sphincter (bird beak appearance). The gold standard test to confirm achalasia cardia is esophageal manometry. The findings are impaired lower esophageal sphincter (LES) relaxation and absent peristalsis with elevated LES pressure.

Management includes:

- Surgical esophagomyotomy.
- Non-surgical measures like sublingual nitroglycerin, nitrates, or calcium channel blockers.
- Injection of botulinum toxin
- Peroral endoscopic myotomy (POEM)

Solution to Question 3:

The instruments shown in the image is used to perform percutaneous endoscopic gastrostomy.

A standard endoscopy is performed to ensure that there are no contraindications to gastrostomy insertion. The stomach is insufflated with air and direct percutaneous needle puncture made at a point where the stomach abuts the abdominal wall. Lignocaine is infused on withdrawal. A trocar is inserted and a wire is passed into the stomach, which can be caught with a snare. The scope is withdrawn, pulling the wire out through the mouth, at which point it is attached to the gastrostomy tube. The gastrostomy is pulled through into the stomach and out through the track created by the trocar insertion.

The insertion of a percutaneous endoscopic gastrostomy (PEG) tube improves nutritional and functional outcomes in patients who cannot maintain oral nutritional intake. PEG insertion is often a preliminary step in the treatment of complex orofacial malignancies and can also be used to support nutrition in patients with various malignant, degenerative, or inflammatory diseases.

Procedure risks include bleeding, perforation, aspiration risk, dislodgment with peritoneal contamination, wound site infection and granulation.

Solution to Question 4:

Estrogen receptor (ER) is used as a prognostic marker.

The term prognostic marker generally is used to describe molecular markers that predict disease-free survival, disease-specific survival, and overall survival, whereas the term predictive marker often is used in the context of predicting response to certain therapies.

Other options

Option A: A diagnostic biomarker detects or confirms the presence of a disease or condition of interest, or identifies an individual with a subtype of the disease.

Option C: When a biomarker can be measured serially to assess the status of a disease or medical condition for evidence of exposure to a medical product or environmental agent, or to detect an effect of a medical product or biological agent, it is a monitoring biomarker

Option D: The most common and serious error in evaluating biomarkers is assuming that a correlation between a biomarker's level and a clinical outcome means the biomarker is a valid surrogate. However, for a biomarker to be a valid surrogate, it must not only be correlated with the outcome but also explain the change in the clinical outcome.

Solution to Question 5:

The given clinical scenario and microscopic image showing coffin lid shaped crystals are suggestive of triple phosphate stones. They are also known as staghorn or struvite stones.

This is caused due to infection with urea-splitting organisms, such as *Proteus*, *Klebsiella*, *Serratia*, or *Enterobacter* species. They are usually formed in alkaline urine.

Staghorn calculi are radio-opaque stones that grow in a branching pattern along the pelvicalyceal system. The above organisms metabolize the urea into ammonium and bicarbonate. Increased metabolism of urea results in alkalinity of the urine, and precipitation of magnesium, ammonium, and phosphate - resulting in the formation of struvite stones.

Complete removal of a staghorn calculus is essential, as any remaining fragments after treatment can lead to a rapid recurrence and continued presence of bacteriuria. Following the successful removal of the infection stone, long-term chemoprophylaxis is required for several months.

Solution to Question 6:

Option C is not true regarding Wilms tumour. The standard treatment of Wilm's tumour involves a combination of chemotherapy and surgery.

Wilms tumor is the most common primary malignant renal tumor in children, typically affecting children less than 5 years (Option B). After neuroblastoma, it is the second most common malignant abdominal tumor in childhood. It most commonly presents as an asymptomatic abdominal mass. (Option A) Other findings include hypertension, abdominal pain, gross painless hematuria, and constitutional symptoms such as fever and weight loss.

CECT abdomen is used to identify the extent of the tumor, integrity of the other kidney, and metastasis. Wilms tumor is metabolically active and can concentrate fluorodeoxyglucose therefore PET/CT is also helpful to identify regional spread and metastatic lesions. A biopsy is discouraged as it may lead to disease upstaging.

Treatment: Surgery and chemotherapy, with or without radiotherapy, are the mainstays of treatment.

- Radical nephrectomy is the standard surgical procedure done. Routine sampling of the para-aortic, hilar, and paracaval lymph nodes should be done.
- Nephron-sparing nephrectomy is done for children with bilateral disease or with a solitary kidney. Preoperative chemotherapy can be used in these cases to induce tumor shrinkage.
- The classical chemotherapy includes vincristine and dactinomycin with the addition of Adriamycin(Doxorubicin).

Despite certain adverse risk factors such as metastases, unfavorable histology and recurrent disease, the prognosis for most children with Wilms' tumor remains very positive. The overall survival rate for children with Wilms' tumor is over 90%.

The tumour can extend into the renal vein and vena cava and metastasises to lymph nodes and lung. But metastasis to lungs occur late in the disease. (Option D)

Solution to Question 7:

Red is the colour code for patients with life threatening conditions who need immediate medical care.

Triage includes rapid classification of the injured based on their injuries and the chances of survival with prompt medical intervention. It provides maximum benefit to the greatest number of injured in a disaster situation. It lessens the burden on medical facilities.

The internationally accepted four-color coded system is used commonly. It includes:

- Red - High priority attention or transfer; Critical, but likely to survive if treatment is given early
- Yellow - Medium priority; Critical, likely to survive if treatment is given within hours
- Green - Ambulatory patients; Stable, likely to survive even if treatment is delayed for hours to days
- Black - Dead or moribund patients; Not breathing, pulseless, so severely injured that no medical care is likely to help

Solution to Question 8:

While all the given options are associated with the causation of cholangiocarcinoma, hepatitis B in particular has an association with intrahepatic cholangiocarcinoma and rarely extrahepatic while all the other options are associated with both intra and extrahepatic cholangiocarcinoma. Hence it is not as significantly associated with cholangiocarcinoma compared to the other options

Cholangiocellular carcinomas (CCCs) or cholangiocarcinomas are mucin-producing adenocarcinomas of the biliary duct system that may originate in the liver and extrahepatic bile ducts. Patients present with signs of obstructive jaundice associated with weight loss and anorexia. Initial workup by ultrasound shows dilation of the intrahepatic biliary tree, but

definitive diagnosis requires visualisation of the biliary tree and MRCP or ERCP (biopsy possible) is preferred

Risk factors for cholangiocarcinoma:

- Chronic inflammatory conditions- PSC, Oriental cholangiohepatitis, Hepatitis C infection
- Parasitic infections- *Opisthorchis viverrini*, *C. sinensis*
- Congenital - Choledochal cysts, Caroli's disease
- Chemical agents- Thorium dioxide (Thorotrast), Vinyl chloride, Dioxin, Asbestos

Solution to Question 9:

The patient presenting with ulcerative lesion with inflammatory signs can be classified as Stage IIIb. Inflammatory breast carcinoma (stage T4d) is characterized by the skin changes of brawny induration, erythema with a raised edge, and edema (peau d'orange) involving >1/3rd of the breast.

T4d in the TNM staging of carcinoma breast points toward a diagnosis of inflammatory breast carcinoma. It shows extensive invasion and proliferation of tumor cells. Inflammatory breast cancer is the most aggressive subtype of breast cancer. The hallmark of this condition is the overexpression of e-cadherin. This results in increased tumor cohesion and the formation of tumoral clumps which can result in emboli. It has a particularly poor prognosis.

In addition to anatomical staging, the eighth edition of the AJCC TNM staging system includes the histological grade, ER, PR, HER2/neu and Ki-67 assessment, multigene testing with Oncotype DX and the response to neoadjuvant chemotherapy to refine the prognostic information.

Solution to Question 10:

The given scenario of a man with a head injury, unconsciousness, nausea, dilated and fixed pupils, and the CT finding of a lenticular hemorrhage also showing a mid-line shift is characteristic of an extradural or epidural hemorrhage. Urgent surgical evacuation of hematoma by craniotomy is the management.

Epidural hemorrhage/hematoma is an intracranial hemorrhage in which the hemorrhage is between the dura and the skull. It classically occurs due to a tear of the middle meningeal artery, following a fracture of the temporal bone. Management is by surgical evacuation of the hematoma or by ligation of the damaged middle meningeal artery.

Clinically, patients with extradural hemorrhage typically present with an initial transient loss of consciousness. A lucid interval is then seen, which may be minutes to hours during which the patient usually presents. Neurological deterioration then occurs following the lucid interval as the hematoma expands.

As the hematoma expands, compensation is exhausted with rapid deterioration. There is contralateral hemiparesis, a reduced conscious level, and ipsilateral pupillary dilatation, the cardinal signs of brain compression and herniation. The prognosis for promptly evacuated

extradural hematoma, without associated primary brain injury is excellent.

Solution to Question 11:

The patient presenting with hoarseness of voice post thyroidectomy can be due to injury to external laryngeal nerve or unilateral recurrent laryngeal nerve. The step in thyroidectomy where it is least to happen is during division of strap muscles.

Recurrent laryngeal nerve is related to the posterior branch of the inferior thyroid artery near the lobes of the thyroid gland. Therefore, during surgery, Inferior thyroid artery is ligated close to the gland to avoid injury to the recurrent laryngeal nerve.

Thyroid surgery may cause injury to both external and recurrent laryngeal nerves (RLN). During thyroidectomy surgeries, there might be unilateral or bilateral recurrent laryngeal nerve injury. Unilateral RLN injury may be asymptomatic but in a few may cause hoarseness of voice.

Bilateral recurrent laryngeal nerve palsy can also occur postoperatively. It causes dyspnea and stridor but the voice is good. These cases required emergency tracheostomy. External laryngeal nerve injury is more common and could result in hoarseness of voice.

Other options

Option B: The RLN frequently passes immediately medial to the tubercle of Zuckerkandl. In order to find and protect RLN during thyroid surgery, a careful, bloodless, and meticulous dissection should be carried out around the tubercle.

Option C: Beahrs' triangle is bounded by common carotid artery as base, inferior thyroid artery superiorly and RLN as the lower arm of triangle. Recognizing the boundaries of the triangle posterior to thyroid after retracting the ipsilateral thyroid lobe medially, helps in avoiding RLN injury.

Option D: The external branch of the superior laryngeal nerve courses along the cricothyroid muscle just medial to the superior pole vessels. To avoid injury to this nerve, the superior pole vessels are divided individually as close as possible to the point where they enter the thyroid gland

Solution to Question 12:

The most common hernia in females is an indirect inguinal hernia.

Inguinal hernia is the most common hernia in men and women. It is classified into :

- Indirect hernia - protrusion of the contents occurs through the deep inguinal ring, along with the spermatic cord, lateral to the inferior epigastric artery. It is the most common subtype of groin hernia in both males and females.
- Direct inguinal hernia - Occurs through the posterior wall of the inguinal canal through the Hasselbach's triangle. The sac is medial to the inferior epigastric artery.

Other options:

Option C: Femoral hernia occurs when the hernial sac descends vertically through the femoral canal up to the saphenous opening and then escapes out into the loose areolar tissue. It is more common in women than in men; however, it is less common than inguinal hernia.

Option D: In an obturator hernia, the contents pass through the obturator canal. It is more commonly seen in females than males; however, it is less common than inguinal hernia.

Note: Inguinal hernias are five times more common than femoral hernias.

Solution to Question 13:

NCCT brain is not done during the primary survey in trauma management. (Option C)

Endotracheal intubation is part of airway management in the primary survey. Intercostal drainage for hemothorax or pneumothorax and chest X-ray especially AP view is part of breathing component assessment of the primary survey in trauma management.

NCCT brain is a part of the secondary survey of trauma management and is performed after initial resuscitation and stabilisation. A CT scan of the head is performed within 1 hour if the GCS is ≤ 14 at the initial assessment. Other indications include a tense fontanelle, suspicion of an open/depressed or base of skull fracture, abnormal pupillary response, abnormal posturing, and a focal neurological defect. A head CT scan should also be performed if there are three or more vomiting episodes, a witnessed loss of consciousness or amnesia for ≥ 5 minutes.

Solution to Question 14:


The next immediate step in management of upper gastrointestinal bleeding is administration of intravenous fluids and electrolyte replacement.

USG findings and history of alcoholism are suggestive of cirrhosis leading to portal hypertension. The upper GI endoscopy provides a definitive diagnosis and aids in predicting the risk of bleeding with an option to treat via band ligation or sclerotherapy. (Option C)

The management of esophageal variceal bleeding involves:

- Resuscitation with IV fluids, airway protection(intubation to protect airway and prevent aspiration), and oxygen
- Blood transfusion to maintain Hb between 7-9g/dL
- Antibiotic prophylaxis with ceftriaxone before endoscopy (Option A)
- An attempt at hemostasis with octreotide or terlipressin or vasopressin
- Upper GI endoscopy within 12 hours after fluid resuscitation to confirm the source of the bleed
- Endoscopic variceal band ligation or sclerotherapy
- In case, of failure or rebleeding - Transjugular intrahepatic portosystemic shunt (TIPS)

Management of acute variceal bleeding



Solution to Question 15:

The clinical scenario along with biopsy showing cells arranged in nests along with amyloid deposits is suggestive of medullary carcinoma of the thyroid. Medullary carcinoma of the thyroid (MTC) arises from the parafollicular C cells.

Histopathology shows polygonal to spindle-shaped cells which form nests, trabeculae, or follicles. Amyloid deposits that are derived from the altered calcitonin molecules are present in the stroma. Calcitonin in the tumor cells and stroma is easily demonstratable by immunohistochemistry.

Features of MTC:


- Painful neck mass with cervical lymphadenopathy
- Dysphagia, dysphonia, or dyspnea if there is local invasion
- Diarrhea due to increased intestinal motility
- Cushingoid habitus due to ectopic ACTH production
- Blood-borne metastasis

Diagnosis is by careful family history, physical examination, raised calcitonin and CEA levels, and fine-needle aspiration biopsy of the mass.

Other options:

Option C and D: A Hurthle cell is a cell in the thyroid that is often associated with Hashimoto's thyroiditis as well as benign and malignant tumors (Hurthle cell adenoma and Hurthle cell carcinoma). Ashkanazy cells are another name given to the cells with Hurthle bodies in Hashimoto's thyroiditis.

Medullary Carcinoma of Thyroid



Solution to Question 16:

The lab reports of this patient, associated with a thyroid mass indicate primary hyperparathyroidism due to parathyroid adenoma. MIBI followed by focused parathyroidectomy to be done in this condition.

The patient presents with hypercalcemia (serum calcium 12.8 mg/dL) and elevated parathyroid hormone (PTH), which strongly indicates primary hyperparathyroidism (PHPT).

The ultrasonographic finding of a mass in the right lower lobe of the thyroid is consistent with a parathyroid adenoma, the most common cause of PHPT.

^{99m}Tc-Sestamibi (MIBI) scan is commonly used in conjunction with ultrasonography to localise the hyperfunctioning parathyroid gland before surgery.

Focused parathyroidectomy is the treatment of choice in such cases when a single adenoma is identified. This approach is minimally invasive and guided by the imaging findings.

Other options

Option A: A normal DEXA scan does not negate the need for surgery in a patient with severe hypercalcemia like this. Severe hypercalcemia (>12 mg/dL) is an absolute indication for surgery, irrespective of the results of the DEXA scan.

Option C: Observation is only appropriate for patients with mild hypercalcemia, no symptoms, and no target organ damage. Severe hypercalcemia (>12 mg/dL) is an absolute indication for surgery.

Option D: Creatinine clearance < 60 ml/min mandates surgery.

Solution to Question 17:

NCCT KUB (Non-contrast CT of the kidney, ureters, and bladder) is the best diagnostic modality for the clinical evaluation of ureteric stones.

NCCT is the preferred investigation for evaluation of:

- Renal stones except for indinavir stones
- Ureteric stones
- Head injury
- Salivary gland stones

Solution to Question 18:

The marked structure represents Corpora amylacea.

Corpora Amylacea are small, rounded, eosinophilic structures composed of glycoproteins and carbohydrates, commonly found in glandular tissues such as:

- Prostate: Seen in benign prostatic hyperplasia (BPH), within prostatic acini.
- Astrocytic processes of the CNS, associated with aging or neurodegenerative diseases.


Clinical Significance:

- Generally benign and age-related in both the prostate and CNS.
- In CNS, linked to diseases like multiple sclerosis or Alzheimer's.

Other Options:

Option B: Corpora arenacea: The pineal gland has a structure called the Corpora arenacea, located within the gland. These bodies are prone to an increase in calcification with advancing age. They are visible on x-ray and can be used as landmarks.


Pineal gland showing corpora arenacea



Option C: Lewy body: aggregates of insoluble fibrillar proteins called alpha-synuclein (specific) and ubiquitin in the neural cells. They are typically seen in Parkinson's disease, Lewy body

dementia, and a few variants of Alzheimer's disease.

Option D: Psammoma bodies : calcified structures seen in areas of cellular degeneration. They have a concentric laminated, or onion-like, appearance. Diagnostic feature of papillary thyroid carcinoma (PTC), the most common form of thyroid cancer.



Solution to Question 19:

CT abdomen and pelvis are not an adjunct to the primary survey. CT scan is a part of a secondary survey.

The primary survey identifies and manages the most immediately life-threatening pathologies first.

The secondary survey includes investigations that help identify all other injuries, excluding the apparent ones. This includes X-rays of the spine and extremities (limbs) and a CT of the abdomen, head, chest, pelvis and spine.

Solution to Question 20:

The patient presents with features suggestive of acute pancreatitis (epigastric pain, fever, history of cholelithiasis, guarding/rigidity, raised serum amylase and lipase, and leukocytosis). For acute pancreatitis, serum lipase is more specific than serum amylase.

Lipase has a higher specificity for diagnosing acute pancreatitis, as elevated amylase can occur in other conditions such as salivary gland disorders, intestinal obstruction, or renal failure. Lipase also remains elevated longer than amylase.

The diagnostic criteria for acute pancreatitis require at least two of the following three features:

- Characteristic Abdominal Pain:

Acute onset of severe epigastric pain, often radiating to the back.

Pain may worsen after eating and is associated with nausea and vomiting.

- Serum Lipase and/or Amylase Elevation:

At least 3 times the upper limit of normal (ULN).

Lipase is preferred due to its higher specificity and longer half-life.

- Imaging Findings:

Characteristic findings of pancreatitis on imaging (CT scan, MRI, or ultrasound), such as:

Pancreatic enlargement

Peripancreatic fluid collections


Inflammatory changes in and around the pancreas

Other Options:

Option A: Amylase levels normalize within 3-5 days after an episode of acute pancreatitis.

Option C: Hypocalcemia is more commonly observed, due to fat saponification in severe cases.

Option D: The degree of elevation of serum lipase does not correlate with the severity of acute pancreatitis. Severity is assessed using clinical scores like Ranson's criteria, APACHE II, or CT severity index.



Solution to Question 21:

TNM staging of the tumour is T3 N2 Mo.

In the given scenario:

- 6 x 4 cm mass - T3
- Ipsilateral Level I and Level II matted axillary lymph nodes- N2

- No distant metastasis - Mo.

In tumour size assessment, the greatest dimension is taken into consideration. For example, in a 6 x 4 cm lump, 6 cm is taken into account, and the tumour is classified as T₃ (>5 cm). If 2 lumps are present in the same breast, staging is according to the bigger lump. If lumps are present in bilateral breasts, separate staging is done for both lumps.

Any contralateral lymph node involvement is considered as distant metastasis.

Solution to Question 22:

The pressure sore is of grade 2 in this case.

Grade 2 of bed sore corresponds to partial-thickness skin loss involving the epidermis and dermis.

Pressure sores, also known as bedsores, pressure ulcers, or decubitus ulcers, are defined as tissue necrosis with ulceration due to prolonged pressure. When the external pressure exceeds the capillary occlusive pressure, blood flow ceases leading to tissue anoxia, necrosis, and ulceration. An external pressure of >30 mmHg leads to the formation of pressure sores.

The common sites of pressure sores in decreasing frequency are the ischium, greater trochanter, sacrum, heel, malleolus (lateral followed by medial), and occiput.

Given below are images showing the stages of pressure sore.



Prevention is the best treatment for pressure sores. They can be prevented with good skincare, the use of air-fluidized beds, special pressure-dispersion cushions, or foams. Fecal or urinary diversion can be considered in special cases.

The position of the bedridden patients should be changed once every 2 hours and wheel-chair bound patients should be taught to get off their chairs for 10 seconds every 10 minutes.

Surgical management of pressure sores follows the same principles involved in acute wound treatment. Prior to the operation, the pressure sores should be adequately debrided, and vacuum-assisted closure (VAC) can be used to promote the growth of granulation tissue required

for surgical closure. Large skin flaps that have muscle with sensory innervation are best for reconstruction.

VAC is done by applying intermittent negative pressure of approximately -125 mmHg to hasten debridement and promote granulation tissue formation. Negative pressure decreases edema, removes interstitial fluid, and increases blood flow. This decreases bacterial growth and increases cell proliferation.

Solution to Question 23:

Dysphagia of solids followed by liquids is the most likely presentation in a patient with ca esophagus.

Initially, patients have difficulty swallowing solids. Later, as the tumor grows and narrows the esophageal lumen, dysphagia progresses to liquids. This progression occurs because solid foods require a wider lumen to pass through, and the narrowing caused by the tumor becomes more critical as it advances.

Carcinoma esophagus is one of the most common malignancies manifesting in mid to late adulthood. It can be a squamous cell or adenocarcinoma.

Squamous cell carcinoma (SCC) affects the upper two-thirds of the esophagus while adenocarcinoma affects the lower third. Tobacco smoking and alcohol are the risk factors for SCC. Obesity and gastroesophageal reflux disease increase the risk of adenocarcinoma.

Dysphagia can be the most common presenting feature but may present late. Other features such as regurgitation, weight loss, vomiting, and odynophagia can be seen. Lymph node involvement indicates a poor prognosis.

Endoscopy is the first line of investigation done. CT is useful to rule out metastasis. Endoscopic ultrasonography is used to assess the depth of tumor invasion through the wall of the esophagus, adjacent organs, and lymph node metastasis.

Management options include radical esophagectomy, neoadjuvant chemotherapy, and palliative chemoradiation or endoscopic treatments.

Other options

Option B: Dysphagia of liquids followed by solids is typically seen in motility disorders, such as achalasia cardia, where there is functional obstruction rather than a mechanical obstruction.

Option C: Chest pain may occur in the CA esophagus, but it is not the most common presenting symptom. It usually arises in advanced cases or due to tumor invasion into surrounding structures.

Option D: A neck mass might occur in metastatic disease (e.g., supraclavicular lymphadenopathy), but not the typical initial presentation.

Solution to Question 24:

This oral cavity tumour is stage T4a.

In the given image, the oral cancer involves the floor of the mouth as well as the lip (adjacent structures). When adjacent structures are involved, the cancer meets the criteria for T4a disease. (resectable)

T4a is defined as "moderately advanced local disease" and includes tumors that invade floor of the mouth or the lip, and is resectable. These structures involve:

- Cortical bone (e.g., mandible or maxilla)
- Inferior alveolar nerve
- Floor of the mouth
- Skin of the face (chin or nose)

The T staging for oral cavity tumors, based on the AJCC 8th edition, is as follows:

- T1: Tumor $\leq$ 2 cm in greatest dimension, no invasion of adjacent structures.
- T2: Tumor $>$ 2 cm but $\leq$ 4 cm in greatest dimension, no invasion of adjacent structures. (Option A)
- T3: Tumor $>$ 4 cm in greatest dimension or minor invasion into structures such as the deep tongue muscle. (Option C)
- T4a (Moderately advanced local disease): Tumor invades adjacent structures like cortical bone, inferior alveolar nerve, floor of the mouth, or skin (chin or nose).
- T4b (Very advanced local disease): Tumor invades masticator space, pterygoid plates, skull base, or encases the internal carotid artery. (Option B)

The gold standard investigation for oral cancer includes an incisional biopsy. An MRI is the imaging modality of choice for assessing primary extent. Surgical management with chemotherapy and adjuvant radiotherapy is the mainstay of management.

Solution to Question 25:


The above clinical scenario of abdominal pain, ulcers in the jejunum and ileum, gastric pH $<$ 2, and elevated gastrin levels are suggestive of gastrinoma. It is commonly located in a triangle formed by the cystic duct, junction of D2-D3, neck and body of pancreas.

Gastrinoma is a neuroendocrinal tumour characterized by gastrin hypersecretion, resulting in gastrointestinal tract ulcers. It occurs more commonly in a sporadic pattern but can be associated with MEN-1 (multiple endocrine neoplasia-1) syndromes in the familial pattern.

Gastrinoma is most commonly located in the gastrinoma triangle (Passaro's triangle) bounded by the lines connecting the:

- cystic duct
- the junction between the second and third parts of the duodenum
- the junction between the neck and the body of the pancreas

Passaro's triangle (Gastrinoma triangle)



©Marrow

The most common presentations of gastrinoma are epigastric pain, diarrhoea, and peptic ulcers. The ulcers occur at typical locations (proximal duodenum) and atypical locations (distal duodenum, jejunum). Duodenal ulcers are the most common. Gastrinoma if associated with marked gastric acid production and recurrent peptic ulcer is called Zollinger-Ellison syndrome.

Diagnosis:


The gastric pH <2 , basal acid output >15 mEq/hr and serum gastrin concentration more than 1000 pg/ml (normal is <100 pg/ml) confirm the diagnosis of gastrinoma. When the diagnosis is in doubt (like gastrin levels 100-1000 pg/ml), then a secretin test is performed. A secretin test is considered positive when serum gastrin levels are more than 200 pg/ml on secretin administration and confirms the diagnosis of gastrinoma.

For localization of the tumour for surgical management, CT or MRI scans should be done first. If the tumour is not localized on these scans, then somatostatin receptor scintigraphy or endoscopic ultrasound is done for better localization.

Solution to Question 26:

The angle between the camera port and the working port is known as Azimuth's angle. The angle must be a minimum of 30° to avoid any clashing between these instruments.

Angles in laparoscopy



©Marrow

Other options

Option B: Manipulation angle is the angle between two working ports. The minimum angle should be 60° .

Option C: Crossing angle refers to the intersection of instruments inside the abdominal cavity.

Option D: The elevation angle is the angle between the working port and the horizontal line. The minimum angle must be 60° to avoid colliding the instrument with the abdominal wall.

Solution to Question 27:

Wound healing occurs by tertiary intention in this case. The wound is cleaned and left open for a few days. This allows soft tissues to drain and also ensures no infection is present before surgical closure can be done.

Wound closure is categorized into primary, secondary, or tertiary:

Primary closure is done for clean wounds with minimal tissue loss that are amenable to approximation of wound edges. Healing occurs by primary intention. Primary closure is ideally performed within 6 to 8 hrs of trauma, but wounds in vascular areas (e.g. face) may be delayed to up to 24 hours after trauma.

Secondary closure allows healing by secondary intention. The wound is left open to heal largely by the formation of granulation tissue and contraction. These wounds often have significant tissue loss precluding tension-free approximation of edges, with devitalized edges. Wound dressings are changed daily to aid in the formation of granulation tissue and subsequent contraction.

Tertiary closure, also known as delayed primary closure, results in a combination of healing by primary and secondary intention. The wound is thoroughly irrigated, debrided, and left open for 2 to 5 days before surgical closure. It is used for wounds that are grossly contaminated without significant tissue loss.

Solution to Question 28:

Total thyroidectomy is performed in patients with radiation-induced differentiated thyroid cancers irrespective of their size.

Management of differentiated thyroid cancers (DTC)

Differentiated thyroid cancers include papillary thyroid cancer, follicular thyroid cancer, and hurtle cell cancer.

Hemithyroidectomy is performed for DTCs between 1-4 cm without extrathyroidal extension or evidence of metastatic disease.

However, total thyroidectomy is performed in the following scenarios

- Tumor at least 4 cm
- Gross extrathyroidal extension
- Evidence of metastatic disease
- Radiation-induced DTC
- Familial nonmedullary thyroid cancer
- Multifocal bilateral DTC.

Central neck dissection is done only in case of involvement of level VI nodes.

If level VI plus other lymph nodes are enlarged - then total thyroidectomy + central neck dissection + modified radical neck dissection is done.

In case of tumor size ≥ 4 cm (T3 & T4), prophylactic central neck dissection is done.

Below is the TNM staging of differentiated thyroid carcinoma.

Papillary carcinoma is the most common thyroid carcinoma. It is most common in iodine-sufficient areas, among children, and in those exposed to ionizing radiation. It is more common in women than men (2:1). Dysphagia, dysphonia, and dyspnea are associated with local invasion. It usually presents as a cold nodule on the scan.

Lymph node metastasis is common. Lateral aberrant thyroid denotes a cervical lymph node invaded by papillary carcinoma. The most common sites of metastasis: lung & liver & bone & brain. It has an excellent prognosis, with a 10-year survival rate in excess of 95%.

Primary Tumor (T)TX Primary cannot be assessedTo No primary tumorT1a ≤1 cm, intrathyroidalT1b 1-2 cm, intrathyroidalT2 2-4 cm, intrathyroidalT3 >4 cm limited to the thyroid or any size with minimal extrathyroidal extension (muscle, soft tissue)T3a- >4cm limited to the thyroidT3b- gross extrathyroidal extension involving only the strap muscle.T4a Invades subcutaneous tissue, larynx, trachea, esophagus, or RLN T4b Invades prevertebral fascia or encases carotid or mediastinal vessels	Regional Lymph Nodes (N)NX Nodes cannot be assessedNo Negative nodesN1aPositive in levelsVI & VIIIN1bPositive unilateral, bilateral, contralateral lateral neck lymph nodes.Distant Metastasis (M)M0No distant metastasis M1Distant metastasis
---	--

Solution to Question 29:

Prolonged fasting is not necessary for enhanced recovery after surgery.

Early oral hydration and nourishment and avoidance of unnecessary IV fluid administration favors the Enhanced Recovery After Surgery (ERAS) protocol.

ERAS is an approach designed to speed the clinical recovery of the patient and reduce the cost and length of stay in the patient. It is achieved by optimizing the patient's health before surgery and then delivering evidence-based best care in the perioperative period.

It includes:

- Preference of minimally invasive procedures (e.g. laparoscopic) over open procedures.
- Minimal perioperative starvation.
- Preference for local or regional anesthesia, and adequate pain control with patient-controlled analgesia.
- Early removal of Ryles tube and early feeding techniques, rather than IV fluid administration.
- Early mobilisation planned physiotherapy.
- Discharge planning is started before the patient is admitted to hospital and involves support from stoma care nurses, physiotherapists etc.

Other Options:

Option B: Epidural anaesthesia is recommended in laparoscopic surgery for pain management.

Option C: Reduction of the hospital stay by 2-3 weeks: ERAS protocols do reduce hospital stay so as to plan for early discharge.

Option D: Beta-blockers are given after major surgery done under general anaesthesia to improve outcome: Beta-blockers may be used postoperatively in certain cases to control heart rate, manage perioperative stress, and improve outcomes, especially in high-risk cardiac patients.

Solution to Question 30:

In intermittent claudication, pain develops after walking some distance and is relieved by rest.

Intermittent(vascular) claudication is most commonly felt in the calf muscles because of the involvement of the superficial femoral artery in chronic vascular occlusive disease.

Intermittent claudication is the debilitating cramp-like pain felt in the leg muscles due to reduced blood flow and anaerobic metabolism brought on by peripheral arterial disease. It usually signifies mild to moderate chronic vascular occlusive disease. The pain is brought on by walking, not present on taking the first step. Relieved by rest in both the standing and sitting positions, usually within 5 minutes (option A). The distance that a patient can walk without stopping is known as claudication distance. It varies only slightly from day to day. The superficial femoral artery (70%) is the most commonly affected artery by claudication and manifests as pain in the calf muscles. Aorto-iliac occlusion may cause thigh or buttock claudication. Leriche's syndrome occurs due to aortoiliac occlusion and presents with a triad of claudication, impotence, and absent femoral pulses.

Boyd's classification of claudication:

- Grade I: Patient complains of pain after walking. If the patient continues to walk, due to increased blood flow in muscle and opening of collaterals metabolites causing pain are washed away and the pain subsides.
- Grade II: Pain persists on continuing to walk; but can walk with effort.
- Grade III: The patient has to take rest to relieve the pain.

Other options

Option A: Rest pain occurs when the occlusive disease progresses in severity. Though this pain is present during the day, it is exacerbated on lying down at night due to loss of gravity-dependent perfusion of the foot. This may be relieved by hanging the foot out of the bed to restore the gravitational effects.

Option B: The development of pain on taking the first step of walking is a classic sign of osteoarthritis.

Option C: Pain aggravated by standing and relieved by bending forward is seen in neurogenic claudication. It may be seen in cases of nerve compression from lumbar intervertebral disc prolapse or spinal stenosis.

Solution to Question 31:

The given clinical scenario of hemodynamic instability, decreased oxygen saturation (85%), GCS of 8/15 and absent breath sounds on the right side of the chest, with a history of a road traffic accident are suggestive of tension pneumothorax. The next step of management in this case is to secure the airway, give oxygen, insert the ICD, and secure the IV line.

According to the latest ATLS guidelines, the order of primary survey in a trauma patient is ABCDEF following the principle of 'what kills first should be managed first' -

- Airway
- Breathing
- Circulation with control of haemorrhage
- Disability(neurological)
- Exposure to environmental control
- Fluid resuscitation

After securing the airway, breathing can be assessed by auscultating for breath sounds and a chest X-ray to look for a pneumothorax or hemothorax. Administering oxygen through a bag and mask is useful during desaturation only if the airway is patent. Therefore, securing the airway is the most important next step for this patient.

Tension pneumothorax is mainly managed by inserting a wide-bore needle in the 5th intercostal space (ICS) followed by placing an intercostal drain in the 5th ICS. Once the airway and breathing are secured, hemodynamic instability must be corrected by securing an IV line to maintain circulation.

Other investigations such as NCCT head and CT head are done later on during further evaluation in the secondary survey.

Solution to Question 32:

The given clinical scenario of foul-smelling breath and dysphagia, and the given barium swallow X-ray showing the collection of contrast in a posterior pharyngeal pouch-like structure, are suggestive of Zenker's diverticulum.

It often presents with the following:

- Dysphagia
- Regurgitation
- Halitosis
- Gurgling sound on swallowing
- Aspiration

A barium swallow study may show a diverticulum arising from the midline of the posterior wall of the distal pharynx near the pharyngoesophageal junction. The diverticulum is typically noted at the C5-6 level. The outpouching may be transient, sometimes referred to as a pharyngeal pouch. The pouch is best identified during swallowing and is best visualized on the lateral view.

Solution to Question 33:

Truelove and Witts classification is used for acute severe ulcerative colitis.

Given below is the Truelove and Witts severity index for ulcerative colitis.

	Mild	Moderate	Severe
Bowel movements	< 4 per day	4 – 6 per day	> 6 per day
Blood in stool	Small	Moderate	Severe
Fever	None	< 37.5°C	> 37.5°C
Tachycardia	None	< 90 mean pulse	>90 mean pulse
Anemia	Mild	Hb > 75% of normal Hb	Hb < 75% of normal Hb
ESR	< 30mm	< 30mm	> 30mm
Endoscopic appearance	ErythemaDecreased vascular markingsFine granularity	Marked erythemaAbsent vascular markingsCoarse granularityContact bleedingNo ulcerations	Spontaneous bleedingUlcerations

Surgery NEET 2024

Question 1:


A male presented with discharge as shown in the image. What is the diagnosis?



- a) Fistula in ano
- b) Fissure in ano
- c) External Haemorrhoids
- d) Pilonidal sinus

Question 2:


A man presented to the emergency department after RTA with complaints of chest pain and breathing difficulty. An X-ray was done and is given below. What type of breathing pattern is observed in this patient?



- a) Cheyne stokes
- b) Paradoxical
- c) Biot
- d) Kussmaul

Question 3:

A mother brought her 3-month-old male child to the hospital, who is currently unable to pass urine. A similar issue was reported 15 days ago. The catheter could be passed without difficulty. The image of the micturating cystourethrogram is given below. What is the most appropriate management for this condition?



- a) Partial resection of verumontanum
- b) Endoscopic dilation of bladder neck
- c) Indwelling catheter
- d) Destruction of valves

Question 4:

What is the condition seen here?



- a) Gastric Ulcers
- b) Oesophageal varices
- c) Candidiasis
- d) Barret's esophagus

Question 5:


The baby was born with this deformity. What is the earliest age to correct this defect?



- a) 3 months
- b) 1 year
- c) 6 months
- d) 18 months

Question 6:

A patient with a history of tuberculosis presents with dysuria and sterile pyuria. The cystoscopy image of the patient is provided below. Based on the patient's history and the cystoscopic findings, which of the following conditions is most likely associated?



- a) Linear ulcer of the bladder

- b) Golf hole ureter
- c) Ureterocele
- d) Urethral stricture

Question 7:

What instrument is required for the following procedure?



- a) Ureteroscope
- b) Cystourethroscope
- c) Nephroscope
- d) Ultrasound guided scope

Question 8:


A patient with a history of umbilical swelling, initially asymptomatic for many years, now complains of pain over the swelling. The swelling was reducible initially but is now non-reducible. What is the appropriate management?



- a) Umbilectomy with herniorrhaphy
- b) Umbilectomy
- c) Conservative management with antibiotics
- d) Incision and drainage

Question 9:

A baby was brought in with complaints of intermittent feculent discharge from the umbilicus. What is the likely diagnosis?



- a) Umbilical polyp

- b) Ileal diverticulum
- c) Urachus
- d) Urachal cyst

Question 10:

A patient presents with renal colic and recurrent urinary tract infections. A CT scan reveals no significant findings. The patient undergoes a percutaneous nephrolithotomy and his urine analysis is as shown. Based on the clinical presentation and the urine analysis findings, what are these crystals made up of?



- a) Struvite
- b) Oxalate
- c) Uric acid
- d) Cystine

Question 11:

What is the treatment plan for this condition?



- a) Full thickness skin grafting
- b) Split thickness skin grafting
- c) Vacuum-assisted closure
- d) Flap

Question 12:

A patient with oral cancer underwent radical surgery and PMMC flap reconstruction as shown. What is the most important indicator for considering adjuvant radiotherapy in this patient?




- a) Close margin ≤ 5 mm

- b) Extra nodal extension
- c) Multiple lymph nodes
- d) Perineural invasion

Question 13:


Identify the stage of bed sore in the sacral region.



- a) Stage 1
- b) Stage 2
- c) Stage 3
- d) Stage 4

Question 14:

Based on the constituents, identify the fluid given to post-operative patients.



- a) Haemaccel
- b) Ringer lactate
- c) Isolyte P
- d) Isolyte M

Question 15:

A female patient presented with a breast mass, as shown in the image below, with concomitant liver metastasis. What is the recommended management plan for this patient?



- a) Simple mastectomy

- b) Neo adjuvant chemotherapy followed by surgery
- c) Modified radical mastectomy
- d) Radical mastectomy

Question 16:

An elderly male who presented with the following lesion as shown, is diagnosed with stage T3 carcinoma with no clinically palpable inguinal lymph nodes. What is the management strategy for this patient?



- a) Penectomy alone
- b) Penectomy + Orchidectomy
- c) Penectomy + Inguinal lymph node dissection
- d) Penectomy + Superficial inguinal lymph node dissection

Question 17:

The most commonly occurring complication of TPN in a sepsis patient in ICU:

- a) Hyperlipidemia
- b) Hypertriglyceridemia
- c) Reactive Hypoglycemia
- d) Hyperglycemia

Question 18:

A CT scan of a patient with a hydatid cyst is shown below. What is the type of hydatid cyst as per the Gharbi classification?



- a) Type 4
- b) Type 3
- c) Type 2
- d) Type 1

Question 19:


Identify the image:



- a) Ileal diverticulum
- b) Bladder exstrophy
- c) Omphalocele
- d) Gastroschisis

Question 20:

Identify the following:




- a) Fluoroscopy
- b) X-ray of alkali ingestion

- c) Barium swallow
- d) X-ray of acid ingestion

Question 21:


Identify pseudomembranous colitis from the given image.



- a) 3
- b) 4
- c) 1
- d) 2

Question 22:

What is the structure marked with the yellow arrow in the intraoperative image of spermatic cord of a patient with a congenital inguinal hernia?



- a) Obturator vein
- b) Femoral vein
- c) Inferior epigastric vein
- d) Testicular vein

Question 23:

A chronic tobacco chewer presented with an ulcerative-nodular lesion of 4x5 cm over the lip, as shown below. Which investigation would most likely confirm his diagnosis?



- a) Superficial incisional biopsy along with normal tissue

- b) Deep incisional biopsy of Ulcer along with normal tissue
- c) Excisional biopsy along with normal tissue
- d) Brush biopsy

Question 24:


A patient presented with recurrent swelling in his lower lip for the past 6 months as shown in the image. What is the diagnosis and treatment option for the patient?



- a) Plunging ranula ; Incision and drainage
- b) Mucous retention cyst ; Excision
- c) Ranula ; Incision and Drainage
- d) Mucocele ; Aspiration

Question 25:

A male child experienced acute scrotal pain while playing. He was taken to the emergency department and a USG doppler was performed, followed by immediate surgical exploration. The following finding was seen. What is the probable diagnosis?



- a) Testicular hematoma
- b) Strangulated inguinal hernia
- c) Torsion of testicular appendage with Gangrene
- d) Testicular torsion with Gangrene

Question 26:

Identify the procedure done in the given image.



- a) Bilateral Orchidectomy with bilateral inguinal node dissection
- b) Hernia repair

- c) Sentinel Lymph node biopsy
- d) Total penectomy with inguinal node dissection

Question 27:

The patient presented with headaches, repeated episodes of elevated blood pressure, and tachycardia. Urine shows elevated levels of vanillylmandelic acid. Her CT scan revealed a retroperitoneal tumor in the right lumbar region. Which among the following statements is true regarding the condition?

- a) It can be associated with MEN 2A syndrome
- b) It is commonly seen as a bilateral condition
- c) It is most commonly malignant
- d) It is more commonly seen in children

Question 28:


What is the ideal pressure exerted by the method of dressing for a non-healing leg ulcer shown below?



- a) 40-50 mmHg
- b) 60-70 mmHg
- c) 110-130 mmHg
- d) 90-100 mmHg

Question 29:

A patient is diagnosed with Stanford type B aortic dissection. What is the most appropriate initial management of this patient?



- a) Administer beta blocker and intensive blood pressure monitoring
- b) Bypass graft
- c) Percutaneous endovascular stent
- d) Balloon dilation

Question 30:

A patient presented with dysuria and increased frequency of micturition. He has a history of recurrent urinary tract infections with *Proteus* species. The radiographic image is given below. Identify the calculi shown.



- a) Cystine
- b) Calcium Phosphate
- c) Xanthine
- d) Calcium oxalate

Question 31:

Identify the given drain.



- a) Underwater seal drain
- b) Closed suction drain

- c) Penrose drain
- d) Nelaton drain

Question 32:

Identify the ulcer shown in the image.



- a) Arterial ulcer
- b) Venous ulcer
- c) Callous ulcer
- d) Neuropathic ulcer

Question 33:

A patient underwent surgery for a leg ulcer due to burns. What is the mechanism by which a graft receives its nutritional supply on postoperative day 3?



- a) Inosculation
- b) Imbibition
- c) Diffusion
- d) Neovascularisation

Question 34:

A woman noticed a neck swelling 2 months back. Clinical examination reveals a solitary nodule on the right side as shown. What is the most appropriate initial investigation for this patient?



- a) Radioisotope I-131 assay

- b) PET Scan
- c) Radioisotope I-123 assay
- d) Serum TSH, T₃ and T₄

Question 35:

A 76 year old elderly man presents with confusion. He is on antihypertensive medication, and on aspirin after a recent MI. He recalls a trivial fall from chair 3 weeks ago. Other neurological examination was normal. What is the likely diagnosis?



- a) Epidural hemorrhage
- b) Subarachnoid hemorrhage
- c) Subdural hemorrhage
- d) Normal

Question 36:

A 45-year-old male presents with acute retention of urine which is drained via the suprapubic route. A retrograde urethrogram was done and is as shown below. What is the most appropriate management?



- a) Serial dilatation
- b) Urethral repair with graft from buccal mucosa
- c) Metallic catheter placement
- d) Endoscopic urethrotomy

Question 37:

A patient diagnosed with GERD has no improvement after long-term medical therapy. He underwent surgical treatment for GERD. Identify the procedure.



- a) Toupet's fundoplication

- b) Dor's fundoplication
- c) Watson's fundoplication
- d) Nissen's fundoplication

Answer Key

Question No.	Correct Option
1	a
2	b
3	d
4	b
5	c
6	b
7	b
8	a
9	b
10	d
11	b
12	b
13	d
14	b
15	a
16	d
17	d
18	c
19	b
20	c
21	a
22	d
23	b
24	b
25	d
26	d
27	a
28	c

29	a
30	b
31	b
32	b
33	a
34	d
35	c
36	b
37	d

Detailed Explanations

Solution to Question 1:

The diagnosis, considering the purulent discharge shown in the image, is a fistula in ano.


A fistula-in-ano is an abnormal hollow tract or cavity that is lined with granulation tissue and that connects a primary opening inside the anal canal to a secondary opening in the perianal skin. The most common cause is a perianal abscess. Other causes include tuberculosis or Crohn's disease.

Symptoms:

- Perianal pain
- Discharge- intermittent purulent discharge (which may be bloody)

Goodsall's rule can be used as a guide in determining the location of the internal opening of the fistula.

- Fistulas with an external opening anteriorly connect to the internal opening by a short, radial tract.
- Fistulas with an external opening posteriorly track in a curvilinear fashion to the posterior midline.



Exceptions to Goodsall's Rule

However, exceptions to this rule often occur: Fistulas with an anterior external opening greater than 3 cm from the anal margin usually track to the posterior midline.

Investigations:

The investigations done for fistula-in-ano are as follows:

- Fistulogram
- Endoanal ultrasound
- MRI
- Computed tomography (CT) scan
- Endocoil MRI
- Hydrogen peroxide-enhanced 3-D endoanal ultrasonography

MRI is the gold standard investigation for fistula imaging. It helps us to identify the secondary extensions of the track, which when missed may cause persistence of the fistula.

Treatment:

- Control of constipation
- Sitz bath
- Fistulotomy
- Fistulectomy
- Ligation of intersphincteric fistula tract (LIFT)
- Setons
- Advancement flaps

Other options:

Option B: A fissure in ano is a linear ulcer usually found in the midline, distal to the dentate line. It is one of the most common causes of anal pain and bleeding. They typically start with a tear in the anoderm within the distal half of the anal canal which then triggers cycles of recurring anal pain and bleeding, which lead to the development of a chronic anal fissure. The exposed internal sphincter muscle within the bed of the fissure frequently spasms, which not only contributes to severe pain but also can restrict blood flow to the fissure, preventing its healing.

Option C: Hemorrhoids are cushions of submucosal tissue containing venules, arterioles, and smooth muscle fibers located in the anal canal. External hemorrhoids are distal to the dentate line and are covered with anoderm; these may engorge, causing pain. Thrombosis of these external hemorrhoids causes severe pain.

Option D: Pilonidal sinus disease is commonly seen in the natal cleft, more in men, usually after puberty. Drivers and hairdressers are more commonly affected. The pilonidal sinus opening is usually seen in the midline between the sacrococcygeal joint's level and the coccyx's tip. The other sites that can be affected include the interdigital cleft and axilla.

Solution to Question 2:


The type of breathing pattern observed in this patient with the X-ray showing contiguous rib fractures on the right side suggestive of a flail chest is paradoxical respiration.

A flail chest usually results from blunt trauma associated with 3 or more rib fractures in 2 or more places. It arises when these injuries cause a part of the chest wall to move independently of the rest of the chest wall. It is diagnosed clinically by observing paradoxical respiration. On inspiration, the loose segment of the chest wall gets displaced inwards, and on expiration, it moves outwards. They are at higher risk of developing pneumothorax/hemothorax. CT scan is the gold standard investigation.


Management includes oxygen administration, adequate analgesia (including opiates) or epidural analgesia, and physiotherapy. Intermittent positive pressure ventilation (IPPV) is reserved for patients developing respiratory failure despite adequate analgesia and oxygen

Treatment algorithm for treatment of flail chest with respiratory failure:

Treatment algorithm of flail chest with respiratory failure



Different patterns of breathing observed are shown below:



Solution to Question 3:

The given image of a micturating cystourethrogram (MCU) shows dilated prostatic urethra and transverse filling defects corresponding with the posterior urethral valves and the most appropriate management for this condition is destruction of valves.

Posterior urethral valves are the most common cause of severe obstructive uropathy in children. The pathological valves are membranes that have a small posterior slit within them, which typically lie just distal to the verumontanum. They act as flap valves and are obstructive to antegrade urinary flow.


Diagnosis is most commonly made antenatally with ultrasound, which demonstrates bilateral hydronephrosis above a distended bladder in fetus along with oligohydramnios and pulmonary hypoplasia.

If the diagnosis is not made antenatally, then babies typically present with urinary infections in the neonatal period or with uremia and renal failure. Clinically a poor urinary stream, dribbling, and palpable bladder (due to diverticula) can be present. The investigation of choice is micturating cystourethrography.


The initial step in treatment is to insert a urinary catheter to drain the urine and decompress the bladder and upper urinary tracts. A small polyethene feeding tube (No. 5 or No. 8 French) is inserted into the bladder, which can be difficult because the tube's tip can coil in the prostatic urethra. A Foley catheter should not be used because the balloon can cause severe bladder spasms, which can result in severe ureteral obstruction.

Endoscopic valve ablation and symptomatic management of urinary tract infections are the definitive treatments.

The below image shows the micturating cystourethrogram (MCU) with dilated prostatic urethra and transverse filling defects (constriction) corresponding with the posterior urethral valves.



The ultrasound shows the typical 'key-hole' sign in which the bladder and the proximal urethra are dilated.



Solution to Question 4:

The given endoscopic image of a patient suggests esophageal varices.

Esophageal varices are tortuous dilated veins within the mucosa and submucosa of the distal esophagus and proximal stomach. Variceal hemorrhage is an emergency that can be treated medically by inducing splanchnic vasoconstriction or endoscopically by sclerotherapy, balloon tamponade, or variceal ligation.

Other options:


Option A: Gastric ulcers are often found near the lesser curvature of the stomach, especially in the antral area. Although duodenal ulcers are more common overall, gastric ulcers are frequently located near the lesser curvature, within a few centimetres of the pylorus. A significant cause of these ulcers is infection with *Helicobacter pylori*.

The following is an endoscopic image of Gastric ulcer:



Option C: Esophageal candidiasis is a fungal infection of the esophagus caused by the overgrowth of yeast, commonly *Candida albicans*. This condition is often seen in individuals with weakened immune systems, such as those with HIV/AIDS, diabetes, or those undergoing immunosuppressive treatments.

The following is an endoscopic image of esophageal candidiasis:



Option D: Barrett's esophagus is a type of metaplasia at the gastro-esophageal junction due to chronic gastroesophageal reflux disease (GERD). The squamous esophageal mucosa is converted to a columnar/intestinal type that is better suited to handle acid reflux. It is associated with an increased risk of esophageal adenocarcinoma.

The following shows an endoscopic image of Barrett's esophagus:



Solution to Question 5:

The earliest age to correct this defect in a child with bilateral cleft lip and palate is 6 months.


Cleft lip repair is done between 3-6 months of age and palate repair is done between 6-12 months of age. As this child's deformity needs to be corrected by palatal repair, the best answer in this scenario is 6 months.

A cleft lip is due to the failure of the fusion of maxillary prominence with medial nasal prominence. It can be unilateral or bilateral. It can vary in severity from a small notch in the upper lip to a double cleft extending into both nostrils. There is also associated underdevelopment of the part of the maxilla that bears the incisors. Millard cleft lip repair is done by rotating the local nasolabial flaps. Cleft lip repair is preferably done between 3-6 months of age.

A cleft palate can occur in the primary palate (anterior cleft) or the secondary palate (posterior palate). A Complete cleft palate involves both the primary and secondary palate. An incomplete cleft palate involves only the secondary palate.

In the bilateral complete cleft palate, a Y-shaped cleft is seen with gaps between the primary and secondary palate and between the two halves of the secondary palate. In addition, there is a bilateral cleft upper lip.

The different types of facial clefts are shown below:



- Unilateral cleft lip with alveolar involvement
- Bilateral cleft lip with alveolar involvement
- Unilateral cleft lip associated with cleft palate
- Bilateral cleft lip and palate
- Cleft palate only

Note: Palate repair is done between 6-12 months of age. The palatal repair is done maintaining the balance between speech outcomes and mid-facial growth. The generally agreed timeline is before speech development. However, the exact timeline is still debated and no uniform consensus is available.

Solution to Question 6:

The cystoscopic findings are suggestive of a "golf hole ureter", which is most likely associated with genitourinary tuberculosis (TB).


Genitourinary TB is caused by the dissemination of the organism through the bloodstream and is thus always a secondary TB. There is either reinfection or reactivation of old TB. It can affect the kidneys, ureters, urinary bladder, prostate, seminal vesicles, epididymis, and vas deferens.

Lesions caused by tuberculosis in the kidney include:

- Golf hole ureteric orifice - This is due to fibrosis causing the rigid withdrawn dilated ureteric orifice to look like a golf hole (cystoscopic finding).
- Papillary ulcer - earliest lesion
- Pyonephrosis - caused by multiple scattered papillary ulcers undergoing caseous necrosis, which later become filled with pus
- Perinephric abscess - by extension of pyonephrosis or tuberculous renal abscess


- Pseudocalculi - occurs when calcium gets deposited in a papillary ulcer healed with fibrosis
- Ghost calyces - caused by the destruction of the calyces
- Thimble bladder -the entire urinary bladder gets fibrosed, stiff, and unable to dilate and accommodate urine (IV urogram finding).
- Putty/cement kidney - amorphous dystrophic calcification eventually involves the entire kidney (seen in X-ray and CT)

The following X-ray KUB shows a thimble bladder:



Ureteric lesions can also be seen in genitourinary tuberculosis. They include:

- Kerr's kink - kinking of the ureteropelvic junction (UPJ)
- Ureteral stricture
- Beaded ureter
- Shortening of the ureter
- Golf hole ureter - Longitudinal fibrosis leads to the shortening of the ureter and pulling up the orifice as a gaping hole. Reimplantation of ureter is done.



Solution to Question 7:

The instrument required for placing the Double J stent (DJ stent), as shown in the radiograph is a cystourethroscope.

Cystourethroscopy allows for the examination and access of the bladder and urethra, which is crucial for guiding the DJ stent from the bladder into the ureter. This procedure involves inserting a flexible or rigid endoscope through the urethra and into the bladder, where the stent can be manoeuvred into the ureter.

While ureteroscopy is typically used to visualize and treat issues directly within the ureter, cystourethroscopy provides the necessary access and visualization to position the DJ stent correctly. This approach is effective for ensuring the stent is placed accurately, particularly in cases where access to the bladder is essential.

Other options:

Option A: A ureteroscope is used to examine and treat the ureters (the tubes connecting the kidneys to the bladder). It is commonly used for removing kidney stones or evaluating issues in the ureter.

Option C: A nephroscope is a type of scope used to examine the kidneys directly. It is often used in procedures like percutaneous nephrolithotomy (PCNL) to remove large kidney stones.

Option D: An ultrasound-guided scope refers to an endoscope or medical device used in conjunction with ultrasound imaging to guide procedures. It's often used in various parts of the body, including the gallbladder, liver, or kidney to help visualize structures during minimally invasive surgeries or biopsies.

Solution to Question 8:

The appropriate management for this patient with a diagnosis of umbilical hernia with painful, non-reducible swelling is umbilectomy with umbilical herniorrhaphy.

An umbilical hernia occurs when a defect in the umbilical or median raphe area allows abdominal contents to protrude. In adults, these hernias often present as a swelling near the umbilicus that can become painful and non-reducible over time. Conditions that cause stretching and thinning of the midline raphe, such as obesity, pregnancy, or liver cirrhosis with ascites, increase the risk of such complications.

The management of painful, non-reducible umbilical hernia typically involves surgical intervention due to the risk of strangulation and bowel obstruction. Umbilical herniorrhaphy with umbilectomy is the standard approach, addressing both the hernia and the associated defect.

In children, the umbilical defect is usually present at birth and typically closes as the umbilical cord stump heals within the first week. If this closure process is delayed, it can result in herniation during the neonatal period. Additionally, in adults, the umbilical ring can stretch and reopen, leading to similar herniation.


Solution to Question 9:

The presence of feculent discharge at the umbilicus with the persistence of the vitellointestinal duct is suggestive of vitelline fistula which occurs due to an ileal diverticulum.

The given clinical image is that of a congenital umbilical anomaly such as patent urachus, urachal cyst, and vitelline cyst (resulting from the persistence of omphalomesenteric duct). The next best step is the USG abdomen to diagnose if the anomaly is cystic, fibrous or fistula and its connections.

Vitello intestinal duct or omphalomesenteric duct is the connection between the yolk sac and the midgut during embryonic life, which eventually gets obliterated. The persistence of the vitellointestinal duct leads to various abnormalities depending on which part of the duct remains patent.


- Persistence of the proximal intestinal end of the duct leads to the Ileal diverticula/Meckel's diverticula.
- Persistence of the distal end near the umbilicus leads to the umbilical polyp or sinus.
- Persistence of the distal end with mucus retention leads to the cyst, which may be associated with the intestine or the umbilicus by a fibrous band.
- A fully patent vitellointestinal duct can lead to an enterocutaneous fistula.



Patent urachus occurs due to the failure of urachus to obliterate, resulting in a connection between the bladder and the umbilicus.

Urachal cyst occurs when the urachus fails to obliterate but there is no connection between the bladder and the umbilicus.

The image below shows a urachal cyst (UC) attached to Meckel's diverticulum (MD) on one side and a urinary bladder (UB) on the other side with a patent tract- urachus (U).



Solution to Question 10:

The image shows a hexagonal crystal, suggesting cystine stones. These stones are a recurrent feature in patients with cystinuria.

Cystinuria is the most common inborn error of amino acid transport. It occurs due to defective basic amino acid transporter in the proximal tubules of the kidney. This leads to the excretion of cystine(dimer of cysteine), ornithine, lysine and arginine, abbreviated as COLA.

The disease manifests as precipitation of cystine stones in the urine leading to renal colic. Oral hydration is the mainstay of treatment.

Crystals in urine are usually associated with kidney stones. Crystals form in the urine depending upon a variety of factors including:

- The degree of concentration of constituent molecules
- The urine pH
- The presence of inhibitors of crystallisation

Calcium oxalate and uric acid crystals can sometimes be found in normal urine that has been left to stand, due to crystal formation *ex vivo*.

Solution to Question 11:

The given clinical image is suggestive of partial thickness burns and the treatment plan for this condition is split thickness skin grafting.

Split-thickness skin graft (Thiersch graft): Partial-thickness skin grafts consisting of the epidermis and a portion of the dermis are called split-thickness skin grafts. Less dermis in the split-thickness skin graft translates into less primary contraction (the degree to which a graft shrinks in surface area after harvesting and before grafting) and more secondary contraction (the degree to which a graft shrinks during healing). Humby knife is used to harvest split-thickness skin grafts.

Advantages of split skin graft over full-thickness skin graft:

- Large grafts can be harvested.
- In the case of thinner split-thickness grafts, the donor site heals more rapidly and can be reused in severe cases such as burns.
- Uptake is good.
- Meshing can be done, i.e., small splits are made in graft and any exudates collected can come out through this, improving graft take-up. It also allows the graft to expand.

Other options

Option A: Full-thickness skin grafts (Wolfe graft) include the epidermis and entire dermis, with portions of the sweat glands, sebaceous glands, and hair follicles. More dermis in the full-thickness skin graft translates into more primary contraction and less secondary contraction.

Option C: VAC is a negative pressure dressing. An occlusive dressing is placed over the sore and it is connected to VAC that generates -125mmHg of pressure. It is used in chronic non-healing ulcers, venous ulcers, diabetic ulcers, burn wounds, and bed sores.

Venous ulcers are due to chronic venous insufficiency and are particularly seen in the lower ■ of the leg above the ankle on the medial aspect, known as the gaiter zone. Ambulatory venous

hypertension is the most acceptable theory in the pathophysiology of venous ulcers. Venous ulcers have a sloping edge and the floor contains granulation tissue covered by slough and exudate. Marjolin's ulcer is a type of squamous cell carcinoma seen in chronic long-standing venous ulcers.



Option D: Flaps are tissues with a blood supply, providing vascularity to a new area. Types include:

- Random Flaps: Rectangular, with no specific blood supply, length-to-breadth ratio $\leq 1.5:1$.
- Axial Flaps: Long flaps based on known blood vessels, suitable for large distances.
- Pedicled/Isolated Flaps: Axial supply allows them to be moved on a stalk or fully isolated.
- Free Flaps: Blood supply disconnected and reconnected via microsurgery.
- Composite Flaps: Combine tissues like skin with bone or muscle.
- Perforator Flaps: Use small vessels from major blood vessels to supply the surface.

Solution to Question 12:

The most important indicator for considering adjuvant radiotherapy in this patient is extranodal extension (ENE).

For patients with oral cavity cancer, ENE is a key indicator for adjuvant radiotherapy. Adjuvant therapy, typically involving external beam radiotherapy, is recommended based on pathological features such as ENE, close margins (1–4.9 mm), or involved margins (≤ 1 mm), lymphovascular invasion (LVI), and perineural invasion (PNI).

The presence of one major criterion - ENE or margins ≤ 1 mm or two minor criteria - close margins (1–4.9 mm), multiple involved nodes, largest node ≥ 3 cm, LVI/PNI or T3/4, guides the decision for additional treatment.

Major criteria

- Extranodal invasion

- Lymphovascular invasion
- Perineural invasion
- T3/T4

Minor criteria

- Close margins (1–4.9 mm)
- Multiple involved nodes
- Largest node ≥ 3 cm

Landmark trials have demonstrated that high-dose cisplatin added to radiotherapy enhances outcomes in high-risk patients, particularly those with ENE and positive margins.

Note: The PMMC flap is the most commonly used flap for soft tissue reconstruction in head and neck surgery. The pectoralis flap can be harvested as a musculocutaneous flap or muscle-only flap and is based on the pectoral branch of the thoracoacromial artery. Following flap harvest, a subcutaneous tunnel is created from the donor site, over the clavicles, to the defect. Division of the pectoral nerve branches ensures atrophy of the muscle to reduce the bulge over the clavicle over time.

Solution to Question 13:

The given image shows a pressure sore with extensive tissue destruction, including damage to the muscle layer and thus it belongs to stage 4.

Pressure sores also known as bed sores, pressure ulcers and decubitus ulcers are ulcers that occur due to prolonged pressure on certain sites thereby leading to tissue necrosis. The commonest sites involved are ischium, greater trochanter, sacrum, heel, malleolus (lateral $\&$ medial), and occiput. Paraplegic patients, elderly patients, and severely ill patients are at a higher risk of developing pressure sores.

Management is primarily proper skin care with pressure dispersion cushions, air-fluidized beds, and urinary or fecal diversion if required. Patients who are bed-bound should be turned once in 2 hours. Those in a wheelchair should lift themselves off the seat for 10 seconds every 10 minutes.

Surgery is the last resort and is done after a proper initial debridement and vacuum-assisted closure. Vacuum-assisted closure is a wound closure technique in which negative pressure is applied to the wound which leads to a decrease in edema. This hastens the debridement and formation of granulation tissue. Skin flaps with a muscle are also used for closure.

Solution to Question 14:

The fluid shown in the image is ringer lactate or Hartmann's solution.

Ringer lactate, or Hartmann's solution, is a crystalloid made up of sodium 131 mEq, chloride 112 mEq, potassium 5 mEq, bicarbonate (as lactate) 29 mEq, and 2 mEq of calcium. It is devoid of glucose.

It is useful in the following conditions:

- Correction of severe hypovolemia
- Metabolic acidosis
- Postoperative period, burns, diarrhoea
- Correction of hypokalemia
- Diabetic ketoacidosis.

It should be avoided in severe shock, liver diseases, vomiting, nasogastric aspiration, and metabolic alkalosis. It cannot be used along with blood transfusion since calcium in the ringer lactate can bind with citrate in the transfused blood leading to clot formation.

Other options:

Option A: Haemaccel is a colloid and contains 145 mmol of sodium, 6.25 mmol of calcium, and 5.1 mmol of potassium. It is mainly used for rapid volume replacement in circulatory shock, burns, and trauma. These release histamine and hence, an H₂ blocker has to be supplemented. It is also used in heart-lung machines.

Options C: Isolyte P is a crystalloid. Isolyte P is commonly used in fluid therapy for paediatrics. One litre contains 50 grams of glucose, 25 mEq of sodium, 22 mEq of potassium, 3 mEq of magnesium, and 23 mEq of acetate. Hyponatremia, renal failure, hyperkalemia, and hypovolemic shock are some of the contraindications.

Option D: Isolyte M is a crystalloid. Isolyte M is used as a maintenance fluid. It mainly contains 50 grams of glucose, 40 mEq sodium, 38 mEq chloride, 35 mEq potassium, 15 mEq phosphate, and 20 mEq acetate. It is useful in correcting hypokalemia in patients with good renal function, and acidosis. It is contraindicated in hyponatremia.

Solution to Question 15:

The image of a large breast mass with an unevenly bosselated surface, overlying skin ulceration, and concomitant liver metastasis is indicative of a malignant phyllodes tumor which is managed by simple mastectomy.

Phyllodes tumors spread primarily through the bloodstream (hematogenous). Lymph node metastases are uncommon. Large or malignant phyllodes tumors may require a simple mastectomy, but axillary dissection is not recommended (Option C and D) due to the rare involvement of axillary lymph nodes. The role of adjuvant chemotherapy (Option B) in malignant phyllodes tumors remains uncertain and is generally not recommended as a standard treatment.

Phyllodes tumor also known as a sero-cystic disease of Brodie or cystosarcoma phyllodes, originates from hormonally responsive stromal cells of the breast. It is usually a benign condition and affects women > 40 years of age, but can be seen in younger women. Sometimes, may develop malignant transformation.

The clinical features include large, sometimes massive, with an unevenly bosselated surface, ulceration of overlying skin occurs because of pressure necrosis and it remains mobile on the chest wall.

There is no role of FNAC in phyllodes tumor. Histopathology of the lump biopsy shows increased stromal cellularity, overgrowth, and typical leaflike architecture. Tumors are treated by excision with margins of at least 2 cm to prevent local recurrence.

Solution to Question 16:


The most appropriate management for this patient with stage T3 carcinoma of the penis with no clinically palpable inguinal lymph nodes is penectomy with superficial inguinal lymph node dissection.

In penile carcinoma, total amputation of the penis is done if the stump is ≤ 2 cm after resection of the tumour and partial penectomy is done if the stump is > 2 cm. Carcinoma of the penis is commonly a squamous cell carcinoma. It arises from the skin of the glans penis or the prepuce. The lesion can be flat or warty in appearance.

- T1- confined to the skin
- T2- invades the corpus spongiosum or cavernosum.
- T3- invades the urethra
- T4- invades adjacent structures.

The first lymph node to be involved is the superficial inguinal lymph node and thus, superficial inguinal lymph node dissection or sentinel lymph node biopsy (SLNB) helps evaluate the spread of penile cancer.

Tumours of low-grade and low-stage are managed by Mohs' surgery. Tumors involving the glans penis require glansectomy, and in advanced cases require partial penectomy.



Solution to Question 17:

The most commonly occurring complication of TPN in a sepsis patient in the ICU is hyperglycemia.

To monitor or prevent complications in patients on TPN, the following steps may be taken:

- Infections
- Monitor the development of glucose intolerance as it is the earliest sign of systemic sepsis. (Note: glucose intolerance- glucose not processed effectively, leading to hyperglycemia)
- Use subclavian vein preferably
- Metabolic complications
- BUN and LFTs weekly
- Serum electrolytes every 2-3 days
- Intestinal atrophy
- Provide at least some nutrients enterally if feasible

TPN also referred to as central parenteral nutrition is used for patients who can't absorb nutrients through their digestive tract, often due to conditions like short bowel syndrome from extensive intestinal resection or fistulas. It can also be temporary until other feeding methods, like a nasojejunal tube, are established.

TPN typically comes in a 3-liter bag with a lipid emulsion, amino acids, glucose, electrolytes, and vitamins. The energy provided is about 150–250 kcal per gram of protein nitrogen, with 30–50% from fat. Folic acid is added weekly, with other vitamins given daily, and long-term TPN may require vitamin B12 supplementation.

Solution to Question 18:

The given CT scan showing the hydatid cyst with detached membranes(Water Lily sign) is classified as CE-3 according to the WHO classification and Type 2 according to the Gharbi classification.

Classification of Hepatic Hydatid Cysts					
WHO-IWGE 2001	Gharbi 1981	Description	Stage		
CE 1	Type I	Unilocular anechoic cystic lesion with double-line sign	Active		
CE 2	Type III	Multiseptate, rosette-like, honeycomb cyst	Active		
CE 3a	Type II	Cyst with detached membranes (Water-lily sign)	Transitional		
CE 3b	Type III	Cyst with daughter cysts in a solid matrix	Transitional		
CE 4	Type IV	Cyst with heterogeneous hypoechoic/hyperechoic contents. No daughter cysts	Inactive		
CE 5	Type V	Solid cyst with a calcified wall	Inactive		

	CE1	CE2	CE3a	CE3b	CE4	CE5
US						
MRI						
CT						


©Marrow

Hydatid Cyst is a zoonotic infection caused by the Echinococcus species(dog tapeworm). Dogs are the definitive hosts of E. granulosus. Sheep are the usual intermediate hosts, but humans are accidental intermediate hosts. Humans are end-stage hosts of the parasite since there is no human-to-human transmission.

The clinical presentation of a hydatid cyst is largely asymptomatic until complications occur. The most common presenting symptoms are abdominal pain, dyspepsia, and vomiting. The most frequent sign is hepatomegaly.

Approximately 75% of hydatid cysts are located in the right lobe of the liver and are solitary.

CT scan is the best imaging modality for hydatid cysts. It shows the presence of a space-occupying lesion with a smooth outline with septa. Diagnosis is suggested by the finding of a multiloculated cyst on ultrasound and is further supported by the finding of a floating membrane within the cysts on a CT scan. Active cysts contain a large number of smaller daughter cysts (multi-vesicular rosette-like cystic lesions with daughter cysts) which is characteristic of hydatid cysts of the liver.




If there is a disruption of the inner layers of the cyst, the following signs may be noted:

- Water lily sign/ camalote sign - air-fluid level with a floating membrane (as shown below)
- Rising sun sign/ serpent sign - a double-wall, dry cyst with crumpled membranes lying at its bottom
- Empty cyst sign - when all the contents are removed




The ultrasound of an active hydatid cyst shows multiple septations and a rosette or honeycomb appearance (as shown below)



The serpent sign is seen as wavy membranes within the cyst, on CT imaging.



A ball of wool sign is seen on ultrasound when the contents settle in the dependent portion of the cyst, as seen below.



The water-lily sign is seen in hydatid infections when there is detachment of the endocyst membrane which results in floating membranes within the pericyst that mimic the appearance of a water lily, on USG.



Solution to Question 19:

The given image showing a midline defect situated in the lower abdominal region below the umbilicus with an exposed bladder is suggestive of ectopia vesicae/ bladder exstrophy.

Ectopia vesicae/ bladder exstrophy is a congenital abdominal wall defect wherein the bladder protrudes from the lower abdominal wall and its mucosa is exposed. This results in the separation of the pubic rami and the rectus muscles.

The condition can be associated with complete epispadias with dorsal chordee, undescended testes, and inguinal hernias in males. Females may also have epispadias and separation of the two halves of the clitoris (bifid clitoris). The condition can be associated with rectal prolapse and short stature in both sexes.

Untreated exstrophy can cause urinary incontinence and is associated with an increased incidence of bladder cancer. Management involves a staged surgical reconstruction of the bladder and genitalia.

Other options:

Option A: The primitive gut is connected to the definitive yolk sac through a narrow stalk-like communication known as the vitellointestinal duct. In about 2% of the population, a small section of the vitellointestinal duct persists to form an outpouching of the ileum called the Meckel diverticulum or ileal diverticulum.

Option C: Gastroschisis is a condition where the intestinal loops herniate through the defect in the abdominal wall. The defect usually lies on one side of the umbilicus (Right side commonly). The loops of the bowel are not covered by amnion. It is associated with elevated alpha-fetoprotein levels.

Option D: Omphalocele is another congenital midline abdominal wall defect at the umbilicus causing herniation of the abdominal contents into the base of the umbilical cord. The herniated viscera is covered with peritoneum without any overlying skin. This condition can be associated

with Beckwith-Wiedemann syndrome which is associated with embryonal tumors such as Wilms tumor.

Solution to Question 20:

The given image shows a barium swallow. It is a type of imaging where the patient ingests barium, a contrast agent, to visualize the esophagus and detect abnormalities. The described indentations in the esophageal lumen caused by anterior osteophytes (bony outgrowths) are characteristic findings that can be seen with a barium swallow.

Fluoroscopy (option A) is a technique that provides real-time X-ray images and is often used during procedures like a barium swallow, but the term is more general. X-ray of alkali ingestion (Option B) and X-ray of acid ingestion (option D) might show damage to the esophagus in barium swallow which is not seen in the image.

Solution to Question 21:

The image 3 given above shows yellow plaques and nodules on the mucosa are suggestive of pseudomembranous colitis. It is usually caused by *Clostridium difficile*. The pathogenesis is mediated by the enterotoxins - toxin A and toxin B.

Clostridium difficile is a gram-positive anaerobic bacillus that causes *Clostridioides* (formerly *Clostridium*) *difficile* infection after ingestion of spores of the organism. The more severe form of the condition is known as pseudomembranous colitis (PMC). Enzyme immunoassay to detect glutamate dehydrogenase is less specific compared to toxin detection for diagnosis. So a combination of the two can be used for diagnostic confirmation.

The condition is most commonly associated with prolonged antimicrobial use, which interferes with the commensal colonic microflora. The most commonly implicated antibiotics are cephalosporins, ampicillin, and clindamycin. The spores are spread by contact with the feces of affected individuals or asymptomatic carriers. It can persist on surfaces for months. Risk factors include poor hand hygiene, old age, and gastrointestinal surgery.

The spores germinate in the small intestine and produce two toxins—toxin A (enterotoxin) and toxin B (cytotoxin). These interfere with the epithelial-cell barrier function and result in diarrhea and pseudomembrane formation. The pseudomembranes appear as small whitish-yellow plaques initially which later coalesce to involve the entire colon wall. It contains necrotic leukocytes, fibrin, and cellular debris. The anamnestic IgG response determines which patients develop the disease or remain asymptomatic.

Symptoms include diarrhea, fever, and abdominal pain. The diagnosis is based on a combination of clinical criteria and detection of toxin or the organism in stool samples. This can be done by nucleic acid amplification testing (NAAT) methods such as polymerase chain reaction (PCR) or by culture.

Discontinuation of any ongoing antimicrobials and adequate hydration are general measures that are recommended. Oral fidaxomicin or vancomycin is the first-line treatment. Oral metronidazole can be used when these are not available.

Other options:

Option C: Familial Adenomatosis Polyposis (image 1) is caused due to a germline loss of function mutation of the tumor suppressor gene, APC (adenomatous polyposis coli) on the 5th chromosome leading to multiple adenomas in the GI tract with 100% risk of malignant conversion associated with congenital retinal pigment epithelium hypertrophy. Variants of FAP are Gardner and Turcot syndromes. Its features are described in the pearl attached.

Option B: Esophageal varices (image 4) are tortuous dilated veins within the mucosa and submucosa of the distal esophagus and proximal stomach. Variceal hemorrhage is an emergency that can be treated medically by inducing splanchnic vasoconstriction or endoscopically by sclerotherapy, balloon tamponade, or variceal ligation. It commonly occurs due to portal hypertension in patients with cirrhosis.

Option D: Ulcerative colitis (UC) (image 2) which is characterized by continuous inflammation of the colon, pseudopolyps, biopsy findings of crypt distortion, ulceration, and infiltration of inflammatory cells until the submucosal layer of the intestinal wall. Diagnosis can be made by colonoscopy followed by a biopsy.

UC is a type of inflammatory bowel disease that primarily affects the rectum and extends proximally to involve the colon. Patients typically present with diarrhea, rectal bleeding, tenesmus, crampy abdominal pain, and mucus discharge. They may also present with anorexia, nausea, vomiting, fever, and weight loss.

The presentation of UC differs from that of Crohn's disease, which is characterized by transmural inflammation of the intestine. Differences are mentioned in the pearl.

Solution to Question 22:

The given clinical image is an open inguinal hernia repair of a child with a structure marked as the spermatic cord whose content includes the testicular vein.

Normally the funicular process of the peritoneum becomes obliterated after the testis has reached the scrotum. The scrotal part of the process remains patent and acquires the name 'tunica vaginalis'. In case of congenital hernia, the whole process remains patent. With the increase in abdominal pressure, abdominal contents come out through the patent peritoneal process. So it reaches the bottom of the scrotum very quickly. It may so happen that the funicular process remains patent up to the top of the testis. So the hernia stops at the top of the testis and is known as a congenital funicular hernia.

Indirect hernia comprises more than 80% of inguinal hernias which are common in women and children. It occurs earlier than a direct hernia and is often complete. The two forms often seen under indirect are congenital and acquired. Unlike congenital, acquired hernias do not become complete at once. They progress gradually. Clinically a congenital hernia may present at a later period.

The pearls attached below describe the inguinal canal and the differences between direct and indirect inguinal hernia.

Other options are not the contents of the spermatic cord. Contents are mentioned in the pearl below.

Solution to Question 23:

The given clinical scenario along with the image showing ulcero-proliferative mass is suggestive of lip cancer which is a type of oral carcinoma. Histopathological diagnosis in the form of a deep incisional biopsy of an ulcer along with normal tissue is the gold standard for diagnosis.

Incisional biopsy is taken from the edge of the ulcer along with normal tissue as the centre of the ulcer is usually necrotic. A narrow deep biopsy is better than a shallow broad biopsy. (Option A)

Excisional biopsy is done for smaller benign lesions. It is not done in oral cancers as they are usually bigger in size. An excisional biopsy may increase the spread of cancer or cause functional impairment. (Option C)

Techniques like brush biopsy and exfoliative biopsy are less sensitive. Hence, they are not used for diagnosis. (Option D)

Solution to Question 24:

The given scenario of recurrent swelling on the lip for 6 months and the image showing soft, and transparent cystic swelling points to a diagnosis of a mucous retention cyst. It has to be managed by an excision.

Mucous retention cysts result from duct obstruction by periductal scars or sialolithiasis (salivary gland stone formation) These cysts are mainly found in the ductal system of a minor salivary gland. The cyst cavity may contain fragments of a sialolith or mucous material. Some may regress spontaneously but best management is done by an excision.

Other Options:

Option A: Cervical or plunging ranula- is a rare type of retention cyst arising from the sublingual gland. The mucous passes through the mylohyoid muscle diaphragm to enter the neck. Clinically present as intraoral and cervical swelling. USG or MRI makes the diagnosis. Management includes excision of the sublingual gland + aspiration of saliva out of the sac.



Option C: Oral ranula - It is a benign cystic lesion formed by extravasation of mucus, involving the sublingual gland. Clinically present as cystic swelling in the floor of the mouth. On examination fluctuation test and transillumination test is positive. It is conservatively managed by marsupialisation. Definitive management is the excision of the cyst. Some may resolve spontaneously.



Option D: Mucocoele occurs when an obstruction in the ducts of minor salivary glands, such as those in the labial or buccal areas, leads to swelling and aspiration is not a definitive treatment as it only provides temporary relief and does not prevent recurrence.

Solution to Question 25:

The given clinical scenario along with the intraoperative specimen is suggestive of testicular torsion with gangrene.

Testicular torsion is the twisting of the spermatic cord and its contents such that the testicular blood supply becomes compromised. It is a surgical emergency and if left untreated, the testes may become gangrenous. Early surgery undertaken to untwist the testis gives a better outcome.

The following are anatomical predispositions to testicular torsion:

- High investment in the tunica vaginalis causes the testis to hang within the tunica like a clapper in a bell. This is the most common cause in adolescents and is typically a bilateral abnormality
- Inversion of the testis. The testis is rotated so that it lies transversely or upside down.
- Separation of the epididymis from the body of the testis. This allows torsion of the testis on the pedicle that connects the testis with the epididymis.

Males in the age group of 10-25 years present with acute scrotal or groin pain with associated nausea and vomiting. On examination, the scrotum appears swollen and tender.

Torsion Testis has to be differentiated from Acute epididymo-orchitis. This is done by Prehn's sign. In the case of acute epididymo-orchitis, the pain gets relieved on the elevation of the testis but not in the torsion of the testes.

Other Signs seen in torsion of the testis are:

- Deming's sign: Spasm of the cremaster muscle and the twisting of the cord cause the affected testis to be positioned higher
- Angell's sign: The testis on the opposite side lies horizontally because of the bilaterality of the mesorchium between the testis and epididymis.
- Cremastic reflex- absent on the side of the affected testis

The morphologic changes range from intense congestion to widespread hemorrhage to testicular infarction depending on the duration of torsion. In advanced stages, the testis is markedly enlarged and consists entirely of soft, necrotic, hemorrhagic tissue.

A Doppler ultrasound may be taken to assess the blood supply but surgical exploration is still indicated as a clinical decision. If the testis is infarcted after the cord is untwisted, an orchidectomy must be undertaken and the opposite testis must be fixed to the scrotum to preserve fertility and prevent torsion. If the affected testis is viable after the cord is untwisted, both the affected and the contralateral testes are fixed to the scrotum (bilateral orchiopexy) using non-absorbable sutures.

Other Options :


Option A: Testicular hematoma will usually be associated with trauma to the testis.



Option B: Strangulated hernia will have a history of inguinal swelling and the presence of cough impulse over the swelling. the following image shows ischemic and necrotic bowel herniated into the sack.



Option C: Testicular appendage torsion is torsion of vestigial appendage present along the testis. It has no function but torsion of appendage can be painful. It presents with a blue dot sign on clinical examination. It is a self-limiting condition and conservative management is done.




Solution to Question 26:

The given post-operative image suggests that this patient underwent Total penectomy with inguinal lymph node dissection.

Total penectomy is done in cases of penile carcinoma if the stump is ≤ 2 cm after resection of the tumour and partial penectomy is done if the stump is > 2 cm. Carcinoma of the penis is commonly a squamous cell carcinoma. It arises from the skin of the glans penis or the prepuce. The lesion can be flat or warty in appearance.

The first lymph node to be involved is the superficial inguinal lymph node and thus, superficial inguinal lymph node dissection or sentinel lymph node biopsy (SLNB) helps evaluate the spread of penile cancer.

Tumours of low-grade and low-stage are managed by Mohs' surgery. Tumors involving the glans penis require glansectomy, and in advanced cases require partial penectomy.



Other options:

Option A: Testis is visible in the given image, therefore it is unlikely to be a specimen of Bilateral Orchidectomy with bilateral inguinal node dissection.

Option B: Hernia repair would be done higher up at the inguinal region.

Option C: SLNB will be done through a smaller incision.

Solution to Question 27:

The given clinical features along with elevated urinary levels of vanillylmandelic acid, and her CT scan showing a retroperitoneal tumor in the right lumbar region is suggestive of pheochromocytoma whose familial forms are associated with MEN2A syndrome.


Patients commonly present with a classical triad of palpitation, headache, and profuse sweating.

Pheochromocytoma is associated with:

- MEN-2: caused by germline mutations of the RET proto-oncogene
- Familial paraganglioma syndrome: caused by germline mutations within the succinate dehydrogenase complex subunit B (SDHB), SDHD, and SDHC genes.
- Von Hippel-Lindau syndrome: caused by a germline mutation in the VHL gene.
- Neurofibromatosis type 1: caused by a germline mutation in the NF1 gene.

Histopathology: Cells with salt and pepper chromatin. Other tumors with similar findings include medullary thyroid carcinoma, small cell cancer lung, and extra-adrenal counterpart of pheochromocytoma- paragangliomas. Given below is the histopathological image of pheochromocytoma.

Zellballen Pattern - Pheochromocytoma and Paraganglioma



Other Options:

Option B: Only 10% of the cases are bilateral.

Option C: Only 10% of the cases are malignant. Pheochromocytomas associated with MEN2A are rarely malignant(3-5%), but often malignant with SDHB gene mutation(50%).

Option D: It is a tumor of adulthood.

Solution to Question 28:

The pressure used in vacuum-assisted closure for wound healing is around -125 mmHg. That lies in the range of 110-130 mmHg

Applying negative pressure appears to hasten debridement and the formation of granulation tissue in chronic wounds and ulcers.

It acts by decreasing edema, by removing interstitial fluid, and as a result, helps in decreasing bacterial counts and cell proliferation increases.

Solution to Question 29:

The initial management of patients with all types of aortic dissection is beta blocker medication to control hypertension along with intense blood pressure monitoring and pain management.

Aortic dissection occurs when a defect or flap occurs in the intima of the aorta. Predisposing factors include mainly age, hypertension, Marfan syndrome, and pregnancy. The most typical presentation of aortic dissection is a tearing interscapular pain. The dissection may also result in aortic wall rupture into the pericardium (cardiac tamponade) or mediastinum (left haemothorax).


Contrast-enhanced CT is currently the most preferred imaging modality for aortic dissection.

Double-lumen aorta is seen on CECT. Transesophageal echocardiography is preferred for patients

who are unstable for transportation.

Acute aortic dissection can be classified by the original DeBakey system, based on the location of the proximal tear and distal extent of the dissection flap, or the Stanford system which uses the location of the proximal tear alone.


Site of dissection	DeBakey	Stanford
Both ascending and descending aorta	Type I	Type A
Only the ascending aorta and arch of the aorta	Type II	Type A
Descending aorta distal to the subclavian artery	Type IIIa Limited to the thoracic aorta	Type B
Type IIIb Extends to the abdominal aorta		



Management of aortic dissection:

The initial management of patients with all types of aortic dissection is beta-blocker along with pain management. The preferred drug is esmolol. Hemodynamic stability is an important consideration.

- Stanford Type A (DeBakey Types I and II) requires surgical intervention once the patient's blood pressure is stabilized.
- Stanford Type B (DeBakey type III) is best managed by antihypertensive drugs. For patients with complicated type B dissection, percutaneously placed endovascular stents(option C)are currently preferred. Surgery (options B and D) is reserved for the rare patients not suitable for stenting.



Solution to Question 30:

The given clinical scenario and X-ray KUB showing a homogenous huge density with the smooth edge seen in the bladder is suggestive of struvite calculi, which is formed due to the precipitation of calcium, magnesium, and ammonium phosphate.

Struvite stones (triple phosphate stones) are usually seen in a setting of infection with urease-producing bacteria like *Proteus*, *Pseudomonas*, and *Staphylococcus*. They can grow very large and form a cast of the renal pelvis and calyces resulting in staghorn calculi. Production of urease enzyme by *Proteus*, which hydrolyses urea to ammonia causes alkalinization of urine. Alkalinization of urine leads to precipitation of organic and inorganic compounds which contributes to the formation of Struvite stones.

Almost 90% of the renal stones are radio-opaque and the remaining 10% are radiolucent. The investigation of choice for renal stones is NCCT.

Struvite stones are usually managed with percutaneous nephrolithotomy (PCNL) as it is usually larger in size and associated with Urinary tract infections.

Extracorporeal shock wave lithotripsy (ESWL) is a non-invasive method that generates shockwaves outside the body, which are focused on the stones. It is the most common method of treating urinary tract stones nowadays. Stones up to approximately 1.5 cm in size are suitable for this form of treatment.

PCNL is indicated in:

- Staghorn calculi - Stone $\geq$ 2 cm
- Multiple stones - $\geq$ 1 cm
- Proximal ureteral stone - $\geq$ 1 cm

Other options:

Option A: Patients with cystinuria (an autosomal recessive disorder) are at risk for Cystine stones. These stones are hexagonal.

Option C: Xanthine stones are seen in inborn errors of metabolism like Lesch Nyhan syndrome and cases of hyperuricemia. It is a radiolucent stone.

Option D: Calcium oxalate stones are seen in conditions causing high urine calcium such as primary hyperparathyroidism and chronic acidemia from distal renal tubular acidosis. Also seen in the condition of high urine oxalate which is seen during a low-calcium diet and in malabsorption syndromes (small bowel disease (eg, Crohn's disease), including a jejunioileal bypass for obesity (bariatric surgery), surgical bowel resection or diversion, or cystic fibrosis). Calcium oxalate monohydrate stones are usually dumbbell-shaped and Calcium oxalate dihydrate stones are usually envelope-shaped.

Solution to Question 31:

The Romo Vac drain, depicted in the image, is a closed suction drain commonly used after mastectomy.

Drains are placed at surgical sites to freely collect the air or fluids that may accumulate. This enhances wound healing by preventing the formation of unnecessary cavities. Drains are prophylactic in elective surgeries and therapeutic in emergency surgeries. Drains can be -

- Open - They are based on the principle of gravity
- Semi-open - Their action is based on capillary effect
- Closed - They carry out their actions through suction

Suction drains are commonly used in chest, breast, abdominal, thyroid, and plastic surgeries. They remove intraperitoneal secretions, anastomotic leaks, and blood from hemorrhage at the surgical site. Romo Vac drain is a negative suction drain closed drain kept in the subcutaneous tissue. Romo Vac drain is indicated in modified radical mastectomy, subcutaneous swelling excision, and amputations.

Special types of drains:

- Chest drains (option A) are connected to an underwater seal to aid the exit of air from the pleural cavity. They are used in pneumothorax, pleural effusion, and hemothorax.
- After surgical exploration of the bile duct, the sphincter of Oddi goes into spasm. To facilitate drainage of bile, a T tube may be used.

Drain insertion site care: The site of insertion of the drain is kept clean by drying and patting the drain insertion site with a clean towel after bath. The dressing is changed daily with folded gauze pieces placed below and above the drain site.

Removal of drains: The catheter at the site is milked for residual collections. It is then clamped and the suture holding the catheter is cut. The tube is then gently pulled out and any clots at the perforation end are removed. The time of removal of drains for various conditions is given below.


- Drains used in surgeries like thyroidectomy are usually removed within 24 hours. Their main purpose is to cover postoperative bleeding.

- In a mastectomy, a drain is placed for 5 days to collect serous collections.
- Drains used to cover the colorectal anastomoses are placed for 5-7 days.
- Common bile duct T-tube drains are usually placed for 10 days.
- Drains for infections are removed when the infection subsides or when the content of the drain is minimal.

After the drain is removed, a folded gauze is placed over the drain insertion area. The dressing can be removed after 24 hours. The patient may be given an antibiotic ointment to apply over the area for a week.

Other options:


Option A: The following image shows an underwater seal drain.



Option C: A Penrose drain is a flexible, soft, latex tube used to drain fluid from a surgical site after surgery. It's a type of passive drain.



Option D: Nelaton drain/catheter is commonly used as an intermittent urinary catheter. Like other intermittent catheters, it is also attached temporarily for draining the bladder and then removed.



Solution to Question 32:

The image of a healing ulcer with a sloping edge in the Gaiter's area indicates a venous ulcer.

Venous ulcers due to chronic venous insufficiency are most commonly seen in the lower ■ of the leg and foot. They usually lie just proximal to the medial or lateral malleolus.

Ambulatory venous hypertension is the most acceptable theory in the pathophysiology. They have a sloping edge and the floor contains granulation tissue covered by slough and exudate. Marjolin's ulcer is a type of squamous cell carcinoma seen in chronic long-standing venous ulcers. These are accompanied by lipodermatosclerosis and hemosiderosis.

Management of venous ulcers:

1) Bisgaard's regimen (4Es)

- Education of the patient
- Elevation of the affected limb to reduce the edema
- Elastic compression stockings - The ideal interface pressure in venous ulceration is 35-40 mmHg.
- Evaluation

2) 4-layer bandaging system comprising of:

- Orthopaedic wool
- Cotton crepe
- Elastic bandage - first compressive layer

- Cohesive bandage - second compressive layer
- 3) Antibiotics are given only if the wound is infected.
 - 4) Regular dressing.
 - 5) Medical management by oral pentoxifylline is considered which helps in reducing the viscosity and improving circulation.
 - 6) Surgical management by ulcer excision with pinch mesh grafting has a good outcome and helps in preventing recurrence.

Note: Vacuum-assisted closure (VAC) is a negative pressure dressing. An occlusive dressing is placed over the sore and it is connected to VAC that generates -125mmHg of pressure. It is used in chronic non-healing ulcers, venous ulcers, diabetic ulcers, burn wounds, and bed sores.

Other options:

Option A: The following image depicts an arterial ulcer

Type of Ulcer	Ischaemic/Arterial	Venous	Neuropathic	Tuberculous
Underlying condition	Peripheral vascular disease mainly atherosclerosis, thromboangitis obliterans, and Raynaud's phenomenon	Varicose veins, previous deep vein thrombosis	Diabetes, peripheral neuropathy	Tuberculosis lymph nodes, cold abscess from bone and joint tuberculosis
Location	Toes, feet, legs, or even upper limb digits	Lower limb, commonly on the medial side of the ankle, never above the middle third of the leg	The plantar surface of the foot and toes	Neck, axilla, and groin
Characteristics	Deep, penetrates the deep fascia exposing the tendons, muscles, and underlying bone. The ulcer is painful, tender, and hyperaesthetic.	Vertically oval, and painless. The floor may have pale to no granulation tissue, induration, and tenderness at the base.	Punched-out with regular calloused margins.	Edges are thin, reddish blue, and undermined. Pale granulation tissue, scanty discharge on the floor, and slight induration at the base
Other features	Features of ischemia in adjacent areas, like pallor, dry skin, brittle nails, loss of hair	Inguinal lymph nodes are enlarged usually the vertical group. The ulcer is recurrent.	Dry, cracked, insensate	Regional lymph nodes are enlarged, non-tender, and matted



Option C: A callous ulcer is a type of foot ulcer that develops from a pre-ulcerative callus. It's a chronic, non-healing wound that can last for months or years. Callous ulcers are characterized by a hard base, rigid walls, pale tissue inside the ulcer, and a variety of shapes and sizes.

Option D: The following image is that of a neuropathic ulcer



Solution to Question 33:

The phase of skin graft uptake in which the graft receives its nutritional supply on postoperative day 3 is known as inosculation. It is the period during which the graft is most at risk for failure.

Split skin graft survives the transplantation surgery by the following process:

- The graft survives the first 48 hours because of the imbibition (option B) of plasma from the wound bed.

- After 48 hours, fine anastomotic connections are made, which leads to the inosculation of blood.
- Neovascularization (option D) is the process by which new blood vessels form, directly invading the graft and restoring the pink color of the skin. There is no step of diffusion (option C) in the process of graft transplantation.

Skin grafts can be broadly divided into two categories:

Split-thickness skin graft (Thiersch graft): Partial-thickness skin grafts consisting of the epidermis and a portion of the dermis are called split-thickness skin grafts. Less dermis in the split-thickness skin graft translates into less primary contraction (the degree to which a graft shrinks in surface area after harvesting and before grafting) and more secondary contraction (the degree to which a graft shrinks during healing).

Full-thickness skin grafts (Wolfe graft) include the epidermis and entire dermis, with portions of the sweat glands, sebaceous glands, and hair follicles. More dermis in the full-thickness skin graft translates into more primary contraction and less secondary contraction.

Advantages of split skin graft over full-thickness skin graft:

- Large grafts can be harvested.
- In the case of thinner split-thickness grafts, the donor site heals more rapidly and can be reused in severe cases such
- as burns.
- Uptake is good.
- Meshing can be done, i.e., small splits are made in graft and any exudates collected can come out through this, improving graft take-up. It also allows the graft to expand.


Advantages of full-thickness skin graft over split skin graft:

- It has a lesser degree of post-graft contraction
- Best cosmetic appearance as they include the highest number of skin appendage elements.

Solution to Question 34:

This patient is presenting with thyroid swelling along with image given is suggestive of Solitary Thyroid Nodule. Assessing the biochemical function of the thyroid, i.e. thyroid function test (TFT) which includes serum TSH, T3, and T4 is the first investigation done on any thyroid condition.

Approach to thyroid nodules



Investigations in a patient with Solitary thyroid nodule

- **Thyroid function tests-** Thyroid stimulating hormone (TSH) levels are the first step in evaluating suspected thyroid disease. Based on TSH levels, further investigations is decided.
- **Ultrasonography** is a non-invasive and inexpensive way to stratify the risk of the nodule into benign or malignant tissue.
- **FNAC:** Although other investigations can assist in evaluating a nodule, FNA cytology provides a definitive diagnosis.
- **Radioiodine Thyroid scan:** It helps to differentiate between a hot nodule and a cold nodule. Thyroid nodules that absorb radioiodine are called hot nodules. These nodules produce excess thyroid hormones and are functional nodules. Whereas, the thyroid nodules that do not absorb radioiodine are called cold nodules and they indicate non-functional nodules. It is mostly done by I-123 radioisotope. (Option C)

Other options:

Option A: I-131 radioisotope is more commonly used to radio-ablate the thyroid tissue.

Option B: PET scan has no role in the evaluation of Solitary thyroid nodules.

Solution to Question 35:

The given clinical vignette along with the CT image showing a crescent-shaped hyperdense lesion that crosses the suture lines is indicative of chronic subdural hematoma/hemorrhage.

Subdural hemorrhage (SDH) is a collection of blood between the dura and the arachnoid layers of the meninges. Most commonly, it results from the rupture of cortical bridging veins. Non-contrast CT is the investigation of choice.

Radiographic features of subdural hemorrhages:

- Typically unilateral (85%)

- Crescent-like distribution (concavo-convex) around the periphery of the brain.
- They are not limited by sutures and can fill dural reflection
- Crescent-shaped hypodense lesion on axial CT: Chronic subdural hemorrhage
- Hyperdense collections on CT: Acute subdural hemorrhage.

CT image showing acute subdural hemorrhage



CT image showing acute on chronic subdural hemorrhage.



Other Options:

Option A: Epidural haemorrhage will be usually lens-shaped and is limited by sutures.

Option B: Subarachnoid haemorrhage will show Hyperdensity in the subarachnoid space.

Solution to Question 36:

The given clinical scenario with a long-segment urethral stricture which will be best managed by urethral repair with a graft from buccal mucosa.


The management of a urethral stricture depends on the site, the length, and the cause of the stricture. They include:

- Endoscopic urethrotomy - The stricture is released using a knife passed through a rigid urethroscope. This is done for a short and incomplete stricture. (Option D)
- Urethroplasty - Excision of the stenosed length of the urethra and re-anastomosis is done. End-end anastomoses are done for strictures that are short and complete, while for a long complete stricture, a buccal mucosa graft can be used to bridge the gap. It is considered when a stricture is due to trauma or recurred following endoscopic treatment.
- Urethral dilation - The urethra is stretched using urethral dilators. This is a blind procedure and it can cause injury. Hence, it is not done frequently. (Option A)
- Intermittent self-dilatation - In patients with recurrent strictures who cannot undergo urethroplasty, intermittent self-dilatation with soft hydrophilic catheters is done daily.
- Metallic catheter placement - This is usually done for palliative care in end stage malignant diseases and recurrent cases. (Option C)

Solution to Question 37:

The procedure in the given image shows a 360-degree wrap, which is known as Nissen's fundoplication; this procedure is done for GERD. Nissen's fundoplication is a complete fundoplication procedure.

Fundoplication is the procedure in which the fundus of the stomach is wrapped around the lower end of the esophagus. In the Nissen's fundoplication procedure, the fundus is wrapped the entire 360 degrees around the esophagus.



The most common complication of Nissen's fundoplication is the gas-bloat syndrome. In this condition, patients are unable to belch gas from the stomach into the oesophagus. This causes increased gas in the stomach, resulting in a bloating sensation, abdominal discomfort, and flatulence.

Other options:

Option A: The Toupet procedure is the re-creation of a sphincter around the EGJ done by wrapping the fundus around the oesophagus. A posterior 180-270-degree wrap is used.

Option B: The Dor procedure is fundoplication, most commonly used in the setting of the esophageal myotomy. It consists of an anterior 180-degree wrap.

Option C: Watson's fundoplication is a 90-degree anterior fundoplication. It is reported to achieve good reflux control while permitting burping and active vomiting and reducing gas bloat.

