

# CS315: DATABASE SYSTEMS QUERY OPTIMIZATION

Arnab Bhattacharya

`arnabb@cse.iitk.ac.in`

Computer Science and Engineering,  
Indian Institute of Technology, Kanpur



`http://web.cse.iitk.ac.in/~cs315/`

2<sup>nd</sup> semester, 2022-23

Tue 10:30-11:45, Thu 12:00-13:15


# Evaluation Plan

- Equivalent expressions provide alternate ways of executing a query




# Evaluation Plan

- Equivalent expressions provide alternate ways of executing a query



- An **evaluation plan** also specifies the algorithms



- **Cost-based query optimization**

# Equivalent Expressions

- Two relational algebra expressions are **equivalent** if they generate the same set of output tuples on every *legal* input relation
  - Order of tuples does not matter
  - For SQL, same multiset of output tuples

# Equivalent Expressions

- Two relational algebra expressions are **equivalent** if they generate the same set of output tuples on every *legal* input relation
  - Order of tuples does not matter
  - For SQL, same multiset of output tuples
- **Equivalence rule**: specifies which expressions are equivalent
- Equivalent expressions are systematically generated by repeatedly applying equivalence rules and replacing one form by another

# Equivalent Expressions

- Two relational algebra expressions are **equivalent** if they generate the same set of output tuples on every *legal* input relation
  - Order of tuples does not matter
  - For SQL, same multiset of output tuples
- **Equivalence rule**: specifies which expressions are equivalent
- Equivalent expressions are systematically generated by repeatedly applying equivalence rules and replacing one form by another
- Evaluation plans must account for all algorithms used
  - Merge join may be costlier than hash join, but since it provides a sorted output, a higher level aggregation will be faster

# Equivalence Rules

1  $\sigma_{\theta_1 \wedge \theta_2}(E) = \sigma_{\theta_1}(\sigma_{\theta_2}(E))$

# Equivalence Rules

- 1  $\sigma_{\theta_1 \wedge \theta_2}(E) = \sigma_{\theta_1}(\sigma_{\theta_2}(E))$
- 2  $\sigma_{\theta_1}(\sigma_{\theta_2}(E)) = \sigma_{\theta_2}(\sigma_{\theta_1}(E))$



# Equivalence Rules

- 1  $\sigma_{\theta_1 \wedge \theta_2}(E) = \sigma_{\theta_1}(\sigma_{\theta_2}(E))$
- 2  $\sigma_{\theta_1}(\sigma_{\theta_2}(E)) = \sigma_{\theta_2}(\sigma_{\theta_1}(E))$
- 3  $\Pi_{L_1}(\Pi_{L_2}(\dots \Pi_{L_n}(E))) = \Pi_{L_1}(E)$

# Equivalence Rules

- 1  $\sigma_{\theta_1 \wedge \theta_2}(E) = \sigma_{\theta_1}(\sigma_{\theta_2}(E))$
- 2  $\sigma_{\theta_1}(\sigma_{\theta_2}(E)) = \sigma_{\theta_2}(\sigma_{\theta_1}(E))$
- 3  $\Pi_{L_1}(\Pi_{L_2}(\dots \Pi_{L_n}(E))) = \Pi_{L_1}(E)$
- 4  $\sigma_{\theta}(E_1 \times E_2) = E_1 \bowtie_{\theta} E_2$

# Equivalence Rules

- 1  $\sigma_{\theta_1 \wedge \theta_2}(E) = \sigma_{\theta_1}(\sigma_{\theta_2}(E))$
- 2  $\sigma_{\theta_1}(\sigma_{\theta_2}(E)) = \sigma_{\theta_2}(\sigma_{\theta_1}(E))$
- 3  $\Pi_{L_1}(\Pi_{L_2}(\dots \Pi_{L_n}(E))) = \Pi_{L_1}(E)$
- 4  $\sigma_{\theta}(E_1 \times E_2) = E_1 \bowtie_{\theta} E_2$
- 5  $\sigma_{\theta_1}(E_1 \bowtie_{\theta_2} E_2) = E_1 \bowtie_{\theta_1 \wedge \theta_2} E_2$

# Equivalence Rules

- 1  $\sigma_{\theta_1 \wedge \theta_2}(E) = \sigma_{\theta_1}(\sigma_{\theta_2}(E))$
- 2  $\sigma_{\theta_1}(\sigma_{\theta_2}(E)) = \sigma_{\theta_2}(\sigma_{\theta_1}(E))$
- 3  $\Pi_{L_1}(\Pi_{L_2}(\dots \Pi_{L_n}(E))) = \Pi_{L_1}(E)$
- 4  $\sigma_{\theta}(E_1 \times E_2) = E_1 \bowtie_{\theta} E_2$
- 5  $\sigma_{\theta_1}(E_1 \bowtie_{\theta_2} E_2) = E_1 \bowtie_{\theta_1 \wedge \theta_2} E_2$
- 6  $E_1 \bowtie_{\theta} E_2 = E_2 \bowtie_{\theta} E_1$

# Equivalence Rules

- 1  $\sigma_{\theta_1 \wedge \theta_2}(E) = \sigma_{\theta_1}(\sigma_{\theta_2}(E))$
- 2  $\sigma_{\theta_1}(\sigma_{\theta_2}(E)) = \sigma_{\theta_2}(\sigma_{\theta_1}(E))$
- 3  $\Pi_{L_1}(\Pi_{L_2}(\dots \Pi_{L_n}(E))) = \Pi_{L_1}(E)$
- 4  $\sigma_{\theta}(E_1 \times E_2) = E_1 \bowtie_{\theta} E_2$
- 5  $\sigma_{\theta_1}(E_1 \bowtie_{\theta_2} E_2) = E_1 \bowtie_{\theta_1 \wedge \theta_2} E_2$
- 6  $E_1 \bowtie_{\theta} E_2 = E_2 \bowtie_{\theta} E_1$
- 7  $(E_1 \times E_2) \times E_3 = E_1 \times (E_2 \times E_3)$

# Equivalence Rules

- 1  $\sigma_{\theta_1 \wedge \theta_2}(E) = \sigma_{\theta_1}(\sigma_{\theta_2}(E))$
- 2  $\sigma_{\theta_1}(\sigma_{\theta_2}(E)) = \sigma_{\theta_2}(\sigma_{\theta_1}(E))$
- 3  $\Pi_{L_1}(\Pi_{L_2}(\dots \Pi_{L_n}(E))) = \Pi_{L_1}(E)$
- 4  $\sigma_{\theta}(E_1 \times E_2) = E_1 \bowtie_{\theta} E_2$
- 5  $\sigma_{\theta_1}(E_1 \bowtie_{\theta_2} E_2) = E_1 \bowtie_{\theta_1 \wedge \theta_2} E_2$
- 6  $E_1 \bowtie_{\theta} E_2 = E_2 \bowtie_{\theta} E_1$
- 7  $(E_1 \times E_2) \times E_3 = E_1 \times (E_2 \times E_3)$
- 8  $(E_1 \bowtie E_2) \bowtie E_3 = E_1 \bowtie (E_2 \bowtie E_3)$

# Equivalence Rules

- 1  $\sigma_{\theta_1 \wedge \theta_2}(E) = \sigma_{\theta_1}(\sigma_{\theta_2}(E))$
- 2  $\sigma_{\theta_1}(\sigma_{\theta_2}(E)) = \sigma_{\theta_2}(\sigma_{\theta_1}(E))$
- 3  $\Pi_{L_1}(\Pi_{L_2}(\dots \Pi_{L_n}(E))) = \Pi_{L_1}(E)$
- 4  $\sigma_{\theta}(E_1 \times E_2) = E_1 \bowtie_{\theta} E_2$
- 5  $\sigma_{\theta_1}(E_1 \bowtie_{\theta_2} E_2) = E_1 \bowtie_{\theta_1 \wedge \theta_2} E_2$
- 6  $E_1 \bowtie_{\theta} E_2 = E_2 \bowtie_{\theta} E_1$
- 7  $(E_1 \times E_2) \times E_3 = E_1 \times (E_2 \times E_3)$
- 8  $(E_1 \bowtie E_2) \bowtie E_3 = E_1 \bowtie (E_2 \bowtie E_3)$
- 9  $(E_1 \bowtie_{\theta_1} E_2) \bowtie_{\theta_2 \wedge \theta_3} E_3 = E_1 \bowtie_{\theta_1 \wedge \theta_3} (E_2 \bowtie_{\theta_2} E_3)$ 
  - $\theta_2$  involves attributes from only  $E_2$  and  $E_3$

# Equivalence Rules

- 1  $\sigma_{\theta_1 \wedge \theta_2}(E) = \sigma_{\theta_1}(\sigma_{\theta_2}(E))$
- 2  $\sigma_{\theta_1}(\sigma_{\theta_2}(E)) = \sigma_{\theta_2}(\sigma_{\theta_1}(E))$
- 3  $\Pi_{L_1}(\Pi_{L_2}(\dots \Pi_{L_n}(E))) = \Pi_{L_1}(E)$
- 4  $\sigma_{\theta}(E_1 \times E_2) = E_1 \bowtie_{\theta} E_2$
- 5  $\sigma_{\theta_1}(E_1 \bowtie_{\theta_2} E_2) = E_1 \bowtie_{\theta_1 \wedge \theta_2} E_2$
- 6  $E_1 \bowtie_{\theta} E_2 = E_2 \bowtie_{\theta} E_1$
- 7  $(E_1 \times E_2) \times E_3 = E_1 \times (E_2 \times E_3)$
- 8  $(E_1 \bowtie E_2) \bowtie E_3 = E_1 \bowtie (E_2 \bowtie E_3)$
- 9  $(E_1 \bowtie_{\theta_1} E_2) \bowtie_{\theta_2 \wedge \theta_3} E_3 = E_1 \bowtie_{\theta_1 \wedge \theta_3} (E_2 \bowtie_{\theta_2} E_3)$ 
  - $\theta_2$  involves attributes from only  $E_2$  and  $E_3$
- 10  $\sigma_{\theta_1}(E_1 \bowtie_{\theta_2} E_2) = (\sigma_{\theta_1}(E_1)) \bowtie_{\theta_2} E_2$ 
  - $\theta_1$  involves attributes from only  $E_1$



# Equivalence Rules

- 1  $\sigma_{\theta_1 \wedge \theta_2}(E) = \sigma_{\theta_1}(\sigma_{\theta_2}(E))$
- 2  $\sigma_{\theta_1}(\sigma_{\theta_2}(E)) = \sigma_{\theta_2}(\sigma_{\theta_1}(E))$
- 3  $\Pi_{L_1}(\Pi_{L_2}(\dots \Pi_{L_n}(E))) = \Pi_{L_1}(E)$
- 4  $\sigma_{\theta}(E_1 \times E_2) = E_1 \bowtie_{\theta} E_2$
- 5  $\sigma_{\theta_1}(E_1 \bowtie_{\theta_2} E_2) = E_1 \bowtie_{\theta_1 \wedge \theta_2} E_2$
- 6  $E_1 \bowtie_{\theta} E_2 = E_2 \bowtie_{\theta} E_1$
- 7  $(E_1 \times E_2) \times E_3 = E_1 \times (E_2 \times E_3)$
- 8  $(E_1 \bowtie E_2) \bowtie E_3 = E_1 \bowtie (E_2 \bowtie E_3)$
- 9  $(E_1 \bowtie_{\theta_1} E_2) \bowtie_{\theta_2 \wedge \theta_3} E_3 = E_1 \bowtie_{\theta_1 \wedge \theta_3} (E_2 \bowtie_{\theta_2} E_3)$ 
  - $\theta_2$  involves attributes from only  $E_2$  and  $E_3$
- 10  $\sigma_{\theta_1}(E_1 \bowtie_{\theta_2} E_2) = (\sigma_{\theta_1}(E_1)) \bowtie_{\theta_2} E_2$ 
  - $\theta_1$  involves attributes from only  $E_1$
- 11  $\sigma_{\theta_1 \wedge \theta_2}(E_1 \bowtie_{\theta_3} E_2) = (\sigma_{\theta_1}(E_1)) \bowtie_{\theta_3} (\sigma_{\theta_2}(E_2))$ 
  - $\theta_1$  and  $\theta_2$  involve attributes from only  $E_1$  and  $E_2$  respectively

# Equivalence Rules (contd.)

- 12  $\Pi_{L_1 \cup L_2}(E_1 \bowtie_{\theta} E_2) = \Pi_{L_1}(E_1) \bowtie_{\theta} \Pi_{L_2}(E_2)$
- $L_1$  and  $L_2$  involve attributes from only  $E_1$  and  $E_2$  respectively
  - $\theta$  involves only  $L_1$  and  $L_2$

# Equivalence Rules (contd.)

- 12  $\Pi_{L_1 \cup L_2}(E_1 \bowtie_{\theta} E_2) = \Pi_{L_1}(E_1) \bowtie_{\theta} \Pi_{L_2}(E_2)$
- $L_1$  and  $L_2$  involve attributes from only  $E_1$  and  $E_2$  respectively
  - $\theta$  involves only  $L_1$  and  $L_2$
- 13  $\Pi_{L_1 \cup L_2}(E_1 \bowtie_{\theta} E_2) = \Pi_{L_1 \cup L_2}((\Pi_{L_1 \cup L_3}(E_1)) \bowtie_{\theta} (\Pi_{L_2 \cup L_4}(E_2)))$
- $L_1$  and  $L_2$  involve attributes from only  $E_1$  and  $E_2$  respectively
  - $L_3$  involves attributes from  $E_1$  but is disjoint with  $L_1$
  - $L_4$  involves attributes from  $E_2$  but is disjoint with  $L_2$
  - $\theta$  involves  $L_3$  and  $L_4$

# Equivalence Rules (contd.)

- 12  $\Pi_{L_1 \cup L_2}(E_1 \bowtie_{\theta} E_2) = \Pi_{L_1}(E_1) \bowtie_{\theta} \Pi_{L_2}(E_2)$ 
  - $L_1$  and  $L_2$  involve attributes from only  $E_1$  and  $E_2$  respectively
  - $\theta$  involves only  $L_1$  and  $L_2$
- 13  $\Pi_{L_1 \cup L_2}(E_1 \bowtie_{\theta} E_2) = \Pi_{L_1 \cup L_2}((\Pi_{L_1 \cup L_3}(E_1)) \bowtie_{\theta} (\Pi_{L_2 \cup L_4}(E_2)))$ 
  - $L_1$  and  $L_2$  involve attributes from only  $E_1$  and  $E_2$  respectively
  - $L_3$  involves attributes from  $E_1$  but is disjoint with  $L_1$
  - $L_4$  involves attributes from  $E_2$  but is disjoint with  $L_2$
  - $\theta$  involves  $L_3$  and  $L_4$
- 14  $E_1 \oplus E_2 = E_2 \oplus E_1$ 
  - $\oplus$  is any of  $\cup$  and  $\cap$

# Equivalence Rules (contd.)

- 12  $\Pi_{L_1 \cup L_2}(E_1 \bowtie_{\theta} E_2) = \Pi_{L_1}(E_1) \bowtie_{\theta} \Pi_{L_2}(E_2)$ 
  - $L_1$  and  $L_2$  involve attributes from only  $E_1$  and  $E_2$  respectively
  - $\theta$  involves only  $L_1$  and  $L_2$
- 13  $\Pi_{L_1 \cup L_2}(E_1 \bowtie_{\theta} E_2) = \Pi_{L_1 \cup L_2}((\Pi_{L_1 \cup L_3}(E_1)) \bowtie_{\theta} (\Pi_{L_2 \cup L_4}(E_2)))$ 
  - $L_1$  and  $L_2$  involve attributes from only  $E_1$  and  $E_2$  respectively
  - $L_3$  involves attributes from  $E_1$  but is disjoint with  $L_1$
  - $L_4$  involves attributes from  $E_2$  but is disjoint with  $L_2$
  - $\theta$  involves  $L_3$  and  $L_4$
- 14  $E_1 \oplus E_2 = E_2 \oplus E_1$ 
  - $\oplus$  is any of  $\cup$  and  $\cap$
- 15  $(E_1 \oplus E_2) \oplus E_3 = E_1 \oplus (E_2 \oplus E_3)$ 
  - $\oplus$  is any of  $\cup$  and  $\cap$

# Equivalence Rules (contd.)

- 12  $\Pi_{L_1 \cup L_2}(E_1 \bowtie_{\theta} E_2) = \Pi_{L_1}(E_1) \bowtie_{\theta} \Pi_{L_2}(E_2)$ 
  - $L_1$  and  $L_2$  involve attributes from only  $E_1$  and  $E_2$  respectively
  - $\theta$  involves only  $L_1$  and  $L_2$
- 13  $\Pi_{L_1 \cup L_2}(E_1 \bowtie_{\theta} E_2) = \Pi_{L_1 \cup L_2}((\Pi_{L_1 \cup L_3}(E_1)) \bowtie_{\theta} (\Pi_{L_2 \cup L_4}(E_2)))$ 
  - $L_1$  and  $L_2$  involve attributes from only  $E_1$  and  $E_2$  respectively
  - $L_3$  involves attributes from  $E_1$  but is disjoint with  $L_1$
  - $L_4$  involves attributes from  $E_2$  but is disjoint with  $L_2$
  - $\theta$  involves  $L_3$  and  $L_4$
- 14  $E_1 \oplus E_2 = E_2 \oplus E_1$ 
  - $\oplus$  is any of  $\cup$  and  $\cap$
- 15  $(E_1 \oplus E_2) \oplus E_3 = E_1 \oplus (E_2 \oplus E_3)$ 
  - $\oplus$  is any of  $\cup$  and  $\cap$
- 16  $\sigma_{\theta}(E_1 \oplus E_2) = \sigma_{\theta}(E_1) \oplus \sigma_{\theta}(E_2)$ 
  - $\oplus$  is any of  $\cup$ ,  $\cap$  and  $-$

# Equivalence Rules (contd.)

- 12  $\Pi_{L_1 \cup L_2}(E_1 \bowtie_{\theta} E_2) = \Pi_{L_1}(E_1) \bowtie_{\theta} \Pi_{L_2}(E_2)$ 
  - $L_1$  and  $L_2$  involve attributes from only  $E_1$  and  $E_2$  respectively
  - $\theta$  involves only  $L_1$  and  $L_2$
- 13  $\Pi_{L_1 \cup L_2}(E_1 \bowtie_{\theta} E_2) = \Pi_{L_1 \cup L_2}((\Pi_{L_1 \cup L_3}(E_1)) \bowtie_{\theta} (\Pi_{L_2 \cup L_4}(E_2)))$ 
  - $L_1$  and  $L_2$  involve attributes from only  $E_1$  and  $E_2$  respectively
  - $L_3$  involves attributes from  $E_1$  but is disjoint with  $L_1$
  - $L_4$  involves attributes from  $E_2$  but is disjoint with  $L_2$
  - $\theta$  involves  $L_3$  and  $L_4$
- 14  $E_1 \oplus E_2 = E_2 \oplus E_1$ 
  - $\oplus$  is any of  $\cup$  and  $\cap$
- 15  $(E_1 \oplus E_2) \oplus E_3 = E_1 \oplus (E_2 \oplus E_3)$ 
  - $\oplus$  is any of  $\cup$  and  $\cap$
- 16  $\sigma_{\theta}(E_1 \oplus E_2) = \sigma_{\theta}(E_1) \oplus \sigma_{\theta}(E_2)$ 
  - $\oplus$  is any of  $\cup$ ,  $\cap$  and  $-$
- 17  $\sigma_{\theta}(E_1 \oplus E_2) = \sigma_{\theta}(E_1) \oplus E_2$ 
  - $\oplus$  is any of  $\cap$  and  $-$

# Equivalence Rules (contd.)

- 12  $\Pi_{L_1 \cup L_2}(E_1 \bowtie_{\theta} E_2) = \Pi_{L_1}(E_1) \bowtie_{\theta} \Pi_{L_2}(E_2)$
- $L_1$  and  $L_2$  involve attributes from only  $E_1$  and  $E_2$  respectively
  - $\theta$  involves only  $L_1$  and  $L_2$
- 13  $\Pi_{L_1 \cup L_2}(E_1 \bowtie_{\theta} E_2) = \Pi_{L_1 \cup L_2}((\Pi_{L_1 \cup L_3}(E_1)) \bowtie_{\theta} (\Pi_{L_2 \cup L_4}(E_2)))$
- $L_1$  and  $L_2$  involve attributes from only  $E_1$  and  $E_2$  respectively
  - $L_3$  involves attributes from  $E_1$  but is disjoint with  $L_1$
  - $L_4$  involves attributes from  $E_2$  but is disjoint with  $L_2$
  - $\theta$  involves  $L_3$  and  $L_4$
- 14  $E_1 \oplus E_2 = E_2 \oplus E_1$
- $\oplus$  is any of  $\cup$  and  $\cap$
- 15  $(E_1 \oplus E_2) \oplus E_3 = E_1 \oplus (E_2 \oplus E_3)$
- $\oplus$  is any of  $\cup$  and  $\cap$
- 16  $\sigma_{\theta}(E_1 \oplus E_2) = \sigma_{\theta}(E_1) \oplus \sigma_{\theta}(E_2)$
- $\oplus$  is any of  $\cup$ ,  $\cap$  and  $-$
- 17  $\sigma_{\theta}(E_1 \oplus E_2) = \sigma_{\theta}(E_1) \oplus E_2$
- $\oplus$  is any of  $\cap$  and  $-$
- 18  $\Pi_L(E_1 \cup E_2) = \Pi_L(E_1) \cup \Pi_L(E_2)$



# Example Schema

- branch (bname, bcity, assets)
- customer (cname, cstreet, ccity)
- account (ano, bname, bal)
- loan (lno, bname, amt)
- depositor (cname, ano)
- borrower (cname, lno)

# Example

- Find names of customers having an account at “Kanpur” city

# Example

- Find names of customers having an account at “Kanpur” city
  - $\Pi_{cname}(\sigma_{bcity="Kanpur"}(branch \bowtie (account \bowtie depositor)))$

# Example

- Find names of customers having an account at “Kanpur” city
  - $\Pi_{cname}(\sigma_{bcity="Kanpur"}(branch \bowtie (account \bowtie depositor)))$
  - $\Pi_{cname}(\sigma_{bcity="Kanpur"}(branch) \bowtie (account \bowtie depositor))$

# Example


- Find names of customers having an account at “Kanpur” city
  - $\Pi_{cname}(\sigma_{bcity="Kanpur"}(branch \bowtie (account \bowtie depositor)))$
  - $\Pi_{cname}(\sigma_{bcity="Kanpur"}(branch) \bowtie (account \bowtie depositor))$
- Find names of customers with an account in “Kanpur” with balance more than 1000

# Example

- Find names of customers having an account at “Kanpur” city
  - $\Pi_{cname}(\sigma_{bcity="Kanpur"}(branch \bowtie (account \bowtie depositor)))$
  - $\Pi_{cname}(\sigma_{bcity="Kanpur"}(branch) \bowtie (account \bowtie depositor))$
- Find names of customers with an account in “Kanpur” with balance more than 1000
  - $\Pi_{cname}(\sigma_{bcity="Kanpur" \wedge bal > 1000}(branch \bowtie (account \bowtie depositor)))$





# Example

- Find names of customers having an account at “Kanpur” city
  - $\Pi_{cname}(\sigma_{bcity="Kanpur"}(branch \bowtie (account \bowtie depositor)))$
  - $\Pi_{cname}(\sigma_{bcity="Kanpur"}(branch) \bowtie (account \bowtie depositor))$
- Find names of customers with an account in “Kanpur” with balance more than 1000
  - $\Pi_{cname}(\sigma_{bcity="Kanpur" \wedge bal > 1000}(branch \bowtie (account \bowtie depositor)))$
  - $\Pi_{cname}(\sigma_{bcity="Kanpur"}(branch) \bowtie \sigma_{bal > 1000}(account)) \bowtie depositor$



# Join Trees

- **Left-deep join tree**
  - Right side of each join is a single relation, and not an intermediate result of join of two or more relations
- Similarly, **right-deep join trees** can be defined
- A tree where at least one child of an internal node is a single relation is called a **zig-zag tree**
- A general tree is also called a **bushy tree**





# Number of Join Orders

- Number of join orders is number of ways relations (leaves) can be placed times the number of different tree configurations

# Number of Join Orders

- Number of join orders is number of ways relations (leaves) can be placed times the number of different tree configurations
- Number of ways leaves can be placed
  - If order does not matter, it is  $n!$
  - If order matters, it is 1

# Number of Join Orders

- Number of join orders is number of ways relations (leaves) can be placed times the number of different tree configurations
- Number of ways leaves can be placed
  - If order does not matter, it is  $n!$
  - If order matters, it is 1
- Number of tree configurations
  - Left-deep and right-deep trees:

# Number of Join Orders

- Number of join orders is number of ways relations (leaves) can be placed times the number of different tree configurations
- Number of ways leaves can be placed
  - If order does not matter, it is  $n!$
  - If order matters, it is 1
- Number of tree configurations
  - Left-deep and right-deep trees: 1
  - Zig-zag trees:

# Number of Join Orders

- Number of join orders is number of ways relations (leaves) can be placed times the number of different tree configurations
- Number of ways leaves can be placed
  - If order does not matter, it is  $n!$
  - If order matters, it is 1
- Number of tree configurations
  - Left-deep and right-deep trees: 1
  - Zig-zag trees:  $2^{n-2}$

# Number of Join Orders

- Number of join orders is number of ways relations (leaves) can be placed times the number of different tree configurations
- Number of ways leaves can be placed
  - If order does not matter, it is  $n!$
  - If order matters, it is 1
- Number of tree configurations
  - Left-deep and right-deep trees: 1
  - Zig-zag trees:  $2^{n-2}$ 
    - Place 2 leaves; rest can be either right or left
  - Bushy trees:

# Number of Join Orders

- Number of join orders is number of ways relations (leaves) can be placed times the number of different tree configurations
- Number of ways leaves can be placed
  - If order does not matter, it is  $n!$
  - If order matters, it is 1
- Number of tree configurations
  - Left-deep and right-deep trees: 1
  - Zig-zag trees:  $2^{n-2}$ 
    - Place 2 leaves; rest can be either right or left
  - Bushy trees:  $(n-1)^{\text{th}}$  **Catalan number**  $\frac{(2n-2)!}{n!(n-1)!}$ 
    - $C(1) = 1$ ;  $C(n) = \sum_{i=1}^{n-1} C(i).C(n-i)$



# Algorithm for Join Order

- Consider set  $S$  as the join of  $n$  relations
- $S$  can be represented as  $S' \bowtie (S - S')$  for any non-empty proper subset  $S' \subset S$
- Choose  $S'$  that minimizes the overall cost
- Recursively choose the best plan for  $S'$  and  $(S - S')$



# Algorithm for Join Order

- Consider set  $S$  as the join of  $n$  relations
- $S$  can be represented as  $S' \bowtie (S - S')$  for any non-empty proper subset  $S' \subset S$
- Choose  $S'$  that minimizes the overall cost
- Recursively choose the best plan for  $S'$  and  $(S - S')$
- Employ dynamic programming solution
- Store solutions of subsets

# Algorithm for Join Order

- Consider set  $S$  as the join of  $n$  relations
- $S$  can be represented as  $S' \bowtie (S - S')$  for any non-empty proper subset  $S' \subset S$
- Choose  $S'$  that minimizes the overall cost
- Recursively choose the best plan for  $S'$  and  $(S - S')$
- Employ dynamic programming solution
- Store solutions of subsets
- **Interesting sort order**: Useful order of records
  - Example: Merge join produces tuples in sorted order which makes later merge joins faster
  - Example: Sorted order makes later grouping and aggregation faster
- Algorithm should find the best subset for *each* interesting sort order

# Heuristics in Query Optimization

- Perform selections early
  - Reduces size of relations

# Heuristics in Query Optimization

- Perform selections early
  - Reduces size of relations
- Perform projections early
  - Reduces number of attributes

# Heuristics in Query Optimization

- Perform selections early
  - Reduces size of relations
- Perform projections early
  - Reduces number of attributes
- Perform joins earlier than cross products
  - Produces smaller relations

# Heuristics in Query Optimization

- Perform selections early
  - Reduces size of relations
- Perform projections early
  - Reduces number of attributes
- Perform joins earlier than cross products
  - Produces smaller relations
- Perform only left-deep (or right-deep) joins
  - Number of trees is less
  - Easier to incorporate other operations
  - Hybrid hash joins can be used

# Heuristics in Query Optimization

- Perform selections early
  - Reduces size of relations
- Perform projections early
  - Reduces number of attributes
- Perform joins earlier than cross products
  - Produces smaller relations
- Perform only left-deep (or right-deep) joins
  - Number of trees is less
  - Easier to incorporate other operations
  - Hybrid hash joins can be used
- Perform semantic optimizations
  - Find all employees earning more than their manager
  - May use domain knowledge to return empty result directly

- For each relation  $r$ 
  - Number of tuples  $n_r$
  - Number of blocks  $b_r$
  - **Blocking factor**, i.e., number of tuples that fit in a block  $f_r = \lfloor n_r / b_r \rfloor$
  - Size of a tuple  $l_r$



- For each relation  $r$ 
  - Number of tuples  $n_r$
  - Number of blocks  $b_r$
  - **Blocking factor**, i.e., number of tuples that fit in a block  $f_r = \lfloor n_r / b_r \rfloor$
  - Size of a tuple  $l_r$
- For each attribute  $A$  of a relation  $r$ 
  - Number of distinct values, i.e., size of  $\Pi_A(r)$ :  $v_r(A)$
  - Minimum  $\min_r(A)$  and maximum  $\max_r(A)$
  - Histogram of values: equi-width and equi-depth

- For each relation  $r$ 
  - Number of tuples  $n_r$
  - Number of blocks  $b_r$
  - **Blocking factor**, i.e., number of tuples that fit in a block  $f_r = \lfloor n_r / b_r \rfloor$
  - Size of a tuple  $l_r$
- For each attribute  $A$  of a relation  $r$ 
  - Number of distinct values, i.e., size of  $\Pi_A(r)$ :  $v_r(A)$
  - Minimum  $\min_r(A)$  and maximum  $\max_r(A)$
  - Histogram of values: equi-width and equi-depth
- For each index
  - Number of levels of the index
  - Number of leaves

# Estimating Size of Selections

- $\sigma_{A=x}(r)$

# Estimating Size of Selections

- $\sigma_{A=x}(r)$ 
  - $n_r/v_r(A)$
  - 1 if key

# Estimating Size of Selections

- $\sigma_{A=x}(r)$ 
  - $n_r/v_r(A)$
  - 1 if key
- $\sigma_{A \leq x}(r)$

# Estimating Size of Selections

- $\sigma_{A=x}(r)$ 
  - $n_r/v_r(A)$
  - 1 if key
- $\sigma_{A \leq x}(r)$ 
  - 0 if  $x < \min_r(A)$  and  $n_r$  if  $x > \max_r(A)$
  - $n_r \cdot \frac{x - \min_r(A)}{\max_r(A) - \min_r(A)}$  assuming uniform distribution
  - Estimate is better when histograms are available

# Estimating Size of Selections

- $\sigma_{A=x}(r)$ 
  - $n_r/v_r(A)$
  - 1 if key
- $\sigma_{A \leq x}(r)$ 
  - 0 if  $x < \min_r(A)$  and  $n_r$  if  $x > \max_r(A)$
  - $n_r \cdot \frac{x - \min_r(A)}{\max_r(A) - \min_r(A)}$  assuming uniform distribution
  - Estimate is better when histograms are available
- $\sigma_{A < x}(r)$

# Estimating Size of Selections

- $\sigma_{A=x}(r)$ 
  - $n_r/v_r(A)$
  - 1 if key
- $\sigma_{A \leq x}(r)$ 
  - 0 if  $x < \min_r(A)$  and  $n_r$  if  $x > \max_r(A)$
  - $n_r \cdot \frac{x - \min_r(A)}{\max_r(A) - \min_r(A)}$  assuming uniform distribution
  - Estimate is better when histograms are available
- $\sigma_{A < x}(r)$ 
  - $\text{size}(\sigma_{A \leq x}(r)) - \text{size}(\sigma_{A=x}(r))$



# Estimating Size of Selections

- $\sigma_{A=x}(r)$ 
  - $n_r/v_r(A)$
  - 1 if key
- $\sigma_{A \leq x}(r)$ 
  - 0 if  $x < \min_r(A)$  and  $n_r$  if  $x > \max_r(A)$
  - $n_r \cdot \frac{x - \min_r(A)}{\max_r(A) - \min_r(A)}$  assuming uniform distribution
  - Estimate is better when histograms are available
- $\sigma_{A < x}(r)$ 
  - $\text{size}(\sigma_{A \leq x}(r)) - \text{size}(\sigma_{A=x}(r))$
- $\sigma_{A \geq x}(r)$

# Estimating Size of Selections

- $\sigma_{A=x}(r)$ 
  - $n_r/v_r(A)$
  - 1 if key
- $\sigma_{A \leq x}(r)$ 
  - 0 if  $x < \min_r(A)$  and  $n_r$  if  $x > \max_r(A)$
  - $n_r \cdot \frac{x - \min_r(A)}{\max_r(A) - \min_r(A)}$  assuming uniform distribution
  - Estimate is better when histograms are available
- $\sigma_{A < x}(r)$ 
  - $\text{size}(\sigma_{A \leq x}(r)) - \text{size}(\sigma_{A=x}(r))$
- $\sigma_{A \geq x}(r)$ 
  - $n_r - \text{size}(\sigma_{A < x}(r))$

# Estimating Size of Selections

- $\sigma_{A=x}(r)$ 
  - $n_r/v_r(A)$
  - 1 if key
- $\sigma_{A \leq x}(r)$ 
  - 0 if  $x < \min_r(A)$  and  $n_r$  if  $x > \max_r(A)$
  - $n_r \cdot \frac{x - \min_r(A)}{\max_r(A) - \min_r(A)}$  assuming uniform distribution
  - Estimate is better when histograms are available
- $\sigma_{A < x}(r)$ 
  - $\text{size}(\sigma_{A \leq x}(r)) - \text{size}(\sigma_{A=x}(r))$
- $\sigma_{A \geq x}(r)$ 
  - $n_r - \text{size}(\sigma_{A < x}(r))$
- $\sigma_{A > x}(r)$

# Estimating Size of Selections

- $\sigma_{A=x}(r)$ 
  - $n_r / v_r(A)$
  - 1 if key
- $\sigma_{A \leq x}(r)$ 
  - 0 if  $x < \min_r(A)$  and  $n_r$  if  $x > \max_r(A)$
  - $n_r \cdot \frac{x - \min_r(A)}{\max_r(A) - \min_r(A)}$  assuming uniform distribution
  - Estimate is better when histograms are available
- $\sigma_{A < x}(r)$ 
  - $\text{size}(\sigma_{A \leq x}(r)) - \text{size}(\sigma_{A=x}(r))$
- $\sigma_{A \geq x}(r)$ 
  - $n_r - \text{size}(\sigma_{A < x}(r))$
- $\sigma_{A > x}(r)$ 
  - $n_r - \text{size}(\sigma_{A \leq x}(r))$

# Estimating Size of Selections

- $\sigma_{A=x}(r)$ 
  - $n_r/v_r(A)$
  - 1 if key
- $\sigma_{A \leq x}(r)$ 
  - 0 if  $x < \min_r(A)$  and  $n_r$  if  $x > \max_r(A)$
  - $n_r \cdot \frac{x - \min_r(A)}{\max_r(A) - \min_r(A)}$  assuming uniform distribution
  - Estimate is better when histograms are available
- $\sigma_{A < x}(r)$ 
  - $\text{size}(\sigma_{A \leq x}(r)) - \text{size}(\sigma_{A=x}(r))$
- $\sigma_{A \geq x}(r)$ 
  - $n_r - \text{size}(\sigma_{A < x}(r))$
- $\sigma_{A > x}(r)$ 
  - $n_r - \text{size}(\sigma_{A \leq x}(r))$
- $\sigma_{A \neq x}(r)$

# Estimating Size of Selections

- $\sigma_{A=x}(r)$ 
  - $n_r / v_r(A)$
  - 1 if key
- $\sigma_{A \leq x}(r)$ 
  - 0 if  $x < \min_r(A)$  and  $n_r$  if  $x > \max_r(A)$
  - $n_r \cdot \frac{x - \min_r(A)}{\max_r(A) - \min_r(A)}$  assuming uniform distribution
  - Estimate is better when histograms are available
- $\sigma_{A < x}(r)$ 
  - $\text{size}(\sigma_{A \leq x}(r)) - \text{size}(\sigma_{A=x}(r))$
- $\sigma_{A \geq x}(r)$ 
  - $n_r - \text{size}(\sigma_{A < x}(r))$
- $\sigma_{A > x}(r)$ 
  - $n_r - \text{size}(\sigma_{A \leq x}(r))$
- $\sigma_{A \neq x}(r)$ 
  - $n_r - \text{size}(\sigma_{A=x}(r))$

# Estimating Size of Complex Selections

- **Selectivity** of a condition  $\theta$  is the probability that a tuple in  $r$  satisfies  $\theta$
- If  $s$  number of tuples satisfy, selectivity is  $s/n_r$

# Estimating Size of Complex Selections

- **Selectivity** of a condition  $\theta$  is the probability that a tuple in  $r$  satisfies  $\theta$
- If  $s$  number of tuples satisfy, selectivity is  $s/n_r$
- Conjunction:  $\sigma_{\theta_1 \wedge \dots \wedge \theta_k}(r)$ 
  - If conditions are independent, estimate is



# Estimating Size of Complex Selections

- **Selectivity** of a condition  $\theta$  is the probability that a tuple in  $r$  satisfies  $\theta$
- If  $s$  number of tuples satisfy, selectivity is  $s/n_r$
- Conjunction:  $\sigma_{\theta_1 \wedge \dots \wedge \theta_k}(r)$ 
  - If conditions are independent, estimate is

$$n_r \times \frac{s_1 \times \dots \times s_k}{n_r^k}$$

# Estimating Size of Complex Selections

- **Selectivity** of a condition  $\theta$  is the probability that a tuple in  $r$  satisfies  $\theta$
- If  $s$  number of tuples satisfy, selectivity is  $s/n_r$
- Conjunction:  $\sigma_{\theta_1 \wedge \dots \wedge \theta_k}(r)$ 
  - If conditions are independent, estimate is

$$n_r \times \frac{s_1 \times \dots \times s_k}{n_r^k}$$

- Disjunction:  $\sigma_{\theta_1 \vee \dots \vee \theta_k}(r)$ 
  - If conditions are independent, estimate is

# Estimating Size of Complex Selections

- **Selectivity** of a condition  $\theta$  is the probability that a tuple in  $r$  satisfies  $\theta$
- If  $s$  number of tuples satisfy, selectivity is  $s/n_r$
- Conjunction:  $\sigma_{\theta_1 \wedge \dots \wedge \theta_k}(r)$ 
  - If conditions are independent, estimate is

$$n_r \times \frac{s_1 \times \dots \times s_k}{n_r^k}$$

- Disjunction:  $\sigma_{\theta_1 \vee \dots \vee \theta_k}(r)$ 
  - If conditions are independent, estimate is

$$n_r \times \left( 1 - \frac{(n_r - s_1) \times \dots \times (n_r - s_k)}{n_r^k} \right)$$

# Estimating Size of Complex Selections

- **Selectivity** of a condition  $\theta$  is the probability that a tuple in  $r$  satisfies  $\theta$
- If  $s$  number of tuples satisfy, selectivity is  $s/n_r$
- Conjunction:  $\sigma_{\theta_1 \wedge \dots \wedge \theta_k}(r)$ 
  - If conditions are independent, estimate is

$$n_r \times \frac{s_1 \times \dots \times s_k}{n_r^k}$$

- Disjunction:  $\sigma_{\theta_1 \vee \dots \vee \theta_k}(r)$ 
  - If conditions are independent, estimate is

$$n_r \times \left( 1 - \frac{(n_r - s_1) \times \dots \times (n_r - s_k)}{n_r^k} \right)$$

- Negation:  $\sigma_{\neg \theta}(r)$ 
  - Estimate is

# Estimating Size of Complex Selections

- **Selectivity** of a condition  $\theta$  is the probability that a tuple in  $r$  satisfies  $\theta$
- If  $s$  number of tuples satisfy, selectivity is  $s/n_r$
- Conjunction:  $\sigma_{\theta_1 \wedge \dots \wedge \theta_k}(r)$ 
  - If conditions are independent, estimate is

$$n_r \times \frac{s_1 \times \dots \times s_k}{n_r^k}$$

- Disjunction:  $\sigma_{\theta_1 \vee \dots \vee \theta_k}(r)$ 
  - If conditions are independent, estimate is

$$n_r \times \left( 1 - \frac{(n_r - s_1) \times \dots \times (n_r - s_k)}{n_r^k} \right)$$

- Negation:  $\sigma_{\neg \theta}(r)$ 
  - Estimate is

$$n_r - \text{size}(\sigma_{\theta}(r))$$

# Estimating Size of Joins

- Size of Cartesian product is  $n_r.n_s$  tuples
- If  $R \cap S = \Phi$ , then  $size(r \bowtie_{\theta} s) =$

# Estimating Size of Joins

- Size of Cartesian product is  $n_r.n_s$  tuples
- If  $R \cap S = \Phi$ , then  $size(r \bowtie_{\theta} s) = size(r \times s)$
- If  $R \cap S$  is a key for  $R$ ,

# Estimating Size of Joins

- Size of Cartesian product is  $n_r \cdot n_s$  tuples
- If  $R \cap S = \Phi$ , then  $size(r \bowtie_{\theta} s) = size(r \times s)$
- If  $R \cap S$  is a key for  $R$ , then each tuple of  $s$  will (natural) join with at most one tuple of  $r$ , resulting in at most  $n_s$  joined tuples
- If  $R \cap S$  is a foreign key in  $S$  referencing  $R$ ,



# Estimating Size of Joins

- Size of Cartesian product is  $n_r \cdot n_s$  tuples
- If  $R \cap S = \Phi$ , then  $size(r \bowtie_{\theta} s) = size(r \times s)$
- If  $R \cap S$  is a key for  $R$ , then each tuple of  $s$  will (natural) join with at most one tuple of  $r$ , resulting in at most  $n_s$  joined tuples
- If  $R \cap S$  is a foreign key in  $S$  referencing  $R$ , then it is exactly  $n_s$

# Estimating Size of Joins

- Size of Cartesian product is  $n_r.n_s$  tuples
- If  $R \cap S = \Phi$ , then  $size(r \bowtie_{\theta} s) = size(r \times s)$
- If  $R \cap S$  is a key for  $R$ , then each tuple of  $s$  will (natural) join with at most one tuple of  $r$ , resulting in at most  $n_s$  joined tuples
- If  $R \cap S$  is a foreign key in  $S$  referencing  $R$ , then it is exactly  $n_s$
- Every tuple of  $r$  can join with  $n_s/v_s(A)$  when joining attribute is  $A$ , thereby producing  $n_r.n_s/v_s(A)$  joined tuples
- Reversing  $r$  and  $s$ , estimate becomes  $n_s.n_r/v_r(A)$
- Lower size is the better estimate
- Histograms can improve the estimates

# Estimating Size of Other Operations

- Projection:  $\text{size}(\Pi_A(r)) =$

# Estimating Size of Other Operations

- Projection:  $size(\Pi_A(r)) = v_r(A)$
- Aggregation:  $size({}_A\mathcal{G}_F(r)) =$

# Estimating Size of Other Operations

- Projection:  $size(\Pi_A(r)) = v_r(A)$
- Aggregation:  $size({}_A\mathcal{G}_F(r)) = v_r(A)$
- Set operations
  - Union:  $size(r \cup s) =$

# Estimating Size of Other Operations

- Projection:  $size(\Pi_A(r)) = v_r(A)$
- Aggregation:  $size(\mathcal{A}\mathcal{G}_F(r)) = v_r(A)$
- Set operations
  - Union:  $size(r \cup s) = n_r + n_s$
  - Intersection:  $size(r \cap s) =$

# Estimating Size of Other Operations

- Projection:  $size(\Pi_A(r)) = v_r(A)$
- Aggregation:  $size(\mathcal{A}\mathcal{G}_F(r)) = v_r(A)$
- Set operations
  - Union:  $size(r \cup s) = n_r + n_s$
  - Intersection:  $size(r \cap s) = \min\{n_r, n_s\}$
  - Set difference:  $size(r - s) =$

# Estimating Size of Other Operations

- Projection:  $size(\Pi_A(r)) = v_r(A)$
- Aggregation:  $size(\mathcal{A}\mathcal{G}_F(r)) = v_r(A)$
- Set operations
  - Union:  $size(r \cup s) = n_r + n_s$
  - Intersection:  $size(r \cap s) = \min\{n_r, n_s\}$
  - Set difference:  $size(r - s) = n_r$
- Set operation estimates are *upper bounds*



# Estimating Size of Other Operations

- Projection:  $size(\Pi_A(r)) = v_r(A)$
- Aggregation:  $size({}_A\mathcal{G}_F(r)) = v_r(A)$
- Set operations
  - Union:  $size(r \cup s) = n_r + n_s$
  - Intersection:  $size(r \cap s) = \min\{n_r, n_s\}$
  - Set difference:  $size(r - s) = n_r$
- Set operation estimates are *upper bounds*
- Outer join
  - Left outer join:  $size(r \rhd s) =$

# Estimating Size of Other Operations

- Projection:  $size(\Pi_A(r)) = v_r(A)$
- Aggregation:  $size(\mathcal{A}\mathcal{G}_F(r)) = v_r(A)$
- Set operations
  - Union:  $size(r \cup s) = n_r + n_s$
  - Intersection:  $size(r \cap s) = \min\{n_r, n_s\}$
  - Set difference:  $size(r - s) = n_r$
- Set operation estimates are *upper bounds*
- Outer join
  - Left outer join:  $size(r \bowtie\!\!\!\lhd s) = size(r \bowtie s) + size(r)$
  - Right outer join:  $size(r \bowtie\!\!\!\rhd s) =$

# Estimating Size of Other Operations

- Projection:  $size(\Pi_A(r)) = v_r(A)$
- Aggregation:  $size({}_A\mathcal{G}_F(r)) = v_r(A)$
- Set operations
  - Union:  $size(r \cup s) = n_r + n_s$
  - Intersection:  $size(r \cap s) = \min\{n_r, n_s\}$
  - Set difference:  $size(r - s) = n_r$
- Set operation estimates are *upper bounds*
- Outer join
  - Left outer join:  $size(r \rhd\bowtie s) = size(r \bowtie s) + size(r)$
  - Right outer join:  $size(r \bowtie\lhd s) = size(r \bowtie s) + size(s)$
  - Full outer join:  $size(r \rhd\bowtie\lhd s) =$

# Estimating Size of Other Operations

- Projection:  $size(\Pi_A(r)) = v_r(A)$
- Aggregation:  $size({}_A\mathcal{G}_F(r)) = v_r(A)$
- Set operations
  - Union:  $size(r \cup s) = n_r + n_s$
  - Intersection:  $size(r \cap s) = \min\{n_r, n_s\}$
  - Set difference:  $size(r - s) = n_r$
- Set operation estimates are *upper bounds*
- Outer join
  - Left outer join:  $size(r \bowtie\!\!\!\lrcorner s) = size(r \bowtie s) + size(r)$
  - Right outer join:  $size(r \bowtie\!\!\!\rceil s) = size(r \bowtie s) + size(s)$
  - Full outer join:  $size(r \bowtie\!\!\!\bowtie s) = size(r \bowtie s) + size(r) + size(s)$
- Outer join estimates are *upper bounds*