

Problem Set 3

Problems marked **(T)** are for discussions in Tutorial sessions.

1. Draw and illustrate in \mathbb{R}^2 .

$$(a) \mathbf{e}_1 + \{n\mathbf{e}_2 | n \in \mathbb{N}\}.$$

$$(b) \mathbf{e}_1 + \{\alpha\mathbf{e}_2 | \alpha \in \mathbb{R}\}.$$

2. In \mathbb{R}^2 , Is $\{\alpha\mathbf{e}_1 | \alpha \in \mathbb{R}\} + \{\alpha\mathbf{e}_2 | \alpha \in \mathbb{R}\} = \mathbb{R}^2$? What about $\{\alpha\mathbf{e}_1 | \alpha \in \mathbb{R}\} + \{\alpha \begin{bmatrix} 1 \\ 1 \end{bmatrix} | \alpha \in \mathbb{R}\} = \mathbb{R}^2$?

3. In \mathbb{R}^3 prove that $\left\{ \alpha \begin{bmatrix} 2 \\ 1 \\ 1 \end{bmatrix} | \alpha \in \mathbb{R} \right\} + \left\{ \alpha \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} | \alpha \in \mathbb{R} \right\} + \left\{ \alpha \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} | \alpha \in \mathbb{R} \right\} = \mathbb{R}^3$. Do you use Gauss-Jordan Elimination (GJE) method somewhere?

4. Let L_1 and L_2 be two nonparallel lines passing through origin in \mathbb{R}^3 . What is $L_1 + L_2$?

5. **(T)** Fix a non-negative integer n and let $\mathbb{R}[x; n] = \left\{ \sum_{i=0}^n c_i x^i : c_0, c_1, \dots, c_n \in \mathbb{R} \right\}$. Show that $\mathbb{R}[x; n]$ is a real vector space with respect to the usual addition and scalar multiplication.

6. Recall that $\mathbb{M}_n(\mathbb{R})$ is the real vector space of all $n \times n$ real matrices. Now, prove the following:

(a) $\mathbb{S} = \{A \in \mathbb{M}_n(\mathbb{R}) : A^T = A\}$ is a subspace of $\mathbb{M}_n(\mathbb{R})$.

(b) Fix $A \in \mathbb{M}_n(\mathbb{R})$. Define $\mathbb{U} = \{B \in \mathbb{M}_n(\mathbb{R}) : AB = BA\}$. Then, \mathbb{U} is a subspace of $\mathbb{M}_n(\mathbb{R})$.

(c) Let $\mathbb{W} = \{a_0 I + a_1 A + \dots + a_m A^m : m \text{ is a non-negative integer, } a_i \in \mathbb{R}\}$. Then, \mathbb{W} is a subspace of \mathbb{U} .

7. In \mathbb{R} , define $x \oplus y = x + y - 1$ and $a \odot x = a(x - 1) + 1$. Show that \mathbb{R} is a real vector space with respect to these operations with additive identity 1 (note that 0 is NOT the additive identity).

8. **(T)** Which of the following are subspaces of \mathbb{R}^3 :

(a) $\{(x, y, z) | x \geq 0\}$,

(b) $\{(x, y, z) | x + y = z\}$,

(c) $\{(x, y, z) | x = y^2\}$.

9. Find the condition on $a, b, c, d \in \mathbb{R}$ so that $S = \{(x, y, z) | ax + by + cz = d\}$ is a subspace of \mathbb{R}^3 .

10. **(T)** Show that $S = \{(x_1, x_2, x_3, x_4) : x_4 - x_3 = x_2 - x_1\} = \text{LS}(\{(1, 0, 0, -1), (0, 1, 0, 1), (0, 0, 1, 1)\})$.

11. **(T)** Let W_1 and W_2 be subspaces of a vector space V such that $W_1 \cup W_2$ is also a subspace. Prove that one of the spaces W_i , $i = 1, 2$ is contained in the other.

12. Let $S = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a subset of a real vector space V . Define **linear span** of S as

$$\text{LS}(\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}) = \{c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \dots + c_n \mathbf{v}_n : c_1, c_2, \dots, c_n \in \mathbb{R}\},$$

i.e., the set of all linear combinations of $\mathbf{v}_1, \dots, \mathbf{v}_n$. Then $\text{LS}(\{\mathbf{v}_1, \dots, \mathbf{v}_n\})$ is a subspace of V .

13. Suppose S and T are two subspaces of a vector space V . Define the **sum**

$$S + T = \{\mathbf{s} + \mathbf{t} : \mathbf{s} \in S, \mathbf{t} \in T\}.$$

Show that $S + T$ satisfies the requirements for a vector space. Moreover, $LS(S \cup T) = S + T$.

14. (T) Find all the subspaces of \mathbb{R}^2 .

15. (T) Let $A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$, with $a_{ij} \in \mathbb{C}$. Then the 4 fundamental subspaces are:

- (a) The column space of A :

$$\text{col}(A) = \{A\mathbf{x} : \mathbf{x} \in \mathbb{C}^n\} = \text{LS}(A[:, 1], \dots, A[:, n]) = \text{LS} \left(\left\{ \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix}, \dots, \begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix} \right\} \right)$$

- (b) The column space of A^* :

$$\text{col}(A^*) = \text{LS}(A^*[1, :], \dots, A^*[m, :]) = \{A^*\mathbf{x} : \mathbf{x} \in \mathbb{C}^m\}.$$

- (c) The null space of A :

$$\text{Null Space}(A) = \mathcal{N}(A) = \{\mathbf{x} \in \mathbb{C}^n : A\mathbf{x} = \mathbf{0}\}.$$

- (d) The null space of A^* :

$$\text{Null Space}(A^*) = \mathcal{N}(A^*) = \{\mathbf{x} \in \mathbb{C}^m : A^*\mathbf{x} = \mathbf{0}\}.$$

Important: In case $A \in \mathbb{M}_{m,n}(\mathbb{R})$, the spaces $\text{col}(A^*)$ and $\text{Null Space}(A^*)$ are called the **row-space** of A and the **left-null space** of A , respectively

Now, determine the above 4 mentioned fundamental spaces for the following matrices.

$$(i) A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 2 & 6 & 8 \\ 2 & 8 & 10 \end{bmatrix} \quad (ii) B = \begin{bmatrix} 1 & 2 & 0 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 1 & 2 \\ 2 & 0 & 0 & 1 \end{bmatrix}$$

- (iii) Suppose $B, C \in \mathbb{M}_{m,n}(\mathbb{C})$ and $S = \text{col}(B)$, $T = \text{col}(C)$. Determine $M \in \mathbb{M}_{m,n}(\mathbb{C})$ such that $\text{col}(M) = S + T$.

16. Construct A such that $[1 \ 1 \ 1]^T \in \text{col}(A)$ and $\mathcal{N}(A) = \text{LS}([1 \ 1 \ 1 \ 1]^T)$.

17. (T) Suppose A is an m by n matrix of rank r .

- (a) If $A\mathbf{x} = \mathbf{b}$ has a solution for every right side \mathbf{b} , what is the column space of A ?

- (b) In part (a), what are all the relations between the numbers m , n and r ?
- (c) Give a specific example of a 3 by 2 matrix A of rank 1 with first row $[2 \ 5]$. Describe the column space, $\text{col}(A)$, and the null space, $N(A)$, completely.
- (d) Suppose the right side \mathbf{b} is same as the first column in your example (part c). Find the complete solution to $A\mathbf{x} = \mathbf{b}$.

18. Suppose $R = \text{RREF}(A)$, where $A = \begin{bmatrix} 1 & 2 & 1 & b \\ 2 & a & 1 & 8 \\ \text{(row 3)} \end{bmatrix}$ and $R = \begin{bmatrix} 1 & 2 & 0 & 3 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$.

- (a) What can you say immediately about row 3 of A ?
- (b) What are the numbers a and b ?
- (c) Describe all solutions of $R\mathbf{x} = \mathbf{0}$. Which among row spaces, column spaces and null spaces are the same for A and for R .

19. Let $A \in \mathbb{M}_n(\mathbb{R})$. Show that $\mathcal{N}(A) \subset \mathcal{N}(A^2) \subset \mathcal{N}(A^3) \cdots$. What if $A = \begin{bmatrix} 0 & 1 & -1 & 7 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$?

20. **(T)** Let $A \in \mathbb{M}_{m,n}(\mathbb{R})$. If $\text{RREF}(A) = \begin{pmatrix} I_r & F \\ 0 & 0 \end{pmatrix}$ then describe $\text{col}(A)$ and $\mathcal{N}(A)$.

21. **(T)** Let $W_1 = \text{span} \left\{ [1 \ 1 \ 0]^T, [-1 \ 1 \ 0]^T \right\}$ and $W_2 = \text{span} \left\{ [1 \ 0 \ 2]^T, [-1 \ 0 \ 4]^T \right\}$. Show that $W_1 + W_2 = \mathbb{R}^3$. Give an example of a vector $\mathbf{v} \in \mathbb{R}^3$ such that \mathbf{v} can be written in two different ways in the form $\mathbf{v} = \mathbf{v}_1 + \mathbf{v}_2$, where $\mathbf{v}_1 \in W_1, \mathbf{v}_2 \in W_2$.