Paediatric Equipment and Drug Doses

- For children, the doses of drugs, defibrillation energy and fluid volumes are based on weight (when this is known).
- If the weight is not known, it can be estimated from the child's age using formulae, noting that the formulae are a guide only and tend to underestimate true weight.
- Children may require a different sized SGA, ETT or length at lips than that predicted by the formulae.

Estimated weight (kg)

Under 1 year old	5
1 - 10 years	(Age in years + 4) x 2
11 - 14 years	Age in years x 3

Cuffed endotracheal tube (ETT) size (mm)

Newborn to 1 year	3 - 4
1 year and over	(Age in years ÷ 4) + 3.5

Endotracheal tube length at lips (cm)

Newborn	6 + weight in kg
Under 1 year	ETT size x 3
1 year and over	(Age in years ÷ 2) + 12

Defibrillation energy

Initial and subsequent	4 J/kg

Paediatric drug doses

- The following pages contain tables of paediatric drug doses. They are based on rounding the child's weight up to 5, 10, 20, 30 or 40 kg. For example, a child known or estimated to weigh between 11 and 20 kg receives a dose based on 20 kg, and a child known or estimated to weigh between 31 and 40 kg receives a dose based on 40 kg.
- Children have a higher proportion of body water than adults and require higher doses of most drugs per kg, to achieve the same effect.
- Children weighing greater than 40 kg can usually be administered adult doses, but clinical judgement must be used if the dose may be potentially harmful, for example when administering paracetamol. If

there is any doubt, a paediatric dose should be administered.

- The tables do not incorporate all administration information and should be read in conjunction with the information within the **'medicines'** section of the comprehensive edition.
- The tables indicate the calculated dose of each drug for a given rounded weight, the concentration of drug that should be drawn up, and the volume that should be administered from that solution.
- Drug doses for rapid sequence intubation (RSI) are not contained within these tables, but are contained within the **'RSI'** guideline.
- Some of the doses within the tables differ slightly from the internationally recommended dose. This is to ensure a practical approach, and to help reduce administration errors. Where there are differences between the dose in the tables and the internationally recommended dose, these differences are not clinically significant.
- The usual recommended defibrillation energy for a child is 4 J/kg, but this may not be available for all defibrillators. If the recommended defibrillation energy level is not available, select the next highest level.

Drug dilution and administration

1. Adrenaline 1:10,000 (0.1 mg/ml)

- Draw up 1 ml of adrenaline from a 1 mg/ml ampoule.
- Dilute to a total volume of 10 ml using 0.9% sodium chloride.
- Administer the volume from the tables as a IV bolus.

2. Adrenaline IV 1:1,000,000 (0.001 mg/ml)

- Use a 1 litre bag of 0.9% sodium chloride.
- Add 1 ml of adrenaline from a 1 mg/ml ampoule.
- Shake well and label.
- Administer the volume from the tables as a IV bolus.
- An IV infusion may be administered if aged 5 years or older.

3. Cefazolin IV 100 mg/ml

- Add approximately 4 ml of 0.9% sodium chloride to a 1 g ampoule of cefazolin powder and shake until dissolved.

- Draw up the contents of the ampoule.
- Dilute to a total volume of 10 ml using 0.9% sodium chloride.
- Discard unrequired drug and administer the volume from the tables as a slow IV push over 1-2 minutes, preferably into a running line.

4. Ceftriaxone IV 200 mg/ml

- Add approximately 4ml of 0.9% sodium chloride to a 2 g ampoule of ceftriaxone powder and shake until dissolved.
- Draw up the contents of the ampoule.
- Dilute to a total volume of 10 ml using 0.9% sodium chloride.
- Discard unrequired drug and administer the volume from the tables as a slow IV push over 1-2 minutes, preferably into a running line.

5. Ceftriaxone IM 400 mg/ml

- Add 4 ml of 0.9% sodium chloride to a 2 g ampoule of ceftriaxone powder and shake until dissolved. The total volume will be 5 ml.
- Draw up 2.5 ml in each of two syringes.
- Discard unrequired drug volume and administer one syringe into each lateral thigh. If this site is not suitable use each lateral upper arm.

6. Fentanyl IV 10 mcg/ml (weight rounded up to 30 kg or more)

- Draw up 2 ml of fentanyl from a 100 mcg/2 ml ampoule.
- Dilute to a total volume of 10 ml using 0.9% sodium chloride.
- Administer the volume from the tables as an IV bolus.

7. Fentanyl IV 1 mcg/ml (weight rounded up to 20 kg or less)

- Draw up 2 ml of fentanyl from a 100 mcg/2 ml ampoule.
- Place in a 100 ml bag of 5% glucose.
- Shake well and label.
- Administer the volume from the tables as an IV bolus.

8. Fentanyl and ketamine IV for post intubation sedation

- Draw up 200 mcg of fentanyl (two 100 mcg/2 ml ampoules) and 200 mg of ketamine (one 200 mg/2 ml ampoule), in the same syringe.
- Dilute to a total volume of 20 ml. This solution contains 10 mcg/ml of fentanyl and 10 mg/ml of ketamine.
- Administer the volume from the tables as an IV bolus.

9. Hydrocortisone IV 10 mg/ml

- Draw up 100 mg of hydrocortisone from a 100 mg/2 ml vial.
- Dilute to a total volume of 10 ml using 0.9% sodium chloride.
- Administer the volume from the tables IV over 1-2 minutes.

10. Ketamine IV for analgesia 1 mg/ml

- Draw up 1 ml of ketamine from a 200 mg/2 ml ampoule.
- Place into a 100 ml bag of 5% glucose. This solution contains 1 mg/ml.
- Draw up the volume from the tables and dilute this further to a total volume of 20 ml using 0.9% sodium chloride.
- Administer IV over approximately 15 minutes.

11. Ketamine IV for dissociation 10 mg/ml

- Draw up 1 ml of ketamine from a 200 mg/2 ml ampoule.
- Dilute to a total volume of 10 ml using 0.9% sodium chloride.
- Administer the volume from the tables as an IV bolus.

12. Levetiracetam

- Draw up the volume described in the tables from ampoule(s) containing 500 mg/5 ml.
- Administer IV over 5 minutes, preferably into a running line.

13. Magnesium IV 1 mmol/ml

- Draw up 5 ml of magnesium from a 10 mmol/5 ml ampoule.
- Dilute to a total volume of 10 ml using 0.9% sodium chloride.

- Discard unrequired drug and administer the volume from the tables IV over approximately 15 minutes, preferably into a running line.

14. Metaraminol IV 0.1 mg/ml

- Draw up 1 ml of metaraminol from a 10 mg/1 ml ampoule.
- Place into a 100 ml bag of 5% glucose. This solution contains 0.1 mg/ml.
- Shake well and label.
- Administer the volume from the tables as an IV bolus.

15. Midazolam IV 1 mg/ml

- Draw up 1 ml of midazolam from a 15 mg/3 ml ampoule.
- Dilute to a total volume of 5 ml using 0.9% sodium chloride.
- Discard unrequired drug and administer the volume from the tables as an IV bolus.

16. Morphine IV 1 mg/ml

- Draw up 1 ml of morphine from a 10 mg/1 ml ampoule.
- Dilute to a total volume of 10 ml using 0.9% sodium chloride.
- Administer the volume from the tables as an IV bolus.

17. Naloxone IV 0.1 mg/ml

- Draw up 1 ml of naloxone from a 0.4 mg/1 ml ampoule.
- Dilute to a total volume of 4 ml using 0.9% sodium chloride.
- Administer the volume from the tables as an IV bolus.

5 kg / less than one year

Cardiac arrest	Dose	Volume	Dilution number
Adrenaline IV	0.05 mg	0.5 ml (1:10,000)	1
Amiodarone IV	25 mg	0.5 ml (undiluted)	-
Manual defibrillation	20-25	Joules	-

LMA	Size 1 (< 5 kg)	Cuff inflation 4 ml	-
ETT (cuffed)	Size 3	9 cm length at lips	-
Other drugs			**Dilution number**
Adrenaline IV (not cardiac arrest)	0.001 mg	1 ml (1:1,000,000)	2
Adrenaline IM	0.05 mg	0.5 ml (1:10,000)	1
Adrenaline IN (epistaxis)	-	-	-
Cefazolin IV	150 mg	1.5 ml (100 mg/ml)	3
Ceftriaxone IV	300 mg	1.5 ml (200 mg/ml)	4
Ceftriaxone IM	300 mg	0.75 ml (400 mg/ml)	5
Fentanyl IV (analgesia)	1-5 mcg	1-5 ml (1 mcg/ml)	7
Fentanyl IM	5 mcg	0.1 ml (undiluted)	-
Fentanyl IN	5 mcg	0.1 ml (undiluted)	-
Fentanyl/ketamine IV (post intubation)	5 mcg/5 mg	0.5 ml	8
Glucagon IM	0.5 mg	0.5 ml (undiluted)	-
10% glucose IV	2 ml/kg	10 ml	-
Hydrocortisone IV	10 mg	1 ml (10 mg/ml)	9
Hydrocortisone IM	10 mg	0.2 ml (undiluted)	-
Ibuprofen liquid PO	50 mg	2.5 ml (20 mg/ml)	-
Ibuprofen tablet PO	-	-	-
Ketamine IV (analgesia)	-	-	-
Ketamine IV (dissociation)	-	-	-
Ketamine IM/PO (analgesia)	-	-	-
Ketamine IM (dissociation)	-	-	-
Levetiracetam IV	250 mg	2.5 ml (100 mg/ml)	12
1% lignocaine IO	5 mg	0.5 ml (undiluted)	-

1% lignocaine SC	15 mg (max)	1.5 ml (max)	-
Loratadine PO	-	-	-
Magnesium IV	-	-	-
Metaraminol IV	0.05-0.1 mg	0.5-1 ml (0.1 mg/ml)	14
Midazolam IV (seizures)	0.5 mg	0.5 ml (1 mg/ml)	15
Midazolam IM (seizures)	1 mg	0.2 ml (undiluted)	-
Morphine IV	0.25 mg	0.25 ml (1 mg/ml)	16
Morphine IM/SC	1 mg	0.1 ml (undiluted)	-
Naloxone IV	0.05 mg	0.5 ml (0.1 mg/ml)	17
Naloxone IM	0.1 mg	0.25 ml (undiluted)	-
Ondansetron IV	-	-	-
Ondansetron IM	-	-	-
Paracetamol liquid PO	75 mg	1.5 ml (50 mg/ml)	-
Paracetamol tablet PO	-	-	-
Prednisolone PO	5 mg	1 ml (5 mg/ml)	-
Prednisone PO	-	-	-
Rocuronium IV	5 mg	0.5 ml (undiluted)	-
0.75% ropivacaine SC	11.25 mg (max)	1.5 ml (max)	-
0.9% sodium chloride IV	20 ml/kg	100 ml	-
8.4% sodium bicarbonate	10 mmol	10 ml (1 mmol/ml)	-
Tranexamic acid IN (epistaxis)	-	-	-
Tranexamic acid IV (non-traumatic bleeding)	100 mg	1 ml (undiluted)	-
Tranexamic acid IV (bleeding from trauma)	200 mg	2 ml (undiluted)	-

10 kg / 1 year

Cardiac arrest	Dose	Volume	Dilution number
Adrenaline IV	0.1 mg	1 ml (1:10,000)	1
Amiodarone IV	50 mg	1 ml (undiluted)	-
Manual defibrillation	40-50	Joules	-
LMA	Size 2 (10-20 kg)	Cuff inflation 10 ml	-
ETT (cuffed)	Size 3 or 4	12 cm length at lips	-
Other drugs			**Dilution number**
Adrenaline IV (not cardiac arrest)	0.002 mg	2 ml (1:1,000,000)	2
Adrenaline IM	0.1 mg	0.1 ml (undiluted)	-
Adrenaline IN (epistaxis)	-	-	-
Cefazolin IV	250 mg	2.5 ml (100 mg/ml)	3
Ceftriaxone IV	500 mg	2.5 ml (200 mg/ml)	4
Ceftriaxone IM	500 mg	1.25 ml (400 mg/ml)	5
Fentanyl IV (analgesia)	2-10 mcg	2-10 ml (1 mcg/ml)	7
Fentanyl IM	10 mcg	0.2 ml (undiluted)	-
Fentanyl IN	15 mcg	0.3 ml (undiluted)	-
Fentanyl/ketamine IV (post intubation)	10 mcg/10 mg	1 ml	8
Glucagon IM	0.5 mg	0.5 ml (undiluted)	-
10% glucose IV	2 ml/kg	20 ml	-
Hydrocortisone IV	25 mg	2.5 ml (10 mg/ml)	9
Hydrocortisone IM	25 mg	0.5 ml (undiluted)	-
Ibuprofen liquid PO	100 mg	5 ml (20 mg/ml)	-
Ibuprofen tablet PO	100 mg	½ tablet	-
Ketamine IV (analgesia)	2 mg	2 ml (1 mg/ml)	10

Ketamine IV (dissociation)	10 mg	1 ml (10 mg/ml)	11
Ketamine IM/PO (analgesia)	5 mg	0.05 ml (undiluted)	-
Ketamine IM (dissociation)	50 mg	0.5 ml (undiluted)	-
Levetiracetam IV	500 mg	5 ml (100 mg/ml)	12
1% lignocaine IO	10 mg	1 ml (undiluted)	-
1% lignocaine SC	30 mg (max)	3 ml (max)	-
Loratadine PO	10 mg	1 tablet	-
Magnesium IV	-	-	-
Metaraminol IV	0.1-0.2 mg	1-2 ml (0.1 mg/ml)	14
Midazolam IV (seizures)	1 mg	1 ml (1 mg/ml)	15
Midazolam IM (seizures)	2 mg	0.4 ml (undiluted)	-
Morphine IV	0.5 mg	0.5 ml (1 mg/ml)	16
Morphine IM/SC	2 mg	0.2 ml (undiluted)	-
Naloxone IV	0.1 mg	1 ml (0.1 mg/ml)	17
Naloxone IM	0.2 mg	0.5 ml (undiluted)	-
Ondansetron IV	2 mg	1 ml (undiluted)	-
Ondansetron IM	1 mg	0.5 ml (undiluted)	-
Paracetamol liquid PO	150 mg	3 ml (50 mg/ml)	-
Paracetamol tablet PO	-	-	-
Prednisolone PO	10 mg	2 ml (5 mg/ml)	-
Prednisone PO	10 mg	½ tablet	-
Rocuronium IV	10 mg	1 ml (undiluted)	-
0.75% ropivacaine SC	22.5 mg (max)	3 ml (max)	-
0.9% sodium chloride IV	20 ml/kg	200 ml	-
8.4% sodium bicarbonate	20 mmol	20 ml (1 mmol/ml)	-

Tranexamic acid IN (epistaxis)	-	-	-
Tranexamic acid IV (non-traumatic bleeding)	200 mg	2 ml (undiluted)	-
Tranexamic acid IV (bleeding from trauma)	400 mg	4 ml (undiluted)	-

20 kg / 2-5 years

Cardiac arrest	Dose	Volume	Dilution number
Adrenaline IV	0.2 mg	2 ml (1:10,000)	1
Amiodarone IV	100 mg	2 ml (undiluted)	-
Manual defibrillation	80-100	Joules	-
LMA	Size 2 (10-20 kg)	Cuff inflation 10 ml	-
ETT (cuffed)	Size 4 or 5	15 cm length at lips	-
Other drugs			**Dilution number**
Adrenaline IV (not cardiac arrest)	0.004 mg	4 ml (1:1,000,000)	2
Adrenaline IM	0.2 mg	0.2 ml (undiluted)	-
Adrenaline IN (epistaxis)	0.1 mg	1 ml (1:10,000)	1
Cefazolin IV	500 mg	5 ml (100 mg/ml)	3
Ceftriaxone IV	1 g	5 ml (200 mg/ml)	4
Ceftriaxone IM	1 g	2.5 ml (400 mg/ml)	5
Fentanyl IV (analgesia)	5-20 mcg	5-20 ml (1 mcg/ml)	7
Fentanyl IM	20 mcg	0.4 ml (undiluted)	-
Fentanyl IN	30 mcg	0.6 ml (undiluted)	-
Fentanyl/ketamine IV (post intubation)	20 mcg/20 mg	2 ml	8
Glucagon IM	0.5-1 mg	0.5-1 ml (undiluted)	-
10% glucose IV	2 ml/kg	40 ml	-
Hydrocortisone IV	50 mg	5 ml (10 mg/ml)	9

Hydrocortisone IM	50 mg	1 ml (undiluted)	-
Ibuprofen liquid PO	150 mg	7.5 ml (20 mg/ml)	-
Ibuprofen tablet PO	100 mg	½ tablet	-
Ketamine IV (analgesia)	4 mg	4 ml (1 mg/ml)	10
Ketamine IV (dissociation)	20 mg	2 ml (10 mg/ml)	11
Ketamine IM/PO (analgesia)	10 mg	0.1 ml (undiluted)	-
Ketamine IM (dissociation)	100 mg	1 ml (undiluted)	-
Levetiracetam IV	1000 mg	10 ml (100 mg/ml)	12
1% lignocaine IO	20 mg	2 ml (undiluted)	-
1% lignocaine SC	60 mg (max)	6 ml (max)	-
Loratadine PO	10 mg	1 tablet	-
Magnesium IV	4 mmol	4 ml (1 mmol/ml)	13
Metaraminol IV	0.2-0.4 mg	2-4 ml (0.1 mg/ml)	14
Midazolam IV (seizures)	2 mg	2 ml (1 mg/ml)	15
Midazolam IM (seizures)	4 mg	0.8 ml (undiluted)	-
Morphine IV	1 mg	1 ml (1 mg/ml)	16
Morphine IM/SC	4 mg	0.4 ml (undiluted)	-
Naloxone IV	0.1-0.2 mg	1-2 ml (0.1 mg/ml)	17
Naloxone IM	0.4 mg	1 ml (undiluted)	-
Ondansetron IV	4 mg	2 ml (undiluted)	-
Ondansetron IM	2 mg	1 ml (undiluted)	-
Paracetamol liquid PO	250 mg	5 ml (50 mg/ml)	-
Paracetamol tablet PO	250 mg	½ tablet	-
Prednisolone PO	20 mg	4 ml (5 mg/ml)	-
Prednisone PO	20 mg	1 tablet	-

Rocuronium IV	20 mg	2 ml (undiluted)	-
0.75% ropivacaine SC	45 mg (max)	6 ml (max)	-
0.9% sodium chloride IV	20 ml/kg	400 ml	-
8.4% sodium bicarbonate	40 mmol	40 ml (1 mmol/ml)	
Tranexamic acid IN (epistaxis)	100 mg	1 ml (undiluted)	-
Tranexamic acid IV (non-traumatic bleeding)	400 mg	4 ml (undiluted)	-
Tranexamic acid IV (bleeding from trauma)	800 mg	8 ml (undiluted)	-

30 kg / 6-10 years

Cardiac arrest	Dose	Volume	Dilution number
Adrenaline IV	0.3 mg	3 ml (1:10,000)	1
Amiodarone IV	150 mg	3 ml (undiluted)	-
Manual defibrillation	120-150	Joules	-
LMA	Size 3 (30-50 kg)	Cuff inflation 20 ml	-
ETT (cuffed)	Size 5 or 6	17 cm length at lips	-
Other drugs			**Dilution number**
Adrenaline IV (not cardiac arrest)	0.006 mg	6 ml (1:1,000,000)	2
Adrenaline IM	0.3 mg	0.3 ml (undiluted)	-
Adrenaline IN (epistaxis)	0.1 mg	1 ml (1:10,000)	1
Cefazolin IV	750 mg	7.5 ml (100 mg/ml)	3
Ceftriaxone IV	1.5 g	7.5 ml (200 mg/ml)	4
Ceftriaxone IM	1.5 g	3.75 ml (400 mg/ml)	5
Fentanyl IV (analgesia)	10-30 mcg	1-3 ml (10 mcg/ml)	6
Fentanyl IM	30 mcg	0.6 ml (undiluted)	-
Fentanyl IN	45 mcg	0.9 ml (undiluted)	-

Fentanyl/ketamine IV (post intubation)	30 mcg/30 mg	3 ml	8
Glucagon IM	1 mg	1 ml (undiluted)	-
10% glucose IV	2 ml/kg	60 ml	-
Hydrocortisone IV	75 mg	7.5 ml (10 mg/ml)	9
Hydrocortisone IM	75 mg	1.5 ml (undiluted)	-
Ibuprofen liquid PO	200 mg	10 ml (20 mg/ml)	-
Ibuprofen tablet PO	200 mg	1 tablet	-
Ketamine IV (analgesia)	6 mg	6 ml (1 mg/ml)	10
Ketamine IV (dissociation)	30 mg	3 ml (10 mg/ml)	11
Ketamine IM/PO (analgesia)	15 mg	0.15 ml (undiluted)	-
Ketamine IM (dissociation)	150 mg	1.5 ml (undiluted)	-
Levetiracetam IV	1500 mg	15 ml (100 mg/ml)	12
1% lignocaine IO	30 mg	3 ml (undiluted)	-
1% lignocaine SC	90 mg (max)	9 ml (max)	-
Loratadine PO	10 mg	1 tablet	-
Magnesium IV	6 mmol	6 ml (1 mmol/ml)	13
Metaraminol IV	0.3-0.6 mg	3-6 ml (0.1 mg/ml)	14
Midazolam IV (seizures)	3 mg	3 ml (1 mg/ml)	15
Midazolam IM (seizures)	6 mg	1.2 ml (undiluted)	-
Morphine IV	1.5 mg	1.5 ml (1 mg/ml)	16
Morphine IM/SC	6 mg	0.6 ml (undiluted)	-
Naloxone IV	0.1- 0.3 mg	1- 3 ml (0.1 mg/ml)	17
Naloxone IM	0.6 mg	1.5 ml (undiluted)	-
Ondansetron IV	6 mg	3 ml (undiluted)	-
Ondansetron IM	3 mg	1.5 ml (undiluted)	-

Paracetamol liquid PO	375 mg	7.5 ml (50 mg/ml)	-
Paracetamol tablet PO	250 mg	½ tablet	-
Prednisolone PO	30 mg	6 ml (5 mg/ml)	-
Prednisone PO	30 mg	1 ½ tablets	-
Rocuronium IV	30 mg	3 ml (undiluted)	-
0.75% ropivacaine SC	67.5 mg (max)	9 ml (max)	-
0.9% sodium chloride IV	20 ml/kg	600 ml	-
8.4% sodium bicarbonate	60 mmol	60 ml (1 mmol/ml)	
Tranexamic acid IN (epistaxis)	100 mg	1 ml (undiluted)	-
Tranexamic acid IV (non-traumatic bleeding)	600 mg	6 ml (undiluted)	-
Tranexamic acid IV (bleeding from trauma)	1200 mg	12 ml (undiluted)	-

40 kg / 11-13 years

Cardiac arrest	Dose	Volume	Dilution number
Adrenaline IV	0.4 mg	4 ml (1:10,000)	1
Amiodarone IV	200 mg	4 ml (undiluted)	-
Manual defibrillation	160-200	Joules	-
LMA	Size 3 (30-50 kg)	Cuff inflation 20 ml	-
ETT (cuffed)	Size 6 or 7	19 cm length at lips	-
Other drugs			**Dilution number**
Adrenaline IV (not cardiac arrest)	0.008 mg	8 ml (1:1,000,000)	2
Adrenaline IM	0.4 mg	0.4 ml (undiluted)	-
Adrenaline IN (epistaxis)	0.2 mg	2 ml (1:10,000)	1
Cefazolin IV	1000 mg	10 ml (100 mg/ ml)	3
Ceftriaxone IV	2 g	10 ml (200 mg/ml)	4

Ceftriaxone IM	2 g	5 ml (400 mg/ml)	5
Fentanyl IV (analgesia)	10-40 mcg	1-4 ml (10 mcg/ml)	6
Fentanyl IM	40 mcg	0.8 ml (undiluted)	-
Fentanyl IN	60 mcg	1.2 ml (undiluted)	-
Fentanyl/ketamine IV (post intubation)	40 mcg/40 mg	4 ml	8
Glucagon IM	1 mg	1 ml (undiluted)	-
10% glucose IV	2 ml/kg	80 ml	-
Hydrocortisone IV	100 mg	10 ml (10 mg/ml)	9
Hydrocortisone IM	100 mg	2 ml (undiluted)	-
Ibuprofen liquid PO	300 mg	15 ml (20 mg/ml)	-
Ibuprofen tablet PO	300 mg	1 ½ tablets	-
Ketamine IV (analgesia)	8 mg	8 ml (1 mg/ml)	10
Ketamine IV (dissociation)	40 mg	4 ml (10 mg/ml)	11
Ketamine IM/PO (analgesia)	20 mg	0.2 ml (undiluted)	-
Ketamine IM (dissociation)	200 mg	2 ml (undiluted)	-
Levetiracetam IV	2000 mg	20 ml (100 mg/ml)	12
1% lignocaine IO	40 mg	4 ml (undiluted)	-
1% lignocaine SC	120 mg (max)	12 ml (max)	-
Loratadine PO	20 mg	2 tablets	-
Magnesium IV	8 mmol	8 ml (1 mmol/ml)	13
Metaraminol IV	0.4-0.8 mg	4-8 ml (0.1 mg/ml)	14
Midazolam IV (seizures)	4 mg	4 ml (1 mg/ml)	15
Midazolam IM (seizures)	8 mg	1.6 ml (undiluted)	-
Morphine IV	2 mg	2 ml (1 mg/ml)	16
Morphine IM/SC	8 mg	0.8 ml (undiluted)	-

Naloxone IV	0.1-0.4 mg	1- 4 ml (0.1 mg/ml)	17
Naloxone IM	0.8 mg	2 ml (undiluted)	-
Ondansetron IV	8 mg	4 ml (undiluted)	-
Ondansetron IM	4 mg	2 ml (undiluted)	-
Paracetamol liquid PO	500 mg	10 ml (50 mg/ml)	-
Paracetamol tablet PO	500 mg	1 tablet	-
Prednisolone PO	40 mg	8 ml (5 mg/ml)	-
Prednisone PO	40 mg	2 tablets	-
Rocuronium IV	40 mg	4 ml (undiluted)	-
0.75% ropivacaine SC	90 mg (max)	12 ml (max)	-
0.9% sodium chloride IV	20 ml/kg	800 ml	-
8.4% sodium bicarbonate	80 mmol	80 ml (1 mmol/ml)	
Tranexamic acid IN (epistaxis)	200 mg	2 ml (undiluted)	-
Tranexamic acid IV (non-traumatic bleeding)	800 mg	8 ml (undiluted)	-
Tranexamic acid IV (bleeding from trauma)	1600 mg	16 ml (undiluted)	-

Paediatric vital signs

Age	Heart rate	Respiratory rate	Blood pressure (systolic)
<1	110-160	30-40	70-90
1-2 years	100-150	25-35	80-95
2-5 years	95-140	25-30	80-100
5-12 years	80-120	20-25	90-110
> 12 years	60-100	15-20	100-120

Infant Glasgow Coma Scale

Best eye opening	
Spontaneously	4
To voice or touch	3
To pain or pressure	2
None	1
Best verbal response	
Smiles, babbles, coos	5
Cries normally	4
Cries only to pain or pressure	3
Moans or grunts	2
None	1
Best motor response	
Normal spontaneous movement	6
Localises or has purposeful movement	5
Withdraws from pain or normal flexion	4
Abnormal (spastic) flexion	3
Extension or rigid	2
None	1

More Information

Version history

28 October 2024 1.0.5

Update: Paediatric vital signs updates. Changes to loratadine, TXA, cefazolin and 8.4% sodium bicarbonate.

11 September 2023 1.0.4

Update: Tables have been updated to include cefazolin and levetiracetam and removal of valproate.

22 November 2022 1.0.1

Update: Defibrillation energy, metaraminol dilution. Added morphine, removed blood volume from vital signs.