Ticagrelor

Mechanism of action

- Ticagrelor has antiplatelet activity. It produces reversible inhibition of the ADP receptor on platelets, decreasing platelet aggregation.

Authorised practice levels

- Paramedics, ICPs and CCPs.

Indications

- STEMI, in conjunction with the primary PCI pathway, when requested by the accepting clinician.

Contraindications

- Known severe allergy.

Cautions

- Clinically significant bleeding. Ticagrelor will increase bleeding.
- At risk of bleeding.
- Pregnancy.

Use in pregnancy or when breastfeeding

- Safety has not been demonstrated during pregnancy. The likelihood of STEMI occurring in a woman who is pregnant is so low that personnel must seek clinical advice prior to administration.
- May be administered if the patient is breastfeeding. Advise the patient to stop breastfeeding and seek further advice from their lead maternity carer or GP.

Dosage

- 180 mg for an adult.

Ticagrelor

Administration

- Administer PO.

Common adverse effects

- Increased bleeding.

Usual onset of effect

- 30-60 minutes.

Usual duration of effect

- 24 hours.

Usual preparation

- 90 mg tablet.

Pharmacokinetics

- Ticagrelor is metabolised in the liver and excreted predominantly in the bile.
- There are no significant effects from liver impairment on acute administration.

Common interactions

- The risk of bleeding will be increased if the patient is taking an anticoagulant.

More Information

Version history

11 September 2023 1.0.4
New: Guideline released.