GLUCO-BETA-EXT: 52-Week Phase III Extension of Glucoren 15 mg

SYNTHETIC DEMONSTRATION DATA. This document is fictional and was generated for testing a document retrieval system. It contains no real patient information and does not report genuine research findings. Do not use it for clinical decisions.

Document ID	MDA-CT-011
Sponsor	Northbridge Clinical Research Unit (fictional)
Document type	Clinical trial summary
Study identifier	GLUCO-BETA-EXT (NB-T2D-3205)
Investigational product	Glucoren (fictional SGLT2 inhibitor), 15 mg once daily
Report date	2026-05-27

Background

Following the GLUCO-BETA Phase II results, this fictional Phase III study assessed long-term efficacy and safety of Glucoren 15 mg added to metformin.

Cohort and methods

640 adults with type 2 diabetes (baseline HbA1c 8.1%) were randomized 1:1 to Glucoren 15 mg or placebo for 52 weeks.

Results

HbA1c decreased by 1.1% with Glucoren versus 0.3% with placebo at week 52. Body weight decreased by 3.1 kg versus 0.5 kg. Systolic blood pressure decreased by 4.6 mmHg versus 1.2 mmHg.

Safety

Genital mycotic infections occurred in 4.5% versus 1.3%. Diabetic ketoacidosis was reported in one Glucoren patient (0.3%). Discontinuation due to adverse events was 3.4% versus 2.5%.

Summary table

Week 52	Glucoren 15 mg (n=320)	Placebo (n=320)
Change in HbA1c	-1.1%	-0.3%
Change in body weight	-3.1 kg	-0.5 kg
Genital mycotic infection	4.5%	1.3%