Onkatinib Therapeutic Drug Monitoring: Plasma Trough Concentrations

SYNTHETIC DEMONSTRATION DATA. This document is fictional and was generated for testing a document retrieval system. It contains no real patient information and does not report genuine research findings. Do not use it for clinical decisions.

Document ID	MDA-LAB-012
Sponsor	Northbridge Clinical Research Unit (fictional)
Document type	Laboratory report
Report identifier	LAB-ONC-2107-PK
Parent study	ONCO-DELTA (NB-ONC-2107)
Assay	LC-MS/MS, lower limit of quantification 2 ng/mL

Purpose

To describe steady-state plasma trough concentrations (Ctrough) of Onkatinib 150 mg and their relationship with toxicity in 96 ONCO-DELTA patients.

Results

Median Ctrough at day 15 was 212 ng/mL (interquartile range 158 to 279). Patients with Ctrough above 300 ng/mL had a higher incidence of grade 3 or higher rash (21% vs 5%) and ALT elevation (11% vs 2%). Ctrough was not associated with objective response.

Recommendation

A target trough range of 150 to 300 ng/mL is proposed. Dose reduction to 100 mg may be considered when Ctrough exceeds 300 ng/mL with grade 2 or higher toxicity.

Summary table

Ctrough group (day 15)	Patients	Grade >=3 rash	Grade >=3 ALT elevation
<=300 ng/mL	78	5%	2%
>300 ng/mL	18	21%	11%