Neurolix Adverse Event Review: Pooled Analysis of Migraine Prevention Studies

SYNTHETIC DEMONSTRATION DATA. This document is fictional and was generated for testing a document retrieval system. It contains no real patient information and does not report genuine research findings. Do not use it for clinical decisions.

Document ID	MDA-AE-005
Sponsor	Northbridge Clinical Research Unit (fictional)
Document type	Adverse event review
Analysis identifier	AER-NLX-POOL-02
Product	Neurolix (fictional CGRP receptor antagonist), 60 mg oral daily
Report date	2025-08-09

Scope

Pooled safety data from three fictional 12-week migraine prevention studies (NLX-201, NLX-301, NLX-302) including 560 Neurolix-treated and 548 placebo-treated adults with episodic migraine.

Most common adverse events

Nausea was the most frequent adverse event (11.0% vs 4.7% placebo), followed by dizziness (6.2% vs 3.1%), constipation (5.4% vs 1.8%) and fatigue (4.1% vs 3.3%). Most events were mild and resolved within two weeks.

Serious adverse events

Serious adverse events occurred in 1.3% of Neurolix patients and 1.1% of placebo patients. Discontinuation due to adverse events was 4.6% vs 2.0%. ALT elevation of grade 3 or higher was observed in 0.3% of Neurolix patients.

Conclusion

Neurolix was generally well tolerated; gastrointestinal events were the main tolerability concern.

Summary table

Adverse event	Neurolix (n=560)	Placebo (n=548)
Nausea	11.0%	4.7%
Dizziness	6.2%	3.1%
Constipation	5.4%	1.8%
Discontinuation due to AE	4.6%	2.0%