Cross-Product Hepatic Safety Signal Review: Onkatinib and Neurolix

SYNTHETIC DEMONSTRATION DATA. This document is fictional and was generated for testing a document retrieval system. It contains no real patient information and does not report genuine research findings. Do not use it for clinical decisions.

Document ID	MDA-AE-009
Sponsor	Northbridge Clinical Research Unit (fictional)
Document type	Adverse event / signal review
Review identifier	SIG-HEP-2025-04
Products	Onkatinib (fictional TKI); Neurolix (fictional CGRP antagonist)
Report date	2025-10-30

Purpose

To review drug-induced liver injury (DILI) signals for two fictional products in the Northbridge portfolio using trial data and spontaneous reports.

Liver enzyme findings

Grade 3 or higher ALT elevation occurred in 4.2% of Onkatinib-treated patients in ONCO-DELTA versus 0.3% of Neurolix-treated patients in the pooled migraine studies. Median time to onset was 6 weeks for Onkatinib and could not be estimated for Neurolix due to few events.

Hy's law cases

One potential Hy's law case (ALT above 3x ULN with bilirubin above 2x ULN) was identified with Onkatinib; none were identified with Neurolix.

Recommendation

Liver function tests every 2 weeks for the first 3 months of Onkatinib therapy are recommended. No additional hepatic monitoring is recommended for Neurolix.

Summary table

Hepatic parameter	Onkatinib	Neurolix
Grade >=3 ALT elevation	4.2%	0.3%
Potential Hy's law cases	1	0
Monitoring recommendation	LFTs every 2 weeks x 3 months	Routine only