GLUCO-BETA: Phase II Dose-Finding Trial of Glucoren in Type 2 Diabetes

SYNTHETIC DEMONSTRATION DATA. This document is fictional and was generated for testing a document retrieval system. It contains no real patient information and does not report genuine research findings. Do not use it for clinical decisions.

Document ID	MDA-CT-003
Sponsor	Northbridge Clinical Research Unit (fictional)
Document type	Clinical trial summary
Study identifier	GLUCO-BETA (NB-T2D-2204)
Investigational product	Glucoren (fictional SGLT2 inhibitor), 5 mg and 15 mg
Report date	2024-11-02

Background

Glucoren is a fictional sodium-glucose cotransporter-2 (SGLT2) inhibitor intended as add-on therapy to metformin.

Cohort and methods

312 adults with type 2 diabetes inadequately controlled on metformin (baseline HbA1c 8.2%) were randomized to Glucoren 5 mg (n=104), Glucoren 15 mg (n=104) or placebo (n=104) for 24 weeks. The primary endpoint was the change in HbA1c at week 24.

Results

HbA1c decreased by 0.6% with Glucoren 5 mg, 0.9% with Glucoren 15 mg and 0.2% with placebo. Body weight decreased by 2.4 kg with the 15 mg dose. Fasting plasma glucose decreased by 1.7 mmol/L with 15 mg. The 15 mg dose was selected for Phase III development.

Safety

Genital mycotic infections occurred in 3.8% (15 mg), 2.9% (5 mg) and 1.0% (placebo). Urinary tract infections were reported in 4.8%, 4.8% and 3.8%. No diabetic ketoacidosis was observed.

Summary table

Week 24	Glucoren 5 mg	Glucoren 15 mg	Placebo
Change in HbA1c	-0.6%	-0.9%	-0.2%
Change in body weight	-1.6 kg	-2.4 kg	-0.3 kg
Genital mycotic infection	2.9%	3.8%	1.0%