RESP-GAMMA: Phase III Trial of Inhaled Pulmavir for Adult RSV Infection

SYNTHETIC DEMONSTRATION DATA. This document is fictional and was generated for testing a document retrieval system. It contains no real patient information and does not report genuine research findings. Do not use it for clinical decisions.

Document ID	MDA-CT-007
Sponsor	Northbridge Clinical Research Unit (fictional)
Document type	Clinical trial summary
Study identifier	RESP-GAMMA (NB-RSV-3302)
Investigational product	Pulmavir (fictional inhaled antiviral), 20 mg twice daily for 5 days
Report date	2025-12-05

Background

Respiratory syncytial virus (RSV) causes significant morbidity in older adults. Pulmavir is a fictional inhaled fusion inhibitor.

Cohort and methods

1,020 adults aged 50 years or older with laboratory-confirmed RSV infection and symptom onset within 72 hours were randomized 1:1 to Pulmavir or placebo. The primary endpoint was time to sustained symptom resolution.

Results

Median time to symptom resolution was 5.1 days with Pulmavir versus 7.4 days with placebo (hazard ratio 1.38; p<0.001). Viral load at day 3 was reduced by 1.2 log10 copies/mL more than placebo. Hospitalization within 28 days occurred in 2.4% vs 4.9% of patients.

Safety

Bronchospasm occurred in 1.6% vs 0.6%, and dysgeusia (taste disturbance) in 3.5% vs 0.4%. No deaths were attributed to study drug.

Summary table

Endpoint	Pulmavir (n=510)	Placebo (n=510)
Median time to resolution	5.1 days	7.4 days
28-day hospitalization	2.4%	4.9%
Bronchospasm	1.6%	0.6%