JOINT-LANTERN: NBX-202 in Adults with Knee Osteoarthritis

SYNTHETIC DEMONSTRATION DATA. This document is fictional and was generated for testing a document retrieval system. It contains no real patient information and does not report genuine research findings. Do not use it for clinical decisions.

Document ID	MDA-CT-014
Sponsor	Northbridge Clinical Research Unit (fictional)
Document type	Synthetic randomized trial summary
Study identifier	JOINT-LANTERN
Investigational product	NBX-202 (fictional)
Report date	2026-06-15

Purpose

This fictional study evaluated NBX-202 for knee osteoarthritis. All participants, sites, outcomes and numbers are invented demonstration data. The report is intended for testing retrieval and citations, not clinical use.

Cohort and methods

310 adults aged 18 to 75 were randomized equally to NBX-202 or placebo. Treatment lasted 16 weeks. The primary endpoint was womac pain score change from baseline, except response proportions which were measured at the endpoint visit. Outcome assessors were blinded.

Primary result

At week 16, womac pain score change was -18.4 points with NBX-202 and -10.1 points with placebo. These values are the prespecified primary analysis of the randomized population. No pediatric results were collected.

Safety

Dyspepsia occurred in 6.5% versus 3.2%; discontinuation due to adverse events was 3.2% versus 1.9%.

Interpretation and limits

The results describe this synthetic 16-week adult study only. Long-term effects, pregnancy outcomes, approved dosing and outcomes in children are not established by this document. No treatment or prescribing recommendation is made.

Summary table

Outcome	NBX-202	Placebo
WOMAC pain score change	-18.4 points	-10.1 points
Participants	155	155