

# On asymptotic power of the new test for equality of two distributions

Viatcheslav Melas and Dmitrii Salnikov

St. Petersburg State University  
Department of Mathematics  
St. Petersburg, Russia

**Abstract.** The paper introduces a new test for equality of two distributions in a class of models. We proved analytically and by stochastic simulation that the test possesses high efficiency. For the case of normal and Cauchy distributions that differ only by shift the asymptotic power of the test appear to be approximately the same as for Wilcoxon-Mann-Whitney and Kolmogorov-Smirnov tests. But if the distributions differ by scale parameters the power of the new test is considerable better.

**Keywords:** Test for equality of two distributions, Asymptotic power, Cauchy distribution, Normal distribution

## 1 Formulation of the problem

Let us consider the classical problem of testing hypothesis on the equality of two distributions

$$H_0 : F_1 = F_2 \quad (1)$$

against the alternative

$$H_1 : F_1 \neq F_2 \quad (2)$$

In the case of two independent samples  $X = (X_1, \dots, X_n)$  and  $Y = (Y_1, \dots, Y_m)$  with the distributions functions  $F_1$  and  $F_2$  respectively.

It is well known (see e.g. [1]) that in the case when both distributions differ only by the means and are normal the classical Student test has a few optimal properties. If the distributions are not normal but still differs only by means a widely popular Wilcoxon-Mann-Whitney (WMW) U-statistic is often used instead. However, it can be shown that if two normal populations differ only in variances, the power of WMW test is very low. If distributions are arbitrary there are some universal techniques such as tests by Kolmogorov - Smirnov and Cramer-von Mises (see [4]) that can be applied but in many cases these tests can be not powerful.

Recently [2] suggested the test basing on U-statistics with the logarithmic kernel and provided its numerical justification for one and many dimensional cases in comparison with a few alternative techniques. However, to the best authors knowledge there are no analytical results about its asymptotic power. Here we introduce a similar but different test and provide a few analytical results on its power.

## 2 The new test and its statistical motivation

Assume that the distribution functions  $F_1$  and  $F_2$  belongs to the class of distribution functions of random values  $\xi$ , such that

$$E[\ln(1 + \xi^2)] < \infty. \quad (3)$$

Many distributions and, in particular, the Cauchy distribution have this property.

Among all distributions with given parameters of shift and scale having this property the Cauchy's one have the maximum entropy.

Consider the following test

$$\Phi_A = \frac{1}{n(n-1)} \sum_{1 \leq i < j \leq n} g(X_i - X_j), \Phi_B = \frac{1}{m(m-1)} \sum_{1 \leq i < j \leq m} g(Y_i - Y_j), \quad (4)$$

$$\Phi_{AB} = -\frac{1}{nm} \sum_{i=1}^n \sum_{j=1}^m g(X_i - Y_j), \Phi_{nm} = \Phi_A + \Phi_B + \Phi_{AB}, \quad (5)$$

where

$$g(u) = -\ln(1 + |u|^2)$$

is under a constant term precision the logarithm of the density of the standard Cauchy distribution. (Note that Zech and Aslan (2005) took  $g(u) = \ln(|u|)$ ).

We would like to have a test that is appropriate for two distributions that differ only by shift and scale parameters and belong to a rather general class of distributions.

In particular, we consider the class of distributions satisfying (3), but the approach can be generalized for other classes of distributions.

Consider the class of distributions given by the property (3). Note that if the parameters are known the test basing on likelihood ratio is the most powerful among tests with given parameters.

The test suggested above can be considered as an approximation of logarithm of this ratio for the Cauchy distribution.

## 3 The analytical study of asymptotic power

Let us consider the case of two distributions having the property (3) and, in particular, the two that differ only by the shift parameters. To simplify notations assume that  $m = n$ . The case  $m \neq n$  is similar. Now the criterion (4) - (5) assumes the form

$$T_n = \Phi_{nn} = \frac{1}{n^2} \sum_{i,j=1}^n \ln(1 + (X_i - Y_j)^2) - \frac{1}{n(n-1)} \sum_{1 \leq i < j \leq n} \ln(1 + (X_i - X_j)^2) \quad (6)$$

$$-\frac{1}{n(n-1)} \sum_{1 \leq i < j \leq n} \ln(1 + (Y_i - Y_j)^2). \quad (7)$$

Denote by  $C(u, v)$  the Cauchy distribution with the density function

$$v/(\pi(v^2 + (x - u)^2)).$$

Let  $f_1(x)$  denotes the density of  $F_1$  and  $f_2(x)$  denotes the density of  $F_2$ . Denote

$$J_h = \int_R g(x - y - |h|/\sqrt{n}) f_1(x) f_2(y) dx dy,$$

If there exists the limit

$$\lim_{n \rightarrow \infty} n(J_h - J_0) \quad (8)$$

denote it by  $J^*(h)$ .

The basic analytical result of the present paper is the following

**Theorem 1.** *Consider the problem of testing hypothesis on the equality of two distributions (1)-(2) where both functions have the property (3). Then (i) under the condition  $n \rightarrow \infty$  the distribution function of  $nT_n$  converges under  $H_0$  to that of the random value*

$$(aZ + b)^2, \quad (9)$$

where  $Z$  has the normal distribution with zero expectation and variance equal to 1,  $a^2 = J_0/3, b = 0$ .

(ii) Let  $F_1 = F(\nu, \mu), F_2 = F(\nu + \theta, \mu)$ , where  $\nu$  and  $\mu$  be the shift and scale parameters of the distribution  $F$ ,  $F$  is arbitrary distribution with property (3) and  $\theta = h/\sqrt{n}, h$  is an arbitrary given number. Then the distribution function of  $nT_n$  converges under  $H_1$  to that of the random value

$$(aZ + b)^2,$$

where  $a^2 = J_0/3, b = 0$  for the case of  $H_0$  and  $a^2 = J_0/3, b^2 = J^*(h)$  for  $H_1$ . In this case the power of the criterion  $T_n$  with significance  $\alpha$  is asymptotically equal to that is given by the formula

$$Pr\{Z \geq z_{1-\alpha/2} - \sqrt{\frac{3J^*(h)}{J_0}}\} + Pr\{Z \leq -z_{1-\alpha/2} - \sqrt{\frac{3J^*(h)}{J_0}}\}.$$

(iii) If  $F_1 = C(\nu, 1), F_2 = C(\nu + \theta)$  then in the part (ii)  $a^2 = (2/3) \ln 3, b = h/3$  and

$$Pr\{Z \geq z_{1-\alpha/2} - (1/\sqrt{6 \ln 3})h\} + Pr\{Z \leq -z_{1-\alpha/2} - (1/\sqrt{6 \ln 3})h\}.$$

The proof of the theorem is given in the Appendix.

## 4 Simulation results

We found by a stochastic simulation that the formula presents an approximation of the power of the test  $T_n$  with a good accuracy (see tables 1-3 below).

At the tables 1-12 results for cases  $n = 100, 500, 1000$  and different values of  $h$  with  $\alpha = 0.05$  are given for normal and Cauchy distributions that differ either by shift or by scale parameters. The critical values were calculated in two way: by simulation of the initial distribution and by random permutations (we used 800 random permutation in all cases). It worth to be noted that the results are very similar. Since the permutation technics is more universal, it can be recommended for practical applications.

Note that in all these cases when the distributions differ only in the scale parameters the power of  $T_n$  and that of the Wilcoxon-Mann-Whitney and the Kolmogorov - Smirnov tests were approximately equal to each other. It can pointed out also that if the variances are not standard but are known we should simply make a the corresponding normalisation. But for the cases where the distributions differ in scale parameters the Wilcoxon-Mann-Whitney is not appropriate at all and the power of the Kolmogorov - Smirnov test is considerably lower.

**Table 1.** Cauchy distribution,  $X \sim C(0, 1)$ ,  $Y \sim C(h/\sqrt{n}, 1)$ ,  $n = 100$

$h$	$T_{n,perm}$	$T_{n,sim}$	$formulae$	$wilcox.test$	$ks.test$	$ad.test$
1	6.4	6.3	6.8	6.6	6.3	7.1
2	10.1	10.6	12.2	11.9	11.1	11.6
3	19.6	20.3	21.5	20.5	20.2	20.7
5	50.9	50.5	49.5	48.5	53.1	52.2
7	82	82.3	77.8	77.2	83.6	80.7
9	96.7	96.8	93.9	91.5	96.5	95.2

**Table 2.** Cauchy distribution,  $X \sim C(0, 1)$ ,  $Y \sim C(h/\sqrt{n}, 1)$ ,  $n = 500$

$h$	$T_{n,perm}$	$T_{n,sim}$	$formula$	$wilcox.test$	$ks.test$	$ad.test$
1	5.8	6.1	6.8	6.4	6.4	7.1
2	11.6	11.6	12.2	12.6	13.9	12.2
3	21	21.8	21.5	22.2	24.3	22.8
5	50.9	51	49.5	48	57.9	50.3
7	82.2	82.4	77.8	75.6	85.9	81.1
9	96.2	96.5	93.9	93.2	97.2	96.0

**Table 3.** Cauchy distribution,  $X \sim C(0, 1)$ ,  $Y \sim C(h/\sqrt{n}, 1)$ ,  $n = 1000$ 

$h$	$T_{n,perm}$	$T_{n,sim}$	$formula$	$wilcox.test$	$ks.test$	$ad.test$
1	6.3	6	6.8	6.8	8.1	6.8
2	11.4	11.9	12.2	12.9	13.4	12.9
3	21	20.9	21.5	22.8	26.2	22.2
5	53.6	53.6	49.5	50.8	59.6	54.2
7	84	84.5	77.8	79.5	87.6	84.4
9	96.6	96.6	93.9	93.2	98.3	96.3

**Table 4.** Cauchy distribution,  $X \sim C(0, 1)$ ,  $Y \sim C(0, 1 + h/\sqrt{n})$ ,  $n = 100$ 

$h$	$T_{n,perm}$	$T_{n,sim}$	$wilcox.test$	$ks.test$	$ad.test$
2	10.6	11.9	5.4	5.4	6.9
4	27.6	29.8	5.5	8.7	11.3
6	49.4	53.6	5.5	15.9	22.2
8	68.8	73.5	5.5	25	37.7
10	84.2	87.1	5.2	36.4	55.4

**Table 5.** Cauchy distribution,  $X \sim C(0, 1)$ ,  $Y \sim C(0, 1 + h/\sqrt{n})$ ,  $n = 500$ 

$h$	$T_{n,perm}$	$T_{n,sim}$	$wilcox.test$	$ks.test$	$ad.test$
2	9.4	10	4.5	6.3	6.2
4	28.5	30.6	4.8	14	12.3
6	54.5	56.5	5	26.1	29.7
8	79.5	80.5	5.2	43.3	51.0
10	93	94	5.2	62.2	74.2

**Table 6.** Cauchy distribution,  $X \sim C(0, 1)$ ,  $Y \sim C(0, 1 + h/\sqrt{n})$ ,  $n = 1000$ 

$h$	$T_{n,perm}$	$T_{n,sim}$	$wilcox.test$	$ks.test$	$ad.test$
2	10.2	10.5	5	7.6	7.3
4	32.4	33.8	5.2	13.8	14.9
6	61.1	62.8	5.2	27.9	32.8
8	84.8	85.6	5.2	47.4	59.7
10	96.1	97.1	5.4	67.9	82.8

**Table 7.** Normal distribution,  $X \sim N(0, 1)$ ,  $Y \sim N(h/\sqrt{n}, 1)$ ,  $n = 100$ 

$h$	$T_{n,perm}$	$T_{n,sim}$	$wilcox.test$	$ks.test$	$ad.test$
1	11.1	11.3	12.5	9.5	12.2
2	29.3	29	31.1	20.5	29.6
3	52.4	53.4	55.8	42	55
4	77.5	77.5	80.6	64.9	78.9
5	91.9	92.5	93.1	84.7	93.1

**Table 8.** Normal distribution,  $X \sim N(0, 1)$ ,  $Y \sim N(h/\sqrt{n}, 1)$ ,  $n = 500$ 

$h$	$T_{n,perm}$	$T_{n,sim}$	$wilcox.test$	$ks.test$	$ad.test$
1	9.2	8.9	9.6	8.3	9.0
2	23.9	23.9	26.3	20.6	25.4
3	47.3	48.9	51.7	41.4	49.7
4	75.3	75.1	77.8	66.9	76.9
5	91.1	91	92.8	86.1	92.6

**Table 9.** Normal distribution,  $X \sim N(0, 1)$ ,  $Y \sim N(h/\sqrt{n}, 1)$ ,  $n = 1000$ 

$h$	$T_{n,perm}$	$T_{n,sim}$	$wilcox.test$	$ks.test$	$ad.test$
1	11	11.3	11.5	10	11.6
2	26.4	27.4	28.5	22	27.7
3	51.3	51.6	54.2	44.6	52.9
4	76.7	77	79.3	68.9	77.9
5	91.6	91.2	92.7	86.6	92.1

**Table 10.** Normal distribution,  $X \sim N(0, 1)$ ,  $Y \sim N(0, 1 + h/\sqrt{n})$ ,  $n = 100$ 

$h$	$T_{n,perm}$	$T_{n,sim}$	$wilcox.test$	$ks.test$	$ad.test$
1	8.1	8.7	6.4	5.3	7.3
2	15	17.4	6.3	7.2	12.7
3	30.5	34.2	6.6	10.7	24.0
4	50.6	57.1	6.7	16.7	39.9
5	70.8	76.7	6.5	24.8	59.9

**Table 11.** Normal distribution,  $X \sim N(0, 1)$ ,  $Y \sim N(0, 1 + h/\sqrt{n})$ ,  $n = 500$ 

$h$	$T_{n,perm}$	$T_{n,sim}$	$wilcox.test$	$ks.test$	$ad.test$
1	8.3	8.4	5	7.4	7.7
2	15.4	16.7	5.1	10.3	12.8
3	33.2	34.7	5.4	16.4	28.3
4	60	63.3	5.6	25.3	52.6
5	83.1	86.3	5.5	40.4	78.1

**Table 12.** Normal distribution,  $X \sim N(0, 1)$ ,  $Y \sim N(0, 1 + h/\sqrt{n})$ ,  $n = 1000$ 

$h$	$T_{n,perm}$	$T_{n,sim}$	$wilcox.test$	$ks.test$	$ad.test$
1	6.7	6.9	5.4	6	6.7
2	15.1	16.4	5.5	9.9	13.1
3	33.2	36	5.4	16.1	30.6
4	62.2	64	5.6	27.5	56.8
5	84.6	86.6	5.4	43.6	81.1

## 5 Conclusion

In this paper we suggested a new test for equality of two distributions. Its asymptotic power was analytically established for the case of distributions that differ only by shift. For the case of Cauchy distribution this formula was presented in a closed form since it proved to be possible to calculate the corresponding integrals analytically.

By stochastic simulation we found that for the Normal and Cauchy distributions that differ only by shift the power of the new test is approximately equal to that of the Wilcoxon-Mann-Whitney and the Kolmogorov - Smirnov tests. However if the distributions differ by the scale parameter our simulations show that the new test is considerably better than the Kolmogorov - Smirnov test. And in this case the the Wilcoxon-Mann-Whitney test is not appropriate at all.

## Acknowledgments

The authors are indebted to professor Yakov Nikitin for the help in calculating the integrals. Work of Viatcheslav Melas was supported by RFBR (grant N 20-01-00096).

## 6 Appendix

Proof of Theorem 1.

Let us begin with studying the asymptotic behaviour of the magnitude  $E(nT_n)^2$ .

**Lemma 1.** (1) If hypothesis  $H_0$  is satisfied and  $F_1$  possesses property (3) then there exists a finite limit of  $d = \lim E(nT_n)^2$  with  $n \rightarrow \infty$ .

(2) If hypothesis  $H_1$  is satisfied for  $F_2(x) = F_1(x - \theta, \theta = h/\sqrt{n})$  and  $F_1$  and  $F_2$  possess property (3) then there exists a finite limit of  $E(nT_n)^2$  for  $n \rightarrow \infty$  and it is given by the formula

$$d + 2J^*(h)J_0 + J^*(h)^2.$$

Proof of the lemma.

Note that  $(nT_n)^2$  is equal to

$$\begin{aligned} n^2 \left[ \frac{1}{n^2} \sum_{i,j=1}^n [g(X_i - Y_j) - J_0] - \frac{1}{n(n-1)} \sum_{i < j, i,j=1}^n [g(X_i - X_j) - J_0] - \right. \\ \left. \frac{1}{n(n-1)} \left[ \sum_{i < j, i,j=1}^n [g(Y_i - Y_j) - J_0] \right]^2 \right], \end{aligned}$$

where  $g(z) = \ln(1 + z^2)$ .

The idea of the proof consists in the splitting the three squares of three sums/

including in this sum and three pairwise products into peculiar sums of the identical structure. Then to each peculiar sum either law of large numbers or central limit theorem is applied.

The square of the first sum,

$$n^2 \left\{ \frac{1}{n^2} \sum_{i,j=1}^n [g(X_i - Y_j) - J_0] \right\}^2,$$

can be represented by sum of the following peculiar sums.

$$1) \ n^2 \left( \frac{1}{n^2} \right)^2 \sum_{i,j=1}^n [(g(X_i - Y_j) - J_1)]^2,$$

$$2a) \ n^2 \left( \frac{1}{n^2} \right)^2 \sum_{i,j=1, i \neq j}^n \sum_{k=1}^n [(g(X_i - Y_j) - J_0)][(g(X_k - Y_j) - J_0)],$$

$$2b) \ n^2 \left( \frac{1}{n^2} \right)^2 \sum_{i,j=1, i \neq j}^n \sum_{k=1}^n [g(Y_k - X_i) - J_0][g(Y_k - X_j) - J_0],$$

$$3) \ n^2 \left( \frac{1}{n^2} \right)^2 \sum_{i,j=1}^n \sum_{l,k=1, (l \neq k) \text{ or } (i \neq j)}^n [g(X_i - Y_j) - J_0][g(X_l - Y_k) - J_0].$$

Similar expression can be obtained for each of the two other squares and three pairwise products. Note the limit with  $n \rightarrow \infty$  for the peculiar sums of the type 1) is finite due to the law of large numbers.

The peculiar sums of type 3) consist of multiplications of independent terms with zero expectation. Therefore for any  $n$  the expectation of these peculiar sums is zero and the limit is also equal to 0.

Consider the peculiar sum  $ES_{xy,2a}^2$ . It can be written as  $I_1 - I_2$ ,

$$I_1 = \frac{1}{n} \sum_{k=1}^n \left\{ \left[ \sum_{i=1}^n (g(x_k - y_i) - J_0) \right] / \sqrt{n} \left[ \sum_{j=1}^n (g(x_k - y_j) - J_0) \right] / \sqrt{n} \right\},$$

$$I_2 = \frac{1}{n^2} \sum_{i,j=1}^n (g(x_i - y_j))^2.$$

Note that  $I_2$  tends with  $n \rightarrow \infty$  to a finite limit due to the law of large numbers. And due to the central limit theorem under fixed  $X_k$  the random value

$$\sum_{i=1}^n [g(x_k - y_i) - J_1] / \sqrt{n}$$



tends to normal random value with zero expectation and a finite variance.

Others sum of type 2) have a similar behaviour. Thus under  $H_0$  there exists a final limit

$$\lim_{n \rightarrow \infty} E(nT_n)^2.$$

Denote this limit by  $d$ .

Thus the first part of the lemma is proved.

Let now  $H_1$  holds with  $|h| > 0$ . In this case only the behavior of the following sums

$$\begin{aligned} n^2 \left\{ \frac{1}{n^2} \sum_{i,j=1}^n [g(X_i - Y_j) - J_0] \right\}^2, \\ S_{xy,xx,2} = n^2 \left( \frac{1}{2n(n-1)n^2} \right) \sum_{k=1}^n \sum_{i,j=1, i \neq j, k, j \neq k}^n [(\ln(1 + (X_k - Y_i)^2) - J_1) [(\ln(1 + (X_k - X_j)^2) - J_1)], \\ S_{xy,xx,3} = n^2 \left( \frac{1}{2n(n-1)n^2} \right) \sum_{i,j=1}^n \sum_{k=1, l=1, l \neq k}^n [(\ln(1 + (X_k - Y_i)^2) - J_1) [(\ln(1 + (X_l - X_j)^2) - J_1)]. \end{aligned}$$

and  $S_{xy,yy,2}, S_{xy,xy,3}$  that are determined in a similar way that  $S_{xy,xx,2}, S_{xy,xx,3}$ . By a direct calculation we will obtain that

$$\lim_{n \rightarrow \infty} E(nT_n)^2 = d + 2J^*(h)J_0 + J^*(h)^2.$$

Lemma is proved.

**Lemma 2.** For  $g(x) = x^2$  the following identity holds

$$\Phi_{nm} = (\bar{x} - \bar{y})^2,$$

where

$$\bar{x} = \left( \sum_{i=1}^n X_i \right) / n, \bar{y} = \left( \sum_{i=1}^m Y_i \right) / m.$$

The proof follows from the known formula [see e.g. [3], p.296]

$$\frac{1}{n(n-1)} \sum_{1 \leq i < j \leq n} (X_i - X_j)^2 = \frac{1}{(n-1)} \sum_{i=1}^n (X_i - \bar{x})^2.$$

by direct calculations.

Assume that  $H_0$  holds. Let  $C$  be an arbitrary positive number,

$$\tilde{X} = (\tilde{X}_1, \dots, \tilde{X}_n), \quad \tilde{Y} = (\tilde{Y}_1, \dots, \tilde{Y}_n),$$

where  $\tilde{X}_i = X_i$ , if  $|X_i| \leq C$  and  $\tilde{X}_i = C$  if  $X_i > 0$ ,  $\tilde{X}_i = -C$  if  $X_i < 0$  otherwise. And  $\tilde{Y}_i$  are determined similarly. Note that  $0 \leq \ln(1 + x^2) \leq x^2$ . Therefore there exists a value  $t$  that depends from  $\tilde{X}$  and  $\tilde{Y}$  such that

$$n \left\{ \frac{1}{n^2} \sum_{i,j=1}^n \ln(1 + (\tilde{X}_i - \tilde{Y}_j)^2) - \frac{1}{n(n-1)} \sum_{i < j} \ln(1 + (\tilde{X}_i - \tilde{X}_j)^2) - \right. \quad (10)$$

$$\frac{1}{n(n-1)} \sum_{i < j} \ln(1 + (\tilde{Y}_i - \tilde{Y}_j)^2) = t \left( \sum_{i=1}^n \tilde{X}_i / \sqrt{n} - \sum_{i=1}^n \tilde{Y}_i / \sqrt{n} \right)^2. \quad (11)$$

For constructing the right hand side we applied Lemma 2. Note that for distributions  $F_1$  and  $F_2$  satisfying (3) it follows from Lemma 1 that the variance of the left hand side is finite. Therefore the variance of the right hand side is also finite for arbitrary  $C$ . Passing to the limit with  $n \rightarrow \infty$  we obtain due to the central limit theorem that the right hand side has the limit distribution of the form (9) where  $Z$  has the normal distribution with zero expectation and variance equal to 1. And its variance is equal to the variance of the left hand side of (11). Since  $C$  is arbitrary we obtain that the limiting distribution has the required form for  $H_0$ .

For determining  $a$  and  $b$  in the part (ii) of the theorem we now can use the equality

$$E((aZ + b)^2)^2 = \lim_{n \rightarrow \infty} E(nT_n)^2, \quad (12)$$

that follows from (11).

Since  $EZ^2 = 1$ ,  $EZ^4 = 3$ , we have for the left hand side (12)

$$3a^4 + 6a^2b^2 + b^4. \quad (13)$$

The formula for the left hand side follows from Lemma 1. And the asymptotic behaviour of the power follows from the asymptotic normality of  $\sqrt{n}T_n$ . In order to calculate the right hand side of (12) in (iii) the following result is crucial.

**Lemma 3.** *If  $X$  and  $Y$  are independent random values with the distribution  $C(0, 1)$ , then*

$$E \ln(1 + (X - Y)^2) = \ln 9, \quad E \ln(1 + (X - Y - \theta)^2) - \ln 9 = \ln(1 + \theta^2/9).$$

In order to prove this Lemma we need the following integrals

$$\int_R \frac{\ln(1 + (x - y)^2)}{\pi(1 + y^2)} dy = \ln(4 + x^2),$$

$$\int_R \frac{\ln(4 + x^2)}{\pi(1 + x^2)} dx = \ln 9,$$

([5] 4.296.2 and 4.295.7.)

$$\int_R \frac{\ln(4 + (x + \theta)^2)}{\pi(x^2 + 1)} dx = \ln(9 + \theta^2),$$

[see [6], formula (2.6.14.19)]. Using these integrals we obtain

$$\begin{aligned} E \ln(1 + (X - Y - \theta)^2) - \ln 9 &= 2 \int_R \int_R \frac{\ln(1 + (x - y - \theta)^2)}{\pi^2(1 + x^2)(1 + y^2)} dx dy - \ln 9 \\ &= \int_R \frac{\ln(4 + (y + \theta)^2)}{\pi(1 + y^2)} dy - \ln 9 = \ln(9 + \theta^2) - \ln 9 = \ln(1 + \theta^2/9). \end{aligned}$$

Submitting here  $\theta = 0$  we obtain both formulas of the Lemma. Note that  $\theta^2 = nh^2$  and

$$\lim_{n \rightarrow \infty} n \ln(1 + \theta^2/9) = (1/9)h^2.$$

Therefore we obtain for the right hand side (8) with some algebra

$$3a^4 + \frac{(2 \ln 9)h^2}{9} + \frac{h^4}{81}. \quad (14)$$

From Lemma 3 and (14) we obtain

$$b = \frac{1}{3}h, \quad a^2 = \frac{2}{3} \ln 3.$$

The formula for the power follows from the form of the limiting distribution (9).

## References

1. Lehmann E.: Testing Statistical Hypotheses, Probability and Statistics Series, Wiley (1986).
2. Zech, G., Aslan, B.: New test for the multivariate two-sample problem based on the concept of minimum energy. Journal of Statistical Computation and Simulation 75(2), pp. 109-119 (2005).
3. Wassily Hoeffding: A class of statistics with asymptotically normal distribution. Ann. Math. Statistics 19, pp. 293-325 (1948).
4. Buening, H.: Kolmogorov-Smirnov and Cram'er-von Mises type two-sample tests with various weight functions. Communications in Statistics-Simulation and Computation, 30, pp. 847-865 (2001).
5. Gradshteyn, I.S., Ryzhik, I.M.: Table of Integrals, series and products. Seventh edition AMSTERDAM,BOSTON,HEIDELBERG, LONDON
6. Prudnikov, A.P., Brychkov, Yu.A., and Marichev, O.I.: Integrals and Series. Elementary Functions (Nauka, Moscow, 1981) [in Russian].