

STUDENT'S HANDBOOK GUIDE TO MATH 115

ELEMENTARY CALCULUS II



EVERYTHING A STUDENT NEEDS TO KNOW ABOUT MATH 115

Master the skills of intergration using the techniques listed in this book. From regular and trigonometric subsitutions, partial fractions, and integration by parts. Using the right technique(s) will lead you to a heroic victory against the machination of the professor.

By slaying his assignment minions, defeating his midterm champion(s) and ending the final themselves. The heroic legacy of your achievement will be remembered on your academic transcript.

Disclaimer: Please don't use the integration symbol as a weapon, especially against an Ancient Red Dragon, or a dragon of any kind. If you do they will most certainly mock you for your stupidity before incinerating you.

CONTENTS

CHAPTER 1-5: MATH 114 REVIEW	1
Derivatives	1
Linear Approximation	1
General Rules of Derivatives	2
L'Hospital's Rule	2
Derivatives of inverse functions	2
Inverse Trigonometric Functions	2
Implicit functions	3
Integrals	4
Riemann Sum	4
Fundamental Theorem of Calculus	4
General Rules for Integrals	4
Properties of Integrals	4
CHAPTER 6: APPLICATIONS OF INTEGRATION	5
Areas between curves	5
Volume	5
Example	5
Volume with an offset x-axis	6
Volumes by Cylindrical Shells	6
Average Value of a function	6
Exercise	6
CHAPTER 7: TECHNIQUES OF INTEGRATION	7
Substitution	7
Arctan Integration	7
Arcsin Integration	7
Integration by Parts	8
Trig Substitution	8
Trigonometric Substitution with Angles	9
Partial Fraction	11
Long Division	11
$Q(x)$ is a product of distinct linear factors	11
$Q(x)$ is a product of linear factors, some of which are repeated	12
$Q(x)$ Contains Irreducible Quadratic Factors, None of Which Is Repeated	12
General Formula	13
Improper Integral	13
Type 1	13
Type 2	13
Nonelementary Integral	14
Exercise	15
CHAPTER 8: FURTHER APPLICATIONS OF INTEGRATION	16
Arc Length	16
Surface Area	16
Exercise	17

CHAPTER 9: DIFFERENTIAL EQUATION	18
Initial Value Problem	18
Population Dynamics	18
The Law of Natural Growth	18
Separable Equation	18
Logistic Model	19
Newton Cooling Equation	19
Linear Equations	20
Integrating factor $\mu(t)$	20
Mixing problem	21
CHAPTER 10: PARAMETRIC EQUATION AND POLAR COORDINATES	22
Parametric Curve	22
Position and speed of a particle in orbit	22
Tangent line	23
Concavity	23
Area	25
Arc length	25
Surface Area	26
ellipse	26
Hyperbolas	27
Exercise	27
PROFESSOR'S VILLAINS	28
Tiamat, The Grade Slayer	28
APPENDIX A: UNIT CIRCLE	29
APPENDIX B: OTHER RULES	30
Log rules	30
Trigonometric Hyperbolic Functions	30
CREDITS	31
General	31
Art	31
Disclaimer	31

CHAPTER 1-5: MATH 114 REVIEW

DERIVATIVES

DERIVATIVE DEFINITION


Let a function $f(x)$ be defined on an open interval $I = (a, b)$. The function $f(x)$ is differentiable at $x_0 \in I$ if the following limit exists

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}.$$

An equivalent formula is derived if we take $x - x_0 = h$

$$f'(x_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}.$$

Geometrical Interpretation The interpretation is shown in the following figure



Physical Interpretation If $x = f(t)$ denote the position of a mass particle on the x -axis, where t stands for time, the value $f'(t_0)$ measures the velocity of the particle at t_0 .

If $f'(t_0) > 0$ the particle moves forward, if $f'(t_0) < 0$ the particle moves backward.

What is the physical interpretation of $f''(t)$?

LINEAR APPROXIMATION

In the definition of derivative, if we relax the limit, we can write

$$f'(x_0) \cong \frac{f(x) - f(x_0)}{x - x_0},$$

or equivalently

$$f(x) \cong f(x_0) + f'(x_0)(x - x_0).$$

The right hand side of the above equality is just the equation of the tangent line,

$$T(x) = f(x_0) + f'(x_0)(x - x_0),$$

thus for x close to x_0 we obtain the following approximation

$$f(x) \cong T(x).$$

EXAMPLE

Use the linear approximation to approximate $\sqrt{66}$.

SOLUTION

Take $f(x) = \sqrt{x}$ and thus if we take $x_0 = 64$, we can write

$$\begin{aligned} f(66) &\cong f(64) + f'(64)(66 - 64) \\ &= 8 + \frac{1}{16}(2) = 8 + \frac{1}{8} = \frac{65}{8} \cong 8.125 \end{aligned}$$

We see the approximate value is close to the real value of $f(66) = 8.1240384$.

EXERCISE

Use linear approximation and approximate $\log_2(2.1)$.

GENERAL RULES OF DERIVATIVES

THE RULES FOR DERIVATIVES

These are the general rules for derivatives.

$$y = x^n \Rightarrow y' = nx^{n-1} dx \quad (1)$$

$$y = \sin(x) \Rightarrow y' = \cos(x) dx \quad (2)$$

$$y = \cos(x) \Rightarrow y' = -\sin(x) dx \quad (3)$$

$$y = e^x \Rightarrow y' = e^x dx \quad (4)$$

$$y = \ln|x| \Rightarrow y' = \frac{1}{x} dx \quad (5)$$

$$(fg)' = f'g + fg' \quad (6)$$

$$\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2} \quad (7)$$

$$(f(g(x)))' = f'(g(x)) \cdot g'(x) \quad (8)$$

$$(a^{f(x)})' = f'(x) \ln(a) \quad (9)$$

$$y = (f(x)^{g(x)}) \Rightarrow y' = y \cdot (g(x) \cdot \ln(f(x)))' \quad (10)$$

$$y = \log_a f(x) \Rightarrow y' = \frac{f(x)'}{f(x) \cdot \ln(a)} \quad (11)$$

$$\sinh(x)' = \cosh(x) \quad (12)$$

$$\cosh(x)' = \sinh(x) \quad (13)$$

L'HOSPITAL'S RULE

L'HOSPITAL'S RULE

Suppose f and g are differentiable and $g'(x) \neq 0$ on an open interval I that contains a (except possibly at a). Suppose that

$$\lim_{x \rightarrow a} f(x) = 0 \text{ and } \lim_{x \rightarrow a} g(x) = 0$$

or that

$$\lim_{x \rightarrow a} f(x) = \pm\infty \text{ and } \lim_{x \rightarrow a} g(x) = \pm\infty$$

Then

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

if the limit on the right side exists (or is ∞ or $-\infty$).

DERIVATIVES OF INVERSE FUNCTIONS

THE GENERAL FORMULA

If we have an expression like this: $y = f^{-1}(x)$. This is equivalent to:

$$y = f^{-1}(x) \Leftrightarrow f(y) = x \quad (14)$$

The derivative of y is then:

$$(f(y))' = (x)'$$

Using the chain rule we get

$$f'(y) \cdot y' = 1$$

$$y' = \frac{1}{f'(y)}$$

$$y' = \frac{1}{f'(f^{-1}(x))}$$

INVERSE TRIGONOMETRIC FUNCTIONS

INVERSE TRIGONOMETRIC FUNCTIONS

These are the standard trigonometric functions, their inverse, and their domains.

$$\theta = \cos^{-1}(x) \Leftrightarrow x = \cos(\theta) \quad (15)$$

$$\theta \in [0, \pi], x \in [-1 \leq x \leq 1]$$

$$\theta = \sin^{-1}(x) \Leftrightarrow x = \sin(\theta) \quad (16)$$

$$\theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right], x \in [-1 \leq x \leq 1]$$


$$\theta = \tan^{-1}(x) \Leftrightarrow x = \tan(\theta) \quad (17)$$

$$\theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right], x \in (-\infty, \infty)$$

When you have an expression in the form like this $f(g^{-1}(x))$ where $f(x)$, and $g^{-1}(x)$ are trigonometric functions then using the following equation above we get these trigonometric identities.

COS INVERSE

If you have $y = \theta = \cos^{-1}(x)$. Then, representing the trigonometric function using a triangle is shown below




$$\sin(\cos^{-1}(x)) = \sqrt{1 - x^2} \quad (18)$$

$$\tan(\cos^{-1}(x)) = \frac{\sqrt{1 - x^2}}{x} \quad (19)$$

SIN INVERSE

If you have $y = \theta = \sin^{-1}(x)$. Then, representing the trigonometric function using a triangle is shown below




$$\cos(\sin^{-1}(x)) = \sqrt{1 - x^2} \quad (20)$$

$$\tan(\sin^{-1}(x)) = \frac{x}{\sqrt{1 - x^2}} \quad (21)$$

TAN INVERSE

If you have $y = \theta = \tan^{-1}(x)$. Then, representing the trigonometric function using a triangle is shown below



$$\sin(\tan^{-1}(x)) = \frac{x}{\sqrt{1 + x^2}} \quad (22)$$

$$\cos(\tan^{-1}(x)) = \frac{1}{\sqrt{1 + x^2}} \quad (23)$$

EXAMPLE

$$y = \tan^{-1}(x)$$

What is y' ?

$$\tan^{-1}(x) = y \Rightarrow \tan(y) = x$$

$$\sec^2(y) \cdot y' = 1$$

$$y' = \frac{1}{\sec^2(y)}$$

$$y' = \frac{1}{\sec^2(\tan^{-1}(x))}$$

From the Tan inverse box we

know that $\sec(\theta) = \sqrt{1 + x^2}$.

$$y' = \frac{1}{\sec^2(\sqrt{1 + x^2})}$$

$$y' = \frac{1}{1 + x^2} dx$$

IMPLICIT FUNCTIONS

If you have an expression where the variable you are deriving is being used in a function or is not alone. Then you have to derive both sides of the = sign to evaluate the expression. This type of function is called an implicit as the variable in question isn't explicitly defined.

You derive each term with respect to the variable you are deriving for and collect the deriving variable on one side and all other terms to the other,

EXAMPLE

$$e^y + y + x^2 + \sin(x) = 3$$

$$y' = e^y \cdot y' + y' + 2x + \cos(x) = 0$$

$$y'(e^y + 1) + 2x + \cos(x) = 0$$

$$y' = \frac{-2x - \cos(x)}{e^y + 1} dx$$

INTEGRALS


RIEMANN SUM

RIEMANN SUM

Let $y = f(x)$ and $a \leq x \leq b$ is a continuous function. The area under the curve of the function is calculated by the formula

$$A = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(x_k) \Delta x,$$

where $\Delta x = \frac{b-a}{n}$, and $x_k = a + k\Delta x$.



$$\begin{aligned} x_1 &= a + \Delta x \\ x_2 &= a + 2\Delta x \\ \dots &\dots \end{aligned} \quad \begin{aligned} A_1 &= f(x_1) \Delta x \\ A_2 &= f(x_2) \Delta x \\ \dots &\dots \end{aligned}$$

FUNDAMENTAL THEOREM OF CALCULUS

FUNDAMENTAL THEOREM OF CALCULUS

The fundamental theorem of calculus states that

$$A = \int_a^b f(x) dx$$

where \int stands for the anti-derivative operation.

EXAMPLE

For finding the area under the curve $y = x^2$, $0 \leq x \leq 1$, we use the Riemann sum. We have $a = 0$, $b = 1$, and thus $\Delta x = \frac{1-0}{n} = \frac{1}{n}$. Intermediate points are $x_k = a + k\Delta x = \frac{k}{n}$, and thus

$$A = \lim_{n \rightarrow \infty} \sum_{k=1}^n \left(\frac{k}{n} \right)^2 \frac{1}{n} = \lim_{n \rightarrow \infty} \frac{1}{n^3} \sum_{k=1}^n k^2.$$

We know

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6},$$

and thus

$$A = \lim_{n \rightarrow \infty} \frac{(n+1)(2n+1)}{6n^2} = \frac{1}{3}.$$

Alternatively, we can use the fundamental theorem of calculus and write

$$A = \int_0^1 x^2 dx = \frac{1}{3}x^3 \Big|_0^1 = \frac{1}{3}.$$

GENERAL RULES FOR INTEGRALS

THE GENERAL RULES FOR INTEGRALS

There are two different types of integrals: indefinite and definite

INDEFINITE

$$\int 1 dx = x + C \quad (24)$$

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C \quad (25)$$

$$\int \frac{1}{x} dx = \ln|x| + C \quad (26)$$

$$\int \cos(x) dx = \sin(x) + C \quad (27)$$

$$\int \sin(x) dx = -\cos(x) + C \quad (28)$$

$$\int e^x dx = e^x + C \quad (29)$$

$$\int \frac{dx}{1+x^2} = \tan^{-1}(x) + C \quad (30)$$

$$\int \frac{dx}{\sqrt{1-x^2}} = \sin^{-1}(x) + C \quad (31)$$

DEFINITE

$$\int_a^b f(x) dx = F(b) - F(a) \quad (32)$$

PROPERTIES OF INTEGRALS

PROPERTIES OF INTEGRALS


$$\int_b^a f(x) dx = - \int_a^b f(x) dx \quad (33)$$

$$\int_a^b f(x) dx + \int_b^c f(x) dx = \int_a^c f(x) dx \quad (34)$$

CHAPTER 6: APPLICATIONS OF INTEGRATION

AREAS BETWEEN CURVES

To find the area between two curves there exist a general formula to calculate such a question.



AREAS BETWEEN CURVES FORMULA

$$\int_a^b |(f(x) - g(x))| dx = \int_a^b |f(x)| dx - \int_a^b |g(x)| dx \quad (35)$$

VOLUME

To find the volume of a curve about the x axis use this equation below.

$$V = \int_a^b \pi y^2 dx \quad (36)$$

To find the volume of a curve about the y axis use this equation below.

$$V = \int_a^b \pi x^2 dy \quad (37)$$

WHAT DOES dv MEAN?

$$dv = \pi y^2 dx$$

or

$$dv = \pi x^2 dy$$

These are called differential volume of the curve, the sum of which gives us the volume.

GENERAL AREA FORMULA

$$V = V_o - V_i = \int_a^b f(x) dx - \int_a^b g(x) dx \quad (38)$$


$f(x)$ is the function that encompasses $g(x)$.

$g(x)$ is the function that encompasses the hollowed out cylinder

EXAMPLE

QUESTION

Find the volume around the y-axis:



SOLUTION

$$\begin{aligned} V &= \int_0^1 \pi x^2 dy - \int_0^1 \pi x^2 dy \\ &= \pi \int_0^1 y dy - \pi \int_0^1 y^2 dy \\ &= \pi \frac{1}{2} y^2 \Big|_0^1 - \pi \frac{1}{3} y^3 \Big|_0^1 \\ &= \pi \frac{1}{2} - \pi \frac{1}{3} = \frac{\pi}{6} \end{aligned}$$

VOLUME WITH AN OFFSET X-AXIS

If you need to calculate a volume with a cylinder removed due to an offset then you will need to modify the general formula.

OFFSET X-AXIS FORMULA

$$V = \int_a^b \pi(a \pm x)^2 dy - (\pi r^2 * (b - a)) \quad (39)$$

The sign is negative if the offset is positive, and positive if the offset is negative. You have to substitute x with the function whose domain is y.

VOLUMES BY CYLINDRICAL SHELLS

We can also calculate the volume of a function using the cylindrical shells method.

CYLINDRICAL SHELL FORMULA

$$V = \int_a^b 2\pi x f(x) dx \quad \text{where } 0 \leq a < b \quad (40)$$

Where:

$2\pi x$ is the circumference

$f(x)$ is the height between two functions.

dx is the thickness, which is infinitely small.

Note: This is for volumes that are generated around the y axis. If the volume generated is done by the x axis then replace the function that uses y as the domain.

OFFSET

If the volume is generated about an axis that is off origin then substitute x in $2\pi x$ with $2\pi(a - x)$ if the offset is positive, or $2\pi(a + x)$ if the offset is negative. The previous rule apply if the volume is generated about the y-axis.

AVERAGE VALUE OF A FUNCTION

This will calculate the average value of a continuous function from all points a to b when a is less than b.

GENERAL EQUATION

$$f_{ave} = \frac{1}{b-a} \int_a^b f(x) dx \quad (41)$$

This also means that there exist a real number c in between $[a, b]$ such that

$$f(c) = f_{ave} = \frac{1}{b-a} \int_a^b f(x) dx \quad (42)$$

That is,

$$\int_a^b f(x) dx = f(c)(b-a) \quad (43)$$

EXERCISE

- Find the area enclosed by the given curves from $2 \leq x \leq 4$

$$y = \frac{5}{x}, \quad y = \frac{10}{x^2}$$

Answer.

$$5 \ln(2) - \frac{5}{2}$$

- Find the area enclosed by the given curves, determine the domain by yourself.

$$2x + y^2 = 8, \quad x = y$$

Answer. 18.

- Find the volume of a solid obtained by rotating the region about the x-axis

$$y = \sqrt{x-1}, \quad y = 0, \quad x = 5$$

Answer. 8π

- Find the average value of the function below from $[2, 5]$

$$\frac{t}{\sqrt{5+t^2}}$$

Answer.

$$\sqrt{\frac{10}{3}} - 1$$

CHAPTER 7: TECHNIQUES OF INTERGRATION

In this chapter we will learn the 4 techniques of intergration or 80 % of this entire course.

GENERAL INTEGRATION FORMULA

Know them inside and out

$$\int x^n dx \text{ where } n \neq -1 = \frac{x^{n+1}}{n+1} + C \quad (44)$$

$$\int x^{-1} dx = \ln|x| + C \quad (45)$$

$$\int e^x dx = e^x + C \quad (46)$$

$$\int \sin(x) dx = -\cos(x) + C \quad (47)$$

$$\int \cos(x) dx = \sin(x) + C \quad (48)$$

$$\int \frac{dx}{1+x^2} dx = \tan^{-1}(x) + C \quad (49)$$

$$\int \frac{dx}{\sqrt{1-x^2}} dx = \sin^{-1}(x) + C \quad (50)$$

$$\int \tan(x) dx = \ln|\sec(x)| + C \quad (51)$$

SUBSTITUTION

SUBSTITUTION RULE

If $u = g(x)$ is a differentiable function whose range is an interval I and f is continuous I , then

$$\int f(g(x)) g'(x) dx = \int f(u) du \quad (52)$$

You basically "simplify" the expression using a different variable where there is a more obvious integration rule you can use to evaluate the expression. Once the antiderivative is found you can resubstitute the original function back into the expression.

Note: You may need to do this multiple times for some expressions.

If you have an expression that looks like this

$$\int g(x)e^{f(x)}$$

Where $f(x)$ and $g(x)$ are both functions and $f(x)$ is not a linear function. Then try using the subsitution technique to evalute the expression. Note this may not work. You may need to fix this.

ARCTAN INTEGRATION

GENERAL FORMULA FOR ARCTAN INTEGRATION

When you have an integral in the form:

$$\int \frac{dx}{a^2+x^2}$$

There exist a close-form formula that gives you the arctan answer.

$$\frac{1}{a} \tan^{-1}\left(\frac{x}{a}\right) + C \quad (53)$$

You may need to factor out the coefficient in order to satisfy this form.

ARCSIN INTEGRATION

GENERAL FORMULA FOR ARCSIN INTEGRATION

When you have an integral in the form:

$$\int \frac{dx}{\sqrt{a^2-x^2}}$$

There exist a close-form formula that gives you the arcsin answer.

$$\sin^{-1}\left(\frac{x}{a}\right) + C \quad (54)$$

You may need to factor out the coefficient in order to satisfy this form.

INTEGRATION BY PARTS

GENERAL FORMULA FOR INTEGRATION BY PARTS

Here is the general formula for integration by parts:

$$\int u dv = uv - \int v du \quad (55)$$

$$\int_a^b u dv = uv|_a^b - \int_a^b v du \quad (56)$$

SPECIAL "POWER" CASE

If the integration looks like this:

$$\int f(x)^n g(x) dx \quad (57)$$

Then it takes n iterations of intergration by parts.

During subsitution of an intergration you may have to change the domain of the integration by the new domain of the substituted equation.

$$\int_0^1 x \ln(x+1) dx$$

Let $t = x + 1$, then $dt = dx$, and $x = t - 1$

$$\int_1^2 (t-1) \ln(t) dt$$

TRIG SUBSTITUTION

TRIGONOMETRIC RULES

Know these rules well.

$$\cos^2(x) + \sin^2(x) = 1 \quad (58)$$

$$1 + \tan^2(x) = \sec^2(x) \quad (59)$$

$$\cos^2(x) - \sin^2(x) = \cos(2x) \quad (60)$$

$$\cos^2(x) = \frac{1 + \cos(2x)}{2} \quad (61)$$

$$\sin^2(x) = \frac{1 - \cos(2x)}{2} \quad (62)$$

$$2 \sin(x) \cos(x) = \sin(2x) \quad (63)$$

$$\int \sec^2(x) dx = \tan(x) + C \quad (64)$$

$$\int \sec(x) \tan(x) dx = \sec(x) + C \quad (65)$$

$$\int \sec(x) dx = \ln |\sec(x) + \tan(x)| + C \quad (66)$$

$$\int \csc(x) dx = -\ln |\csc(x) + \cot(x)| + C \quad (67)$$

IF YOU HAVE THIS FORM OF INTEGRATION

$$\int \cos^a(x) \sin^b(x) dx \quad (68)$$

Where at least one of the powers are odd, then seperate the odd powered trig function into an even powered trig function.

Let a be odd and greater than 2, then

$$\int \sin^a(x) \cos^b(x) dx$$

$$\int \sin^{a-1}(x) \cos^b(x) \sin(x) dx$$

Let $u = \cos(x)$, then $du = -\sin(x)dx$

$$-\int (1 - \cos^2(x))^{\frac{a-1}{2}} \cos^b(x) \sin(x) dx$$

$$-\int (1 - u^2)^{\frac{a-1}{2}} u^b du$$

You get the idea

TRIGONOMETRIC SUBSTITUTION WITH ANGLES

Let b be odd and greater than 2, then

$$\int \sin^a(x) \cos^b(x) dx$$

$$\int \sin^a(x) \cos^{b-1}(x) \cos(x) dx$$

Let $u = \sin(x)$, then $du = \cos(x)dx$

$$\int \sin^a(x)(1 - \sin^2(x))^{\frac{b-1}{2}} \cos(x) dx$$

$$\int u^a(1 - u^2)^{\frac{b-1}{2}} du$$

You get the idea

BOTH ARE EVEN POWERS

If both powers are the same and are even then you can use this: $2 \sin(x) \cos(x) = \sin(2x)$ to replace the multitrig function into a single trig function.

Let a be an even number

$$\int \sin^a(x) \cos^a(x) dx$$

$$\int \left(\frac{\sin(2x)}{2}\right)^a dx$$

$$\frac{1}{2^a} \int \sin^a(2x) dx$$

You get the idea

FOR TAN AND SEC INTERGRATIONS

For tan and sec integrations there are two rules you can follow.

$$\int \tan^{odd}(x) \sec^n(x) dx$$

↓

$$\int \tan^{odd-1}(x) \sec^{n-1}(x) \tan(x) \sec(x) dx$$

Let $u = \sec(x)$

$$\int (\sec^2(x) - 1)^{\frac{odd-1}{2}} \sec^{n-1}(x) \tan(x) \sec(x) dx$$

$$\int \tan^n(x) \sec^{even}(x) dx$$

↓

$$\int \tan^n(x) \sec^{even-2}(x) \sec^2(x) dx$$

Let $u = \tan(x)$

$$\int \tan^n(x)(\tan^2(x) + 1)^{\frac{even-2}{2}} \sec^2(x) dx$$

TAN SUBSTITUTION WITH ANGLE

For the general intergration shown below:

$$\int \frac{dx}{f(a^2 + b^2 x^2)^\alpha} \quad (69)$$

The standard subsitution is we let $x = \frac{a}{b} \tan(\theta)$. Therefore we get this general formula.

$$x = \frac{a}{b} \tan(\theta)$$

$$dx = \frac{a}{b} \sec^2 \theta d\theta$$

In the denominator

$$a^2 + b^2 \left(\frac{a}{b} \tan(x)\right)^2 = a^2 + b^2 \frac{a^2}{b^2} \tan^2 \theta$$

$$= a^2 + a^2 \tan^2(x)$$


$$= a^2(1 + \tan^2(x))$$

$$= (a^2 \sec^2 \theta)^\alpha$$

So we get this subsitution

$$\frac{a}{a^2 b} \int \frac{\sec^2 \theta d\theta}{(\sec^2 \theta)^\alpha}$$

From here you can manipulate the integral so it easy to work with



Once you intergrated the expression you have to convert the domain from radian to real. Use the figure above to subsitute the trigonometric function with the real expressions.

SIN SUBSTITUTION WITH ANGLE

For the general intergration shown below:

$$\int \frac{dx}{f(a^2 - b^2x^2)^\alpha} \quad (70)$$

The standard subsitution is we let $x = \frac{a}{b} \sin(\theta)$. Therefore we get this general formula.

$$x = \frac{a}{b} \sin(\theta)$$

$$dx = \frac{a}{b} \cos \theta d\theta$$


In the demoninator

$$\begin{aligned} a^2 - b^2 \left(\frac{a}{b} \sin \theta \right)^2 &= a^2 - b^2 \frac{a^2}{b^2} \sin^2 \theta \\ &= a^2 - a^2 \sin^2 \theta \\ &= a^2(1 - \sin^2 \theta) \\ &= (a^2 \cos^2 \theta)^\alpha \end{aligned}$$

So we get this subsitution

$$\frac{a}{a^{2\alpha}b} \int \frac{\cos \theta d\theta}{(\cos^2 \theta)^\alpha}$$

From here you can manipulate the integral so it easy to work with



Once you intergrated the expression you have to convert the domain from radian to real. Use the figure above to subsitute the trigonometric function with the real expressions.

SEC SUBSTITUTION WITH ANGLE

For the general intergration shown below:

$$\int \frac{dx}{f(b^2x^2 - a^2)^\alpha} \quad (71)$$

The standard subsitution is we let $x = \frac{a}{b} \sec(\theta)$. Therefore we get this general formula.

$$x = \frac{a}{b} \sec(\theta)$$

$$dx = \frac{a}{b} \tan \theta \ sec \theta d\theta$$


In the demoninator

$$\begin{aligned} b^2 \left(\frac{a}{b} \sec \theta \right)^2 - a^2 &= b^2 \frac{a^2}{b^2} \sec^2 \theta - a^2 \\ &= a^2 \sec^2 \theta - a^2 \\ &= a^2(\sec^2 \theta - 1) \\ &= (a^2 \tan^2 \theta)^\alpha \end{aligned}$$

So we get this subsitution

$$\frac{a}{a^{2\alpha}b} \int \frac{\tan \theta \ sec \theta d\theta}{(\tan^2 \theta)^\alpha}$$

From here you can manipulate the integral so it easy to work with



Once you intergrated the expression you have to convert the domain from radian to real. Use the figure above to subsitute the trigonometric function with the real expressions.

PARTIAL FRACTION

PARTIAL FRACTION

If you have an integral that look a fraction of polynomials then you have to use partial fractions to evaluate it. Here is an generic example:

$$\int f(x) = \int \frac{P(x)}{Q(x)} \quad (72)$$

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$$

$$Q(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$$

$$a_n \neq 0$$

WORKING WITH PARTIAL FRACTIONS

This will only work if the degree (that is the highest power of the polynomial) of the numerator is less than the degree of the denominator.

If $\text{Deg}(P(x)) \geq \text{Deg}(Q(x))$, then you would have to use long division to make the degree of the numerator be less than the denominator.

LONG DIVISION

LONG DIVISION

If you need to use long division in order to integrate using partial fraction then the overall general equation will look like this.

$$Q(x) \overline{P(x)} = S(x) \quad (73)$$

$$\int f(x) = \int \frac{P(x)}{Q(x)} = \int S(x) + \int \frac{R(x)}{Q(x)} \quad (74)$$

Where:

$S(x)$: Is the quotient

$R(x)$: Is the remainder

Generally $\int S(x)$ is pretty easy to evaluate as you are integrating a polynomial. So the power rule applies.

Once the expression is in the proper format we can continue with our evaluation.

Q(X) IS A PRODUCT OF DISTINCT LINEAR FACTORS

THE "SIMPLE" CASE

If you can factorize $Q(x)$ to

$$Q(x) = (a_1 x + b_1)(a_2 x + b_2) \dots (a_k x + b_k)$$

then you can write the partial fraction like this:

$$\int \frac{R(x)}{Q(x)} = \int \frac{A_1}{(a_1 x + b_1)} + \int \frac{A_2}{(a_2 x + b_2)} + \dots + \int \frac{A_k}{(a_k x + b_k)} \quad (75)$$

To solve for the coefficients one must isolate them by multiplying its denominator with all other terms and $R(x)$, then set x such that the denominator is zero cancelling all other coefficients.

GENERAL EQUATION

$$A_1 = \frac{R(x)(a_1 x + b_1)}{(a_1 x + b_1)(a_2 x + b_2) \dots (a_k x + b_k)}, \quad x = \frac{-b_1}{a_1}$$

$$A_2 = \frac{R(x)(a_2 x + b_2)}{(a_1 x + b_1)(a_2 x + b_2) \dots (a_k x + b_k)}, \quad x = \frac{-b_2}{a_2}$$

$$A_k = \frac{R(x)(a_k x + b_k)}{(a_1 x + b_1)(a_2 x + b_2) \dots (a_k x + b_k)}, \quad x = \frac{-b_k}{a_k}$$

Once the coefficients are found they can be substituted into the expression and be evaluate. In this case its pretty easy as it's very similar to

$$\int \frac{dx}{f(x)} = \ln f(x) + C$$

Therefore

$$\int \frac{A_1}{(a_1 x + b_1)} = \frac{A_1}{a_1} \ln(a_1 x + b_1) + C$$

$$\int \frac{A_2}{(a_2 x + b_2)} = \frac{A_2}{a_2} \ln(a_2 x + b_2) + C$$

⋮

$$\int \frac{A_k}{(a_k x + b_k)} = \frac{A_k}{a_k} \ln(a_k x + b_k) + C$$

Therefore

$$\int \frac{R(x)}{Q(x)} = \frac{A_1}{a_1} \ln(a_1 x + b_1) + \frac{A_2}{a_2} \ln(a_2 x + b_2) + \dots + \frac{A_k}{a_k} \ln(a_k x + b_k) + C$$

Q(x) IS A PRODUCT OF LINEAR FACTORS, SOME OF WHICH ARE REPEATED

REPEATED FACTORS

If you have a factor that is $(a_i x + b_i)^r$ Where: $r > 1$ then the partial fraction decomposition would look like this:

$$\frac{A_1}{a_i x + b_i} + \frac{A_2}{(a_i x + b_i)^2} + \dots + \frac{A_r}{(a_i x + b_i)^r} \quad (76)$$

There would be r terms after the decomposition with each term having a factor with a unique power $[1-r]$.

Depending on the expression you may have to use techniques from both case 1 and 2 in order to integrate the expression.

EXAMPLE

$$\begin{aligned} & \int \frac{x^4 - 2x^2 + 4x + 1}{x^3 - x^2 - x + 1} dx \\ &= \int x + 1 + \frac{4x}{x^3 - x^2 - x + 1} dx \\ &= \int x + 1 + \frac{4x}{(x-1)^2(x+1)} dx \\ &= \frac{4x}{(x-1)^2(x+1)} = \frac{A}{x-1} + \frac{B}{(x-1)^2} + \frac{C}{x+1} \end{aligned}$$

Solving for the coefficients using techniques from case 1 will help with most cases except for A as when $x = 1$ the denominator is zero. Case 1 will help you figure out that $B = 2$ and $C = -1$.

Let $x = 2$, then

$$\begin{aligned} \frac{4(2)}{((2)-1)^2(2+1)} &= \frac{A}{2-1} + \frac{2}{((2)-1)^2} - \frac{1}{2+1} \\ \frac{8}{3} &= A + 2 - \frac{1}{3} \\ A &= 1 \end{aligned}$$

After the coefficients are found we substitute them into the expression and evaluate.

$$\begin{aligned} & \int x + 1 + \frac{1}{x-1} + \frac{2}{(x-1)^2} - \frac{1}{x+1} dx \\ &= \frac{x^2}{2} + x + \ln|x-1| - \frac{2}{x-1} - \ln|x+1| + C \end{aligned}$$

Q(x) CONTAINS IRREDUCIBLE QUADRATIC FACTORS, NONE OF WHICH IS REPEATED

QUADRATIC FACTOR

If $Q(x)$ has the factor $ax^2 + bx + c$, where $b^2 - 4ac < 0$, then, in addition to the partial fractions in equations 58 and 59, the decomposition will have a term of the form:

$$\frac{Ax + B}{ax^2 + bx + c} \quad (77)$$

The term above could be integrated by completing the square (if necessary) and using the formula.

$$ax^2 + bx + c = a(x - h)^2 + k$$

$$h = \frac{b}{2a}$$

$$k = c - \frac{b^2}{4a}$$

$$u = x - h$$

$$\begin{aligned} d &= \frac{k}{a} \\ &= \frac{1}{a(u^2 + d)} \end{aligned}$$

$$\int \frac{dx}{a(u^2 + d)} = \frac{1}{a\sqrt{d}} \tan^{-1}\left(\frac{x}{a\sqrt{d}}\right) + C \quad (78)$$

EXAMPLE

$$\begin{aligned} & \text{Evaluate } \int \frac{2x^2 - x - 4}{x^3 + 4x} dx \\ & \int \frac{2x^2 - x + 4}{x(x^2 + 4)} dx = \int \frac{A}{x} + \frac{Bx + C}{x^2 + 4} \\ & \frac{(2x^2 - x + 4)x(x^2 + 4)}{x(x^2 + 4)} = A(x^2 + 4) + (Bx + C)x \\ & \frac{(2x^2 - x + 4)x(x^2 + 4)}{x(x^2 + 4)} = (A + B)x^2 + Cx + 4A \end{aligned}$$

We get the coefficients: $A + B = 2$, $C = -1$, $4A = 4$

$$A = 1, B = 1, C = -1$$

$$\begin{aligned} & \int \frac{2x^2 - x + 4}{x(x^2 + 4)} dx = \int \frac{1}{x} dx + \int \frac{x-1}{x^2+4} dx \\ &= \int \frac{1}{x} dx + \int \frac{x}{x^2+4} dx - \int \frac{1}{x^2+4} dx \\ &= \ln|x| + \frac{1}{2} \ln|x^2+4| - \frac{1}{2} \tan^{-1}\frac{x}{2} + C \end{aligned}$$

GENERAL FORMULA

THE GENERAL IDEA

1. Convert the fraction such that the polynomial degree of the denominator is greater than the polynomial degree of the numerator (if necessary).
2. Factorize both polynomials.
3. For each factor of the denominator the partial fraction decomposition is a polynomial whose degree is one less than the denominator.

$$\frac{D_n x^{n-1} + D_{n-1} x^{n-2} \dots + D_2 x + D_1}{(c_n x^n + c_{n-1} x^{n-1} + \dots + c_1 x + c_0)} \quad (79)$$

Where c and D are the coefficients

4. Rearrange such that each coefficient is with a variable.
5. Use linear algebra to solve for the coefficients (look at the previous example).

IMPROPER INTEGRAL

When you have infinity as part of the range or the function is discontinuous over the range.

CONVERGENCE VS DIVERGENCE

An integral **converges** if the value of the integral is a real number.

It **diverges** if the value of the integral is $\pm\infty$

TYPE 1

TYPE 1

$$\int_a^{\infty} f(x) dx = \lim_{r \rightarrow \infty} \int_a^r f(x) dx \quad (80)$$

$$\int_{-\infty}^a f(x) dx = \lim_{r \rightarrow -\infty} \int_r^a f(x) dx \quad (81)$$

Where $a \in \mathbb{R}$ / a constant

$$\int_{-\infty}^{\infty} f(x) dx = \int_{-\infty}^0 f(x) dx + \int_0^{\infty} f(x) dx \quad (82)$$

Use equation 77 and 78 to solve equation 79 were $a = 0$.

EXAMPLE

Evaluate this expression:

$$\begin{aligned} & \int_1^{\infty} \frac{dx}{x} \\ & \lim_{r \rightarrow \infty} \int_1^r \frac{dx}{x} \\ &= \ln|x| \Big|_1^r \\ &= \ln|r| - \ln|1| \\ &= \ln|r| - 0 \\ &= \ln|r| = \infty \end{aligned}$$

TYPE 2

TYPE 2

If the function is discontinuous over the range of domain, then you have to split the integration between the point of discontinuity.

$$\int_a^b f(x) dx = \int_a^{c^-} f(x) dx + \int_{c^+}^b f(x) dx \quad (83)$$

Where c is the domain where the range of the function is discontinuous. i.e. $f(c) = \text{DNE/UNKNOWN}$.

HAVING + AND - ON THE POINT OF DISCONTINUITY

It's important to have the appropriate + and - symbols for c as you are evaluating the integrals when they approach c from the right or left side respectively. The direction of approach matter as it could affect the sign of the value (positive or negative).

EXAMPLE

Evaluate this expression:


$$\int_0^3 \frac{dx}{x-1}$$

Because this function is discontinuous when $x = 1$ we have to split the evaluation into two integral.

$$\int_0^{1^-} \frac{dx}{x-1}$$

and

$$\int_{1^+}^3 \frac{dx}{x-1}$$



$$\begin{aligned}
 & \int_0^{1^-} \frac{dx}{x-1} \\
 &= \lim_{r \rightarrow 1^-} \frac{dx}{x-1} \\
 &= \lim_{r \rightarrow 1^-} \ln|r-1| \Big|_0^{r^-} \\
 &= \lim_{r \rightarrow 1^-} \ln|r-1| - \ln|0-1| \\
 &= \lim_{r \rightarrow 1^-} \ln|r-1| - 0 \\
 &= \lim_{r \rightarrow 1^-} \ln|r-1| = -\infty
 \end{aligned}$$

This integration diverges therefore the entire function diverge.

CONDITION OF DIVERGENCE

If one of the integrals leads to a divergence then the entire function is divergent.

NONELEMENTARY INTEGRAL

PROVING DIVERGENCE OR CONVERGENCE OF NONELEMENTARY INTEGRALS

If you have an integral that is nonelementary. i.e. having a summation of trig functions, polynomials, and constant. Then the answer to the expression is impossible to find based on current understanding. One way to solve the expression (somewhat) is by using proofs.

BIG-O

Used in CS and Math to determine the behaviour of a function based on input size (More of a CS definition). The basic idea is that you find a single term expression where the order of expression is as big or bigger than the nonelementary function.

BIG-O

Big-O: Let f be the nonelementary function and g a single term function or something that is easily integratable. Then you find a function g such that.

$$f(x) \leq g(x) \quad \forall x \geq x_0 \in \mathbb{R}$$

or

$$\lim_{x \rightarrow 0} \left| \frac{f(x)}{g(x)} \right| < \infty$$

You can say that Big-O will give the "upper bound" for the growth rate of function f .

There is a "lower bound" equivalent called Big-Omega but in a lot of cases this function can be a constant like 0.

Once we found our boundaries we can integrate them to find if the nonelementary function converges or diverges.

EXAMPLE

Evaluate this expression:

$$\int_1^{\infty} \frac{dx}{x^2 + x^{20}}$$

Because this expression is a nonelementary function we have to limit its range using simpler, elementary functions.

So let 0 be the lower bound as this function will always be greater than 0.

$$0 \leq \int_1^{\infty} \frac{dx}{x^2 + x^{20}}$$

Now we have to find the upper bound.

Let x^2 be our upper bound

$$\frac{1}{x^2 + x^{20}} \leq \frac{1}{x^2} \quad \forall x \geq 1$$

We now have bounded the nonelementary function. Therefore the value of the function must be within values of the two simpler functions.

$$0 \leq \int_1^{\infty} \frac{dx}{x^2 + x^{20}} \leq \int_1^{\infty} \frac{dx}{x^2}$$

Evaluating the upper bound gives this value

$$\lim_{r \rightarrow \infty} \left. \frac{-1}{x} \right|_1^r = \left(\frac{-1}{r} - \frac{-1}{1} \right) = 0 - (-1) = 1$$

Therefore

$$0 \leq \int_1^\infty \frac{dx}{x^2 + x^{20}} \leq 1$$

Therefore

$$\int_1^\infty \frac{dx}{x^2 + x^{20}}$$

Converges to some value between 0 and 1.

CONDITION FOR CONVERGENCE OF NONELEMENTARY FUNCTIONS

An Nonelementary function converges onto a value if and only if the upper **and** lower bounds $\neq \pm\infty$. Otherwise, the function diverges.

Answer.

$$\frac{3\pi}{16}$$

6. Evaluate the integral using the indicated trigonometric substitution

$$\int \frac{x^3}{\sqrt{x^2 + 64}} dx, \quad x = 8 \tan(\theta)$$

Answer.

$$\frac{1}{3}(x^2 - 128)\sqrt{x^2 + 64} + C$$

7. Evaluate the integral

$$\int \frac{x^3}{x-1} dx$$

Answer.

$$\frac{x^3}{3} + \frac{x^2}{2} + x + \ln|x-1| + C$$

8. Evaluate the integral

$$\int \frac{50x+6}{(7x+1)(x-1)} dx$$

Answer.

$$\frac{\ln|7x+1|}{7} + 7 \ln|x-1| + C$$

9. Determine whether the integral is convergent or divergent

$$\int_8^\infty \frac{1}{x^2+x} dx$$

Answer.

$$\ln\left(\frac{9}{8}\right)$$

10. Determine whether the integral is convergent or divergent

$$\int_0^\infty 43 \frac{\arctan(x)}{2+e^x} dx$$

Answer. It converges.

11. Determine whether the integral is convergent or divergent

$$\int_{-\infty}^0 \frac{1}{5-7x} dx$$

Answer. It diverges

12. Determine whether the integral is convergent or divergent

$$\int_{-\infty}^\infty 9xe^{-x^2} dx$$

Answer. 0.

CHAPTER 8: FURTHER APPLICATIONS OF INTEGRATION

ARC LENGTH

Calculate the length of a function given a range from a to b.

EQUATION

$$L = \int_a^b \sqrt{1 + (f'(x))^2} dx \quad (84)$$

PROOF

To prove this equation, let assume that you have an arbitrary function f that is continuous. The distance between two points on the graph can be approximated by the summation of very small slopes.

Let ds be the distance between two points on the graph (rise over run).

$$ds = \sqrt{\Delta x^2 + \Delta y^2}$$

$$ds = \Delta x \sqrt{1 + \frac{\Delta y^2}{\Delta x^2}}$$

$$ds = \Delta x \sqrt{1 + \left(\frac{\Delta y}{\Delta x}\right)^2}$$

$\frac{\Delta y}{\Delta x}$ is the equivalent to $\frac{dy}{dx} = f'(x)$

Δx is equivalent to dx

∴

$$ds = dx \sqrt{1 + (f'(x))^2}$$

To get the total distance we add them together

$$L = \int_a^b \sqrt{1 + (f'(x))^2} dx$$

EXAMPLE

Find the arc length from 0 to $\frac{\pi}{3}$ of this function:

$$y = \ln(\cos(x))$$

$$y = \ln |\cos(x)|$$

$$y' = \frac{-\sin(x)}{\cos(x)} dx$$

$$L = \int_0^{\frac{\pi}{3}} \sqrt{1 + \left(\frac{-\sin(x)}{\cos(x)}\right)^2} dx$$

$$L = \int_0^{\frac{\pi}{3}} \sqrt{1 + \tan^2(x)} dx$$

$$L = \int_0^{\frac{\pi}{3}} \sqrt{\sec^2(x)} dx$$

$$L = \int_0^{\frac{\pi}{3}} \sec(x) dx$$

$$L = \ln |\sec(x) + \tan(x)| \Big|_0^{\frac{\pi}{3}}$$

$$L = \ln \left| \sec\left(\frac{\pi}{3}\right) + \tan\left(\frac{\pi}{3}\right) \right| - \ln \left| \sec(0) + \tan(0) \right|$$

$$L = \ln |2 + \sqrt{3}|$$

SURFACE AREA

SURFACE AREA

$$SA = \int_a^b 2\pi f(x) ds \quad (85)$$

Where ds is the arc length segment.

$$ds = \sqrt{1 + (f'(x))^2} dx$$

$$SA = 2\pi \int_a^b f(x) \sqrt{1 + (f'(x))^2} dx \quad (86)$$

Basically you are adding infinitely many cylinder of size ds together. As the surface area of a cylinder excluding the top and bottom is $2\pi rh$, where h is ds .

IF YOU ARE ROTATING AROUND THE Y-AXIS

If you are rotating around the y axis then you would use this equation

$$L = \int_a^b 2\pi x(y) \sqrt{1 + (x'(y))^2} dy \quad (87)$$

EXAMPLE

Find the surface area of this function around the y-axis from 0 to 1:

$$y = x^2$$

$$x = \sqrt{y}$$

$$dx = \frac{1}{2\sqrt{y}} dy$$

$$SA = 2\pi \int_0^1 \sqrt{y} \sqrt{1 + \left(\frac{1}{2\sqrt{y}}\right)^2} dy$$

$$SA = 2\pi \int_0^1 \sqrt{y} \sqrt{1 + \left(\frac{1}{4y}\right)} dy$$

$$SA = 2\pi \int_0^1 \sqrt{y + \left(\frac{1}{4y}\right)} dy$$

$$SA = 2\pi \int_0^1 \sqrt{y + \left(\frac{1}{4}\right)} dy$$

EXERCISE

- Find the exact length of the curve

$$36y^2 = (x^2 - 4)^3, 4 \leq x \leq 9$$

Answer.

$$\frac{1}{2} \left(225 - \frac{40}{3} \right)$$

- Find the exact length of the curve

$$y = \frac{x^3}{3} + \frac{1}{4x}, 1 \leq x \leq 2$$

Answer.

$$\frac{7}{3} + \frac{1}{8}$$

- Find the exact area of the surface obtained by rotating the curve about the x-axis

$$y = x^3, 0 \leq x \leq 4$$

Answer.

$$\frac{\pi}{27} \left(2305^{\frac{3}{2}} - 1 \right)$$

- Find the exact area of the surface obtained by rotating the curve about the x-axis

$$y = \frac{x^3}{6} + \frac{1}{2x}, \frac{1}{2} \leq x \leq 1$$

Answer.

$$\frac{263\pi}{256}$$

- Find the exact area of the surface obtained by rotating the curve about the y-axis

$$y = \frac{1}{4}x^2 - \frac{1}{2} \ln(x), 2 \leq x \leq 4$$

Answer.

$$\frac{62\pi}{3}$$

CHAPTER 9: DIFFERENTIAL EQUATION

Differential equations may be the most important applications for calculus. Scientists from all fields use differential equation to create models to explain the phenomenon that they are studying. Although it is often impossible to find an explicit formula for the solution of a differential equation, we will see that graphical and numerical approaches provide the information we need.

INITIAL VALUE PROBLEM

Suppose that you have a differential equation y' with a specific initial condition or the "starting" value. Find y with a known constant.

INITIAL VALUE PROBLEM (IVP)

Generally the problem is in this form of expression:

$$\begin{cases} y'(t) = f(t, y(t)) \\ y(t_0) = y_0 \end{cases}$$

What we are trying to find is a function y that satisfy both conditions above. Later we will talk about seperable equation and how it will help solve this problem.

POPULATION DYNAMICS

For example let's use differential equations to explain population dynamics.

THE LAW OF NATURAL GROWTH

This is one model where we assume that the population grows at a rate proportional to the size of the population.

Is this a reasonable assumption? Suppose we have a population of $P = 1,000,000,000$ Skavens and at a certain time it is growing at a rate of $P' = 1,000,000$ Skavens per hour. Now let's take another billion Skavens. Each half of the combined population was previously growing at a rate of one million Skavens per hour. We would expect the total population of two billion to increase at a rate of 2 million Skavens per

hour initially (provided there's enough food, land, and lack of backstabbing). So if we double the size, we double the growth rate. It seems reasonable that the growth rate should be proportional to the size.

Notation Meaning

t	time (the independent variable)
P	# of individuals in the population (the dependent variable)

RATE OF GROWTH OF A POPULATION

$$\frac{dP}{dt} = kP \quad (88)$$

Where:

k is the proportionality constant or growth-rate

This equation is a differential equation because it contains an unknown function P and its derivative $\frac{dP}{dt}$.

$$P'(t) = C(ke^{kt}) = k(Ce^{kt}) = kP(t) \quad (89)$$

SEPARABLE EQUATION

How did we get the population dynamics equation from before? To derive this equation we have to use the property of seperable equation.

SEPARABLE EQUATION

If you have an expression like below.

$$\frac{dy}{dx} = f(x)g(y)$$

Then using the properties of Algebra allows us to do this.

$$h(y) = \frac{1}{f(y)}$$

$$\frac{dy}{dx} = \frac{g(x)}{h(y)}$$

$$h(y) dy = g(x) dx$$

$$\int h(y) dy = \int g(x) dx \quad (90)$$

EXAMPLE

Evaluate this expression:

$$\frac{dy}{dt} = ky$$

Where k is a constant

Solution

$$\begin{aligned}\frac{dy}{dt} &= ky \\ \frac{dy}{y} &= kdt \\ \int \frac{dy}{y} &= \int kdt \\ \ln|y| &= kt + C \\ |y| &= e^{kt} \cdot \pm e^C \\ |y| &= Ce^{kt}\end{aligned}$$

Solve for C : You would set y to the value of y_0 or the initial condition/seed value, and t to t_0 .

LOGISTIC MODEL

The logistic model is a better representation of explaining the dynamics of a population. In our previous model we never considered an upper limit on the size of a population, it just continues to grow. The logistic model considers a carrying capacity M such that the relative growth rate will be negative if P is greater than M and positive if P is less than M .

Meaning

P	population function
k	relative growth rate
M	carrying capacity

LOGISTIC MODEL EQUATION

$$\frac{dP}{dt} = kP\left(1 - \frac{P}{M}\right) \quad (91)$$

Because the equation above is separable we get this

$$\frac{dP}{P(1 - \frac{P}{M})} = k dt$$

This left side expression can be evaluated using partial fractions to get this

$$\int \frac{M dP}{P(M - P)} = \int \frac{dP}{P} + \int \frac{dP}{M - P}$$

$$\int \frac{dP}{P} + \int \frac{dP}{M - P} = \int k dt$$

$$\ln\left|\frac{P}{M - P}\right| = kt + C$$

$$\left|\frac{P}{M - P}\right| = \pm e^C \cdot e^{kt} = Ce^{kt}$$

$$\left|\frac{M - P}{P}\right| = \pm e^{-C} \cdot e^{-kt} = Ce^{-kt}$$

$$\frac{M}{P} - 1 = Ce^{-kt} \Rightarrow \frac{P}{M} = \frac{1}{1 + Ce^{-kt}}$$

$$P = \frac{M}{1 + Ce^{-kt}}$$

$$\frac{M - P_0}{P_0} = Ce^0 = C$$

LOGISTIC EQUATION

$$P(t) = \frac{M}{1 + Ce^{-kt}} \text{ Where } C = \frac{M - P_0}{P_0} \quad (92)$$

$$\lim_{t \rightarrow \infty} P(t) = M$$

M is the terminal value

NEWTON COOLING EQUATION

Lets say you want to model the transfer of heat between an object and its environment. We can use differential equation to model this behaviour.

NEWTON COOLING EQUATION

Notation	Meaning
t	Time t
$T(t)$	Temperature at time t
T_E	Temperature of the environment
T_0	Temperature at $t = 0$
$\frac{dT}{dt}$	The rate of change of the difference in temperature T

$$\frac{dT}{dt} = -k(T_0 - T_E) \quad k > 0 \quad (93)$$

Using techniques discusses previously we get this

$$T(t) = T_E + (T_0 - T_E)e^{-kt} \quad (94)$$

EXAMPLE

Suppose that you have a bowl of hot soup at 60 degrees C, a refrigerator at 4 degrees C, and $k = 0.1$. How long will it take for the soup in the refrigerator to drop down to 50 degrees C?

Solution

$$\begin{aligned} T(t) &= 4 + 56e^{-0.1t} \\ 50 &= 4 + 56e^{-0.1t} \\ 46 &= 56e^{-0.1t} \\ \frac{46}{56} &= e^{-0.1t} \\ \ln\left(\frac{46}{56}\right) &= -0.1t \\ -10\ln\left(\frac{46}{56}\right) &= t \end{aligned}$$

FINAL 2018 EXAMPLE

A pot of soup if initial temperature of 80 degrees C is placed into a refrigerator which is set at 5 degrees C, at $T = 75$ degrees C, the temperature of the pot drop with a rate of change at 2 degrees C per second. Find the time when the temp is at 30 degrees C.

T_0	80 degrees C
T_E	5 degrees C
T	75 degrees C
$\frac{dT}{dt}$	-2 degrees C / s

Solution

We first have to find k .

$$-2 = -k(80 - 5)$$

$$-2 = -k(75)$$

$$k = \frac{2}{75}$$

Once we have found k we can now solve the problem.

$$\begin{aligned} 30 &= 5 + (80 - 5)e^{-\frac{2}{75}(t)} \\ 25 &= 75e^{-\frac{2}{75}(t)} \\ \frac{25}{75} &= e^{-\frac{2}{75}(t)} \\ \frac{1}{3} &= e^{-\frac{2}{75}(t)} \\ \ln\left(\frac{1}{3}\right) &= -\frac{2}{75}(t) \\ -\frac{75}{2} \ln\left(\frac{1}{3}\right) &= t \end{aligned}$$

LINEAR EQUATIONS

FO LINEAR DIFF EQ

A first-order linear differential equation is one that can be put into the form:

$$\begin{cases} y' + p(x)y = r(x) \\ y(0) = y_0 \end{cases} \quad (95)$$

$$\frac{dy}{dx} + p(x)y = Q(x) \quad (96)$$

Its linear as y' and y are both linear have a power of 1.

In order to evaluate this function we need a special function called an **integrating factor** denoted as $\mu(t)$

INTEGRATING FACTOR $\mu(t)$

WHAT IS $\mu(t)$

$\mu(t)$ is a special function that fulfills the condition as an intergrating factor:

$$\mu(t)p(t) = \mu'(t) \quad (97)$$

Multiplying the entire linear differential equation with $\mu(t)$ lead to this expression:

$$\mu(t)y' + \mu'(t)y = \mu(t)r(t)$$

This is the result from the product rule of differentiation.

$$(y(t)\mu(t))' = y'(t)\mu(t) + y(t)\mu'(t)$$

So we can replace the left side with $(y(t)\mu(t))'$. Our expression now is:

$$(y(t)\mu(t))' = \mu(t)r(t)$$

Intergrating both sides:

$$\begin{aligned} \int (y(t)\mu(t))' dt &= \int \mu(t)r(t) dt \\ y(t)\mu(t) + C &= \int \mu(t)r(t) dt \\ y(t)\mu(t) &= \int \mu(t)r(t) dt - C \\ y(t) &= \frac{\int \mu(t)r(t) dt + C}{\mu(t)} \end{aligned}$$

THE UNIVERSAL INTEGRATING FACTOR

If we use this function $\mu(t) = e^{\int p(t)} dt + C$ as our integrating factor we get this expression once we substituted $\mu(t)$ into the equation from before.

$$y(t) = \frac{\int e^{\int p(t)} r(t) dt + C}{e^{\int p(t)}}$$

THE UNIVERSAL EQUATION

The universal equation is shown below:

$$y(t) = \frac{\int e^{\int p(t)} r(t) dt}{e^{\int p(t)}} + \frac{C}{e^{\int p(t)}} \quad (98)$$

EXAMPLE

Evaluate this expression $\begin{cases} y' + \tan(t) y = \sec(t) \\ y(0) = 1 \end{cases}$

Solution

Let: $p(t) = \tan(t)$ $r(t) = \sec(t)$

$$\mu(t) = e^{\int \tan(t)} = e^{\ln |\sec(t)|} = \sec(t)$$

$$e^{\ln |\sec(t)|} y' + e^{\ln |\sec(t)|} \tan(t) = e^{\ln |\sec(t)|} \sec(t)$$

$$\int (e^{\ln |\sec(t)|} y)' dt = \int e^{\ln |\sec(t)|} \sec(t) dt$$

Using the log rule we can get this:

$$\sec(t)y = \int \sec^2(t) dt + C$$

$$y = \frac{\tan(t) + C}{\sec(t)}$$

$$y = \sin(t) + C \cos(t)$$

Evaluate using the initial condition

$$y_0 = 1 = \sin(0) + C \cos(0)$$

$$1 = 0 + C \Rightarrow C = 1$$

$$y(t) = \sin(t) + \cos(t)$$

MIXING PROBLEM

Suppose you have a water tank that contains 1000 liters of pure water at time $t = 0$. A stream of 30 L/min of salty water of a concentration of 0.05 kg/L flows into the tank. The content is kept thoroughly mixed. Simultaneously a stream of 50 l/min flows out of the tank. Find the total amount of salt at time $t = 10$.

To solve this we need a function to measure the rate of change of the salt with respect to time. We also need a function to measure the salt concentration in the tank at time t .

Let:

$c(t) =$ Salt concentration in the tank at time t

$\frac{dC}{dt} =$ incoming salt - outgoing salt

$C(t) = c(t) \cdot 1000$ The total amount of salt at time t

$$\frac{dC}{dt} = (0.05 \text{ kg/L} \times 50 \text{ L/min}) - (50 \text{ L/min} c(t))$$

$$1000 \frac{dc}{dt} = 2.5 - 50 c(t)$$

$$\frac{dc}{dt} = \frac{2.5}{1000} - \frac{50 c(t)}{1000}$$

Using the property of separable equation get us:

$$\frac{dc}{c(t)} = \frac{2.5}{1000} - \frac{50}{1000} dt$$

$$\frac{dc}{c(t)} = 0.0025 - 0.05 dt$$

You get the idea

CHAPTER 10: PARAMETRIC EQUATION AND POLAR COORDINATES

So far we have described curves by giving y as a function of x , i.e. $y = f(x)$ or x as a function of y , i.e. $x = g(y)$, or by giving a relation between x and y that defines y implicitly as a function of x like $y^2 + x^2 = 1$.

Some curves, such as the cycloid, are best handled when both are given in terms of a third variable t called a parameter, i.e. $x = f(t)$, $y = g(t)$.

PARAMETRIC CURVE

NOTATION



$$C = (x(t), y(t)) \quad (99)$$

$x = x(t)$, $y = y(t)$ are functions of parametric t

POSITION AND SPEED OF A PARTICLE IN ORBIT

Lets say that the x position of the particle can be defined as $x = \cos(2t)$ and the y position can be defined as $y = \sin(2t)$. Then we have a parameter curve that is defined as $C = (\cos(2t), \sin(2t))$.

This is what the parameter curve would look like from $0 \leq t \leq 2\pi$:



It

creates a circle.

VELOCITY VECTOR AND SPEED

THE VELOCITY VECTOR EQUATION

The velocity of the particle can be figured out by finding the rate of change on both the x and y axes. This will give us a vector that is the tangential velocity of the particle at a given time t .

$$\vec{V} = < x'(t), y'(t) > \quad (100)$$

To get the speed we take the magnitude of the velocity vector

$$V = \|\vec{V}\| = \sqrt{(x'(t))^2 + (y'(t))^2} \quad (101)$$

For the previous example, the equation for the velocity vector and speed is:

$$\begin{aligned} \vec{V}(t) &= < (\cos(2t))', (\sin(2t))' > \\ \vec{V}(t) &= < -2 \sin(2t), 2 \cos(2t) > \\ V(t) &= \sqrt{(-2 \sin(2t))^2 + (2 \cos(2t))^2} \\ V(t) &= \sqrt{4 \sin^2(2t) + 4 \cos^2(2t)} \\ V(t) &= \sqrt{4 (\sin^2(2t) + \cos^2(2t))} \\ V(t) &= \sqrt{4} \\ V(t) &= 2 \end{aligned}$$

TANGENT LINE

TANGENT LINE EQUATION

$$m = \frac{dy}{dx} = \frac{dy/dt}{dx/dt} \quad (102)$$

Tangent line equation

$$y - y_0 = m(x - x_0) \quad (103)$$

EXAMPLE

Find the tangent of this parametric equation $C = (2 \cos(3t), 2 \sin(3t))$ when $t = \frac{\pi}{3}$ and $t = \frac{\pi}{6}$

First we need to find P_0 and P_1 for the two times.

$$P_1 = 2 \cos\left(3\frac{\pi}{3}\right), 2 \sin\left(3\frac{\pi}{3}\right) \rightarrow P_0 = (-2, 0)$$

$$P_2 = 2 \cos\left(3\frac{\pi}{6}\right), 2 \sin\left(3\frac{\pi}{6}\right) \rightarrow P_1 = (0, 2)$$

Now we need to find the slope.

$$m = \frac{\cancel{\beta} \cos(3t)}{\cancel{\beta} - \sin(3t)} \Rightarrow m = -\frac{\cos(3t)}{\sin(3t)}$$

$$m_1 = -\frac{\cos(3(\pi/3))}{\sin(3(\pi/3))} = \frac{1}{0} = \infty$$


$$m_2 = -\frac{\cos(3(\pi/6))}{\sin(3(\pi/6))} = -\frac{0}{1} = 0$$

Inputting the slopes into our tangent line equation gives us

$$y_1 = \infty(x + 2) \Rightarrow \frac{y_1}{\infty} = x + 2$$

$$0 = x + 2 \Rightarrow x = -2$$

$$y_2 = 0(x - 2)$$



CONCAVITY

To find the concavity of a parametric curve you find the second-order derivative of the curve. This is different than finding the second-order derivative of a regular function.

SECOND-ORDER DERIVATIVE OF A PARAMETRIC CURVE

Because you are trying to find the second-order derivative with respect to x , you have to arrange the expression as such.

$$\begin{aligned} \frac{d^2y}{dx^2} &= \frac{d}{dt} \left(\frac{dy(t)}{dx(t)} \right) \frac{dt}{dx} \\ &= \frac{\frac{dy'(t)}{dx'(t)}}{\frac{dx}{dt}} \end{aligned}$$

Using the quotient rule gives you

$$= \frac{\frac{y''(t)x'(t) - y'(t)x''(t)}{(x'(t))^2}}{x'(t)}$$

$$\frac{d^2y}{dx^2} = \frac{y''(t)x'(t) - y'(t)x''(t)}{(x'(t))^3} \quad (104)$$

EXAMPLE

For this parametric curve $C = (t^2, t^3 - 3t)$.

- Find for all t such that the curve passes through $(3, 0)$
- Find the tangent line that all pass through $(3, 0)$
- Find points where the tangent line is horizontal and vertical
- Find the intervals where it concaves up and down

Solution

- This is simple, find t such that it will satisfy the point.

$$3 = t^2 = \pm\sqrt{3}$$

- Using the slope equation from the last subsection gives us this expression:

$$\frac{3t^2 - 3}{2t} = m(t)$$

Input both values of t into the slope equation

gives us these values for the slope:

$$\frac{3(-\sqrt{3})^2 - 3}{2 \cdot -\sqrt{3}} = -\sqrt{3}$$

$$\frac{3(\sqrt{3})^2 - 3}{2\sqrt{3}} = \sqrt{3}$$

Therefore, the tangent lines are

$$y_1 = \sqrt{3}(x - 3)$$

$$y_2 = -\sqrt{3}(x - 3)$$

c) Set the slope $m = 0$, then you have:

$$0 = \frac{3t^2 - 3}{2t}$$

$$t = \pm 1$$

Input t into the orginal equation

$$P_{horizontal} = ((\pm 1)^2, (\pm 1)^3 - 3(\pm 1))$$

$$P_{horizontal} = \begin{cases} (1, -2) \\ (1, 2) \end{cases}$$

For the vertical slope set $m = \infty$, then you have:

$$\infty = \frac{3t^2 - 3}{2t}$$

$$2t = \frac{3t^2 - 3}{\infty}$$

$$2t = 0$$

$$t = 0$$

Like before input t back into the orginal eqauation which will give you $P_{vertical} = (0, 0)$.

d) To find concavity we use the second-order derivative eqauation from before.





$$\frac{d^2y}{dx^2} = \frac{(6t)(2t) - (2)(3t^2 - 3)}{(2t)^3}$$

$$\frac{d^2y}{dx^2} = \frac{12t^2 - 6t^2 + 6}{8t^3}$$

$$\frac{d^2y}{dx^2} = \frac{6t^2 + 6}{8t^3}$$

The function concaves upward when $t > 0$ and concaves downward when $t < 0$.

Here is the graphical representation of the curve.



AREA

As we discussed before the area underneath a function between two points is defined by the fundamental theorem of calculus i.e., $A = \int_a^b f(x) dx$.

We can use this same theorem to find the area of a parametric curve.

We know that

$$\frac{dx}{dt} = x'(t)$$

Therefore, $dx = x'(t) dt$.

Substitute this into the fundamental theorem of calculus gives us this:

$$A = \int_a^b y x'(t) dt$$

AREA OF A PARAMETRIC CURVE

$$A = \int_a^b y(t)x'(t) dt \quad (105)$$

To find the area on the y-axis then use this equation:


$$A = \int_a^b x(t)y'(t) dt \quad (106)$$

EXAMPLE

Find the area of this parameter curve $C = (t - \sin(t), 1 - \cos(t))$ from $0 \leq t \leq 2\pi$.

Solution.

$$\begin{aligned} x'(t) &= 1 - \cos(t) dt \\ \int_0^{2\pi} (1 - \cos(t))(1 - \cos(t)) dt &= \\ \int_0^{2\pi} (1 - \cos(t))^2 dt &= \\ \int_0^{2\pi} \cos^2(t) - 2\cos(t) + 1 dt &= \\ \int_0^{2\pi} \frac{1 + \cos(2t)}{2} - 2\cos(t) + 1 dt &= \\ \frac{1}{2}t + \frac{1}{4}\sin(2t) - 2\sin(t) + t \Big|_0^{2\pi} &= \\ 2\pi + \pi = 3\pi & \end{aligned}$$



ARC LENGTH

Similar to $L = \int_a^b \sqrt{1 + (f'(x))^2} dx$, but replace dx with $x'(t) dt$ and replace $f'(x)$ with $\frac{dy}{dx}$.

ARC LENGTH OF A PARAMETRIC CURVE

$$L = \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt \quad (107)$$

EXAMPLE

Find the arc length of this parametric curve $C = (e^t - t, 4e^{t/2})$ from 0 to 2.

Solution.

$$\begin{aligned} x'(t) &= e^t - 1, \quad y'(t) = 2e^{t/2} \\ \int_0^2 \sqrt{(e^t - 1)^2 + (2e^{t/2})^2} dt &= \\ \int_0^2 \sqrt{e^{2t} + 2e^t + 1} dt &= \\ \int_0^2 \sqrt{(e^t + 1)^2} dt &= \\ \int_0^2 e^t + 1 dt &= \\ e^t + t \Big|_0^2 &= \\ e^2 + 1 & \end{aligned}$$

SURFACE AREA

Similar to $SA = \int_a^b 2\pi f(x) \sqrt{1 + (f'(x))^2} dx$, but replace dx with $x'(t) dt$ and replace $f'(x)$ with $\frac{dy}{dx}$.

SURFACE AREA OF A PARAMETRIC CURVE

$$SA = \int_a^b 2\pi y(t) \sqrt{(x'(t))^2 + (y'(t))^2} dt \quad (108)$$

EXAMPLE

Prove that the surface area of a sphere with radius a is $4\pi a^2$.

Solution. The parametric curve for a semicircle with a radius a is $C = (a \cos(t), a \sin(t))$ from $0 \leq t \leq \pi$.

$$x'(t) = -a \sin(t), \quad y'(t) = a \cos(t)$$

$$SA = \int_0^\pi 2\pi a \sin(t) \sqrt{(-a \sin(t))^2 + (a \cos(t))^2} dt$$

$$SA = \int_0^\pi 2\pi a \sin(t) \sqrt{a^2 \sin^2(t) + a^2 \cos^2(t)} dt$$

$$SA = \int_0^\pi 2\pi a^2 \sin(t) \sqrt{\sin^2(t) + \cos^2(t)} dt$$

$$SA = \int_0^\pi 2\pi a^2 \sin(t) \sqrt{1} dt$$

$$SA = \int_0^\pi 2\pi a^2 \sin(t) dt$$

$$SA = 2\pi a^2 \int_0^\pi \sin(t) dt$$

$$SA = 2\pi a^2 \left(-\cos(t) \Big|_0^\pi \right)$$

$$SA = 2\pi a^2 (-(-1) - (-1))$$

$$SA = 2\pi a^2 (2)$$

$$SA = 4\pi a^2$$

■

ELLIPSE

ELLIPSE



Are parametrics curves that encompasses all circles.

$$C = (a \cos(t), b \sin(t)) \quad a, b \in \mathbb{R} \quad (109)$$


$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \quad (110)$$

EXAMPLE

This is an ellipse where the parametric equation is $C = (\cos(t), 2 \sin(t))$. As you can see the "radius" of the curve changes from 1 to 2.



HYPERBOLAS



EXERCISE

1. Consider the parametric equation below

$$C = (x^2 - 2, t + 1), \quad -3 \leq t \leq 3$$

Find the arc length of the parametric equation using the domain provided.

2. Consider the following parametric equation

$$C = (\sin\left(\frac{1}{2}\theta\right), \cos\left(\frac{1}{2}\theta\right), -\pi \leq \tau \leq \pi)$$

Find the arc length and the surface area of the parametric equation using the domain above

3. Find an equation of the tangent to the curve at the point corresponding to the given value of the parameter $x = t \cos(t)$, $y = t \sin(t)$; $t = \pi$

4. Find an equation of the tangent to the curve at the given point by both eliminating the parameter and without eliminating the parameter.

$$x = 5 + \ln(t), \quad y = t^2 + 8, \quad (5, 9)$$

5. Find dy/dx and d^2y/dx^2 of this parametric equation

$$x = e^t, \quad y = te^{-t}$$

At which value of t is the curve concave upward?

PROFESSOR'S VILLAINS

WEBASSIGN MINION

Small website, neutral evil

Armor Class Collectively 10

Hit Points 30 (4d10 + 8)

Speed 1 - 2 weeks

STR	DEX	CON	INT	WIS	CHA
7 (-2)	15 (+2)	9 (-1)	8 (-1)	7 (-2)	8 (-1)

Damage Vulnerabilities Calculator, Wolfram Alpha, Google

Senses —

Languages Algebra, Integration, Derivative

Challenge 1/8 (25 XP)

COVERAGE

Everything that is being discussed in class during the existence of the minion.

COMPOSITION

Multiple Choice. Some questions are MC

Numerical. Some questions require a numerical answer.

Mathematical Expression. Some questions require a mathematical expression.

MIDTERM CHAMPION

Huge paper, party slayer, neutral evil

Armor Class 35

Hit Points 70

Speed 75 minutes

STR	DEX	CON	INT	WIS	CHA
27 (+8)	10 (+0)	25 (+7)	16 (+3)	13 (+1)	21 (+5)

Damage Immunities Calculator, Smartphones, any calculating devices

Senses —

Languages Algebra, Integration, Derivative

Challenge 17 (18,000 XP)

COVERAGE

Everything from the beginning of the book to Sec substitution with angles in the chapter: Techniques of

Integration.

COMPOSITION

Multiple Choice. There are 10 MC question on the midterm, each worth 7 marks each.

GRADE SLAYER TIAMAT

Gargantuan paper psychic, Lathrain of grades, chaotic evil

Armor Class 55

Hit Points N > 0

Speed 120 minutes

STR	DEX	CON	INT	WIS	CHA
30 (+10)	10 (+0)	30 (+10)	26 (+8)	26 (+8)	28 (+9)

Damage Immunities Calculator, Smartphones, any calculating devices

Senses —

Languages Algebra, Integration, Derivative

Challenge 30 (155,000 XP)

COVERAGE

Everything

COMPOSITION


Multiple Choice. 30 % of the entire test is MC, in 2018 it was 8 MC

Long Answer. 70 % of the entire test is long answer, in 2018 6 Long answer

TIAMAT, THE GRADE SLAYER

Be aware that the final is going to be absurdly difficult. If his midterm champion didn't tell you anything then you should turn away and drop the class. If however, you truly believe that you can take on the final then you should at least: **Practice, Practice, Practice** - If you think you've practiced enough then your wrong and should practice some more. Moreover, you should **double check your answers**, there is a good chance that you rolled a nat 1 in arithmetic and messed up somewhere.

APPENDIX A: UNIT CIRCLE



APPENDIX B: OTHER RULES

LOG RULES

LOG RULES

$\log_b a$ is only defined if and only if $a > 0$ and $b > 0$.

$$\log_b a = c \Rightarrow a = b^c \quad (111)$$

$$\ln = \log_e \quad (112)$$

$$\log_b 1 = 0 \quad (113)$$

$$\log_b b = 1 \quad (114)$$

$$b^{\log_b a} = a \quad (115)$$

$$\log_b a^n = n \log_b a \quad (116)$$

$$\log_b a + \log_b c = \log_b a \cdot c \quad (117)$$

$$\log_b a - \log_b c = \log_b \frac{a}{c} \quad (118)$$

$$\frac{\log_b a}{\log_b c} = \log_c a \quad (119)$$

TRIGONOMETRIC HYPERBOLIC FUNCTIONS

TRIGONOMETRIC HYPERBOLIC FUNCTIONS

$$\sinh(x) = \frac{e^x - e^{-x}}{2} \quad (120)$$

$$\cosh(x) = \frac{e^x + e^{-x}}{2} \quad (121)$$

$$\sinh(x)' = \cosh(x) \quad (122)$$

$$\cosh(x)' = \sinh(x) \quad (123)$$

$$\cosh^2(x) - \sinh^2(x) = 1 \quad (124)$$

$$e^x = \frac{\cosh(x) + \sinh(x)}{2} \quad (125)$$

$$e^{-x} = \frac{\cosh(x) - \sinh(x)}{2} \quad (126)$$

$\sinh(x)$ is an odd function and $\cosh(x)$ is an even function.

CREDITS

GENERAL

- Created by: Jihoon Og, u/DnD_Notes
- Editor: Mohammad Ali, Niksirat
- Compiled on Monday 15th April, 2019 at 12:59
- Typesetting engine: [L^AT_EX](#)
- Dungeon and Dragon (5e) LaTeX [Template](#)

ART

- Red Dragon for the cover art is from [D&D Beyond](#)
- Integral symbol is from [here](#)
- The stick figure that is attacking the dragon is from [xkcd](#)
- Andy the D&D Ampersand is from [Dungeon and Dragons](#)
- Cover art formatting and design done in Photoshop CC 2019

DISCLAIMER

This document is completely unofficial and in no way endorsed by Wizards of the Coast or Games Workshop. All associated marks, names, races, race insignia characters, locations, illustrations and images from Dungeons and Dragons, and Warhammer are either ®, ©, TM and/or Copyright Wizards of the Coast Ltd 2012-2018. All used without permission. No challenge to their status intended. All Rights Reserved to their respective owners.