

<b>SDG Goal</b>	<b>3</b>	<b>Good health and well-being</b>
<b>SDG Target</b>	<b>3.5</b>	<b>Strengthen the prevention and treatment of substance abuse, including narcotic drug abuse and harmful use of alcohol</b>
<b>SDG Indicator</b>	<b>3.5.1</b>	<b>Coverage of treatment interventions (pharmacological, psychosocial and rehabilitation and aftercare services) for substance use disorders</b>

1. Name of data series	
<b>Inpatient treatment of dependence syndrom due to psychoactive substance use, total</b>	
Compliant with SDG metadata: <b>yes, partly</b>	<a href="#">SDG Metadata</a>
2. Definition of indicator	
The indicator “Inpatient treatment of dependence syndrom due to psychoactive substance use, total” is defined as the number of patients treated in hospitals and the number of patients treated in prevention and rehabilitation facilities with the diagnosis: F10.2-F19.2 (dependence syndrome).	
3. Comparison with SDG metadata (as of 20/09/2019)	
The indicator describes the SDG indicator 3.5.1 only partly, because it only covers inpatient treatment.	
4. Data description	
<p>The data on “inpatient treatment of dependence syndrom due to psychoactive substance use, total” in hospitals and in prevention and rehabilitation facilities are derived from the hospital statistics. Both statistics are conducted by the Federal Statistical Office.</p> <p>The indicator is defined as the number of patients treated in hospitals and the number of patients treated in prevention and rehabilitation facilities with the diagnosis F10.2-F19.2 (dependence syndrome). According to ICD-10 the following codes are included:</p> <ul style="list-style-type: none"> <li>• F10.2 Mental and behavioral disorders due to use of alcohol</li> <li>• F11.2 Mental and behavioural disorders due to use of opioids</li> <li>• F12.2 Mental and behavioural disorders due to use of cannabinoids</li> <li>• F13.2 Mental and behavioural disorders due to use of sedatives or hypnotics</li> <li>• F14.2 Mental and behavioural disorders due to use of cocaine</li> <li>• F15.2 Mental and behavioural disorders due to use of other stimulants, including caffeine</li> <li>• F16.2 Mental and behavioural disorders due to use of hallucinogens</li> <li>• F17.2 Mental and behavioural disorders due to use of tobacco</li> <li>• F18.2 Mental and behavioural disorders due to use of volatile solvents</li> <li>• F19.2 Mental and behavioural disorders due to multiple drug use and use of other psychoactive substances</li> </ul> <p>ICD-10 is the International Statistical Classification of Diseases and Related Health Problems 10th Revision (German Modification).</p>	
5. Calculation method	
$\text{Inpatient treatment of dependence syndrom due to psychoactive substance use, total} = \frac{\sum_{\text{ICD-Codes}} \text{Patients treated in hospitals (number)} + \text{Patients treated in prevention and rehabilitation facilities (number)}}{1,000}$	
6. Unit of measure	1,000

7. Timeliness	8. Frequency
t + 12 months	Annual
9. Last regular revision	10. Revised period
Not applicable	Not applicable

11. Accessibility of source data
<p>“Diagnostic data of the hospitals” – GBE  <a href="http://www.gbe-bund.de/gbe10/i?i=702:1018E">http://www.gbe-bund.de/gbe10/i?i=702:1018E</a></p> <p>and</p> <p>“Diagnostic data of the prevention or rehabilitation facilities” – GBE  <a href="http://www.gbe-bund.de/gbe10/i?i=902:1020E">http://www.gbe-bund.de/gbe10/i?i=902:1020E</a></p>
12. Metadata on source data
<p>Quality report of Hospital statistics - diagnostic data of the hospital patients (only available in German):  “Qualitätsbericht – Diagnosen der Krankenhauspatientinnen und –patienten”  <a href="https://www.destatis.de/DE/Methoden/Qualitaet/Qualitaetsberichte/Gesundheit/diagnose-krankenhauspatienten.pdf?__blob=publicationFile">https://www.destatis.de/DE/Methoden/Qualitaet/Qualitaetsberichte/Gesundheit/diagnose-krankenhauspatienten.pdf?__blob=publicationFile</a></p> <p>and</p> <p>Quality report of Hospital statistics - diagnostic data of the prevention and rehabilitation facilities patients (only available in German):  “Qualitätsbericht - Diagnosen der Vorsorge- und Rehabilitationseinrichtungen”  <a href="https://www.destatis.de/DE/Methoden/Qualitaet/Qualitaetsberichte/Gesundheit/diagnose-vorsorge-rehaeinrichtungen.pdf?__blob=publicationFile">https://www.destatis.de/DE/Methoden/Qualitaet/Qualitaetsberichte/Gesundheit/diagnose-vorsorge-rehaeinrichtungen.pdf?__blob=publicationFile</a></p>

13. Related SDG data series (duplicate indicators or sub-indicators to same indicator)
3.5.1 Inpatient treatment of alcohol dependence

For more information please contact: <https://www.destatis.de/EN/Service/Contact/Contact.html>

<b>SDG Goal</b>	<b>3</b>	<b>Good health and well-being</b>
<b>SDG Target</b>	<b>3.5</b>	<b>Strengthen the prevention and treatment of substance abuse, including narcotic drug abuse and harmful use of alcohol</b>
<b>SDG Indicator</b>	<b>3.5.1</b>	<b>Coverage of treatment interventions (pharmacological, psychosocial and rehabilitation and aftercare services) for substance use disorders</b>

1. Name of data series
<b>Inpatient treatment of alcohol dependence</b>
Compliant with SDG metadata: <b>yes, partly</b> <a href="#">SDG Metadata</a>

2. Definition of indicator
The indicator “Inpatient treatment of alcohol dependence” is defined as the number of patients treated in hospitals and the number of patients treated in prevention and rehabilitation facilities with the diagnosis: F10.2.
3. Comparison with SDG metadata (as of 20/09/2019)
The indicator describes the SDG indicator 3.5.1 only partly, because it only covers inpatient treatment.

4. Data description	
The data on “inpatient treatment of alcohol dependence” in hospitals and in prevention and rehabilitation facilities are derived from the hospital statistics. Both statistics are conducted by the Federal Statistical Office. The indicator is defined as the number of patients treated in hospitals and the number of patients treated in prevention and rehabilitation facilities with the diagnosis F10.2 Mental and behavioral disorders due to use of alcohol. ICD-10 is the International Statistical Classification of Diseases and Related Health Problems 10th Revision (German Modification).	
5. Calculation method	
$\text{Inpatient treatment of alcohol dependence} = \frac{\sum_{\text{ICD-Code}} \text{Patients treated in hospitals (number)} + \text{Patients treated in prevention and rehabilitation facilities (number)}}{1,000}$	
6. Unit of measure	1,000

7. Timeliness	8. Frequency
t + 12 months	Annual
9. Last regular revision	10. Revised period
Not applicable	Not applicable

11. Accessibility of source data
<p>“Diagnostic data of the hospitals” – GBE  <a href="http://www.gbe-bund.de/gbe10/i?i=702:1018E">http://www.gbe-bund.de/gbe10/i?i=702:1018E</a></p> <p>and</p> <p>“Diagnostic data of the prevention or rehabilitation facilities” – GBE  <a href="http://www.gbe-bund.de/gbe10/i?i=902:1020E">http://www.gbe-bund.de/gbe10/i?i=902:1020E</a></p>
12. Metadata on source data
<p>Quality report of Hospital statistics - diagnostic data of the hospital patients (only available in German):  “Qualitätsbericht – Diagnosen der Krankenhauspatientinnen und –patienten”  <a href="https://www.destatis.de/DE/Methoden/Qualitaet/Qualitaetsberichte/Gesundheit/diagnose-krankenhauspatienten.pdf?__blob=publicationFile">https://www.destatis.de/DE/Methoden/Qualitaet/Qualitaetsberichte/Gesundheit/diagnose-krankenhauspatienten.pdf?__blob=publicationFile</a></p> <p>and</p> <p>Quality report of Hospital statistics - diagnostic data of the prevention and rehabilitation facilities patients (only available in German):  “Qualitätsbericht - Diagnosen der Vorsorge- und Rehabilitationseinrichtungen”  <a href="https://www.destatis.de/DE/Methoden/Qualitaet/Qualitaetsberichte/Gesundheit/diagnose-vorsorge-rehaeinrichtungen.pdf?__blob=publicationFile">https://www.destatis.de/DE/Methoden/Qualitaet/Qualitaetsberichte/Gesundheit/diagnose-vorsorge-rehaeinrichtungen.pdf?__blob=publicationFile</a></p>
13. Related SDG data series (duplicate indicators or sub-indicators to same indicator)
3.5.1 Inpatient treatment of dependence syndrom due to psychoactive substance use, total
For more information please contact: <a href="https://www.destatis.de/EN/Service/Contact/Contact.html">https://www.destatis.de/EN/Service/Contact/Contact.html</a>