

Designing Oligos for Universal Knock-in Vectors using Type 2 Restriction Enzymes:

Note In the following document when orientation words are used, they are used in the context of the reading frame of your genetic loci of interest. e.g. A 5' Reverse CRISPR means that the target site for the CRISPR is on the reverse strand at your locus and is toward the 5' end of the gene. Upstream Homology Domains are 5' of the CRISPR cut and Downstream Homology Domains are 3' of the cut. Also note: Upper case and lower case bases are the same; they are typed the way they are as a visual marker of the different parts of the homology domains.

For the 5' Homology Domain:

1) Open the sequence file for your gene and identify the CRISPR site. (In this example it is a Reverse CRISPR target in Yellow, the PAM is in Orange)

Copy the 48bp 5' of the CRISPR cut into a new sequence file; this is your 5' homology.

	Sequence	Start	Length	End	ORF	Tm	%GC	Linear
	2391	232<0>	48<0>	279<2>		72°C	48%	<input checked="" type="checkbox"/> Dam/Dcm

	* 10 * 20 * 30 * 40 * 50 * 60 * 70 *
1	CGCTATATGAACCCGACGGCGCACGGGGAGGAGAAAAACGACCCACATGCTGCCAGACTCCGAATGGGTTAATG
76	AAGAGCGTGTCTTTCATCGTCAAAGATAGCTGAGAAATGTGGTGATATTAACGCACCAGAACAACTCTTGCGT
151	AGGACGTAGCTGAGGAAAAGAGTGAAATCTACTCATCGAGGACTGAGACGGTGGTACTTCTTGAAGCACCATGA
226	GCTGGA ^{TTTTCTTACGCGGTTGTTGGATGAAATCTCCAACCACTCCA} ^{CCTTCG} TGGGCAAGATATGGCTCACGT
301	TATTCATCATCTTCCGCATTGTTTGAAGTGTGTGGGGGAGAAATCGATATACTACGATGAACAGAGCAAATTTG
376	TGTGTAATACCCAGCAACCTGGTTGTGAGAACGTTTGCTACGATGCATTTCACCACTCTCTCATGTCCGGTTCT
451	GGGTTTCCAGATCATTTTGATCACAACCCCACTATCATGTACTTGGGATTGCTATGCACAAGATCGCTCGGT


2) Observe the next three bases immediately upstream of your 48bp of homology, and pick a base not present to be the 3bp spacer between your homology and the Universal PAM in the vector. (Here the three bases are “GGA” so I chose “ccc” for my spacer)

Add the spacer to your new file 5' (in front) of your homology.


	* 10 * 20 * 30 * 40 * 50 * 60 * 70 *
1	cccGTTTCTTACGCGGTTGTTGGATGAAATCTCCAACCACTCCA ^{CCTTCG}

3) Determine where the last codon is in your homology. Here the 3' G in the homology domain is the first base in the last codon upstream of the CRISPR cut.


Complete the codon by adding the remaining bases for that codon from your sequence to ensure your integration event will be in frame.




4) Add the BfuAI overhang sequences for cloning, to the ends of your homology domain. 5'GCGG and 3'GGAT (or 3'CTTC for the prism series). (Here we are setting both overhangs to prevent errors in copying sequence for your oligos in the next two steps.)



5) Your 5' Homology Oligo A will be this sequence from the beginning to the end of the last codon. Copy and paste this into a new file and save it. (In this example this oligo sequence is 5'GCGGcccGTTTTCTTACGCGGTTGTTGGATGAAATCTCCAACCACTCCACCTTCGtg3')



6) Your 5' Homology Oligo B will be the reverse complement of this sequence from beginning of the spacer to the end of the sequence. Copy the reverse complement, paste it into a new file, and save it. (In this example this oligo sequence is 5'ATCCcaCGAAGGTGGAGTGGTTGGAGATTTTCATCCAACAACCGCGTAAGAAAACggg3')



Continue to next page – >

For the 3' Homology Domain:

7) Open sequence file for your gene and identify the CRISPR site. (Reverse CRISPR target in Yellow, PAM in Orange)

Copy the 48bp 3' of the CRISPR cut into a new sequence file; this is your 3' homology.

8) Observe the next three bases downstream of your 48bp of homology, and pick a base not present to be the 3bp spacer between your homology and the Universal PAM in the vector. (Here the bases are “CTG” so I chose “aaa” for my spacer.)

Add the spacer to your new file 3' of (after) your homology.


9) Add the BspQI overhang sequences for cloning, to the ends of your homology domain. 5'AAG and 3'CCG. (Here we are setting both overhangs to prevent errors in copying sequence for your oligos in the next two steps.)

10) Your 3' Homology Oligo A will be this sequence from the beginning of the sequence to the end of the spacer. (Here =

5'AAGTGGGCAAGATATGGCTCACGTTATTTCATCATCTTCCGCATTGTTTGAaaa3')

Continue to next page –>

11) Your 3' Homology Oligo B will be the reverse compliment of this sequence from the beginning of the homology to the end of the sequence. (Here = 5'CGGtttTCAAAACAATGCGGAAGATGATGAATAACGTGAGCCATATCTTGCCCA3')



1 AAG TGGGCAAGATATGGCTC ACGTTATTCATCACTTCCGCATTGTTTTGA aaaCCG