

Impacts of halibut bycatch and wastage on halibut coast-wide yield and spawning biomass

Steve Martell

University of British Columbia

martell.steve@gmail.com

April 17, 2012

Introduction

Simulation model

Scenarios

Results

EBio

SBio

YBio

Wastage

Yield Loss Ratio

Summary

Outline

Introduction

Simulation model

Scenarios

Results

EBio

SBio

YBio

Wastage

Yield Loss Ratio

Summary

Introduction

Simulation model

Scenarios

Results

EBio

SBio

YBio

Wastage

Yield Loss Ratio

Summary

Objectives

Provide an alternative investigation into the effects of halibut bycatch and wastage in the GOA and BSAI groundfish fisheries.

Introduction

Simulation model

Scenarios

Results

EBio

SBio

YBio

Wastage

Yield Loss Ratio

Summary

Objectives

Provide an alternative investigation into the effects of halibut bycatch and wastage in the GOA and BSAI groundfish fisheries.

Research Question:

What are the impacts of bycatch reductions on future estimates of halibut **biomass**, **yield**, **spawning biomass** and **wastage** by age–size categories over a fifteen year time horizon?

Objectives

Provide an alternative investigation into the effects of halibut bycatch and wastage in the GOA and BSAI groundfish fisheries.

Research Question:

What are the impacts of bycatch reductions on future estimates of halibut **biomass**, **yield**, **spawning biomass** and **wastage** by age–size categories over a fifteen year time horizon?

To answer this question:

Developed a deterministic sex/age-structured simulation model using IPHC assessment outputs to parameterize the model.

Outline

Introduction

Simulation model

Scenarios

Results

EBio

SBio

YBio

Wastage

Yield Loss Ratio

Summary

Introduction

Simulation model

Scenarios

Results

EBio

SBio

YBio

Wastage

Yield Loss Ratio

Summary

Simulation model overview

- 2 genders, 30 age-classes, 1996–2026 (or longer).

Simulation model overview

- 2 genders, 30 age-classes, 1996–2026 (or longer).
- Inputs: IPHC initial age-composition, natural mortality (by sex), annual recruits, fishing mortality rates, size-selectivity, survey mean length-at-age.

Simulation model overview

- 2 genders, 30 age-classes, 1996–2026 (or longer).
- Inputs: IPHC initial age-composition, natural mortality (by sex), annual recruits, fishing mortality rates, size-selectivity, survey mean length-at-age.
- Not used: empirical catch weight-at-age, age-smearing.

Simulation model overview

- 2 genders, 30 age-classes, 1996–2026 (or longer).
- Inputs: IPHC initial age-composition, natural mortality (by sex), annual recruits, fishing mortality rates, size-selectivity, survey mean length-at-age.
- Not used: empirical catch weight-at-age, age-smearing.
- **Exploitable biomass based on vulnerable biomass in commercial fishery.**

Simulation model overview

- 2 genders, 30 age-classes, 1996–2026 (or longer).
- Inputs: IPHC initial age-composition, natural mortality (by sex), annual recruits, fishing mortality rates, size-selectivity, survey mean length-at-age.
- Not used: empirical catch weight-at-age, age-smearing.
- Exploitable biomass based on vulnerable biomass in commercial fishery.
- Wastage based on joint probability:
 $P(\text{capture}) * (1 - P(\text{retention})) * M_d$.

Introduction

Simulation model

Scenarios

Results

EBio

SBio

YBio

Wastage

Yield Loss Ratio

Summary

Simulation model overview

- 2 genders, 30 age-classes, 1996–2026 (or longer).
- Inputs: IPHC initial age-composition, natural mortality (by sex), annual recruits, fishing mortality rates, size-selectivity, survey mean length-at-age.
- Not used: empirical catch weight-at-age, age-smearing.
- Exploitable biomass based on vulnerable biomass in commercial fishery.
- Wastage based on joint probability:
 $P(\text{capture}) * (1 - P(\text{retention})) * M_d$.
- Apportionment based on 2011 apportionment.

Simulation model overview

- 2 genders, 30 age-classes, 1996–2026 (or longer).
- Inputs: IPHC initial age-composition, natural mortality (by sex), annual recruits, fishing mortality rates, size-selectivity, survey mean length-at-age.
- Not used: empirical catch weight-at-age, age-smearing.
- Exploitable biomass based on vulnerable biomass in commercial fishery.
- Wastage based on joint probability:
 $P(\text{capture}) * (1 - P(\text{retention})) * M_d$.
- Apportionment based on 2011 apportionment.
- Area based CEY's based area specific harvest rate.

Simulation model overview

- 2 genders, 30 age-classes, 1996–2026 (or longer).
- Inputs: IPHC initial age-composition, natural mortality (by sex), annual recruits, fishing mortality rates, size-selectivity, survey mean length-at-age.
- Not used: empirical catch weight-at-age, age-smearing.
- Exploitable biomass based on vulnerable biomass in commercial fishery.
- Wastage based on joint probability:
 $P(\text{capture}) * (1 - P(\text{retention})) * M_d$.
- Apportionment based on 2011 apportionment.
- Area based CEY's based area specific harvest rate.
- Commercial fishery share: $\text{CEY} - (\text{other removals})$.

Simulation model overview

- 2 genders, 30 age-classes, 1996–2026 (or longer).
- Inputs: IPHC initial age-composition, natural mortality (by sex), annual recruits, fishing mortality rates, size-selectivity, survey mean length-at-age.
- Not used: empirical catch weight-at-age, age-smearing.
- Exploitable biomass based on vulnerable biomass in commercial fishery.
- Wastage based on joint probability:
 $P(\text{capture}) * (1 - P(\text{retention})) * M_d$.
- Apportionment based on 2011 apportionment.
- Area based CEY's based area specific harvest rate.
- Commercial fishery share: CEY - (other removals).
- Reduction in bycatch translates to increase in commercial catch.

[Introduction](#)[Simulation model](#)[Scenarios](#)[Results](#)[EBio](#)[SBio](#)[YBio](#)[Wastage](#)[Yield Loss Ratio](#)[Summary](#)

Model outputs

Exploitable Biomass (EBio):

Start of year biomass vulnerable to the commercial gear.

Model outputs

Exploitable Biomass (EBio):

Start of year biomass vulnerable to the commercial gear.

Spawning Biomass (SBio):

Start of year female mature biomass.

Model outputs

Exploitable Biomass (EBio):

Start of year biomass vulnerable to the commercial gear.

Spawning Biomass (SBio):

Start of year female mature biomass.

Commercial Yield (YBio):

Weight of commercial landings (excluding wastage).

Model outputs

Exploitable Biomass (EBio):

Start of year biomass vulnerable to the commercial gear.

Spawning Biomass (SBio):

Start of year female mature biomass.

Commercial Yield (YBio):

Weight of commercial landings (excluding wastage).

Wastage (WBio):

Weight of dead discarded undersized fish in commercial fishery.

Model outputs

Exploitable Biomass (EBio):

Start of year biomass vulnerable to the commercial gear.

Spawning Biomass (SBio):

Start of year female mature biomass.

Commercial Yield (YBio):

Weight of commercial landings (excluding wastage).

Wastage (WBio):

Weight of dead discarded undersized fish in commercial fishery.

Lost Yield (LBio):

Difference between YBio with *no* bycatch & *with* bycatch.

Model outputs

Exploitable Biomass (EBio):

Start of year biomass vulnerable to the commercial gear.

Spawning Biomass (SBio):

Start of year female mature biomass.

Commercial Yield (YBio):

Weight of commercial landings (excluding wastage).

Wastage (WBio):

Weight of dead discarded undersized fish in commercial fishery.

Lost Yield (LBio):

Difference between YBio with *no* bycatch & *with* bycatch.

Yield Loss Ratio (YLR):

Ratio between the yield loss and the bycatch.

Outline

Introduction

Simulation model

Scenarios

Results

EBio

SBio

YBio

Wastage

Yield Loss Ratio

Summary

Introduction

Simulation model

Scenarios

Results

EBio

SBio

YBio

Wastage

Yield Loss Ratio

Summary

Growth & Recruitment Scenarios (states of nature)

1. Poor (60% below average recruitment)
2. Average
3. Good (60% above average recruitment)

Growth:

Growth & Recruitment Scenarios (states of nature)

Growth:


1. Density-independent (using 2011 average length-at-age)
2. Density dependent.

Policy Scenarios

Three alternative policies:

1. Status quo: 2011 BSAI bycatch levels
2. 50% Reduction in BSAI bycatch (4ABCDE)
 - ▶ decrease from 5.535 million lb to 2.765 million lb.
3. 50% Reduction in GULF bycatch (3AB)
 - ▶ decrease from 5.752 million lb to 2.876 million lb.

Summary plots



Outline

Introduction

Simulation model

Scenarios

Results

EBio

SBio

YBio

Wastage

Yield Loss Ratio

Summary

Introduction

Simulation model

Scenarios

Results

EBio

SBio

YBio







Wastage

Yield Loss Ratio

Summary







Decision table: EBio

Effect of bycatch reduction on Exploitable Biomass

Policy	Status quo	50% BSAI	50% GULF
DI Growth			
DD Growth			
Average	633	631.5	631







Decision table: SBio

Effect of bycatch reduction on Female Spawning Biomass

Policy	Status quo	50% BSAI	50% GULF
DI Growth			
DD Growth			
Average	490.5	488	487.5







Decision table: Commercial Yield

Effect of bycatch reduction on Commercial Yield

Policy	Status quo	50% BSAI	50% GULF
DI Growth			
DD Growth			
Average	101	103	103







Decision table: Comm. Wastage

Effect of bycatch reduction on Commercial Wastage

Policy	Status quo	50% BSAI	50% GULF
DI Growth	 <p>2.796</p>	 <p>2.883</p>	 <p>2.891</p>
DD Growth	 <p>2.192</p>	 <p>2.279</p>	 <p>2.287</p>
Average	2.494	2.581	2.589

Decision table: Yield Loss Ratio

Effect of bycatch reduction on Yield Loss Ratio

Policy	Status quo	50% BSAI	50% GULF
DI Growth			
DD Growth			
Average	2.494	2.581	2.589

Outline

Introduction

Simulation model

Scenarios

Results

EBio

SBio

YBio

Wastage

Yield Loss Ratio

Summary

Introduction

Simulation model

Scenarios

Results

EBio

SBio

YBio

Wastage

Yield Loss Ratio

Summary

Summary

- Density-dependent growth dominates the scale of EBio in comparison to recruitment trends.

Summary

- Density-dependent growth dominates the scale of EBio in comparison to recruitment trends.
- Exploitable and spawning coastwide biomass are largely insensitive to BSAI and GULF bycatch.

Summary

- Density-dependent growth dominates the scale of EBio in comparison to recruitment trends.
- Exploitable and spawning coastwide biomass are largely insensitive to BSAI and GULF bycatch.
- Fishing mortality is dominated by the directed commercial fishery.

Introduction

Simulation model

Scenarios

Results

EBio


SBio

YBio

Wastage

Yield Loss Ratio

Summary



Summary

- Density-dependent growth dominates the scale of EBio in comparison to recruitment trends.
- Exploitable and spawning coastwide biomass are largely insensitive to BSAI and GULF bycatch.
- Fishing mortality is dominated by the directed commercial fishery.
- Reducing bycatch in BSAI or Gulf by 50% (≈ 2.7 million lb) results in a ≈ 2 million lb yr^{-1} increase in the directed fishery.

Acknowledgments

IPHC staff
At-sea Processors Association
United Catcher Boats
Pacific Seafood Processors Association
Alaska Groundfish Data Bank
Marine Conservation Alliance
Groundfish Fourm
Alaska Whitefish Trawlers