

Part I: Moving towards a sustainable fisheries
framework for BC herring: data, models &
alternative assumptions.

Part II: Stock assessment and management
advice for BC Herring stocks (2011/2012)

Steven Martell, Jake Schweigert, Jaclyn Cleary, Vivian Haist

University of British Columbia

martell.steve@gmail.com

August 13, 2011



Moving towards the sustainable fisheries framework.

Overview

- ▶ Review of the HCAM model in June 17-18, 2010.
 - ▶ Model parameterization of q .
 - ▶ Parametrization of q , M , and selectivity is confounded.
- ▶ Development of a new integrated Statistical Catch Age Model ($iSCA_M$).
- ▶ Data, assumptions and Analytical methods.
- ▶ Outstanding issues.



Introduction

- ▶ Current harvest control rule for BC herring:
 - ▶ Cutoffs set at $0.25 B_0$
 - ▶ 20% exploitation rate
 - ▶ Estimates of B_0 were last updated in 1996.
- ▶ HCAM model assumed $q = 1$ for the dive survey data.
- ▶ Natural mortality is modelled as a random-walk.
- ▶ Gill net selectivity is a function of weight-at-age.



Harvest Strategy Compliant with Precautionary Approach




Figure: Fisheries management framework consistent with a precautionary approach.



Key elements for the new framework

Reference points

- ▶ Limit Reference Point (LRP) & Upper Stock Reference (USR) requires knowledge of stock productivity and population scale.
- ▶ Removal Rate requires knowledge of stock productivity.
- ▶ MSY-based reference points require *a priori* allocation to different gears.

Risk & Decision making

- ▶ Onus on being able to reliably determine stock status (informative data).



Herring Stock Assessment Model Review

Summary of Panel Recommendations

- ▶ Panel concluded that $q_2 = 1$ was inappropriate.
- ▶ CUTOFFS can be fixed or annually estimated (should be updated if management objective is 25% B_0)
- ▶ A model based approach to estimating B_0 and B_{MSY} is appropriate.
- ▶ Recruitment variation should be estimated within the model rather than fixing it at a pre-specified level.
- ▶ Issues regarding estimating selectivity vs. availability should be explored (data is limited to estimate availability).
- ▶ Science advice should be risk neutral.
- ▶ MSE should explore elements of the Sustainable Fisheries Framework (i.e., ensure that $B_t > 0.4B_{MSY}$ with 95% certainty over two generations.)
- ▶ ...



Input data

The input data for $i\text{SCA}_M$ is the same as HCAM:

- ▶ Catch by gear,
- ▶ Spawn survey index,
- ▶ Age-composition data for all gears,
- ▶ Empirical weight-at-age data.



