

# Math Camp

Justin Grimmer

Professor  
Department of Political Science  
Stanford University

September 18th, 2018

# Questions?

# Questions?

- 1) What is a continuous random variable?

# Questions?

- 1) What is a continuous random variable?
- 2) What does it mean when we say  $X \sim \text{Normal}(\mu, \sigma^2)$

# Questions?

- 1) What is a continuous random variable?
- 2) What does it mean when we say  $X \sim \text{Normal}(\mu, \sigma^2)$
- 3) Explain why the pdf and cdf contain the same information

# Questions?

- 1) What is a continuous random variable?
- 2) What does it mean when we say  $X \sim \text{Normal}(\mu, \sigma^2)$
- 3) Explain why the pdf and cdf contain the same information
- 4) Explain why the height of the pdf isn't a probability

# Questions?

- 1) What is a continuous random variable?
- 2) What does it mean when we say  $X \sim \text{Normal}(\mu, \sigma^2)$
- 3) Explain why the pdf and cdf contain the same information
- 4) Explain why the height of the pdf isn't a probability
- 5) Suppose  $Z \sim \text{Normal}(0, 1)$ . What is  $Y = aZ + b$ ?

# Where We've Been, Where We're Going

## Multivariate Distributions

- 1) Joint Density
- 2) Covariance, Marginalization
- 3) Independence of Random Variables
- 4) Properties of Sums of Random Variables
- 5) The Multivariate Normal Distribution and You

# Continuous Random Variable

## Definition

*X is a continuous random variable if there exists a nonnegative function defined for all  $x \in \mathbb{R}$  having the property for any (measurable) set of real numbers B,*

$$P(X \in B) = \int_B f(x)dx$$

We'll call  $f(\cdot)$  the **probability density function** for X.

## Definition

**Multivariate Distribution** We will say that  $X$  and  $Y$  are *jointly continuous* if, for all  $x \in \mathbb{R}$  and  $y \in \mathbb{R}$ , there exists a function  $f(x, y)$  such that for every set  $C \subset \mathbb{R}^2$ ,

## Definition

**Multivariate Distribution** We will say that  $X$  and  $Y$  are *jointly continuous* if, for all  $x \in \mathbb{R}$  and  $y \in \mathbb{R}$ , there exists a function  $f(x, y)$  such that for every set  $C \subset \mathbb{R}^2$ ,

What is  $C \subset R^2$ ?

## Definition

**Multivariate Distribution** We will say that  $X$  and  $Y$  are *jointly continuous* if, for all  $x \in \mathbb{R}$  and  $y \in \mathbb{R}$ , there exists a function  $f(x, y)$  such that every set  $C \subset \mathbb{R}^2$ ,

What is  $C \subset R^2$ ?

$$- R^2 = R \underbrace{\times}_{\text{Cartesian Product}} R$$

## Definition

**Multivariate Distribution** We will say that  $X$  and  $Y$  are *jointly continuous* if, for all  $x \in \mathbb{R}$  and  $y \in \mathbb{R}$ , there exists a function  $f(x, y)$  such that every set  $C \subset \mathbb{R}^2$ ,

What is  $C \subset R^2$ ?

- $R^2 = R \times R$

  
Cartesian Product

- This is the 2-d plane (your piece of paper)

## Definition

**Multivariate Distribution** We will say that  $X$  and  $Y$  are *jointly continuous* if, for all  $x \in \mathbb{R}$  and  $y \in \mathbb{R}$ , there exists a function  $f(x, y)$  such that every set  $C \subset \mathbb{R}^2$ ,

What is  $C \subset R^2$ ?

- $R^2 = R \times R$

 Cartesian Product


- This is the 2-d plane (your piece of paper)
- $C$  is a subset of the 2-d plane

## Definition

**Multivariate Distribution** We will say that  $X$  and  $Y$  are *jointly continuous* if, for all  $x \in \mathbb{R}$  and  $y \in \mathbb{R}$ , there exists a function  $f(x, y)$  such that every set  $C \subset \mathbb{R}^2$ ,

What is  $C \subset R^2$ ?

- $R^2 = R \times R$   
 $\underbrace{\quad \quad}_{\text{Cartesian Product}}$
- This is the 2-d plane (your piece of paper)
- $C$  is a subset of the 2-d plane
  - $C = \{x, y : x \in [0, 1], y \in [0, 1]\}$




## Definition

**Multivariate Distribution** We will say that  $X$  and  $Y$  are *jointly continuous* if, for all  $x \in \mathbb{R}$  and  $y \in \mathbb{R}$ , there exists a function  $f(x, y)$  such that every set  $C \subset \mathbb{R}^2$ ,

What is  $C \subset R^2$ ?

- $R^2 = R \times R$   
     $\underbrace{\quad \quad}_{\text{Cartesian Product}}$
- This is the 2-d plane (your piece of paper)
- $C$  is a subset of the 2-d plane
  - $C = \{x, y : x \in [0, 1], y \in [0, 1]\}$
  - $C = \{x, y : x^2 + y^2 \leq 1\}$



## Definition


**Multivariate Distribution** We will say that  $X$  and  $Y$  are *jointly continuous* if, for all  $x \in \mathbb{R}$  and  $y \in \mathbb{R}$ , there exists a function  $f(x, y)$  such that every set  $C \subset \mathbb{R}^2$ ,

What is  $C \subset R^2$ ?

- $R^2 = R \times R$

$\underbrace{\quad \times \quad}_{\text{Cartesian Product}}$   $R$

- This is the 2-d plane (your piece of paper)
- $C$  is a subset of the 2-d plane
  - $C = \{x, y : x \in [0, 1], y \in [0, 1]\}$
  - $C = \{x, y : x^2 + y^2 \leq 1\}$
  - $C = \{x, y : x > y, x, y \in (0, 2)\}$



## Definition

**Multivariate Distribution** We will say that  $X$  and  $Y$  are *jointly continuous* if, for all  $x \in \mathbb{R}$  and  $y \in \mathbb{R}$ , there exists a function  $f(x, y)$  such that every set  $C \subset \mathbb{R}^2$ ,

$$P\{(X, Y) \in C\} = \iint_{(x,y) \in C} f(x, y) dx dy$$

What is  $C \subset \mathbb{R}^2$ ?

- $R^2 = R \times R$

 Cartesian Product

- This is the 2-d plane (your piece of paper)
- $C$  is a subset of the 2-d plane
  - $C = \{x, y : x \in [0, 1], y \in [0, 1]\}$
  - $C = \{x, y : x^2 + y^2 \leq 1\}$
  - $C = \{x, y : x > y, x, y \in (0, 2)\}$

## Definition

**Multivariate Distribution** We will say that  $X$  and  $Y$  are *jointly continuous* if, for all  $x \in \mathbb{R}$  and  $y \in \mathbb{R}$ , there exists a function  $f(x, y)$  such that every set  $C \subset \mathbb{R}^2$ ,

$$P\{(X, Y) \in C\} = \iint_{(x,y) \in C} f(x, y) dx dy$$

What is  $C \subset \mathbb{R}^2$ ?

- $R^2 = R \times R$

 Cartesian Product

- This is the 2-d plane (your piece of paper)
- $C$  is a subset of the 2-d plane
  - $C = \{x, y : x \in [0, 1], y \in [0, 1]\}$
  - $C = \{x, y : x^2 + y^2 \leq 1\}$
  - $C = \{x, y : x > y, x, y \in (0, 2)\}$
  - $C = \{x, y : x \in A, y \in B\}$

## Definition

**Multivariate Distribution** We will say that  $X$  and  $Y$  are *jointly continuous* if, for all  $x \in \mathbb{R}$  and  $y \in \mathbb{R}$ , there exists a function  $f(x, y)$  such that for every set  $C \subset \mathbb{R}^2$ ,

$$P\{(X, Y) \in C\} = \int_B \int_A f(x, y) dx dy$$

What is  $C \subset \mathbb{R}^2$ ?

- $\mathbb{R}^2 = \mathbb{R} \times \mathbb{R}$

Cartesian Product

- This is the 2-d plane (your piece of paper)
- $C$  is a subset of the 2-d plane
  - $C = \{x, y : x \in [0, 1], y \in [0, 1]\}$
  - $C = \{x, y : x^2 + y^2 \leq 1\}$
  - $C = \{x, y : x > y, x, y \in (0, 2)\}$
  - $C = \{x, y : x \in A, y \in B\}$

# Examples of Joint PDFs

# Examples of Joint PDFs

- We're going to focus (initially) on pdfs of **two** random variables

# Examples of Joint PDFs

- We're going to focus (initially) on pdfs of **two** random variables
- Consider a function  $f : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$

# Examples of Joint PDFs

- We're going to focus (initially) on pdfs of **two** random variables
- Consider a function  $f : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ 
  - Input: an  $x$  value and a  $y$  value.

# Examples of Joint PDFs


- We're going to focus (initially) on pdfs of **two** random variables
- Consider a function  $f : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ 
  - Input: an  $x$  value and a  $y$  value.
  - Output: a number from the real line

# Examples of Joint PDFs

- We're going to focus (initially) on pdfs of **two** random variables
- Consider a function  $f : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ 
  - Input: an  $x$  value and a  $y$  value.
  - Output: a number from the real line
  - $f(x,y) = a$


# Examples of Joint PDFs

- We're going to focus (initially) on pdfs of **two** random variables
- Consider a function  $f : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ 
  - Input: an  $x$  value and a  $y$  value.
  - Output: a number from the real line
  - $f(x,y) = a$




# Examples of Joint PDFs

- We're going to focus (initially) on pdfs of **two** random variables
- Consider a function  $f : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ 
  - Input: an  $x$  value and a  $y$  value.
  - Output: a number from the real line
  - $f(x,y) = a$




# Examples of Joint PDFs

- We're going to focus (initially) on pdfs of **two** random variables
- Consider a function  $f : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ 
  - Input: an  $x$  value and a  $y$  value.
  - Output: a number from the real line
  - $f(x,y) = a$




# Examples of Joint PDFs

- We're going to focus (initially) on pdfs of **two** random variables
- Consider a function  $f : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ 
  - Input: an  $x$  value and a  $y$  value.
  - Output: a number from the real line
  - $f(x,y) = a$




# Examples of Joint PDFs

- We're going to focus (initially) on pdfs of **two** random variables
- Consider a function  $f : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ 
  - Input: an  $x$  value and a  $y$  value.
  - Output: a number from the real line
  - $f(x,y) = a$




# Equivalently: Contour Plots

Aerial view of probability density function: contour plots




# Equivalently: Contour Plots

Aerial view of probability density function: contour plots



# Equivalently: Contour Plots

Aerial view of probability density function: contour plots




# Example 3D-Contour Plots

Joint distribution of  $X$  and  $Y$ .

# Example 3D-Contour Plots

Joint distribution of  $X$  and  $Y$ .


1)  $f(x, y) = 1$  if  $x \in [0, 1], y \in [0, 1]$ ,  $f(x, y) = 0$



# Example 3D-Contour Plots

Joint distribution of  $X$  and  $Y$ .


1)  $f(x, y) = 1$  if  $x \in [0, 1], y \in [0, 1]$ ,  $f(x, y) = 0$



# Example 3D-Contour Plots

Joint distribution of  $X$  and  $Y$ .


2)  $f(x, y) = 2 \exp(-2y) \exp(-x)$  if  $x \in [0, \infty)$ ,  $y \in [0, \infty)$ ,  $f(x, y) = 0$  otherwise



# Example 3D-Contour Plots

Joint distribution of  $X$  and  $Y$ .


2)  $f(x, y) = 2 \exp(-2y) \exp(-x)$  if  $x \in [0, \infty)$ ,  $y \in [0, \infty)$ ,  $f(x, y) = 0$  otherwise



# Example 3D-Contour Plots

Joint distribution of  $X$  and  $Y$ .


3)  $f(x, y) = x + y$ , if  $x \in [0, 1], y \in [0, 1]$



# Example 3D-Contour Plots

Joint distribution of  $X$  and  $Y$ .

3)  $f(x, y) = x + y$ , if  $x \in [0, 1]$ ,  $y \in [0, 1]$




## Definition

### Multivariate Cumulative Density Function


For jointly continuous random variables  $X$  and  $Y$  define,  $F(b, a)$  as

$$F(b, a) = P\{X \leq b, Y \leq a\} = \int_{-\infty}^a \int_{-\infty}^b f(x, y) dx dy$$

# A Picture




# A Picture



$$F(b, a) = \int_{-\infty}^a \int_{-\infty}^b f(x, y) dx dy$$

# A Picture




Examples:

$$- F(-1, -0.5)$$

$$F(b, a) = \int_{-\infty}^a \int_{-\infty}^b f(x, y) dx dy$$

# A Picture




Examples:

$$- F(-0.5, -1.25)$$

$$F(b, a) = \int_{-\infty}^a \int_{-\infty}^b f(x, y) dx dy$$

# A Picture




Examples:

- $F(0, 0)$

$$F(b, a) = \int_{-\infty}^a \int_{-\infty}^b f(x, y) dx dy$$

# A Picture



Examples:

- $F(1.5, 0.25)$

$$F(b, a) = \int_{-\infty}^a \int_{-\infty}^b f(x, y) dx dy$$

# Marginalization

## Definition

*Moving from Joint Distributions to Univariate PDFs*

Define  $f_X(x)$  as the pdf for  $X$ ,

$$f_X(x) = \int_{-\infty}^{\infty} f(x, y) dy$$

Similarly, define  $f_Y(y)$  as the pdf for  $Y$ ,

$$f_Y(y) = \int_{-\infty}^{\infty} f(x, y) dx$$


# Conditional Probability Distribution Function

## Definition


Suppose  $X$  and  $Y$  are continuous random variables with joint pdf  $f(x, y)$ . Then define the **conditional probability function**  $f(x|y)$  as

$$f(x|y) = \frac{f(x, y)}{f_Y(y)}$$


# A Picture



# A Picture



# A Picture



# Why Does Marginalization Work?

Begin with **discrete** case.

# Why Does Marginalization Work?

Begin with **discrete** case.

Consider jointly distributed discrete random variables,  $X$  and  $Y$ . We'll suppose they have joint pmf,

# Why Does Marginalization Work?

Begin with **discrete** case.

Consider jointly distributed discrete random variables,  $X$  and  $Y$ . We'll suppose they have joint pmf,

$$P(X = x, Y = y) = p(x, y)$$

# Why Does Marginalization Work?

Begin with **discrete** case.

Consider jointly distributed discrete random variables,  $X$  and  $Y$ . We'll suppose they have joint pmf,

$$P(X = x, Y = y) = p(x, y)$$

Suppose that the distribution allocates its mass at  $x_1, x_2, \dots, x_M$  and  $y_1, y_2, \dots, y_N$ .

# Why Does Marginalization Work?

Begin with **discrete** case.

Consider jointly distributed discrete random variables,  $X$  and  $Y$ . We'll suppose they have joint pmf,

$$P(X = x, Y = y) = p(x, y)$$

Suppose that the distribution allocates its mass at  $x_1, x_2, \dots, x_M$  and  $y_1, y_2, \dots, y_N$ .

Define the conditional mass function  $P(X = x | Y = y)$  as,

# Why Does Marginalization Work?

Begin with **discrete** case.

Consider jointly distributed discrete random variables,  $X$  and  $Y$ . We'll suppose they have joint pmf,

$$P(X = x, Y = y) = p(x, y)$$

Suppose that the distribution allocates its mass at  $x_1, x_2, \dots, x_M$  and  $y_1, y_2, \dots, y_N$ .

Define the conditional mass function  $P(X = x | Y = y)$  as,

$$P(X = x | Y = y) \equiv = p(x|y)$$

# Why Does Marginalization Work?

Begin with **discrete** case.

Consider jointly distributed discrete random variables,  $X$  and  $Y$ . We'll suppose they have joint pmf,

$$P(X = x, Y = y) = p(x, y)$$

Suppose that the distribution allocates its mass at  $x_1, x_2, \dots, x_M$  and  $y_1, y_2, \dots, y_N$ .

Define the conditional mass function  $P(X = x | Y = y)$  as,

$$\begin{aligned} P(X = x | Y = y) &\equiv = p(x|y) \\ &= p(x, y)/p(y) \end{aligned}$$

# Why Does Marginalization Work?

Begin with **discrete** case.

Consider jointly distributed discrete random variables,  $X$  and  $Y$ . We'll suppose they have joint pmf,

$$P(X = x, Y = y) = p(x, y)$$

Suppose that the distribution allocates its mass at  $x_1, x_2, \dots, x_M$  and  $y_1, y_2, \dots, y_N$ .

Define the conditional mass function  $P(X = x | Y = y)$  as,

$$\begin{aligned} P(X = x | Y = y) &\equiv = p(x|y) \\ &= p(x, y)/p(y) \end{aligned}$$

Then it follows that:

# Why Does Marginalization Work?

Begin with **discrete** case.

Consider jointly distributed discrete random variables,  $X$  and  $Y$ . We'll suppose they have joint pmf,

$$P(X = x, Y = y) = p(x, y)$$

Suppose that the distribution allocates its mass at  $x_1, x_2, \dots, x_M$  and  $y_1, y_2, \dots, y_N$ .

Define the conditional mass function  $P(X = x | Y = y)$  as,

$$\begin{aligned} P(X = x | Y = y) &\equiv = p(x|y) \\ &= p(x, y)/p(y) \end{aligned}$$

Then it follows that:

$$p(x, y) = p(x|y)p(y)$$

# Why Does Marginalization Work?

Begin with **discrete** case.

Consider jointly distributed discrete random variables,  $X$  and  $Y$ . We'll suppose they have joint pmf,

$$P(X = x, Y = y) = p(x, y)$$

Suppose that the distribution allocates its mass at  $x_1, x_2, \dots, x_M$  and  $y_1, y_2, \dots, y_N$ .

Define the conditional mass function  $P(X = x | Y = y)$  as,

$$\begin{aligned} P(X = x | Y = y) &\equiv = p(x|y) \\ &= p(x, y)/p(y) \end{aligned}$$

Then it follows that:

$$p(x, y) = p(x|y)p(y)$$

Marginalizing **over**  $y$  to get  $p(x)$  is then,

# Why Does Marginalization Work?

Begin with **discrete** case.

Consider jointly distributed discrete random variables,  $X$  and  $Y$ . We'll suppose they have joint pmf,

$$P(X = x, Y = y) = p(x, y)$$

Suppose that the distribution allocates its mass at  $x_1, x_2, \dots, x_M$  and  $y_1, y_2, \dots, y_N$ .

Define the conditional mass function  $P(X = x | Y = y)$  as,

$$\begin{aligned} P(X = x | Y = y) &\equiv = p(x|y) \\ &= p(x, y)/p(y) \end{aligned}$$

Then it follows that:

$$p(x, y) = p(x|y)p(y)$$

Marginalizing **over**  $y$  to get  $p(x)$  is then,

$$p(x_j) = \sum_{i=1}^N p(x_j | y_i)p(y_i)$$

# A Table

	Y = 0	Y = 1	
X = 0	p(0,0)	p(0, 1)	$p_X(0)$
X = 1	p(1,0)	p(1,1)	$p_X(1)$
	$p_Y(0)$	$p_Y(1)$	

# A Table

	Y = 0	Y = 1	
X = 0	0.01	0.05	?
X = 1	0.25	0.69	?
	0.26	0.74	

$$p_X(0) = p(0|y=0)p(y=0) + p(0|y=1)p(y=1)$$

# A Table

	Y = 0	Y = 1	
X = 0	0.01	0.05	?
X = 1	0.25	0.69	?
	0.26	0.74	

$$\begin{aligned} p_X(0) &= p(0|y=0)p(y=0) + p(0|y=1)p(y=1) \\ &= \frac{0.01}{0.26} \times 0.26 + \frac{0.05}{0.74} \times 0.74 \end{aligned}$$

# A Table

	Y = 0	Y = 1	
X = 0	0.01	0.05	?
X = 1	0.25	0.69	?
	0.26	0.74	

$$\begin{aligned} p_X(0) &= p(0|y=0)p(y=0) + p(0|y=1)p(y=1) \\ &= \frac{0.01}{0.26} \times 0.26 + \frac{0.05}{0.74} \times 0.74 \\ &= 0.06 \end{aligned}$$

# A Table

	Y = 0	Y = 1	
X = 0	0.01	0.05	?
X = 1	0.25	0.69	?
	0.26	0.74	

$$\begin{aligned} p_X(0) &= p(0|y=0)p(y=0) + p(0|y=1)p(y=1) \\ &= \frac{0.01}{0.26} \times 0.26 + \frac{0.05}{0.74} \times 0.74 \\ &= 0.06 \end{aligned}$$

$$p_X(1) = p(1|y=0)p(y=0) + p(1|y=1)p(y=1)$$

# A Table

	Y = 0	Y = 1	
X = 0	0.01	0.05	?
X = 1	0.25	0.69	?
	0.26	0.74	

$$\begin{aligned} p_X(0) &= p(0|y=0)p(y=0) + p(0|y=1)p(y=1) \\ &= \frac{0.01}{0.26} \times 0.26 + \frac{0.05}{0.74} \times 0.74 \\ &= 0.06 \end{aligned}$$

$$\begin{aligned} p_X(1) &= p(1|y=0)p(y=0) + p(1|y=1)p(y=1) \\ &= \frac{0.25}{0.26} \times 0.26 + \frac{0.69}{0.74} \times 0.74 \end{aligned}$$

# A Table

	Y = 0	Y = 1	
X = 0	0.01	0.05	?
X = 1	0.25	0.69	?
	0.26	0.74	

$$\begin{aligned} p_X(0) &= p(0|y=0)p(y=0) + p(0|y=1)p(y=1) \\ &= \frac{0.01}{0.26} \times 0.26 + \frac{0.05}{0.74} \times 0.74 \\ &= 0.06 \end{aligned}$$

$$\begin{aligned} p_X(1) &= p(1|y=0)p(y=0) + p(1|y=1)p(y=1) \\ &= \frac{0.25}{0.26} \times 0.26 + \frac{0.69}{0.74} \times 0.74 \\ &= 0.94 \end{aligned}$$

## Move to the Continuous Case

For **jointly distributed continuous** random variables  $X$  and  $Y$  define,

# Move to the Continuous Case

For **jointly distributed continuous** random variables  $X$  and  $Y$  define,

$$f_{X|Y}(x|y) = \frac{f(x,y)}{f_Y(y)}$$

## Move to the Continuous Case

For **jointly distributed continuous** random variables  $X$  and  $Y$  define,

$$f_{X|Y}(x|y) = \frac{f(x,y)}{f_Y(y)}$$

Then, analogously, we can define

## Move to the Continuous Case

For **jointly distributed continuous** random variables  $X$  and  $Y$  define,

$$f_{X|Y}(x|y) = \frac{f(x,y)}{f_Y(y)}$$

Then, analogously, we can define

$$f_X(x) = \int_{-\infty}^{\infty} f_{X|Y}(x|y)f_Y(y)dy$$

## Move to the Continuous Case

For **jointly distributed continuous** random variables  $X$  and  $Y$  define,

$$f_{X|Y}(x|y) = \frac{f(x,y)}{f_Y(y)}$$

Then, analogously, we can define

$$f_X(x) = \int_{-\infty}^{\infty} f_{X|Y}(x|y)f_Y(y)dy$$

In words:

## Move to the Continuous Case

For **jointly distributed continuous** random variables  $X$  and  $Y$  define,

$$f_{X|Y}(x|y) = \frac{f(x,y)}{f_Y(y)}$$

Then, analogously, we can define

$$f_X(x) = \int_{-\infty}^{\infty} f_{X|Y}(x|y)f_Y(y)dy$$

In words:

- Think of  $f_{X|Y}(x|y)$  as the pdf for  $X$  at a value of  $Y$ .

## Move to the Continuous Case

For **jointly distributed continuous** random variables  $X$  and  $Y$  define,

$$f_{X|Y}(x|y) = \frac{f(x,y)}{f_Y(y)}$$

Then, analogously, we can define

$$f_X(x) = \int_{-\infty}^{\infty} f_{X|Y}(x|y)f_Y(y)dy$$

In words:

- Think of  $f_{X|Y}(x|y)$  as the pdf for  $X$  at a value of  $Y$ .
- We average over those pdfs to get the final pdf for  $X$  (want densities where there is lots of area of  $Y$  to receive lots of weight, the densities without much area from  $Y$  should receive little weight)

# A (Simple) Example

Suppose  $X$  and  $Y$  are jointly continuous and that

## A (Simple) Example


Suppose  $X$  and  $Y$  are jointly continuous and that

$$f(x, y) = x + y, \text{ if } x \in [0, 1], y \in [0, 1]$$

# A (Simple) Example

Suppose  $X$  and  $Y$  are jointly continuous and that

$$\begin{aligned}f(x, y) &= x + y, \text{ if } x \in [0, 1], y \in [0, 1] \\&= 0, \text{ otherwise}\end{aligned}$$




# A (Simple) Example

Suppose  $X$  and  $Y$  are jointly continuous and that

$$\begin{aligned}f(x, y) &= x + y, \text{ if } x \in [0, 1], y \in [0, 1] \\&= 0, \text{ otherwise}\end{aligned}$$

We want  $f_X(x)$ . Assume we have  $f_Y(y) = 1/2 + y$ .




## A (Simple) Example

Suppose  $X$  and  $Y$  are jointly continuous and that

$$\begin{aligned}f(x, y) &= x + y, \text{ if } x \in [0, 1], y \in [0, 1] \\&= 0, \text{ otherwise}\end{aligned}$$

We want  $f_X(x)$ . Assume we have  $f_Y(y) = 1/2 + y$ .

Then:  $f(x|y) = \frac{x+y}{1/2+y}$ .




## A (Simple) Example

Suppose  $X$  and  $Y$  are jointly continuous and that

$$\begin{aligned}f(x, y) &= x + y, \text{ if } x \in [0, 1], y \in [0, 1] \\&= 0, \text{ otherwise}\end{aligned}$$

We want  $f_X(x)$ . Assume we have  $f_Y(y) = 1/2 + y$ .

Then:  $f(x|y) = \frac{x+y}{1/2+y}$ .




# A (Simple) Example

Suppose  $X$  and  $Y$  are jointly continuous and that

$$\begin{aligned}f(x, y) &= x + y, \text{ if } x \in [0, 1], y \in [0, 1] \\&= 0, \text{ otherwise}\end{aligned}$$

We want  $f_X(x)$ . Assume we have  $f_Y(y) = 1/2 + y$ .

Then:  $f(x|y) = \frac{x+y}{1/2+y}$ .




## A (Simple) Example

Suppose  $X$  and  $Y$  are jointly continuous and that


$$\begin{aligned}f(x, y) &= x + y, \text{ if } x \in [0, 1], y \in [0, 1] \\&= 0, \text{ otherwise}\end{aligned}$$

We want  $f_X(x)$ . Assume we have  $f_Y(y) = 1/2 + y$ .


Then:  $f(x|y) = \frac{x+y}{1/2+y}$ .  $f(x) = \int_0^1 f(x|y)f(y)dy = 1/2 + x$




# A (Simple) Example



# A (Simple) Example



# A (Simple) Example



# A (Simple) Example

## Example

(Ross, Example 1)

Suppose  $X$  and  $Y$  are jointly distributed with pdf (for all  $x > 0, y > 0$ )

$$f(x, y) = 2 \exp(-x) \exp(-2y)$$

=

# Example

(Ross, Example 1)

Suppose  $X$  and  $Y$  are jointly distributed with pdf (for all  $x > 0, y > 0$ )

$$f(x, y) = 2 \exp(-x) \exp(-2y)$$

- 1) Verify this is a pdf

=

## Example

(Ross, Example 1)

Suppose  $X$  and  $Y$  are jointly distributed with pdf (for all  $x > 0, y > 0$ )

$$f(x, y) = 2 \exp(-x) \exp(-2y)$$

- 1) Verify this is a pdf

$$\int_0^\infty \int_0^\infty f(x, y) dx dy =$$

## Example

(Ross, Example 1)

Suppose  $X$  and  $Y$  are jointly distributed with pdf (for all  $x > 0, y > 0$ )

$$f(x, y) = 2 \exp(-x) \exp(-2y)$$

- 1) Verify this is a pdf

$$\int_0^\infty \int_0^\infty f(x, y) dx dy = 2 \int_0^\infty \int_0^\infty \exp(-x) \exp(-2y) dx dy$$

## Example

(Ross, Example 1)

Suppose  $X$  and  $Y$  are jointly distributed with pdf (for all  $x > 0, y > 0$ )

$$f(x, y) = 2 \exp(-x) \exp(-2y)$$

- 1) Verify this is a pdf

$$\begin{aligned} \int_0^\infty \int_0^\infty f(x, y) dx dy &= 2 \int_0^\infty \int_0^\infty \exp(-x) \exp(-2y) dx dy \\ &= 2 \int_0^\infty \exp(-2y) dy \int_0^\infty \exp(-x) dx \end{aligned}$$

## Example

(Ross, Example 1)

Suppose  $X$  and  $Y$  are jointly distributed with pdf (for all  $x > 0, y > 0$ )

$$f(x, y) = 2 \exp(-x) \exp(-2y)$$

- 1) Verify this is a pdf

$$\begin{aligned} \int_0^\infty \int_0^\infty f(x, y) dx dy &= 2 \int_0^\infty \int_0^\infty \exp(-x) \exp(-2y) dx dy \\ &= 2 \int_0^\infty \exp(-2y) dy \int_0^\infty \exp(-x) dx \\ &= 2 \left( -\frac{1}{2} \exp(-2y) \Big|_0^\infty \right) \left( -\exp(-x) \Big|_0^\infty \right) \end{aligned}$$

## Example

(Ross, Example 1)

Suppose  $X$  and  $Y$  are jointly distributed with pdf (for all  $x > 0, y > 0$ )

$$f(x, y) = 2 \exp(-x) \exp(-2y)$$

- 1) Verify this is a pdf

$$\begin{aligned} \int_0^\infty \int_0^\infty f(x, y) dx dy &= 2 \int_0^\infty \int_0^\infty \exp(-x) \exp(-2y) dx dy \\ &= 2 \int_0^\infty \exp(-2y) dy \int_0^\infty \exp(-x) dx \\ &= 2 \left( -\frac{1}{2} \exp(-2y) \Big|_0^\infty \right) \left( -\exp(-x) \Big|_0^\infty \right) \\ &= 2 \left[ \left( -\frac{1}{2} (\lim_{y \rightarrow \infty} \exp(-2y) - 1) \right) \left( -(\lim_{x \rightarrow \infty} \exp(-x) - 1) \right) \right] \end{aligned}$$

## Example

(Ross, Example 1)

Suppose  $X$  and  $Y$  are jointly distributed with pdf (for all  $x > 0, y > 0$ )

$$f(x, y) = 2 \exp(-x) \exp(-2y)$$

- 1) Verify this is a pdf

$$\begin{aligned} \int_0^\infty \int_0^\infty f(x, y) dx dy &= 2 \int_0^\infty \int_0^\infty \exp(-x) \exp(-2y) dx dy \\ &= 2 \int_0^\infty \exp(-2y) dy \int_0^\infty \exp(-x) dx \\ &= 2 \left( -\frac{1}{2} \exp(-2y) \Big|_0^\infty \right) \left( -\exp(-x) \Big|_0^\infty \right) \\ &= 2 \left[ \left( -\frac{1}{2} (\lim_{y \rightarrow \infty} \exp(-2y) - 1) \right) \left( -(\lim_{x \rightarrow \infty} \exp(-x) - 1) \right) \right] \\ &= 2 \left[ -\frac{1}{2} (-1) \times -1(-1) \right] \end{aligned}$$

## Example

(Ross, Example 1)

Suppose  $X$  and  $Y$  are jointly distributed with pdf (for all  $x > 0, y > 0$ )

$$f(x, y) = 2 \exp(-x) \exp(-2y)$$

- 1) Verify this is a pdf

$$\begin{aligned} \int_0^\infty \int_0^\infty f(x, y) dx dy &= 2 \int_0^\infty \int_0^\infty \exp(-x) \exp(-2y) dx dy \\ &= 2 \int_0^\infty \exp(-2y) dy \int_0^\infty \exp(-x) dx \\ &= 2 \left( -\frac{1}{2} \exp(-2y) \Big|_0^\infty \right) \left( -\exp(-x) \Big|_0^\infty \right) \\ &= 2 \left[ \left( -\frac{1}{2} \left( \lim_{y \rightarrow \infty} \exp(-2y) - 1 \right) \right) \left( -\left( \lim_{x \rightarrow \infty} \exp(-x) - 1 \right) \right) \right] \\ &= 2 \left[ -\frac{1}{2} (-1) \times -1 (-1) \right] \\ &= 1 \end{aligned}$$

## 2) Calculate CDF

## 2) Calculate CDF

$$F(x, y) \equiv P\{X \leq b, Y \leq a\} = 2 \int_0^a \int_0^b \exp(-x) \exp(-2y) dx dy$$

## 2) Calculate CDF

$$\begin{aligned} F(x, y) \equiv P\{X \leq b, Y \leq a\} &= 2 \int_0^a \int_0^b \exp(-x) \exp(-2y) dx dy \\ &= 2 \left( \int_0^a \exp(-2y) dy \right) \left( \int_0^b \exp(-x) dx \right) \end{aligned}$$

## 2) Calculate CDF

$$\begin{aligned}F(x, y) \equiv P\{X \leq b, Y \leq a\} &= 2 \int_0^a \int_0^b \exp(-x) \exp(-2y) dx dy \\&= 2 \left( \int_0^a \exp(-2y) dy \right) \left( \int_0^b \exp(-x) dx \right) \\&= 2 \left[ -\frac{1}{2} (\exp(-2a) - 1) \right] [-(\exp(-b) - 1)]\end{aligned}$$

## 2) Calculate CDF

$$\begin{aligned}F(x, y) \equiv P\{X \leq b, Y \leq a\} &= 2 \int_0^a \int_0^b \exp(-x) \exp(-2y) dx dy \\&= 2 \left( \int_0^a \exp(-2y) dy \right) \left( \int_0^b \exp(-x) dx \right) \\&= 2 \left[ -\frac{1}{2} (\exp(-2a) - 1) \right] [-(\exp(-b) - 1)] \\&= [1 - \exp(-2a)] [1 - \exp(-b)]\end{aligned}$$

3) Calculate  $f_X(x)$  and  $f_Y(y)$

3) Calculate  $f_X(x)$  and  $f_Y(y)$

$$f_X(x) = \int_0^{\infty} 2 \exp(-x) \exp(-2y) dy$$

3) Calculate  $f_X(x)$  and  $f_Y(y)$

$$\begin{aligned}f_X(x) &= \int_0^{\infty} 2 \exp(-x) \exp(-2y) dy \\&= 2 \exp(-x) \int_0^{\infty} \exp(-2y) dy\end{aligned}$$

3) Calculate  $f_X(x)$  and  $f_Y(y)$

$$\begin{aligned}f_X(x) &= \int_0^{\infty} 2 \exp(-x) \exp(-2y) dy \\&= 2 \exp(-x) \int_0^{\infty} \exp(-2y) dy \\&= 2 \exp(-x) \left[ -\frac{1}{2}(0 - 1) \right]\end{aligned}$$

3) Calculate  $f_X(x)$  and  $f_Y(y)$

$$\begin{aligned}f_X(x) &= \int_0^{\infty} 2 \exp(-x) \exp(-2y) dy \\&= 2 \exp(-x) \int_0^{\infty} \exp(-2y) dy \\&= 2 \exp(-x) \left[ -\frac{1}{2}(0 - 1) \right] \\&= \exp(-x)\end{aligned}$$

3) Calculate  $f_X(x)$  and  $f_Y(y)$

$$\begin{aligned}f_X(x) &= \int_0^{\infty} 2 \exp(-x) \exp(-2y) dy \\&= 2 \exp(-x) \int_0^{\infty} \exp(-2y) dy \\&= 2 \exp(-x) \left[ -\frac{1}{2}(0 - 1) \right] \\&= \exp(-x)\end{aligned}$$

$$f_Y(y) = \int_0^{\infty} 2 \exp(-x) \exp(-2y) dx$$

3) Calculate  $f_X(x)$  and  $f_Y(y)$

$$\begin{aligned}f_X(x) &= \int_0^{\infty} 2 \exp(-x) \exp(-2y) dy \\&= 2 \exp(-x) \int_0^{\infty} \exp(-2y) dy \\&= 2 \exp(-x) \left[ -\frac{1}{2}(0 - 1) \right] \\&= \exp(-x)\end{aligned}$$

$$\begin{aligned}f_Y(y) &= \int_0^{\infty} 2 \exp(-x) \exp(-2y) dx \\&= 2 \exp(-2y) \int_0^{\infty} \exp(-x) dx\end{aligned}$$

3) Calculate  $f_X(x)$  and  $f_Y(y)$

$$\begin{aligned}f_X(x) &= \int_0^{\infty} 2 \exp(-x) \exp(-2y) dy \\&= 2 \exp(-x) \int_0^{\infty} \exp(-2y) dy \\&= 2 \exp(-x) \left[ -\frac{1}{2}(0 - 1) \right] \\&= \exp(-x)\end{aligned}$$

$$\begin{aligned}f_Y(y) &= \int_0^{\infty} 2 \exp(-x) \exp(-2y) dx \\&= 2 \exp(-2y) \int_0^{\infty} \exp(-x) dx \\&= 2 \exp(-2y) [-(0 - 1)]\end{aligned}$$

3) Calculate  $f_X(x)$  and  $f_Y(y)$

$$\begin{aligned}f_X(x) &= \int_0^{\infty} 2 \exp(-x) \exp(-2y) dy \\&= 2 \exp(-x) \int_0^{\infty} \exp(-2y) dy \\&= 2 \exp(-x) \left[ -\frac{1}{2}(0 - 1) \right] \\&= \exp(-x)\end{aligned}$$

$$\begin{aligned}f_Y(y) &= \int_0^{\infty} 2 \exp(-x) \exp(-2y) dx \\&= 2 \exp(-2y) \int_0^{\infty} \exp(-x) dx \\&= 2 \exp(-2y) [-(0 - 1)] \\&= 2 \exp(-2y)\end{aligned}$$

## Definition

*Two random variables  $X$  and  $Y$  are independent if for any two sets of real numbers  $A$  and  $B$ ,*

## Definition

*Two random variables  $X$  and  $Y$  are independent if for any two sets of real numbers  $A$  and  $B$ ,*

$$P\{X \in A, Y \in B\} = P\{X \in A\}P\{Y \in B\}$$

## Definition

*Two random variables  $X$  and  $Y$  are independent if for any two sets of real numbers  $A$  and  $B$ ,*

$$P\{X \in A, Y \in B\} = P\{X \in A\}P\{Y \in B\}$$

*Equivalently we will say  $X$  and  $Y$  are independent if,*

## Definition

*Two random variables  $X$  and  $Y$  are independent if for any two sets of real numbers  $A$  and  $B$ ,*

$$P\{X \in A, Y \in B\} = P\{X \in A\}P\{Y \in B\}$$

*Equivalently we will say  $X$  and  $Y$  are independent if,*

$$f(x, y) = f_X(x)f_Y(y)$$

## Definition

*Two random variables  $X$  and  $Y$  are independent if for any two sets of real numbers  $A$  and  $B$ ,*

$$P\{X \in A, Y \in B\} = P\{X \in A\}P\{Y \in B\}$$

*Equivalently we will say  $X$  and  $Y$  are independent if,*

$$f(x, y) = f_X(x)f_Y(y)$$

*If  $X$  and  $Y$  are not independent, we will say they are **dependent***

# Conditional Distribution

If  $X$  and  $Y$  are independent, then

$$\begin{aligned}f_{X|Y}(x|y) &= \frac{f(x,y)}{f_Y(y)} \\&= \frac{f_X(x)f_Y(y)}{f_Y(y)} \\&= f_X(x)\end{aligned}$$

In words: the distribution of  $X$  does not change as levels of  $Y$  change.

# A (Simple) Example of Dependence

Suppose  $X$  and  $Y$  are jointly continuous and that

# A (Simple) Example of Dependence

Suppose  $X$  and  $Y$  are jointly continuous and that

$$f(x, y) = x + y, \text{ if } x \in [0, 1], y \in [0, 1]$$

## A (Simple) Example of Dependence

Suppose  $X$  and  $Y$  are jointly continuous and that

$$\begin{aligned}f(x, y) &= x + y, \text{ if } x \in [0, 1], y \in [0, 1] \\&= 0, \text{ otherwise}\end{aligned}$$

## A (Simple) Example of Dependence

Suppose  $X$  and  $Y$  are jointly continuous and that

$$\begin{aligned}f(x, y) &= x + y, \text{ if } x \in [0, 1], y \in [0, 1] \\&= 0, \text{ otherwise}\end{aligned}$$

$$f_X(x) = \int_0^1 (x + y) dy$$

## A (Simple) Example of Dependence

Suppose  $X$  and  $Y$  are jointly continuous and that

$$\begin{aligned}f(x, y) &= x + y, \text{ if } x \in [0, 1], y \in [0, 1] \\&= 0, \text{ otherwise}\end{aligned}$$

$$\begin{aligned}f_X(x) &= \int_0^1 (x + y) dy \\&= xy + \frac{y^2}{2} \Big|_0^1\end{aligned}$$

# A (Simple) Example of Dependence

Suppose  $X$  and  $Y$  are jointly continuous and that

$$\begin{aligned}f(x, y) &= x + y, \text{ if } x \in [0, 1], y \in [0, 1] \\&= 0, \text{ otherwise}\end{aligned}$$

$$\begin{aligned}f_X(x) &= \int_0^1 (x + y) dy \\&= xy + \frac{y^2}{2} \Big|_0^1 \\&= x + \frac{1}{2}\end{aligned}$$

# A (Simple) Example of Dependence

Suppose  $X$  and  $Y$  are jointly continuous and that

$$\begin{aligned}f(x, y) &= x + y, \text{ if } x \in [0, 1], y \in [0, 1] \\&= 0, \text{ otherwise}\end{aligned}$$

$$\begin{aligned}f_X(x) &= \int_0^1 (x + y) dy \\&= xy + \frac{y^2}{2} \Big|_0^1 \\&= x + \frac{1}{2} \\f_Y(y) &= \frac{1}{2} + y\end{aligned}$$

# A (Simple) Example of Dependence

Suppose  $X$  and  $Y$  are jointly continuous and that

$$\begin{aligned}f(x, y) &= x + y, \text{ if } x \in [0, 1], y \in [0, 1] \\&= 0, \text{ otherwise}\end{aligned}$$

$$f(x, y) = x + y$$

# A (Simple) Example of Dependence

Suppose  $X$  and  $Y$  are jointly continuous and that

$$\begin{aligned}f(x, y) &= x + y, \text{ if } x \in [0, 1], y \in [0, 1] \\&= 0, \text{ otherwise}\end{aligned}$$

$$\begin{aligned}f(x, y) &= x + y \\f_X(x)f_Y(y) &= \left(\frac{1}{2} + x\right)\left(\frac{1}{2} + y\right)\end{aligned}$$

# A (Simple) Example of Dependence

Suppose  $X$  and  $Y$  are jointly continuous and that

$$\begin{aligned}f(x, y) &= x + y, \text{ if } x \in [0, 1], y \in [0, 1] \\&= 0, \text{ otherwise}\end{aligned}$$

$$\begin{aligned}f(x, y) &= x + y \\f_X(x)f_Y(y) &= \left(\frac{1}{2} + x\right)\left(\frac{1}{2} + y\right) \\&= \frac{1}{4} + \frac{x+y}{2} + xy\end{aligned}$$

# A (Simple) Example of Dependence

Suppose  $X$  and  $Y$  are jointly continuous and that

$$\begin{aligned}f(x, y) &= x + y, \text{ if } x \in [0, 1], y \in [0, 1] \\&= 0, \text{ otherwise}\end{aligned}$$

$$\begin{aligned}f(x, y) &= x + y \\f_X(x)f_Y(y) &= \left(\frac{1}{2} + x\right)\left(\frac{1}{2} + y\right) \\&= \frac{1}{4} + \frac{x+y}{2} + xy\end{aligned}$$

Intuition: at different levels of  $X$  the distribution on  $Y$  behaves differently.

# A (Simple) Example of Dependence

Suppose  $X$  and  $Y$  are jointly continuous and that

$$\begin{aligned}f(x, y) &= x + y, \text{ if } x \in [0, 1], y \in [0, 1] \\&= 0, \text{ otherwise}\end{aligned}$$

$$\begin{aligned}f(x, y) &= x + y \\f_X(x)f_Y(y) &= \left(\frac{1}{2} + x\right)\left(\frac{1}{2} + y\right) \\&= \frac{1}{4} + \frac{x+y}{2} + xy\end{aligned}$$

Intuition: at different levels of  $X$  the distribution on  $Y$  behaves differently.  
 **$X$  provides information about  $Y$**

# Expectation

## Definition

For jointly continuous random variables  $X$  and  $Y$  define,

$$E[X] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xf(x, y) dx dy$$

$$E[Y] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} yf(x, y) dx dy$$

$$E[XY] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xyf(x, y) dx dy$$

## Proposition

Suppose  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  (that isn't crazy). Then,

$$E[g(X, Y)] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x, y)f(x, y) dx dy$$

# Covariance

## Definition

*For jointly continuous random variables  $X$  and  $Y$  define, the covariance of  $X$  and  $Y$  as,*

# Covariance

## Definition

*For jointly continuous random variables  $X$  and  $Y$  define, the covariance of  $X$  and  $Y$  as,*

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])]$$

# Covariance

## Definition

*For jointly continuous random variables  $X$  and  $Y$  define, the covariance of  $X$  and  $Y$  as,*

$$\begin{aligned}\text{cov}(X, Y) &= E[(X - E[X])(Y - E[Y])] \\ &= E[XY - E[X]Y - E[Y]X + E[X]E[Y]]\end{aligned}$$

# Covariance

## Definition

*For jointly continuous random variables  $X$  and  $Y$  define, the covariance of  $X$  and  $Y$  as,*

$$\begin{aligned}\text{cov}(X, Y) &= E[(X - E[X])(Y - E[Y])] \\ &= E[XY - E[X]Y - E[Y]X + E[X]E[Y]] \\ &= E[XY] - 2E[X]E[Y] + E[E[X]E[Y]]\end{aligned}$$

# Covariance

## Definition

*For jointly continuous random variables  $X$  and  $Y$  define, the covariance of  $X$  and  $Y$  as,*

$$\begin{aligned}\text{cov}(X, Y) &= E[(X - E[X])(Y - E[Y])] \\ &= E[XY - E[X]Y - E[Y]X + E[X]E[Y]] \\ &= E[XY] - 2E[X]E[Y] + E[E[X]E[Y]] \\ &= E[XY] - E[X]E[Y]\end{aligned}$$

# Covariance

## Definition

*For jointly continuous random variables  $X$  and  $Y$  define, the covariance of  $X$  and  $Y$  as,*

$$\begin{aligned}\text{cov}(X, Y) &= E[(X - E[X])(Y - E[Y])] \\ &= E[XY - E[X]Y - E[Y]X + E[X]E[Y]] \\ &= E[XY] - 2E[X]E[Y] + E[E[X]E[Y]] \\ &= E[XY] - E[X]E[Y]\end{aligned}$$

*Define the correlation of  $X$  and  $Y$  as,*

# Covariance

## Definition

For jointly continuous random variables  $X$  and  $Y$  define, the covariance of  $X$  and  $Y$  as,

$$\begin{aligned}\text{cov}(X, Y) &= E[(X - E[X])(Y - E[Y])] \\ &= E[XY - E[X]Y - E[Y]X + E[X]E[Y]] \\ &= E[XY] - 2E[X]E[Y] + E[E[X]E[Y]] \\ &= E[XY] - E[X]E[Y]\end{aligned}$$

Define the correlation of  $X$  and  $Y$  as,

$$\text{cor}(X, Y) = \frac{\text{cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}$$

# Some Observations

Variance is the covariance of a random variable with itself.

# Some Observations

Variance is the covariance of a random variable with itself.

$$\begin{aligned}\text{cov}(X, X) &= E[XX] - E[X]E[X] \\ &= E[X^2] - E[X]^2\end{aligned}$$

## Some Observations

Variance is the covariance of a random variable with itself.

$$\begin{aligned}\text{cov}(X, X) &= E[XX] - E[X]E[X] \\ &= E[X^2] - E[X]^2\end{aligned}$$

Correlation measures the linear relationship between two random variables

## Some Observations

Variance is the covariance of a random variable with itself.

$$\begin{aligned}\text{cov}(X, X) &= E[XX] - E[X]E[X] \\ &= E[X^2] - E[X]^2\end{aligned}$$

Correlation measures the linear relationship between two random variables

Suppose  $X = Y$

# Some Observations

Variance is the covariance of a random variable with itself.

$$\begin{aligned}\text{cov}(X, X) &= E[XX] - E[X]E[X] \\ &= E[X^2] - E[X]^2\end{aligned}$$

Correlation measures the linear relationship between two random variables

Suppose  $X = Y$

$$\text{cor}(X, Y) = \frac{\text{cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}$$

# Some Observations

Variance is the covariance of a random variable with itself.

$$\begin{aligned}\text{cov}(X, X) &= E[XX] - E[X]E[X] \\ &= E[X^2] - E[X]^2\end{aligned}$$

Correlation measures the linear relationship between two random variables

Suppose  $X = Y$

$$\begin{aligned}\text{cor}(X, Y) &= \frac{\text{cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} \\ &= \frac{\text{Var}(X)}{\text{Var}(X)}\end{aligned}$$

# Some Observations

Variance is the covariance of a random variable with itself.

$$\begin{aligned}\text{cov}(X, X) &= E[XX] - E[X]E[X] \\ &= E[X^2] - E[X]^2\end{aligned}$$

Correlation measures the linear relationship between two random variables

Suppose  $X = Y$

$$\begin{aligned}\text{cor}(X, Y) &= \frac{\text{cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} \\ &= \frac{\text{Var}(X)}{\text{Var}(X)} \\ &= 1\end{aligned}$$

# Some Observations

Variance is the covariance of a random variable with itself.

$$\begin{aligned}\text{cov}(X, X) &= E[XX] - E[X]E[X] \\ &= E[X^2] - E[X]^2\end{aligned}$$

Correlation measures the linear relationship between two random variables

Suppose  $X = Y$

$$\begin{aligned}\text{cor}(X, Y) &= \frac{\text{cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} \\ &= \frac{\text{Var}(X)}{\text{Var}(X)} \\ &= 1\end{aligned}$$

Suppose  $X = -Y$

# Some Observations

Variance is the covariance of a random variable with itself.

$$\begin{aligned}\text{cov}(X, X) &= E[XX] - E[X]E[X] \\ &= E[X^2] - E[X]^2\end{aligned}$$

Correlation measures the linear relationship between two random variables

Suppose  $X = Y$

$$\begin{aligned}\text{cor}(X, Y) &= \frac{\text{cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} \\ &= \frac{\text{Var}(X)}{\text{Var}(X)} \\ &= 1\end{aligned}$$

Suppose  $X = -Y$

$$\text{cor}(X, Y) = \frac{\text{cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}$$

# Some Observations

Variance is the covariance of a random variable with itself.

$$\begin{aligned}\text{cov}(X, X) &= E[XX] - E[X]E[X] \\ &= E[X^2] - E[X]^2\end{aligned}$$

Correlation measures the linear relationship between two random variables

Suppose  $X = Y$

$$\begin{aligned}\text{cor}(X, Y) &= \frac{\text{cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} \\ &= \frac{\text{Var}(X)}{\text{Var}(X)} \\ &= 1\end{aligned}$$

Suppose  $X = -Y$

$$\begin{aligned}\text{cor}(X, Y) &= \frac{\text{cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} \\ &= \frac{-\text{Var}(X)}{\text{Var}(X)}\end{aligned}$$

# Correlation is Between -1 and 1

$$|cor(X, Y)| \leq 1$$

- Proof 1: Variance trick
- Proof 2: Cauchy-Schwartz Inequality
  - “Inner product” of any two vectors  $X$  and  $Y$  is less than or equal to the length of vector  $X$  times the length of vector  $Y$

## Example: $X + Y$

Suppose  $X$  and  $Y$  have pdf  $x + y$  for  $x, y \in [0, 1]$ .

## Example: $X + Y$

Suppose  $X$  and  $Y$  have pdf  $x + y$  for  $x, y \in [0, 1]$ .

$\text{Cov}(X, Y)$

## Example: $X + Y$

Suppose  $X$  and  $Y$  have pdf  $x + y$  for  $x, y \in [0, 1]$ .

$\text{Cov}(X, Y)$

$$E[XY] = \int_0^1 \int_0^1 xy(x + y) dx dy$$

## Example: $X + Y$

Suppose  $X$  and  $Y$  have pdf  $x + y$  for  $x, y \in [0, 1]$ .

$\text{Cov}(X, Y)$

$$\begin{aligned} E[XY] &= \int_0^1 \int_0^1 xy(x+y) dx dy \\ &= \int_0^1 \int_0^1 (x^2y + y^2x) dx dy \end{aligned}$$

## Example: $X + Y$

Suppose  $X$  and  $Y$  have pdf  $x + y$  for  $x, y \in [0, 1]$ .

$\text{Cov}(X, Y)$

$$\begin{aligned} E[XY] &= \int_0^1 \int_0^1 xy(x+y) dx dy \\ &= \int_0^1 \int_0^1 (x^2y + y^2x) dx dy \\ &= \int_0^1 \left(\frac{y}{3} + \frac{y^2}{2}\right) dy \end{aligned}$$

## Example: $X + Y$

Suppose  $X$  and  $Y$  have pdf  $x + y$  for  $x, y \in [0, 1]$ .

$\text{Cov}(X, Y)$

$$\begin{aligned} E[XY] &= \int_0^1 \int_0^1 xy(x+y) dx dy \\ &= \int_0^1 \int_0^1 (x^2y + y^2x) dx dy \\ &= \int_0^1 \left( \frac{y}{3} + \frac{y^2}{2} \right) dy \\ &= \frac{1}{6} + \frac{1}{6} = \frac{1}{3} \end{aligned}$$

## Example: $X + Y$

Suppose  $X$  and  $Y$  have pdf  $x + y$  for  $x, y \in [0, 1]$ .

$\text{Cov}(X, Y)$

$$\begin{aligned} E[XY] &= \int_0^1 \int_0^1 xy(x+y) dx dy \\ &= \int_0^1 \int_0^1 (x^2y + y^2x) dx dy \\ &= \int_0^1 \left( \frac{y}{3} + \frac{y^2}{2} \right) dy \\ &= \frac{1}{6} + \frac{1}{6} = \frac{1}{3} \end{aligned}$$

$$E[X] = \int_0^1 \int_0^1 x(x+y) dx dy$$

## Example: $X + Y$

Suppose  $X$  and  $Y$  have pdf  $x + y$  for  $x, y \in [0, 1]$ .

$\text{Cov}(X, Y)$

$$\begin{aligned} E[XY] &= \int_0^1 \int_0^1 xy(x+y) dx dy \\ &= \int_0^1 \int_0^1 (x^2y + y^2x) dx dy \\ &= \int_0^1 \left( \frac{y}{3} + \frac{y^2}{2} \right) dy \\ &= \frac{1}{6} + \frac{1}{6} = \frac{1}{3} \end{aligned}$$

$$\begin{aligned} E[X] &= \int_0^1 \int_0^1 x(x+y) dx dy \\ &= \frac{7}{12} \end{aligned}$$

## Example: $X + Y$

Suppose  $X$  and  $Y$  have pdf  $x + y$  for  $x, y \in [0, 1]$ .

$\text{Cov}(X, Y)$

$$\begin{aligned} E[XY] &= \int_0^1 \int_0^1 xy(x+y) dx dy \\ &= \int_0^1 \int_0^1 (x^2y + y^2x) dx dy \\ &= \int_0^1 \left( \frac{y}{3} + \frac{y^2}{2} \right) dy \\ &= \frac{1}{6} + \frac{1}{6} = \frac{1}{3} \end{aligned}$$

$$\begin{aligned} E[Y] &= \int_0^1 \int_0^1 y(x+y) dx dy \\ &= \end{aligned}$$

## Example: $X + Y$

Suppose  $X$  and  $Y$  have pdf  $x + y$  for  $x, y \in [0, 1]$ .

$\text{Cov}(X, Y)$

$$\begin{aligned} E[XY] &= \int_0^1 \int_0^1 xy(x+y) dx dy \\ &= \int_0^1 \int_0^1 (x^2y + y^2x) dx dy \\ &= \int_0^1 \left( \frac{y}{3} + \frac{y^2}{2} \right) dy \\ &= \frac{1}{6} + \frac{1}{6} = \frac{1}{3} \end{aligned}$$

$$\begin{aligned} E[Y] &= \int_0^1 \int_0^1 y(x+y) dx dy \\ &= \frac{7}{12} \end{aligned}$$

## Example: $X + Y$

$$\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$$

## Example: $X + Y$

$$\begin{aligned}\text{Cov}(X, Y) &= E[XY] - E[X]E[Y] \\ &= \frac{1}{3} - \frac{49}{144} = -\frac{1}{144}\end{aligned}$$

## Example: $X + Y$

$$\begin{aligned}\text{Cov}(X, Y) &= E[XY] - E[X]E[Y] \\ &= \frac{1}{3} - \frac{49}{144} = -\frac{1}{144}\end{aligned}$$

$$\text{Cor}(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}$$

## Example: $X + Y$

$$\begin{aligned}\text{Cov}(X, Y) &= E[XY] - E[X]E[Y] \\ &= \frac{1}{3} - \frac{49}{144} = -\frac{1}{144}\end{aligned}$$

$$\begin{aligned}\text{Cor}(X, Y) &= \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} \\ &= \frac{-\frac{1}{144}}{\frac{11}{144}}\end{aligned}$$

## Example: $X + Y$

$$\begin{aligned}\text{Cov}(X, Y) &= E[XY] - E[X]E[Y] \\ &= \frac{1}{3} - \frac{49}{144} = -\frac{1}{144}\end{aligned}$$

$$\begin{aligned}\text{Cor}(X, Y) &= \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} \\ &= \frac{-\frac{1}{144}}{\frac{11}{144}} \\ &= \frac{-1}{11}\end{aligned}$$

# Sums of Random Variables

Suppose we have a sequence of random variables  $X_i$ ,  $i = 1, 2, \dots, N$ . Suppose that they have joint pdf,

$$f(\mathbf{x}) = f(x_1, x_2, \dots, x_n)$$

$$1) E[\sum_{i=1}^N X_i] = \sum_{i=1}^N E[X_i]$$

$$2) \text{var}(\sum_{i=1}^N X_i) = \sum_{i=1}^N \text{var}(X_i) + 2 \sum_{i < j} \text{cov}(X_i, X_j)$$

# Sums of Random Variables

## Proposition

Suppose we have a sequence of random variables  $X_i$ ,  $i = 1, 2, \dots, N$ .

Suppose that they have joint pdf,

$$f(\mathbf{x}) = f(x_1, x_2, \dots, x_n)$$

Then

$$E\left[\sum_{i=1}^N X_i\right] = \sum_{i=1}^N E[X_i]$$

Proof.



Proof.

$$E\left[\sum_{i=1}^N X_i\right] = E[X_1 + X_2 + \dots + X_N]$$



Proof.

$$\begin{aligned} E\left[\sum_{i=1}^N X_i\right] &= E[X_1 + X_2 + \dots + X_N] \\ &= \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} (x_1 + x_2 + \dots + x_N) f(x_1, x_2, \dots, x_N) dx_1 dx_2 \dots dx_N \end{aligned}$$



Proof.

$$\begin{aligned} E\left[\sum_{i=1}^N X_i\right] &= E[X_1 + X_2 + \dots + X_N] \\ &= \int_{-\infty}^{\infty} \dots \iint_{-\infty}^{\infty} (x_1 + x_2 + \dots + x_N) f(x_1, x_2, \dots, x_N) dx_1 dx_2 \dots dx_N \\ &= \int_{-\infty}^{\infty} x_1 f_{X_1}(x_1) dx_1 + \int_{-\infty}^{\infty} x_2 f_{X_2}(x_2) dx_2 + \dots + \int_{-\infty}^{\infty} x_N f_{X_N}(x_N) dx_N \end{aligned}$$



Proof.

$$\begin{aligned} E\left[\sum_{i=1}^N X_i\right] &= E[X_1 + X_2 + \dots + X_N] \\ &= \int_{-\infty}^{\infty} \dots \iint_{-\infty}^{\infty} (x_1 + x_2 + \dots + x_N) f(x_1, x_2, \dots, x_N) dx_1 dx_2 \dots dx_N \\ &= \int_{-\infty}^{\infty} x_1 f_{X_1}(x_1) dx_1 + \int_{-\infty}^{\infty} x_2 f_{X_2}(x_2) dx_2 + \dots + \int_{-\infty}^{\infty} x_N f_{X_N}(x_N) dx_N \\ &= E[X_1] + E[X_2] + \dots + E[X_N] \end{aligned}$$



# Sums of Random Variable

## Proposition

Suppose  $X_i$  is a sequence of random variables. Then

$$\text{var}\left(\sum_{i=1}^N X_i\right) = \sum_{i=1}^N \text{var}(X_i) + 2 \sum_{i < j} \text{cov}(X_i, X_j)$$

# Sums of Random Variable

Proof.

Consider two random variables,  $X_1$  and  $X_2$ . Then,

# Sums of Random Variable

Proof.

Consider two random variables,  $X_1$  and  $X_2$ . Then,

$$\text{var}(X_1 + X_2) = E[(X_1 + X_2)^2] - (E[X_1] + E[X_2])^2$$

# Sums of Random Variable

Proof.

Consider two random variables,  $X_1$  and  $X_2$ . Then,

$$\begin{aligned}\text{var}(X_1 + X_2) &= E[(X_1 + X_2)^2] - (E[X_1] + E[X_2])^2 \\ &= E[X_1^2] + 2E[X_1 X_2] + E[X_2^2] \\ &\quad - (E[X_1])^2 - 2E[X_1]E[X_2] - 2E[X_2]^2\end{aligned}$$

# Sums of Random Variable

Proof.

Consider two random variables,  $X_1$  and  $X_2$ . Then,

$$\begin{aligned}\text{var}(X_1 + X_2) &= E[(X_1 + X_2)^2] - (E[X_1] + E[X_2])^2 \\&= E[X_1^2] + 2E[X_1 X_2] + E[X_2^2] \\&\quad - (E[X_1])^2 - 2E[X_1]E[X_2] - (E[X_2])^2 \\&= \underbrace{E[X_1^2] - (E[X_1])^2}_{\text{var}(X_1)} + \underbrace{E[X_2^2] - (E[X_2])^2}_{\text{var}(X_2)} \\&\quad + 2\underbrace{(E[X_1 X_2] - E[X_1]E[X_2])}_{\text{cov}(X_1, X_2)}\end{aligned}$$

# Sums of Random Variable

Proof.

Consider two random variables,  $X_1$  and  $X_2$ . Then,

$$\begin{aligned}\text{var}(X_1 + X_2) &= E[(X_1 + X_2)^2] - (E[X_1] + E[X_2])^2 \\&= E[X_1^2] + 2E[X_1 X_2] + E[X_2^2] \\&\quad - (E[X_1])^2 - 2E[X_1]E[X_2] - (E[X_2])^2 \\&= \underbrace{E[X_1^2] - (E[X_1])^2}_{\text{var}(X_1)} + \underbrace{E[X_2^2] - (E[X_2])^2}_{\text{var}(X_2)} \\&\quad + 2\underbrace{(E[X_1 X_2] - E[X_1]E[X_2])}_{\text{cov}(X_1, X_2)} \\&= \text{var}(X_1) + \text{var}(X_2) + 2\text{cov}(X_1, X_2)\end{aligned}$$



## Definition

Suppose  $\mathbf{X} = (X_1, X_2, \dots, X_N)$  is a vector of random variables. If  $\mathbf{X}$  has pdf

$$f(\mathbf{x}) = (2\pi)^{-N/2} \det(\boldsymbol{\Sigma})^{-1/2} \exp\left(-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu})\right)$$

Then we will say  $\mathbf{X}$  is a *Multivariate Normal Distribution*,

$$\mathbf{X} \sim \text{Multivariate Normal}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

- **Regularly** used for likelihood, Bayesian, and other parametric inferences

# Multivariate Normal Distribution

Consider the (bivariate) special case where  $\mu = (0, 0)$  and

$$\Sigma = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

# Multivariate Normal Distribution

Consider the (bivariate) special case where  $\mu = (0, 0)$  and

$$\Sigma = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

Then

$$f(x_1, x_2) = (2\pi)^{-2/2} 1^{-1/2} \exp \left( -\frac{1}{2} \left( (\mathbf{x} - \mathbf{0})' \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} (\mathbf{x} - \mathbf{0}) \right) \right)$$

# Multivariate Normal Distribution

Consider the (bivariate) special case where  $\mu = (0, 0)$  and

$$\Sigma = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

Then

$$\begin{aligned} f(x_1, x_2) &= (2\pi)^{-2/2} 1^{-1/2} \exp \left( -\frac{1}{2} \left( (\mathbf{x} - \mathbf{0})' \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} (\mathbf{x} - \mathbf{0}) \right) \right) \\ &= \frac{1}{2\pi} \exp \left( -\frac{1}{2} (x_1^2 + x_2^2) \right) \end{aligned}$$

# Multivariate Normal Distribution

Consider the (bivariate) special case where  $\mu = (0, 0)$  and

$$\Sigma = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

Then

$$\begin{aligned} f(x_1, x_2) &= (2\pi)^{-2/2} 1^{-1/2} \exp \left( -\frac{1}{2} \left( (\mathbf{x} - \mathbf{0})' \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} (\mathbf{x} - \mathbf{0}) \right) \right) \\ &= \frac{1}{2\pi} \exp \left( -\frac{1}{2} (x_1^2 + x_2^2) \right) \\ &= \frac{1}{\sqrt{2\pi}} \exp \left( -\frac{x_1^2}{2} \right) \frac{1}{\sqrt{2\pi}} \exp \left( -\frac{x_2^2}{2} \right) \end{aligned}$$

# Multivariate Normal Distribution

Consider the (bivariate) special case where  $\mu = (0, 0)$  and

$$\Sigma = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

Then

$$\begin{aligned} f(x_1, x_2) &= (2\pi)^{-2/2} 1^{-1/2} \exp \left( -\frac{1}{2} \left( (\mathbf{x} - \mathbf{0})' \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} (\mathbf{x} - \mathbf{0}) \right) \right) \\ &= \frac{1}{2\pi} \exp \left( -\frac{1}{2} (x_1^2 + x_2^2) \right) \\ &= \frac{1}{\sqrt{2\pi}} \exp \left( -\frac{x_1^2}{2} \right) \frac{1}{\sqrt{2\pi}} \exp \left( -\frac{x_2^2}{2} \right) \end{aligned}$$

↪ product of univariate standard normally distributed random variables

# Standard Multivariate Normal

## Definition

Suppose  $\mathbf{Z} = (Z_1, Z_2, \dots, Z_N)$  is

$$\mathbf{Z} \sim \text{Multivariate Normal}(\mathbf{0}, \mathbf{I}_N).$$

Then we will call  $\mathbf{Z}$  the standard multivariate normal.

# Properties of the Multivariate Normal Distribution

Suppose  $\mathbf{X} = (X_1, X_2, \dots, X_N)$

$$\begin{aligned}E[\mathbf{X}] &= \boldsymbol{\mu} \\ \text{cov}(\mathbf{X}) &= \boldsymbol{\Sigma}\end{aligned}$$

So that,

$$\boldsymbol{\Sigma} = \begin{pmatrix} \text{var}(X_1) & \text{cov}(X_1, X_2) & \dots & \text{cov}(X_1, X_N) \\ \text{cov}(X_2, X_1) & \text{var}(X_2) & \dots & \text{cov}(X_2, X_N) \\ \vdots & \vdots & \ddots & \vdots \\ \text{cov}(X_N, X_1) & \text{cov}(X_N, X_2) & \dots & \text{var}(X_N) \end{pmatrix}$$

# Independence and Multivariate Normal

## Proposition

*Suppose  $X$  and  $Y$  are independent. Then*

$$\text{cov}(X, Y) = 0$$

Proof.

Suppose  $X$  and  $Y$  are independent.

Proof.

Suppose  $X$  and  $Y$  are independent.

$$\text{cov}(X, Y) = E[XY] - E[X]E[Y]$$

Proof.

Suppose  $X$  and  $Y$  are independent.

$$\text{cov}(X, Y) = E[XY] - E[X]E[Y]$$

Calculating  $E[XY]$

Proof.

Suppose  $X$  and  $Y$  are independent.

$$\text{cov}(X, Y) = E[XY] - E[X]E[Y]$$

Calculating  $E[XY]$

$$E[XY] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xyf(x, y)dxdy$$

Proof.

Suppose  $X$  and  $Y$  are independent.

$$\text{cov}(X, Y) = E[XY] - E[X]E[Y]$$

Calculating  $E[XY]$

$$\begin{aligned} E[XY] &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xyf(x, y)dxdy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xyf_X(x)f_Y(y)dxdy \end{aligned}$$

Proof.

Suppose  $X$  and  $Y$  are independent.

$$\text{cov}(X, Y) = E[XY] - E[X]E[Y]$$

Calculating  $E[XY]$

$$\begin{aligned} E[XY] &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xyf(x, y)dxdy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xyf_X(x)f_Y(y)dxdy \\ &= \int_{-\infty}^{\infty} xf_X(x)dx \int_{-\infty}^{\infty} yf_Y(y)dy \end{aligned}$$

Proof.

Suppose  $X$  and  $Y$  are independent.

$$\text{cov}(X, Y) = E[XY] - E[X]E[Y]$$

Calculating  $E[XY]$

$$\begin{aligned} E[XY] &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xyf(x, y)dxdy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xyf_X(x)f_Y(y)dxdy \\ &= \int_{-\infty}^{\infty} xf_X(x)dx \int_{-\infty}^{\infty} yf_Y(y)dy \\ &= E[X]E[Y] \end{aligned}$$

Proof.

Suppose  $X$  and  $Y$  are independent.

$$\text{cov}(X, Y) = E[XY] - E[X]E[Y]$$

Calculating  $E[XY]$

$$\begin{aligned} E[XY] &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xyf(x, y)dxdy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xyf_X(x)f_Y(y)dxdy \\ &= \int_{-\infty}^{\infty} xf_X(x)dx \int_{-\infty}^{\infty} yf_Y(y)dy \\ &= E[X]E[Y] \end{aligned}$$

Then  $\text{cov}(X, Y) = 0$ .



## Zero covariance does not generally imply Independent

Suppose  $X \in \{-1, 1\}$  with  $P(X = 1) = P(X = -1) = 1/2$ .

Suppose  $Y \in \{-1, 0, 1\}$  with  $Y = 0$  if  $X = -1$  and

$P(Y = 1) = P(Y = -1)$  if  $X = 1$ .

$$\begin{aligned} E[XY] &= \sum_{i \in \{-1, 1\}} \sum_{j \in \{-1, 0, 1\}} ijP(X = i, Y = j) \\ &= -1 \times 0 \times P(X = -1, Y = 0) + 1 \times 1 \times P(X = 1, Y = 1) \\ &\quad -1 \times 1 \times P(X = 1, Y = -1) \\ &= 0 + P(X = 1, Y = 1) - P(X = 1, Y = -1) \\ &= 0.25 - 0.25 = 0 \end{aligned}$$

$$E[X] = 0$$

$$E[Y] = 0$$

## Proposition

Suppose  $\mathbf{X} \sim \text{Multivariate Normal}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ . where  $\mathbf{X} = (X_1, X_2, \dots, X_N)$ .  
If  $\text{cov}(X_i, X_j) = 0$ , then  $X_i$  and  $X_j$  are independent

## Tomorrow

- Changing Coordinates
- Moment Generating Functions
- Famous Inequalities
- Different Notions of Convergence