

MA144 Methods of Mathematical Modelling 2, Assignment 1

Dyson Dyson

Question 1

Consider the polar curve with equation $r = f(\theta)$, where $\theta \in [a, b]$ and f is some function of θ .

Q1 (a)

Using the arc-length formula for parametric curves (in the lecture notes), show that the arc length of the polar curve is given by the integral

$$\int_a^b \sqrt{\left(\frac{df}{d\theta}\right)^2 + f^2} d\theta.$$

The polar curve with equation $r = f(\theta)$ can be parametrised as $\underline{r}(\theta) = (f(\theta) \cos \theta, f(\theta) \sin \theta)$. Then

$$\frac{d\underline{r}}{d\theta} = \left(\frac{df}{d\theta}(\theta) \cos \theta - f(\theta) \sin \theta, \frac{df}{d\theta}(\theta) \sin \theta + f(\theta) \cos \theta \right)$$

Therefore

$$\begin{aligned} \|\underline{r}'(\theta)\| &= \sqrt{(f'(\theta) \cos \theta - f(\theta) \sin \theta)^2 + (f'(\theta) \sin \theta + f(\theta) \cos \theta)^2} \\ &= \sqrt{f'(\theta)^2 \cos^2 \theta - 2f(\theta)f'(\theta) \cos \theta \sin \theta + f(\theta)^2 \sin^2 \theta} \\ &\quad + f'(\theta)^2 \sin^2 \theta + 2f(\theta)f'(\theta) \cos \theta \sin \theta + f(\theta)^2 \cos^2 \theta \\ &= \sqrt{f'(\theta)^2(\cos^2 \theta + \sin^2 \theta) + f(\theta)^2(\cos^2 \theta + \sin^2 \theta)} \\ &= \sqrt{f'(\theta)^2 + f(\theta)^2} \\ &= \sqrt{\left(\frac{df}{d\theta}\right)^2 + f^2} \end{aligned}$$

Therefore the arc length of the curve is

$$s = \int_a^b \sqrt{\left(\frac{df}{d\theta}\right)^2 + f^2} d\theta$$

as required.

Q1 (b)

Sketch the closed curve with polar equation $r = 1 + \cos \theta$. Find its arc length.




Figure 1: A sketch of $r = 1 + \cos \theta$

$\frac{dr}{d\theta} = -\sin \theta$ so the arc length is

$$\begin{aligned} s &= \int_0^{2\pi} \sqrt{\sin^2 \theta + (1 + \cos \theta)^2} d\theta \\ &= \int_0^{2\pi} \sqrt{\sin^2 \theta + 1 + 2 \cos \theta + \cos^2 \theta} d\theta \\ &= \int_0^{2\pi} \sqrt{2 + 2 \cos \theta} d\theta \end{aligned}$$

I don't know how to do this integral but apparently it's 4π .

Q1 (c)

Consider the curve with polar equation $r = 1 + \cos k\theta$, where $\theta \in \mathbb{R}$. Give a value of the constant k such that the curve is not simple. Justify your answer.

The curve given by $r = 1 + \cos k\theta$ is non-simple (self-intersecting) whenever k is not an integer.

Question 2

Consider the curve \mathcal{C} with equation

$$4y^2 - 9x^2 = 1, \quad y > 0$$

Q2 (a)

Parametrise the curve in terms of hyperbolic functions of t . Is this a regular parametrisation of the curve?

Since $\cosh^2 t - \sinh^2 t = 1$, we can let $y = \frac{1}{2} \cosh t$ and $x = \frac{1}{3} \sinh t$. Then we'll have the equation of \mathcal{C} . Therefore we can parametrise \mathcal{C} as $\underline{r}(t) = (\frac{1}{3} \sinh t, \frac{1}{2} \cosh t)$.

Since $\frac{1}{2} \cosh t > 0$ for all $t \in (-\infty, \infty)$, the requirement of $y > 0$ is satisfied by $t \in (-\infty, \infty)$, so that's the range of this parametrisation.

Q2 (b)

Sketch the curve described by your parametrisation (include an arrow to indicate its orientation). Give the equations of any asymptotes.




Figure 2: A sketch of $\underline{r}(t) = (\frac{1}{3} \sinh t, \frac{1}{2} \cosh t)$

There is a turning point at $(0, \frac{1}{2})$ and asymptotes are $y = \frac{3}{2}x$ and $y = -\frac{3}{2}x$.

Q2 (c)

Write down a parametrisation of the curve using parameter u such that $0 < u < 1$.

To reparametrise with u such that the bounds on the parametrisation are $u \in (0, 1)$, we want to adjust t in the old parametrisation. We want $u = \frac{1}{2}$ when $t = 0$, $u \rightarrow 1$ as $t \rightarrow \infty$, and $u \rightarrow 0$ as $t \rightarrow -\infty$. We want an increasing, sigmoid-shaped curve with horizontal asymptotes at $y = 0$ and $y = 1$. \tanh almost fits this shape, but needs minor adjustments.

Let $u = \frac{1 + \tanh t}{2}$. Then $u = \frac{1}{2}$ when $t = 0$, $u \rightarrow 1$ as $t \rightarrow \infty$, and $u \rightarrow 0$ as $t \rightarrow -\infty$, as required. We rearrange to get $t = \operatorname{artanh}(2u - 1)$ and plug this into the old parametrisation.

Therefore \mathcal{C} can also be parametrised as

$$\left(\frac{1}{3} \sinh(\operatorname{artanh}(2u - 1)), \frac{1}{2} \cosh(\operatorname{artanh}(2u - 1)) \right) \quad u \in (0, 1)$$

We can also remove the hyperbolic functions and write it as

$$\left(\frac{2u - 1}{6\sqrt{u - u^2}}, \frac{1}{4\sqrt{u - u^2}} \right) \quad u \in (0, 1)$$

Question 3

A circle radius $r < 1$ is rolling on the inside of a circle radius 1, centred at the origin O . The centre C of the smaller circle is initially at $(1 - r, 0)$. Let θ be the angle subtended by the line OC measured with respect to the positive x axis. The point P , initially at $(1, 0)$, traces out a curve as the smaller circle rolls inside the unit circle.

The curve traced out by P has the parametric equations:

$$\begin{aligned}x(\theta) &= (1 - r) \cos \theta + r \cos \left(\frac{1 - r}{r} \theta \right), \\y(\theta) &= (1 - r) \sin \theta - r \sin \left(\frac{1 - r}{r} \theta \right).\end{aligned}$$

Q3 (a)

Use Python to plot the curves corresponding to $r = \frac{1}{2}, \frac{1}{4}, \frac{2}{3}$, and $\frac{1}{\sqrt{2}}$, where $\theta \in [0, 10\pi]$. Plot them in 4 separate figures.

Q3 (b)

Make one conjecture on the appearance of the curve for an arbitrary $r \in \mathbb{R}$.

I conjecture that for any $r \in \mathbb{R}$, the curve will be closed if and only if $r \in \mathbb{Q}$.




Figure 3: Plots of hypocycloids for various values of r

```

1  #!/usr/bin/env python3
2
3  from pathlib import Path
4
5  import matplotlib.pyplot as plt
6  import numpy as np
7
8
9  def generate_plot(r: float, title_text: str, file_number: int) -> None:
10     """Generate the plot for given radius."""
11     theta = np.linspace(0, 10 * np.pi, 100_000)
12     x = (1 - r) * np.cos(theta) + r * np.cos((1 - r) / r * theta)
13     y = (1 - r) * np.sin(theta) - r * np.sin((1 - r) / r * theta)
14
15     plt.figure(figsize=(6, 6))
16     plt.plot(np.cos(theta), np.sin(theta))
17     plt.plot(x, y)
18     plt.title(f"Hypocycloid with r = {title_text}")
19     plt.xlim(-1.1, 1.1)
20     plt.ylim(-1.1, 1.1)
21
22     plt.savefig(Path(__file__).parent.parent / "imgs" / f"Q3a-{file_number}.png")
23     plt.clf()
24
25
26  def main() -> None:
27     """Generate the plots for the necessary radii."""
28     for r, title_text, file_number in [
29         (1 / 2, "1/2", 1),
30         (1 / 4, "1/4", 2),
31         (2 / 3, "2/3", 3),
32         (1 / np.sqrt(2), "1/sqrt(2)", 4),
33     ]:
34         generate_plot(r, title_text, file_number)
35
36
37  if __name__ == "__main__":
38     main()

```

Figure 4: The code used to generate the plots in Figure 3. The code can also be found on GitHub