

Using eDNA and dissolved oxygen data to identify species-specific responses to hypoxic events



REED
COLLEGE



PMEL
Pacific Marine Environmental Laboratory

11/10/2024

Eleanor (Ella) Crotty (she/her)

Reed College

Environmental Studies/Biology

Pacific Marine Environmental Laboratory

Zachary Gold¹, Shannon Brown^{1,2}, Sean M.

McAllister^{1,2}, Jeannette Waddell³, Kathryn R Hough³,

Simone R. Alin¹, Matthew P. Galaska¹

¹NOAA PMEL

²UW CICOES


³NOAA OCNMS

Background

- **Environmental DNA (eDNA):** Genetic material shed into the environment by organisms
 - We sequence this to detect species in water
- **Hypoxia:** Low oxygen concentration
 - 20% saturation and lower harmful
 - 2 mg/L and lower harmful






Automated sampler deployed at TH042 mooring for summer 2021-24



PMEL
Pacific Marine Environmental Laboratory

eDNA sampling





PMEL
Pacific Marine Environmental Laboratory

Alin et al. 2023 *Oceanography*

Objective: Understand how species presence changes with oxygen levels

Overall Ocean Molecular Ecology (OME) lab project: Understand increasingly severe hypoxic events in the Olympic Coast National Marine Sanctuary (OCNMS) and develop autonomous eDNA monitoring systems



Environmental Data



O₂




OCNMS Mooring

Biodiversity




PPS Sampler



Hypoxia in the Pacific Northwest


Figure: Barth et al.
(2024)



Hypoxic Events

Mooring data

Length & Severity of OCNMS Hypoxic Events




Hypoxic event = period where the daily average oxygen saturation was < 20%

Hypoxic Events

eDNA sample data

Length & Severity of OCNMS Hypoxic Events + eDNA Sampling Dates




Hypoxic event = period where the daily average oxygen saturation was < 20%

Hypoxic Events



eDNA sampling windows

Length & Severity of OCNMS Hypoxic Events + eDNA Sampling Dates



Hypoxic event = period where the daily average oxygen saturation was < 20%


Temperature & Oxygen



Hypoxic event = period where the daily average oxygen saturation was < 20%

Results



- 79 eDNA samples taken in 2021-22
- 2 barcode regions used
- 8,228 unique gene sequences detected
- 674 species detected
- 64 species of interest identified by OCNMS + coastal treaty tribes
- 24 species of interest detected



PMEL
Pacific Marine Environmental Laboratory

Results

Clupea pallasii Pacific herring



PMEL
Pacific Marine Environmental Laboratory

Copepods

Indicator species used in OCNMS

P. mimus: Northern copepod


Paracalanus sp.: Southern copepods

Opposite hypoxic responses




PMEL
Pacific Marine Environmental Laboratory

Pseudocalanus *mimus* Presence vs Oxygen




Paracalanus sp. C AC-2013 Presence vs Oxygen



Results

- Wider strategy
 - Any species with large or significant hypoxia effect
- Identified affected species
 - Pacific sanddab
 - *Aureococcus anophagefferens* (Harmful algal bloom)


Gregarious
jellyfish



Pacific sanddab



A. anophagefferens



Steven Wilhelm





PMEL
Pacific Marine Environmental Laboratory

Results

Scientific Name	Common Name	Change With Hypoxia (<20% saturation)
<i>Paracalanus</i> sp. C AC-2013	Southern Copepod	More common
<i>Pseudocalanus mimus</i>	Northern Copepod	Less common
<i>Aureococcus anophagefferens</i>	Diatom (HAB)	Less common
<i>Citharichthys sordidus</i>	Pacific sanddab	More common
<i>Clytia gregaria</i>	Gregarious jellyfish	More common



Citharichthys sordidus (Pacific sanddab)

Detected more often in hypoxia
Benthic (seafloor resident) fish




Aureococcus anophagefferens (Diatom)

Detected less often in hypoxia
Harmful algal bloom (HAB) causer



Summary

- Combined eDNA and oceanographic data
 - Co-located datasets
- Evidence of hypoxia **tolerance** in two OCNMS priority species
- Identified other species affected by hypoxia



Next Steps

- 2023 data gap
- Community composition
- Relative abundance
- Pseudo-nitzschia (diatom) detection data

Biology senior thesis



Pseudo-nitzschia (NOAA NMFS)

Acknowledgements

- Ocean Molecular Ecology Group
 - Zachary Gold
 - Shannon Brown
 - Sean McAllister
 - Han Weinrich
 - Samantha Setta
- OCNMS crew: Haley Wilson, Andrew Micks, Kathy Hough, and Hatcher Cox
- Evan Howard (PMEL)
- NOAA Library
- Funding: NOAA Ernest F. Hollings Undergraduate Scholarship Program





Citations

- Alin, Simone R., Jan A. Newton, Richard A. Feely, Dana Greeley, Beth Curry, Julian Herndon, and Mark Warner. 2024. "A Decade-Long Cruise Time Series (2008–2018) of Physical and Biogeochemical Conditions in the Southern Salish Sea, North America." *Earth System Science Data* 16 (2): 837–65. <https://doi.org/10.5194/essd-16-837-2024>.
- Barth, John A., Stephen D. Pierce, Brendan R. Carter, Francis Chan, Anatoli Y. Erofeev, Jennifer L. Fisher, Richard A. Feely, et al. 2024. "Widespread and Increasing Near-Bottom Hypoxia in the Coastal Ocean off the United States Pacific Northwest." *Scientific Reports* 14 (1): 3798. <https://doi.org/10.1038/s41598-024-54476-0>.
- Domenici, P, C Lefrançois, and A Shingles. 2007. "Hypoxia and the Antipredator Behaviours of Fishes." *Philosophical Transactions of the Royal Society B: Biological Sciences* 362 (1487): 2105–21. <https://doi.org/10.1098/rstb.2007.2103>.
- Domenici, P., J. F. Steffensen, and R. S. Batty. 2000. "The Effect of Progressive Hypoxia on Swimming Activity and Schooling in Atlantic Herring." *Journal of Fish Biology* 57 (6): 1526–38. <https://doi.org/10.1111/j.1095-8649.2000.tb02229.x>.
- Fisheries, NOAA. 2024. "Local Biological Indicators | NOAA Fisheries." NOAA. West Coast. February 2, 2024. <https://www.fisheries.noaa.gov/west-coast/science-data/local-biological-indicators>.
- Gobler, Christopher J., Dianna L. Berry, Sonya T. Dyhrman, Steven W. Wilhelm, Asaf Salamov, Alexei V. Lobanov, Yan Zhang, et al. 2011. "Niche of Harmful Alga *Aureococcus Anophagefferens* Revealed through Ecogenomics." *Proceedings of the National Academy of Sciences* 108 (11): 4352–57. <https://doi.org/10.1073/pnas.1016106108>.
- Moriarty, Pamela E., Timothy E. Essington, John K. Horne, Julie E. Keister, Lingbo Li, Sandra L. Parker-Stetter, and Mei Sato. 2020. "Unexpected Food Web Responses to Low Dissolved Oxygen in an Estuarine Fjord." *Ecological Applications* 30 (8): e02204. <https://doi.org/10.1002/eap.2204>.
- "Oceanographic Moorings | Olympic Coast National Marine Sanctuary." n.d. Accessed July 19, 2024. <https://olympiccoast.noaa.gov/science/oceanographic-moorings/>.
- Onukwufor, John, and Chris Wood. 2022. "The Osmorespiratory Compromise in Marine Flatfish: Differential Effects of Temperature, Salinity, and Hypoxia on Diffusive Water Flux and Oxygen Consumption of English Sole (*Parophrys Vetulus*) and Pacific Sanddab (*Citharichthys Sordidus*)."*Marine Biology* 169 (April). <https://doi.org/10.1007/s00227-022-04040-z>.
- "Pacific Northwest Dead Zones Linked to Climate Change on JSTOR." n.d. Accessed June 17, 2024. https://www.jstor.org/stable/25595236?searchText=ocean+hypoxia&searchUri=%2Faction%2FdBasicSearch%3FQuery%3Docean%2Bhypoxia%26so%3Drel&ab_segments=0%2Fbasic_search_gsv%2Fcontrol&refreqid=fasty-default%3Afe4bd87fbb5a4d458696e62c77054c6f.
- "Pacific Sanddab | Washington Department of Fish & Wildlife." n.d. Accessed July 23, 2024. <https://wdfw.wa.gov/species-habitats/species/citharichthys-sordidus>.
- Risien, Craig M., Kathryn R. Hough, Jeannette Waddell, Melanie R. Fewings, and Brandy T. Cervantes. 2024. "Hydrographic Shipboard Profile Data Collected within Olympic Coast National Marine Sanctuary, 2005–2023." *Data in Brief* 53 (April):110171. <https://doi.org/10.1016/j.dib.2024.110171>.
- Vaquer-Sunyer, Raquel, and Carlos M. Duarte. 2008. "Thresholds of Hypoxia for Marine Biodiversity." *Proceedings of the National Academy of Sciences* 105 (40): 15452–57. <https://doi.org/10.1073/pnas.0803833105>.
- Zhang, Qing-Chun, Li-Mei Qiu, Ren-Cheng Yu, Fan-Zhou Kong, Yun-Feng Wang, Tian Yan, Christopher J. Gobler, and Ming-Jiang Zhou. 2012. "Emergence of Brown Tides Caused by *Aureococcus Anophagefferens* Hargraves et Sieburth in China." *HARMFUL ALGAE* 19 (September):117–24. <https://doi.org/10.1016/j.hal.2012.06.007>.

Mystery hydrozoan

Detected more often in hypoxia

DNA sequence could still be used as an indicator



Rate Change

Detection rate (detections/number of samples) in hypoxic conditions

-

Detection rate in non-hypoxic conditions

=

"Rate Change"



PMEL
Pacific Marine Environmental Laboratory