



Data Article

Hydrographic shipboard profile data collected within Olympic coast national marine sanctuary, 2005–2023



Craig M. Risien^{a,*}, Kathryn R. Hough^b, Jeannette Waddell^b,
Melanie R. Fewings^a, Brandy T. Cervantes^a

^a College of Earth, Ocean, and Atmospheric Sciences, Oregon State University, Corvallis, OR 97331, USA

^b National Oceanic and Atmospheric Administration, Olympic Coast National Marine Sanctuary, Port Angeles, WA 98362, USA

ARTICLE INFO

Article history:

Received 9 January 2024

Revised 31 January 2024

Accepted 2 February 2024

Available online 7 February 2024

Dataset link:

[Conductivity–Temperature–Depth \(CTD\) and dissolved oxygen profile data from shipboard surveys collected within Olympic Coast National Marine Sanctuary, 2005–2023 \(Original data\)](#)

Keywords:

Seawater temperature

Practical salinity

Density

Dissolved oxygen

CTD

Northern California current

Washington state

ABSTRACT

Olympic Coast National Marine Sanctuary (OCNMS), which was established in 1994 and covers an area of 8257 km², is located along Washington State's remote and rugged outer coast towards the northernmost extent of the California Current System (CCS). In this region, summertime equatorward winds drive seasonal upwelling of cold, nutrient rich waters onto the continental shelf. These waters help fuel a highly diverse and productive ecosystem that includes marine mammal and seabird communities as well as commercially and culturally important fisheries. The sanctuary is located within the boundaries of the legally defined Usual and Accustomed (U&A) fishing grounds of four Coastal Treaty Tribes, the Hoh Tribe, Makah Tribe, Quileute Tribe, and the Quinault Indian Nation, which hold treaty fishing rights and co-manage fisheries and other natural resources within the sanctuary through state, federal, and international partnerships and agreements. This data article describes shipboard hydrographic Conductivity–Temperature–Depth (CTD) and dissolved oxygen profile data that were collected within the sanctuary at fourteen locations during mooring deployment, recovery, and maintenance cruises between the months of May and October from 2005–2023. The 792 CTD profiles

* Corresponding author.

E-mail address: craig.risien@oregonstate.edu (C.M. Risien).

were acquired using Sea-Bird Scientific 19 SeaCAT or 19plus SeaCAT CTD profilers with associated SBE-43 (Sea-Bird Electronics) or Beckman or YSI-type (Yellow Springs Instruments) dissolved oxygen sensors. The data were processed using Sea-Bird Scientific's SBE Data Processing application. These data are needed for improving our understanding of sub-surface oceanographic conditions – including marine heat waves, changes in timing of spring transition to upwelling, seasonal hypoxia, and ocean acidification – in this important but undersampled region, and can be used to help improve the management of marine resources regionally and within the sanctuary. The CTD cast data are available via Zenodo at <https://doi.org/10.5281/zenodo.10466124>.

© 2024 The Author(s). Published by Elsevier Inc.
This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

Specifications Table

Subject	Oceanography			
Specific subject area	Hydrographic data collected within Olympic Coast National Marine Sanctuary (47.13–48.51°N, 124.18–125.68°W)			
Data format	Raw			
Type of data	Filtered			
Data collection	Figure			
	This data collection consists of 792 raw and processed Conductivity-Temperature-Depth (CTD) and dissolved oxygen profiles that were collected within Olympic Coast National Marine Sanctuary during mooring deployment, recovery, and maintenance cruises between the months of May and October from 2005–2023. These data were acquired using Sea-Bird Scientific 19 SeaCAT or 19plus SeaCAT CTD profilers with associated SBE-43 or Beckman or YSI-type dissolved oxygen sensors. They were processed using Sea-Bird Scientific's SBE Data Processing (v7.26.7) application.			
Data source location	Hydrographic data collected within Olympic Coast National Marine Sanctuary			
	Station Name	Latitude	Longitude	Water Depth (m, MLLW)
	<i>Makah Bay (MB)</i>			
	MB015	48.3254°N	124.6768°W	15
	MB042	48.3240°N	124.7354°W	42
	<i>Cape Alava (CA)</i>			
	CA015	48.1663°N	124.7568°W	15
	CA042	48.1660°N	124.8234°W	42
	CA065	48.1659°N	124.8949°W	65
	<i>Teahwhit Head (TH)</i>			
	TH015	47.8761°N	124.6195°W	15
	TH042	47.8762°N	124.7334°W	42
	TH065	47.8767°N	124.7967°W	65
	<i>Kalaloch (KL)</i>			
	KL015	47.6008°N	124.4284°W	15
	KL027	47.5946°N	124.4971°W	27
	KL050	47.5933°N	124.6112°W	50
	<i>Cape Elizabeth (CE)</i>			
	CE015	47.3568°N	124.3481°W	15
	CE042	47.3531°N	124.4887°W	42
	CE065	47.3528°N	124.5669°W	65
Data accessibility	Repository name: Zenodo			
	Data identification number: 10.5281/zenodo.10466124			
	Direct dataset link: https://doi.org/10.5281/zenodo.10466124			
	Zenodo is an open repository operated by the European Organization for Nuclear Research, known as CERN.			

1. Value Of The Data

- The data presented here can be used for model validation analyses [1,2] in this oceanographically and ecologically important but undersampled region.
- This CTD data set can be used as part of the process to quality control associated OCNMS mooring data [3] via cross-sensor calibrations and comparisons, which can correct for sensor drift.
- Using the long time series presented here, researchers can characterize regional ocean change at intra and interannual time scales as well as place extreme events [4] in greater context.

2. Background

The nineteen years (2005–2023) of CTD and dissolved oxygen profile data described in this data article were previously not publicly available. Making these data publicly available helps advance the understanding of subsurface oceanographic conditions within Olympic Coast National Marine Sanctuary and can be used to improve the management of commercially and culturally important marine resources. These profile data have been processed using the same procedures as those used to process CTD data collected along the Newport Hydrographic Line [5] located off the central Oregon coast, which allows for regional comparisons.

3. Data Description

The 792 CTD and dissolved oxygen profiles described here are available via Zenodo at <https://doi.org/10.5281/zenodo.10466124> [6]. The data set consists of the instrument configuration and calibration files (.con and .xmlcon) derived from annual vendor conducted sensor calibrations and the individual cast data files (.hex and .cnv) that contain CTD and dissolved oxygen observations collected between May 2005 and October 2023 at fourteen hydrographic stations located within Olympic Coast National Marine Sanctuary (Figs. 1 and 2). Additionally, the data set includes 792 CF (Climate and Forecast) compliant NetCDF files (Table 1) that con-

Table 1

Names, descriptions, and units of variables included in processed NetCDF files. NetCDF files are named using the convention sitename_yyyymmddThhmmss(_binned).nc with times given in Coordinated Universal Time (UTC). For example, a CTD cast that was conducted at the Cape Elizabeth 42-meter site on 28 August 2013 at 13:10:45 UTC is named CE042_20130828T131045.nc. The associated file that contains 1-dbar bin averaged profiles for this cast is named CE042_20130828T131045_binned.nc.

Variable name	Variable description	Units
Time	CTD cast start time	seconds since 1970-01-01 00:00:00 UTC
Pressure	Seawater pressure	dbar
Depth	Seawater depth calculated from pressure	meters
Temperature	Seawater temperature	°C
Salinity	Seawater practical salinity	1
potential_density	Potential density of seawater calculated from absolute salinity, potential temperature with respect to a reference seawater pressure of 0 dbar	kg m ⁻³
dissolved_oxygen	Seawater dissolved oxygen concentration	ml l ⁻¹
Latitude	Latitude	°N
Longitude ^a	Longitude	°E

^a The Specifications Table above lists the longitudes of the CTD casts in °W, but the processed NetCDF file contents use °E to follow the CF standard.




Fig. 1. A regional map showing GEBCO bathymetry [7] including the 25, 50, 100, and 200-m isobaths offshore of Washington State's Olympic Peninsula. The boundary of Olympic Coast National Marine Sanctuary is shown in magenta. The 14 mooring sites are shown as black circles with the numbers inside the black circles indicating the water depth at MLLW in meters. The inset map shows North America with the geographic bounds of the regional map indicated as a red box off the Washington coast.




Fig. 2. CTD data availability at each of the 14 mooring locations. Each black circle represents an individual CTD cast.

tain the processed (original and 1 dbar vertically bin averaged) temperature, practical salinity, potential density, and dissolved oxygen data for each CTD cast (Fig. 3).

4. Experimental Design, Materials and Methods

The 792 CTD+DO profiles were processed using Sea-Bird Scientific's SBE Data Processing (v7.26.7) application, which includes a series of data processing modules [8]. The data presented here were processed using six of the modules in the following order: *Data Conversion*, *Filter*, *Align CTD*, *Loop Edit*, *Derive*, and *Bin Average*. The *Cell Thermal Mass* (CTM) module was not applied to the data because the SBE Data Processing manual [8] states that the correction is negligible in regions that do not have steep temperature gradients. In regions that do have steep temperature gradients, the correction is about 0.005 PSU. These processing steps, including not applying the CTM module to the data, are the same as those used to process CTD data that make up the Newport Hydrographic Line time series located off the central Oregon coast [5] thus allowing for a direct comparison between the two regions.

The raw cast data files (.hex), which include the upcast and downcast data (time, pressure, conductivity, temperature, and dissolved oxygen) were converted to ASCII files (.cnv) using the *Data Conversion* module and associated instrument configuration and calibration files (.con and .xmlcon). For the years 2005–2007 when Beckman or YSI-type sensors were used to measure dissolved oxygen, data were converted using the Owens-Millard equation [9]. For the period 2008–2023 when SBE-43 oxygen sensors were used to measure dissolved oxygen, data were converted using the SBE-43 equation, which is a modified version of the Owens-Millard equa-




Fig. 3. Example temperature (upper left), practical salinity (upper right), potential density (lower left), and dissolved oxygen (lower right) profiles that were recorded at the 42-m Cape Elizabeth site on 25 July 2013. Full profiles (up and down casts) are shown in black. 1 dbar binned downcast profiles are shown in red. The dashed blue line in the lower right panel shows the 1.4 ml/l hypoxia threshold.

tion [10]. Converted pressure, conductivity and temperature data were then filtered using the *Filter* module. Pressure data collected using SBE 19 SeaCAT or SBE 19plus SeaCAT CTD profilers were filtered using a 2-s or a 1-s low-pass filter, respectively. Conductivity and temperature data were filtered using a 0.5-s low-pass filter. Profiles were then aligned, using the *Align CTD* module, by advancing temperature and conductivity observations by 0.5 and -0.5 s, respectively. Oxygen data were advanced 5 s. Next the *Loop Edit* module, which removes scans associated with pressure slowdowns and reversals, was applied to the aligned data. Scans where the CTD vertical speed was less than 0.25 m s^{-1} were excluded. Additionally, surface (1 m depth) soak data and bad data scans were omitted. After applying the *Loop Edit* module to the data, the *Derive* module was used to calculate depth (meters), practical salinity (PSU), and potential density (kg m^{-3}). Finally, all upcast data were excluded and downcast data were binned to 1-dbar bins using the *Bin Average* module. Bad scans were excluded from the binning process. Fig. 3 shows an example CTD cast that was collected at the 42-m Cape Elizabeth site on 25 July 2013. The temperature, practical salinity, potential density, and dissolved oxygen profiles shown in Fig. 3 have been processed using the SBE Data Processing modules and procedures described above.

Limitations

The processing steps described above did not include correcting for sensor drift within a field season. However, sensors were vendor calibrated annually prior to the start of each field season. Sea-Bird Scientific estimates that the typical drift rates associated with CTD temperature, conductivity and pressure sensors are $0.0002 \text{ }^{\circ}\text{C/month}$, $0.0001 \text{ Siemens/meter/month}$, and $0.0015\text{--}0.004\%$ full scale/month, respectively [11]. The calibration drift rate of the SBE-43 is estimated to be less than 0.5% over 1000 h of operation [12].

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data Availability

Conductivity–Temperature–Depth (CTD) and dissolved oxygen profile data from shipboard surveys collected within Olympic Coast National Marine Sanctuary, 2005–2023 (Original data) (Zenodo).

CRediT Author Statement

Craig M. Risien: Writing – original draft, Data curation, Validation; **Kathryn R. Hough:** Writing – review & editing, Data curation, Validation; **Jeannette Waddell:** Supervision, Writing – review & editing; **Melanie R. Fewings:** Supervision, Writing – review & editing; **Brandy T. Cervantes:** Writing – review & editing.

Ethics Statement

The authors have read and adhered to the ethical requirements for publication in Data in Brief and confirm that the current work does not involve human subjects, animal experiments, or any data collected from social media platforms.

Acknowledgments

We thank all the captains, crews, and technicians who enabled the collection of these data, and NOAA's Office of National Marine Sanctuaries for sustaining this monitoring program for more than two decades. Support for CMR, MRF, and BTC was provided by NOAA's [Climate Program Office](#) (Open Funder Registry/FundRef number [100007298](#)) through Climate Observations and Monitoring Program grant NA22OAR4310559.

References

- [1] P. MacCready, R.M. McCabe, S.A. Siedlecki, M. Lorenz, S.N. Giddings, J. Bos, S. Albertson, N.S. Banas, S. Garnier, Estuarine circulation, mixing, and residence times in the Salish Sea, *J. Geophys. Res.* 126 (2021) e2020JC016738, doi:[10.1029/2020JC016738](#).
- [2] S.A. Siedlecki, I.C. Kaplan, A.J. Hermann, T.T. Nguyen, N.A. Bond, J.A. Newton, G.D. Williams, W.T. Peterson, S.R. Alin, R.A. Feely, Experiments with seasonal forecasts of ocean conditions for the Northern region of the California Current upwelling system, *Sci. Rep.* 6 (2016) 27203, doi:[10.1038/srep27203](#).
- [3] S.R. Alin, S.A. Siedlecki, H. Berger, R.A. Feely, J.E. Waddell, B.R. Carter, J.A. Newton, E.J. Schumacker, D. Ayres, Evaluating the evolving ocean acidification risk to Dungeness crab: time-series observations and modelling on the Olympic Coast, Washington, USA, *Oceanography* 36 (2–3) (2023) 138–147, doi:[10.5670/oceanog.2023.216](#).
- [4] J.A. Koehlinger, J. Newton, J. Mickett, L. Thompson, T. Klinger, Large and transient positive temperature anomalies in Washington's coastal nearshore waters during the 2013–2015 Northeast Pacific marine heatwave, *PLoS One* 18 (2) (2023) e0280646, doi:[10.1371/journal.pone.0280646](#).
- [5] C.M. Risien, M.R. Fewings, J.L. Fisher, J.O. Peterson, C.A. Morgan, Spatially gridded cross-shelf hydrographic sections and monthly climatologies from shipboard survey data collected along the Newport hydrographic Line, 1997–2021, *Data Br.* 41 (2022), doi:[10.1016/j.dib.2022.107922](#).
- [6] C. Risien, K. Hough, J. Waddell, M. Fewings, B. Cervantes, Conductivity–Temperature–Depth (CTD) and dissolved oxygen profile data from shipboard surveys collected within Olympic Coast National Marine Sanctuary, 2005–2023, *Zenodo* v1 (2024), doi:[10.5281/zenodo.10466124](#).
- [7] GEBCO Bathymetric Compilation Group 2021, The GEBCO_2021 Grid - a continuous terrain model of the global oceans and land, NERC EDS British Oceanographic Data Centre NOC, 2021. doi:[10.5285/c6612cbe-50b3-0cff-e053-6c86abc09f8f](#).
- [8] [Sea-Bird ElectronicsInc., SEASOFT V2: SBE Data Processing Manual, Bellevue, Washington, 2010.](#)
- [9] W.B. Owens, R.C. Millard, Jr., A new algorithm for CTD oxygen calibration, *J. Phys. Oceanogr.* 15 (1985) 621–631 <0621:ANAFCO>2.0.CO;2, doi:[10.1175/1520-0485\(1985\)015](#).
- [10] [Sea-Bird Electronics, Inc.Application note no. 64–2: SBE 43 Dissolved Oxygen Sensor Calibration and Data Corrections, Bellevue, Washington, 2012.](#)
- [11] [Sea-Bird Electronics, Inc.Training Module 9 Getting the Highest Accuracy Data: CTD Care and Calibrations, Bellevue, Washington, 2023.](#)
- [12] [Sea-Bird Electronics, Inc.SBE 43 Dissolved Oxygen Sensor Data Sheet, Bellevue, Washington, 2022.](#)