

Geometric Series formulas

Interval	Sum	Condition	Interval	Sum	Condition
Infinite	$\sum_{k=0}^{\infty} a^k = \frac{1}{1-a}$	$ a < 1$	Finite on $[1, N]$	$\sum_{k=1}^N a^k = \frac{a(1-a^{N+1})}{1-a}$	None
Finite on $[0, N]$	$\sum_{k=0}^N a^k = \frac{1-a^{N+1}}{1-a}$	None	Finite on $[N_1, N_2]$	$\sum_{k=N_1}^{N_2} a^k = \frac{a^{N_1} - a^{N_2+1}}{1-a}$	None
Infinite	$\sum_{k=1}^{\infty} a^k = \frac{a}{1-a}$	$ a < 1$	Finite on $[1, N]$	$\sum_{k=1}^N k = \frac{N(N+1)}{2}$	None

Elementary signal classification

Name	Continuous	Discrete	Name	Continuous	Discrete
Unit Step function	$u(t) = \begin{cases} 1, & t \geq 0 \\ 0, & t < 0 \end{cases}$	$u[n] = \begin{cases} 1, & n \geq 0 \\ 0, & n < 0 \end{cases}$	Signum signal	$Sgn(t) = \begin{cases} 1, & t > 0 \\ -1, & t < 0 \end{cases}$	$Sgn[n] = \begin{cases} 1, & n > 0 \\ -1, & n < 0 \end{cases}$
Ramp signal	$r(t) = \begin{cases} t, & t \geq 0 \\ 0, & t < 0 \end{cases}$	$r[n] = nu[n] = \begin{cases} n, & n \geq 0 \\ 0, & n < 0 \end{cases}$	Sinusoidal signal	$x(t) = \sin(2\pi f_0 t + \theta)$	$X[n] = \sin(2\pi f_0 n + \theta)$
Impulse function	$\delta(t) = 0, t \neq 0$	$\delta[n] = \begin{cases} 1, & n = 0 \\ 0, & \text{otherwise} \end{cases}$	Sinc function	$\sin(\omega_0 t) = \frac{\sin(\pi \omega_0 t)}{\pi \omega_0 t}$	$\sin[\omega_0 n] = \frac{\sin(\pi \omega_0 n)}{\pi \omega_0 n}$
Rectangular pulse function	$\Pi\left(\frac{t}{\tau}\right) = \begin{cases} 1, & t \leq \tau/2 \\ 0, & t > \tau/2 \end{cases}$	$\Pi\left[\frac{n}{2N}\right] = \begin{cases} 1, & n \leq N \\ 0, & n > N \end{cases}$	Triangular pulse	$\Lambda\left(\frac{t}{\tau}\right) = \begin{cases} 1 - t /\tau, & t \leq \tau \\ 0, & t > \tau \end{cases}$	$\Lambda\left[\frac{n}{N}\right] = \begin{cases} 1 - n /N, & n \leq N \\ 0, & \text{elsewhere} \end{cases}$

Important Properties of Signals

Name	Properties	Name	Properties
Impulse properties	$\int_{-\infty}^{\infty} \delta(t) dt = 1$ $\delta(\alpha t) = \frac{1}{ \alpha } \delta(t)$ $\delta(\alpha t + b) = \frac{1}{ \alpha } \delta\left(t + \frac{b}{\alpha}\right)$ $\int_{-\infty}^{\infty} \phi(t) \delta(t - \lambda) dt = \phi(\lambda)$ $\phi(t) \delta(t - \lambda) = \phi(\lambda) \delta(t - \lambda)$	Time period of linear combination of two signals	<p>Sum of signals is periodic if $\frac{T_1}{T_2} = \frac{m}{n} = \text{rational number}$</p> <p>The fundamental period of $g(t)$ is given by $nT_1 = mT_2$ provided that the values of m and n are chosen such that the greatest common divisor (gcd) between m and n is 1</p>
Signals in term of unit step and vice versa	$u(t) = \int_{-\infty}^t \delta(\tau) d\tau$ $\delta(t) = \frac{d}{dt} u(t)$ $r(t) = tu(t)$ $u(t) = \frac{d}{dt} r(t)$ $u(t) = \frac{d}{dt} r(t)$ $sgn = u(t) - u(-t)$ $sgn = 2u(t) - 1$ $\Pi\left(\frac{t}{\tau}\right) = u\left(t + \frac{\tau}{2}\right) - u\left(t - \frac{\tau}{2}\right)$	Odd and even & symmetry	$x_e(t) = x_e(-t)$ $x_o(t) = -x_o(-t)$ $x(t) = x_e(t) + x_o(t)$ $x_e(t) = \frac{1}{2} [x(t) + x(-t)]$ $x_o(t) = \frac{1}{2} [x(t) - x(-t)]$
Derivative of impulse (doublet)	$\frac{d}{dt} \delta(t) = \delta'(t) = \begin{cases} \text{undefined}, & t = 0 \\ 0, & \text{otherwise} \end{cases}$ $\delta'(\alpha t) = \frac{1}{\alpha \alpha } \delta'(t)$ $\int_{-\infty}^{\infty} x(t) \delta'(t - \lambda) dt = -x'(\lambda)$ $x(t) \delta'(t) = x(0) \delta'(t) - x'(0) \delta(t)$	Combined operation	$x(t) \Rightarrow Kx(t) + C$ <p>Scale by K then shift by C....</p> $x(t) \Rightarrow x(\alpha t - \beta)$ <p>Shift by β: $[x(t - \beta)]$ Then Compress by α: $[x(t - \beta) \Rightarrow x(\alpha t - \beta)]$</p> <p>OR Compress by α: $[x(t) \Rightarrow x(\alpha t)]$ then Shift by $\frac{\beta}{\alpha}$: $[x(\alpha t) \Rightarrow x(\alpha(t - \frac{\beta}{\alpha})) = x(\alpha t - \beta)]$</p>
Energy and power	Periodic signals have infinite energy hence power type signals.		

Properties of the Continuous-Time Fourier Series

$$x(t) = \sum_{k=-\infty}^{+\infty} a_k e^{jk\omega_0 t} = \sum_{k=-\infty}^{+\infty} a_k e^{jk(2\pi/T)t}$$

$$a_k = \frac{1}{T} \int_T x(t) e^{-jk\omega_0 t} dt = \frac{1}{T} \int_T x(t) e^{-jk(2\pi/T)t} dt$$

Property	Time Domain	Fourier series
	$\left. \begin{array}{l} x(t) \\ y(t) \end{array} \right\} \begin{array}{l} \text{Periodic with period } T \text{ and} \\ \text{fundamental frequency } \omega_0 = 2\pi/T \end{array}$	$\begin{array}{l} a_k \\ b_k \end{array}$
Linearity	$Ax(t) + By(t)$	$Aa_k + Bb_k$
Time-Shifting	$x(t - t_0)$	$a_k e^{-jk\omega_0 t_0} = a_k e^{-jk(2\pi/T)t_0}$
Frequency-Shifting	$e^{jM\omega_0 t} = e^{jM(2\pi/T)t} x(t)$	a_{k-M}
Conjugation	$x^*(t)$	a_{-k}^*
Time Reversal	$x(-t)$	a_{-k}
Time Scaling	$x(\alpha t), \alpha > 0$ (periodic with period T/α)	a_k
Periodic Convolution	$\int_T x(\tau) y(t - \tau) d\tau$	$T a_k b_k$
Multiplication	$x(t) y(t)$	$\sum_{l=-\infty}^{+\infty} a_l b_{k-l}$
Differentiation	$\frac{dx(t)}{dt}$	$jk\omega_0 a_k = jk \frac{2\pi}{T} a_k$
Integration	$\int_{-\infty}^t x(t) dt$ (finite-valued and periodic only if $a_0 = 0$)	$\left(\frac{1}{jk\omega_0} \right) a_k = \left(\frac{1}{jk(2\pi/T)} \right) a_k$
Conjugate Symmetry for Real Signals	$x(t)$ real	$\begin{cases} a_k = a_{-k}^* \\ \Re\{a_k\} = \Re\{a_{-k}\} \\ \Im\{a_k\} = -\Im\{a_{-k}\} \\ a_k = a_{-k} \\ \angle a_k = -\angle a_{-k} \end{cases}$
Real and Even Signals	$x(t)$ real and even	a_k real and even
Real and Odd Signals	$x(t)$ real and odd	a_k purely imaginary and odd
Even-Odd Decomposition of Real Signals	$\begin{cases} x_e(t) = \mathcal{E}v\{x(t)\} & [x(t) \text{ real}] \\ x_o(t) = \mathcal{O}d\{x(t)\} & [x(t) \text{ real}] \end{cases}$	$\begin{cases} \Re\{a_k\} \\ j\Im\{a_k\} \end{cases}$

Parseval's Relation for Periodic Signals

$$\frac{1}{T} \int_T |x(t)|^2 dt = \sum_{k=-\infty}^{+\infty} |a_k|^2$$

TNT

Properties of the Discrete-Time Fourier Series

$$x[n] = \sum_{k=\langle N \rangle} a_k e^{jk\omega_0 n} = \sum_{k=\langle N \rangle} a_k e^{jk(2\pi/N)n}$$

$$a_k = \frac{1}{N} \sum_{n=\langle N \rangle} x[n] e^{-jk\omega_0 n} = \frac{1}{N} \sum_{n=\langle N \rangle} x[n] e^{-jk(2\pi/N)n}$$

Property	Time Domain	Fourier Series
	$\left. \begin{matrix} x[n] \\ y[n] \end{matrix} \right\}$ Periodic with period N and fundamental frequency $\omega_0 = 2\pi/N$	$\left. \begin{matrix} a_k \\ b_k \end{matrix} \right\}$ Periodic with period N
Linearity	$Ax[n] + By[n]$	$Aa_k + Bb_k$
Time shift	$x[n - n_0]$	$a_k e^{-jk(2\pi/N)n_0}$
Frequency Shift	$e^{jM(2\pi/N)n} x[n]$	a_{k-M}
Conjugation	$x^*[n]$	a_{-k}^*
Time Reversal	$x[-n]$	a_{-k}
Time Scaling	$x_{(m)}[n] = \begin{cases} x[n/m] & \text{if } n \text{ is a multiple of } m \\ 0 & \text{if } n \text{ is not a multiple of } m \end{cases}$ (periodic with period mN)	$\frac{1}{m} a_k$ (viewed as periodic with period mN)
Periodic Convolution	$\sum_{r=\langle N \rangle} x[r] y[n - r]$	$N a_k b_k$
Multiplication	$x[n] y[n]$	$\sum_{l=\langle N \rangle} a_l b_{k-l}$
First Difference	$x[n] - x[n - 1]$	$(1 - e^{-jk(2\pi/N)}) a_k$
Running Sum	$\sum_{k=-\infty}^n x[k]$ (finite-valued and periodic only if $a_0 = 0$)	$\left(\frac{1}{1 - e^{-jk(2\pi/N)}} \right) a_k$
Conjugate Symmetry for Real Signals	$x[n]$ real	$\begin{cases} a_k = a_{-k}^* \\ \Re\{a_k\} = \Re\{a_{-k}\} \\ \Im\{a_k\} = -\Im\{a_{-k}\} \\ a_k = a_{-k} \\ \angle a_k = -\angle a_{-k} \end{cases}$
Real and Even Signals	$x[n]$ real and even	a_k real and even
Real and Odd Signals	$x[n]$ real and odd	a_k purely imaginary and odd
Even-Odd Decomposition of Real Signals	$x_e[n] = \mathcal{E}v\{x[n]\}$ $[x[n] \text{ real}]$ $x_o[n] = \mathcal{O}d\{x[n]\}$ $[x[n] \text{ real}]$	$\Re\{a_k\}$ $j\Im\{a_k\}$

Parseval's Relation for Periodic Signals

$$\frac{1}{N} \sum_{n=\langle N \rangle} |x[n]|^2 = \sum_{k=\langle N \rangle} |a_k|^2$$



Properties of the Continuous-Time Fourier Transform

$$x(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(j\omega) e^{j\omega t} d\omega$$

$$X(j\omega) = \int_{-\infty}^{\infty} x(t) e^{-j\omega t} dt$$

Property	Aperiodic Signal	Fourier Transform
	$x(t)$	$X(j\omega)$
	$y(t)$	$Y(j\omega)$
Linearity	$ax(t) + by(t)$	$aX(j\omega) + bY(j\omega)$
Time-shifting	$x(t - t_0)$	$e^{-j\omega t_0} X(j\omega)$
Frequency-shifting	$e^{j\omega_0 t} x(t)$	$X(j(\omega - \omega_0))$
Conjugation	$x^*(t)$	$X^*(-j\omega)$
Time-Reversal	$x(-t)$	$X(-j\omega)$
Time- and Frequency-Scaling	$x(at)$	$\frac{1}{ a } X\left(\frac{j\omega}{a}\right)$
Convolution	$x(t) * y(t)$	$X(j\omega) Y(j\omega)$
Multiplication	$x(t)y(t)$	$\frac{1}{2\pi} X(j\omega) * Y(j\omega)$
Differentiation in Time	$\frac{d}{dt} x(t)$	$j\omega X(j\omega)$
Integration	$\int_{-\infty}^t x(t) dt$	$\frac{1}{j\omega} X(j\omega) + \pi X(0) \delta(\omega)$
Differentiation in Frequency	$tx(t)$	$j \frac{d}{d\omega} X(j\omega)$
Conjugate Symmetry for Real Signals	$x(t)$ real	$\begin{cases} X(j\omega) = X^*(-j\omega) \\ \Re\{X(j\omega)\} = \Re\{X(-j\omega)\} \\ \Im\{X(j\omega)\} = -\Im\{X(-j\omega)\} \\ X(j\omega) = X(-j\omega) \\ \angle X(j\omega) = -\angle X(-j\omega) \end{cases}$
Symmetry for Real and Even Signals	$x(t)$ real and even	$X(j\omega)$ real and even
Symmetry for Real and Odd Signals	$x(t)$ real and odd	$X(j\omega)$ purely imaginary and odd
Even-Odd Decomposition for Real Signals	$x_e(t) = \mathcal{E}v\{x(t)\}$ $[x(t) \text{ real}]$ $x_o(t) = \mathcal{O}d\{x(t)\}$ $[x(t) \text{ real}]$	$\Re\{X(j\omega)\}$ $j\Im\{X(j\omega)\}$

Parseval's Relation for Aperiodic Signals

$$\int_{-\infty}^{+\infty} |x(t)|^2 dt = \frac{1}{2\pi} \int_{-\infty}^{+\infty} |X(j\omega)|^2 d\omega$$

TNT

Basic Continuous-Time Fourier Transform Pairs

Signal	Fourier Transform	Fourier Series Coefficients (if periodic)
$\sum_{k=-\infty}^{+\infty} a_k e^{jk\omega_0 t}$	$2\pi \sum_{k=-\infty}^{+\infty} a_k \delta(\omega - k\omega_0)$	a_k
$e^{j\omega_0 t}$	$2\pi \delta(\omega - \omega_0)$	$a_1 = 1$ $a_k = 0$, otherwise
$\cos \omega_0 t$	$\pi [\delta(\omega - \omega_0) + \delta(\omega + \omega_0)]$	$a_1 = a_{-1} = \frac{1}{2}$ $a_k = 0$, otherwise
$\sin \omega_0 t$	$\frac{\pi}{j} [\delta(\omega - \omega_0) - \delta(\omega + \omega_0)]$	$a_1 = -a_{-1} = \frac{1}{2j}$ $a_k = 0$, otherwise
$x(t) = 1$	$2\pi \delta(\omega)$	$a_0 = 1$, $a_k = 0$, $k \neq 0$ (this is the Fourier series representation for any choice of $T > 0$)
Periodic square wave $x(t) = \begin{cases} 1, & t < T_1 \\ 0, & T_1 < t \leq \frac{T}{2} \end{cases}$ and $x(t+T) = x(t)$	$\sum_{k=-\infty}^{+\infty} \frac{2 \sin k\omega_0 T_1}{k} \delta(\omega - k\omega_0)$	$\frac{\omega_0 T_1}{\pi} \text{sinc}\left(\frac{k\omega_0 T_1}{\pi}\right) = \frac{\sin k\omega_0 T_1}{k\pi}$
$\sum_{n=-\infty}^{+\infty} \delta(t - nT)$	$\frac{2\pi}{T} \sum_{k=-\infty}^{+\infty} \delta\left(\omega - \frac{2\pi k}{T}\right)$	$a_k = \frac{1}{T}$ for all k
$x(t) = \begin{cases} 1, & t < T_1 \\ 0, & t > T_1 \end{cases}$	$\frac{2 \sin \omega T_1}{\omega}$	—
$\frac{\sin Wt}{\pi t}$	$X(j\omega) = \begin{cases} 1, & \omega < W \\ 0, & \omega > W \end{cases}$	—
$\delta(t)$	1	—
$u(t)$	$\frac{1}{j\omega} + \pi \delta(\omega)$	—
$\delta(t - t_0)$	$e^{-j\omega t_0}$	—
$e^{-at} u(t), \Re\{a\} > 0$	$\frac{1}{a + j\omega}$	—
$te^{-at} u(t), \Re\{a\} > 0$	$\frac{1}{(a + j\omega)^2}$	—
$\frac{t^{n-1}}{(n-1)!} e^{-at} u(t), \Re\{a\} > 0$	$\frac{1}{(a + j\omega)^n}$	—

TNT

Properties of the Discrete-Time Fourier Transform

$$x[n] = \frac{1}{2\pi} \int_{2\pi} X(e^{j\omega}) e^{j\omega n} d\omega$$

$$X(e^{j\omega}) = \sum_{n=-\infty}^{+\infty} x[n] e^{-j\omega n}$$

Property	Aperiodic Signals	Fourier Transform
Linearity	$x[n]$ $y[n]$ $ax[n] + by[n]$	$X(e^{j\omega})$ $Y(e^{j\omega})$ $aX(e^{j\omega}) + bY(e^{j\omega})$
Time-Shifting	$x[n - n_0]$	$e^{-j\omega n_0} X(e^{j\omega})$
Frequency-Shifting	$e^{j\omega_0 n} x[n]$	$X(e^{j(\omega - \omega_0)})$
Conjugation	$x^*[n]$	$X^*(e^{-j\omega})$
Time Reversal	$x[-n]$	$X(e^{-j\omega})$
Time Expansions	$x_{(k)}[n] = \begin{cases} x[n/k], & \text{if } n = \text{multiple of } k \\ 0, & \text{if } n \neq \text{multiple of } k \end{cases}$	$X(e^{jk\omega})$
Convolution	$x[n] * y[n]$	$X(e^{j\omega}) Y(e^{j\omega})$
Multiplication	$x[n] y[n]$	$\frac{1}{2\pi} \int_{2\pi} X(e^{j\theta}) Y(e^{j(\omega - \theta)}) d\theta$
Differencing in Time	$x[n] - x[n - 1]$	$(1 - e^{-j\omega}) X(e^{j\omega})$
Accumulation	$\sum_{k=-\infty}^n x[k]$	$\frac{1}{1 - e^{-j\omega}} X(e^{j\omega})$ $+ \pi X(e^{j0}) \sum_{k=-\infty}^{+\infty} \delta(\omega - 2\pi k)$
Differentiation in Frequency	$n x[n]$	$j \frac{dX(e^{j\omega})}{d\omega}$
Conjugate Symmetry for Real Signals	$x[n]$ real	$\begin{cases} X(e^{j\omega}) = X^*(e^{-j\omega}) \\ \Re\{X(e^{j\omega})\} = \Re\{X(e^{-j\omega})\} \\ \Im\{X(e^{j\omega})\} = -\Im\{X(e^{-j\omega})\} \\ X(e^{j\omega}) = X(e^{-j\omega}) \\ \angle X(e^{j\omega}) = -\angle X(e^{-j\omega}) \end{cases}$
Symmetry for Real, Even Signals	$x[n]$ real and even	$X(e^{j\omega})$ real and even
Symmetry for Real, Odd Signals	$x[n]$ real and odd	$X(e^{j\omega})$ purely imaginary and odd
Even-odd Decomposition of Real Signals	$x_e[n] = \mathcal{E}\{x[n]\}$ [x[n] real] $x_o[n] = \mathcal{O}\{x[n]\}$ [x[n] real]	$\Re\{X(e^{j\omega})\}$ $j\Im\{X(e^{j\omega})\}$

Parseval's Relation for Aperiodic Signals

$$\sum_{n=-\infty}^{+\infty} |x[n]|^2 = \frac{1}{2\pi} \int_{2\pi} |X(e^{j\omega})|^2 d\omega$$

TNT

Basic Discrete-Time Fourier Transform Pairs

Signal	Fourier Transform	Fourier Series Co-efficient (if periodic)
$\sum_{k=-\infty}^{+\infty} a_k e^{jk(2\pi/N)n}$	$2\pi \sum_{k=-\infty}^{+\infty} a_k \delta\left(\omega - \frac{2\pi k}{N}\right)$	a_k
$e^{j\omega_0 n}$	$2\pi \sum_{l=-\infty}^{+\infty} \delta(\omega - \omega_0 - 2\pi l)$	(a) $\omega_0 = \frac{2\pi m}{N}$ $a_k = \begin{cases} 1, & k = m, m \pm N, m \pm 2N, \dots \\ 0, & \text{otherwise} \end{cases}$ (b) $\frac{\omega_0}{2\pi}$ irrational \Rightarrow The signal is aperiodic
$\cos \omega_0 n$	$\pi \sum_{l=-\infty}^{+\infty} \{\delta(\omega - \omega_0 - 2\pi l) + \delta(\omega + \omega_0 - 2\pi l)\}$	(a) $\omega_0 = \frac{2\pi m}{N}$ $a_k = \begin{cases} \frac{1}{2}, & k = \pm m, \pm m \pm N, \pm m \pm 2N, \dots \\ 0, & \text{otherwise} \end{cases}$ (b) $\frac{\omega_0}{2\pi}$ irrational \Rightarrow The signal is aperiodic
$\sin \omega_0 n$	$\frac{\pi}{j} \sum_{l=-\infty}^{+\infty} \{\delta(\omega - \omega_0 - 2\pi l) - \delta(\omega + \omega_0 - 2\pi l)\}$	(a) $\omega_0 = \frac{2\pi r}{N}$ $a_k = \begin{cases} \frac{1}{2j}, & k = r, r \pm N, r \pm 2N, \dots \\ -\frac{1}{2j}, & k = -r, -r \pm N, -r \pm 2N, \dots \\ 0, & \text{otherwise} \end{cases}$ (b) $\frac{\omega_0}{2\pi}$ irrational \Rightarrow The signal is aperiodic
$x[n] = 1$	$2\pi \sum_{l=-\infty}^{+\infty} \delta(\omega - 2\pi l)$	$a_k = \begin{cases} 1, & k = 0, \pm N, \pm 2N, \dots \\ 0, & \text{otherwise} \end{cases}$
Periodic square wave $x[n] = \begin{cases} 1, & n \leq N_1 \\ 0, & N_1 < n \leq N/2 \end{cases}$ and $x[n+N] = x[n]$	$2\pi \sum_{k=-\infty}^{+\infty} a_k \delta\left(\omega - \frac{2\pi k}{N}\right)$	$a_k = \frac{\sin[(2\pi k/N)(N_1 + \frac{1}{2})]}{N \sin[2\pi k/2N]}$, $k \neq 0, \pm N, \pm 2N, \dots$ $a_k = \frac{2N_1 + 1}{N}$, $k = 0, \pm N, \pm 2N, \dots$
$\sum_{k=-\infty}^{+\infty} \delta[n - kN]$	$\frac{2\pi}{N} \sum_{k=-\infty}^{+\infty} \delta\left(\omega - \frac{2\pi k}{N}\right)$	$a_k = \frac{1}{N}$ for all k
$a^n u[n]$, $ a < 1$	$\frac{1}{1 - ae^{-j\omega}}$	—
$x[n] = \begin{cases} 1, & n \leq N_1 \\ 0, & n > N_1 \end{cases}$	$\frac{\sin[\omega(N_1 + \frac{1}{2})]}{\sin(\omega/2)}$	—
$\frac{\sin Wn}{\pi n} = \frac{W}{\pi} \text{sinc}\left(\frac{Wn}{\pi}\right)$ $0 < W < \pi$	$X(\omega) = \begin{cases} 1, & 0 \leq \omega \leq W \\ 0, & W < \omega \leq \pi \end{cases}$ $X(\omega)$ periodic with period 2π	—
$\delta[n]$	1	—
$u[n]$	$\frac{1}{1 - e^{-j\omega}} + \sum_{k=-\infty}^{+\infty} \pi \delta(\omega - 2\pi k)$	—
$\delta[n - n_0]$	$e^{-j\omega n_0}$	—
$(n+1)a^n u[n]$, $ a < 1$	$\frac{1}{(1 - ae^{-j\omega})^2}$	—
$\frac{(n+r-1)!}{n!(r-1)!} a^n u[n]$, $ a < 1$	$\frac{1}{(1 - ae^{-j\omega})^r}$	—

TNT

Properties of Laplace Transform

Property	Signal	Transform	ROC
	$x(t)$	$X(s)$	R
	$x_1(t)$	$X_1(s)$	R_1
	$x_2(t)$	$X_2(s)$	R_2
Linearity	$ax_1(t) + bx_2(t)$	$aX_1(s) + bX_2(s)$	At least $R_1 \cap R_2$
Time shifting	$x(t - t_0)$	$e^{-st_0} X(s)$	R
Shifting in the s -Domain	$e^{s_0 t} x(t)$	$X(s - s_0)$	Shifted version of R [i.e., s is in the ROC if $(s - s_0)$ is in R]
Time scaling	$x(at)$	$\frac{1}{ a } X\left(\frac{s}{a}\right)$	“Scaled” ROC (i.e., s is in the ROC if (s/a) is in R)
Conjugation	$x^*(t)$	$X^*(s^*)$	R
Convolution	$x_1(t) * x_2(t)$	$X_1(s)X_2(s)$	At least $R_1 \cap R_2$
Differentiation in the Time Domain	$\frac{d}{dt}x(t)$	$sX(s)$	At least R
Differentiation in the s -Domain	$-tx(t)$	$\frac{d}{ds}X(s)$	R
Integration in the Time Domain	$\int_{-\infty}^t x(\tau)d(\tau)$	$\frac{1}{s}X(s)$	At least $R \cap \{\Re\{s\} > 0\}$

Initial- and Final Value Theorems

If $x(t) = 0$ for $t < 0$ and $x(t)$ contains no impulses or higher-order singularities at $t = 0$, then

$$x(0^+) = \lim_{s \rightarrow \infty} sX(s)$$

If $x(t) = 0$ for $t < 0$ and $x(t)$ has a finite limit as $t \rightarrow \infty$, then

$$\lim_{t \rightarrow \infty} x(t) = \lim_{s \rightarrow 0} sX(s)$$

TNT

Laplace Transform of Elementary Functions

Signal	Transform	Roc
1. $\delta(t)$	1	All s
2. $u(t)$	$\frac{1}{s}$	$\Re\{s\} > 0$
3. $-u(-t)$	$\frac{1}{s}$	$\Re\{s\} < 0$
4. $\frac{t^{n-1}}{(n-1)!}u(t)$	$\frac{1}{s^n}$	$\Re\{s\} > 0$
5. $-\frac{t^{n-1}}{(n-1)!}u(-t)$	$\frac{1}{s^n}$	$\Re\{s\} < 0$
6. $e^{-\alpha t}u(t)$	$\frac{1}{s + \alpha}$	$\Re\{s\} > -\alpha$
7. $-e^{-\alpha t}u(-t)$	$\frac{1}{s + \alpha}$	$\Re\{s\} < -\alpha$
8. $\frac{t^{n-1}}{(n-1)!}e^{-\alpha t}u(t)$	$\frac{1}{(s + \alpha)^n}$	$\Re\{s\} > -\alpha$
9. $-\frac{t^{n-1}}{(n-1)!}e^{-\alpha t}u(-t)$	$\frac{1}{(s + \alpha)^n}$	$\Re\{s\} < -\alpha$
10. $\delta(t - T)$	e^{-sT}	All s
11. $[\cos \omega_0 t]u(t)$	$\frac{s}{s^2 + \omega_0^2}$	$\Re\{s\} > 0$
12. $[\sin \omega_0 t]u(t)$	$\frac{\omega_0}{s^2 + \omega_0^2}$	$\Re\{s\} > 0$
13. $[e^{-\alpha t} \cos \omega_0 t]u(t)$	$\frac{s + \alpha}{(s + \alpha)^2 + \omega_0^2}$	$\Re\{s\} > -\alpha$
14. $[e^{-\alpha t} \sin \omega_0 t]u(t)$	$\frac{\omega_0}{(s + \alpha)^2 + \omega_0^2}$	$\Re\{s\} > -\alpha$
15. $u_n(t) = \frac{d^n \delta(t)}{dt^n}$	s^n	All s
16. $u_{-n}(t) = \underbrace{u(t) * \dots * u(t)}_{n \text{ times}}$	$\frac{1}{s^n}$	$\Re\{s\} > 0$

TNT

Haris H.

Sequence	Transform	ROC
$\delta[n]$	1	All z
$\delta[n - m]$	z^{-m}	$ z > 0, m > 0; z < \infty, m < 0$
$a^n u[n]$	$\frac{1}{1 - az^{-1}}$	$ z > a $
$-a^n u[-n - 1]$	$\frac{1}{1 - az^{-1}}$	$ z < a $
$na^n u[n]$	$\frac{az^{-1}}{(1 - az^{-1})^2}$	$ z > a $
$-na^n u[-n - 1]$	$\frac{az^{-1}}{(1 - az^{-1})^2}$	$ z < a $
$a^n \cos(bn)u[n]$	$\frac{1 - a \cos(b)z^{-1}}{1 - 2a \cos(b)z^{-1} + a^2 z^{-2}}$	$ z > a $
$a^n \sin(bn)u[n]$	$\frac{a \sin(b)z^{-1}}{1 - 2a \cos(b)z^{-1} + a^2 z^{-2}}$	$ z > a $

Signal	Transform	ROC
$\delta(t)$	1	All s
$\delta(t - T)$	e^{-sT}	All s
$e^{-at}u(t)$	$\frac{1}{s + a}$	$\Re\{s\} > -a$
$-e^{-at}u(-t)$	$\frac{1}{s + a}$	$\Re\{s\} < -a$
$\frac{t^{n-1}}{(n-1)!}e^{-at}u(t)$	$\frac{1}{(s + a)^n}$	$\Re\{s\} > -a$
$-\frac{t^{n-1}}{(n-1)!}e^{-at}u(-t)$	$\frac{1}{(s + a)^n}$	$\Re\{s\} < -a$
$e^{-at} \cos(bt)u(t)$	$\frac{s + a}{(s + a)^2 + b^2}$	$\Re\{s\} > -a$
$e^{-at} \sin(bt)u(t)$	$\frac{b}{(s + a)^2 + b^2}$	$\Re\{s\} > -a$

Table 9.1: Laplace transforms for common signals