

# Solutions to EE3210 Tutorial 10 Problems

## Problem 1:

(a) Using the analysis formula of the continuous-time Fourier transform, we have

$$\begin{aligned} X(\omega) &= \int_{-\infty}^{+\infty} x(t)e^{-j\omega t} dt = \int_{-\infty}^{+\infty} \sum_{k=0}^{+\infty} \alpha^k \delta(t - kT) e^{-j\omega t} dt \\ &= \sum_{k=0}^{+\infty} \alpha^k \int_{-\infty}^{+\infty} \delta(t - kT) e^{-j\omega t} dt = \sum_{k=0}^{+\infty} \alpha^k e^{-jk\omega T} = \sum_{k=0}^{+\infty} (\alpha e^{-j\omega T})^k \\ &= \frac{1}{1 - \alpha e^{-j\omega T}}. \end{aligned}$$

(b) Using the analysis formula of the discrete-time Fourier transform, we have

$$\begin{aligned} X[\Omega] &= \sum_{n=-\infty}^{+\infty} x[n]e^{-j\Omega n} = \sum_{n=-\infty}^{+\infty} 4^n u[-n]e^{-j\Omega n} = \sum_{l=-\infty}^{+\infty} 4^{-l} u[l]e^{j\Omega l} = \sum_{l=0}^{+\infty} \left(\frac{1}{4}e^{j\Omega}\right)^l \\ &= \frac{1}{1 - \frac{1}{4}e^{j\Omega}}. \end{aligned}$$

## Problem 2:

(a) Using the time reversal property, we have

$$x(-t) \leftrightarrow X(-\omega).$$

Then, using the time shift property, we have

$$x[-(t-1)] \leftrightarrow e^{-j\omega} X(-\omega)$$

and

$$x[-(t+1)] \leftrightarrow e^{j\omega} X(-\omega).$$

Therefore, using the linearity property, we obtain

$$x(1-t) + x(-1-t) \leftrightarrow e^{-j\omega} X(-\omega) + e^{j\omega} X(-\omega) = 2 \cos \omega X(-\omega).$$

(b) Using the time shift property, we have

$$x(t - 6) \leftrightarrow e^{-j6\omega} X(\omega).$$

Then, using the time scaling property, we have

$$x(3t - 6) \leftrightarrow \frac{1}{3} e^{-j2\omega} X\left(\frac{\omega}{3}\right).$$

(c) Using the differentiation in time property, we have

$$\frac{dx(t)}{dt} \leftrightarrow (j\omega) X(\omega).$$

Applying this property again, we have

$$\frac{d^2x(t)}{dt^2} \leftrightarrow (j\omega)^2 X(\omega) = -\omega^2 X(\omega).$$

Finally, using the time shift property, we have

$$\frac{d^2x(t - 1)}{dt^2} \leftrightarrow -\omega^2 e^{-j\omega} X(\omega).$$

### Problem 3:

(a) Using the synthesis formula of the continuous-time Fourier transform, we have

$$x(t) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} X(\omega) e^{j\omega t} d\omega = \int_{-\infty}^{+\infty} \delta(\omega - \omega_0) e^{j\omega t} d\omega = e^{j\omega_0 t}.$$

*Remark.* In this case,  $x(t) = e^{j\omega_0 t}$  is in fact a periodic signal. Although continuous-time periodic signals in general do not satisfy the absolutely integrable condition, i.e.,

$$\int_{-\infty}^{+\infty} |x(t)| dt < \infty, \quad (1)$$

they have Fourier transforms if impulse functions such as  $\delta(\omega - \omega_0)$  are permitted in the transform. This implies that (1) is a sufficient but not necessary condition for the existence of Fourier transforms. ■

(b) Using the synthesis formula of the continuous-time Fourier transform, we have

$$\begin{aligned} x(t) &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} X(\omega) e^{j\omega t} d\omega \\ &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} 2[\delta(\omega - 1) - \delta(\omega + 1)] e^{j\omega t} d\omega + \frac{1}{2\pi} \int_{-\infty}^{+\infty} 3[\delta(\omega - 2\pi) + \delta(\omega + 2\pi)] e^{j\omega t} d\omega \\ &= \frac{1}{\pi} (e^{jt} - e^{-jt}) + \frac{3}{2\pi} (e^{j2\pi t} + e^{-j2\pi t}) \\ &= \frac{3}{\pi} \cos(2\pi t) + j \frac{2}{\pi} \sin t. \end{aligned}$$