

CITY UNIVERSITY OF HONG KONG
Department of Electronic Engineering

EE 3118 Linear Systems and Signal Analysis


Homework #3

1. Problem 2.29 (a), (b), (c), pp. 144.
2. Problem 2.40, pp. 148.
3. Problem 3.3, pp. 251.
4. Problem 3.22 (b), and (a) for Fig. (b), (e). pp. 255.
5. Problem 3.26, pp. 257.
6. Problem 3.40 (a), (d), (e), pp. 261.
7. Problem 3.43, (a), (b), pp. 262.

Homework #3

Problem 2.29, PP. 144

$$(b) \quad h(t) = e^{-6t} u(3-t)$$



Solution: The LTI system is non-causal, since $h(t) \neq 0$ for $t < 0$.

For stability, check

$$\int_{-\infty}^{\infty} |h(t)| dt = \int_{-\infty}^3 e^{-6t} dt = \infty$$

The system is unstable.

$$(g) \quad h(t) = (2e^{-t} - e^{(t-100)/100}) u(t)$$

It is causal, since $h(t) \equiv 0$ for $t < 0$.

It is unstable since

$$\int_{-\infty}^{\infty} |h(t)| dt = \infty$$

Problem 2.40, PP. 148

(a) Solution: To find the impulse response,

let $x(t) = \delta(t)$. Then,

$$h(t) = \int_{-\infty}^t e^{-(t-\tau)} \delta(\tau-2) d\tau$$

$$= \begin{cases} e^{-(t-2)} & t > 2 \\ 0 & t < 2 \end{cases} = e^{-(t-2)} u(t-2)$$

Prob. 3.3

Using Euler identity

$$\cos \theta = \frac{e^{j\theta} + e^{-j\theta}}{2}, \quad \sin \theta = \frac{e^{j\theta} - e^{-j\theta}}{2j}$$

We ~~write~~ write

$$\begin{aligned} x(t) &= 2 + \cos \frac{2\pi}{3} t + 4 \sin \frac{5\pi}{3} t \\ &= 2 + \frac{e^{j2\frac{\pi}{3}t} + e^{-j2\frac{\pi}{3}t}}{2} + 4 \cdot \frac{e^{j5\frac{\pi}{3}t} - e^{-j5\frac{\pi}{3}t}}{2j} \\ &= \cancel{\frac{2j}{2}} e^{-j5\frac{\pi}{3}t} + \frac{1}{2} e^{-j2\frac{\pi}{3}t} + 2 + \frac{1}{2} e^{j2\frac{\pi}{3}t} \\ &\quad - 2j e^{j5\frac{\pi}{3}t} \end{aligned}$$

\Rightarrow

$$\omega_0 = \frac{\pi}{3}$$

$$a_{-5} = 2j$$

$$a_{-2} = \frac{1}{2}$$

$$a_0 = 2$$


$$a_2 = \frac{1}{2}$$

$$a_5 = -2j$$

The key to this problem is twofold:

- (i) All sinusoids can be expressed using Euler identity. As a result it is unnecessary to go through integration to find a_k .
- (ii) Any Fourier series is unique! Thus, if we can expand $x(t)$ in the standard form (as above), we need to identify only the coefficients.

Prob. 3.22 (a), Fig. (b)



Within one period, $T=6$, (say, $[-2, 4]$), $\omega_0 = \frac{2\pi}{T} = \frac{\pi}{3}$

$$x(t) = \begin{cases} 0 & 2 \leq t \leq 4 \\ 2-t & 1 \leq t \leq 2 \\ 1 & -1 \leq t \leq 1 \\ t+2 & -2 \leq t \leq -1 \end{cases}$$

$$\begin{aligned} x_k &= \frac{1}{T} \int_T x(t) e^{-jk\omega_0 t} dt \\ &= \frac{1}{T} \left[\int_{-2}^{-1} (t+2) e^{-jk\omega_0 t} dt + \int_{-1}^1 e^{-jk\omega_0 t} dt \right. \\ &\quad \left. + \int_1^2 (2-t) e^{-jk\omega_0 t} dt \right] \end{aligned}$$

Use integration by parts

$$\int_a^b u dv = uv \Big|_a^b - \int_a^b v du$$

$$\int_{-2}^{-1} t e^{-jk\omega_0 t} dt = -\frac{1}{jk\omega_0} t e^{-jk\omega_0 t} \Big|_{-2}^{-1} + \int_{-2}^{-1} \frac{1}{jk\omega_0} e^{-jk\omega_0 t} dt$$

$$(u=t, e^{-jk\omega_0 t} dt = dv \Rightarrow v = -\frac{1}{jk\omega_0} e^{-jk\omega_0 t}, du=dt)$$

Prob. 3.26

$$a_k = \begin{cases} 2 & k=0 \\ j(\frac{1}{2})^{|k|} & \text{otherwise} \end{cases}$$

$$\begin{aligned} \Rightarrow x(t) &= \sum_{k=-\infty}^{\infty} a_k e^{jk\omega_0 t} \\ &= \sum_{k=-\infty}^{-1} j(\frac{1}{2})^{|k|} e^{jk\omega_0 t} + 2 + \sum_{k=1}^{\infty} j(\frac{1}{2})^{|k|} e^{jk\omega_0 t} \\ &= \sum_{k=-\infty}^{-1} j(\frac{1}{2})^{-k} e^{jk\omega_0 t} + 2 + \sum_{k=1}^{\infty} j(\frac{1}{2})^k e^{jk\omega_0 t} \\ &\quad \quad \quad (k < 0, |k| = -k) \quad \quad \quad (k > 0, |k| = k) \\ &= \sum_{k=1}^{\infty} j(\frac{1}{2})^k e^{-jk\omega_0 t} + 2 + \sum_{k=1}^{\infty} j(\frac{1}{2})^k e^{jk\omega_0 t} \\ &\quad \quad \quad (\text{Variable change } \times) \\ &= 2 + \sum_{k=1}^{\infty} j(\frac{1}{2})^k \left[e^{jk\omega_0 t} + e^{-jk\omega_0 t} \right] \\ &= 2 + \sum_{k=1}^{\infty} j(\frac{1}{2})^k \cdot 2 \cos k\omega_0 t \\ &\quad \quad \quad (\text{Euler Identity}) \end{aligned}$$

(a) No, $x(t)$ is not real

(b) Yes, $x(t)$ is even, since

$$\begin{aligned} x(-t) &= 2 + 2 \sum_{k=1}^{\infty} j(\frac{1}{2})^k \cos k\omega_0(-t) \\ &= 2 + 2 \sum_{k=1}^{\infty} j(\frac{1}{2})^k \cos k\omega_0 t \\ &= x(t) \end{aligned}$$

$$(c) \frac{dx(t)}{dt} = \sum_{k=1}^{\infty} j \left(\frac{1}{2}\right)^k \cdot 2 \left(-k\omega_0 \sin k\omega_0 t\right)$$

$$= -2j\omega_0 \sum_{k=1}^{\infty} k \left(\frac{1}{2}\right)^k \sin k\omega_0 t$$

Let $y(t) = \frac{dx(t)}{dt}$. Then,

$$y(-t) = -2j\omega_0 \sum_{k=1}^{\infty} k \left(\frac{1}{2}\right)^k \sin k\omega_0 (-t)$$

$$= 2j\omega_0 \sum_{k=1}^{\infty} k \left(\frac{1}{2}\right)^k \sin k\omega_0 t$$

$$= -y(t).$$

No, $\frac{dx(t)}{dt}$ is not even; in fact, it is odd.

1. Prob. 3.40

(a) $x(t-t_0) + x(t+t_0)$

$$x(t-t_0) \longleftrightarrow X_k e^{-jk\omega_0 t_0}$$

$$x(t+t_0) \longleftrightarrow X_k e^{-jk\omega_0(-t_0)} = X_k e^{jk\omega_0 t_0}$$

$$\begin{aligned} x(t-t_0) + x(t+t_0) &\longleftrightarrow X_k e^{-jk\omega_0 t_0} + X_k e^{jk\omega_0 t_0} \\ &= X_k (e^{jk\omega_0 t_0} + e^{-jk\omega_0 t_0}) \\ &= 2X_k \cos k\omega_0 t_0 \end{aligned}$$

(c) $x(3t-1)$

Let $y(t) = x(3t)$, $z(t) = x(3t-1)$. Then,

$$\begin{aligned} z(t) &= x(3t-1) \\ &= x\left[3\left(t-\frac{1}{3}\right)\right] \\ &= y\left(t-\frac{1}{3}\right) \end{aligned}$$

Time-shifting: $z_k = y_k e^{-jk\omega_y \cdot \frac{1}{3}}$

Time-scaling: $y_k = x_k$

$$\Rightarrow z_k = x_k e^{-jk\omega_x}$$

$$\begin{aligned} 3T_y &= T_x \\ \omega_x &= \frac{2\pi}{T_x} = \frac{2\pi}{3T_y} = \frac{1}{3}\omega_y \end{aligned}$$

Prob. 3.43

(i) For an odd harmonic signal, $a_k = 0$ for any even k .
Consider

$$\begin{aligned}
 x\left(t + \frac{T}{2}\right) &= \sum_{k=-\infty}^{\infty} a_k e^{jk \frac{2\pi}{T} \left(t + \frac{T}{2}\right)} \\
 &= \sum_{k=-\infty}^{\infty} a_k e^{jk \frac{2\pi}{T} t} e^{jk\pi} \\
 &= \sum_{k=-\infty}^{\infty} a_k e^{jk \frac{2\pi}{T} t} \cos k\pi \\
 &= \sum_{k=-\infty}^{\infty} a_k e^{jk \frac{2\pi}{T} t} (-1)^k \\
 &= - \sum_{k=-\infty}^{\infty} a_k e^{jk \frac{2\pi}{T} t} \quad \left(\text{since } a_k = 0 \text{ for even } k \text{ and } (-1)^k = -1 \text{ for odd } k \right) \\
 &= -x(t)
 \end{aligned}$$

(ii) If $x(t) = -x\left(t + \frac{T}{2}\right)$, in addition to $x(t) = x(t + T)$
Then

$$x(t) = \sum_{k=-\infty}^{\infty} a_k e^{jk \frac{2\pi}{T} t}$$

where

$$a_k = \frac{1}{T} \int_T x(t) e^{-jk\omega_0 t} dt, \quad \omega_0 = \frac{2\pi}{T}$$

let $\tau = t + \frac{T}{2}$

$$\begin{aligned}
 &= -\frac{1}{T} \int_T x\left(t + \frac{T}{2}\right) e^{-jk\omega_0 t} dt \\
 &= -\frac{1}{T} \int_T x(\tau) e^{-jk\omega_0 \left(\tau - \frac{T}{2}\right)} d\tau
 \end{aligned}$$

$$= -\frac{1}{T} \int_0^T x(\tau) e^{-j k \omega_0 \tau} d\tau$$

$$= -\left(\frac{1}{T} \int_0^T x(\tau) e^{-j k \omega_0 \tau} d\tau\right) \cos k\pi$$

$$= -\frac{1}{T} \int_0^T x(\tau) e^{-j k \omega_0 \tau} d\tau$$

$$= -(-1)^k a_k$$

$$\Rightarrow [1 + (-1)^k] a_k = 0$$

For any even k , this means that

$$2a_k = 0,$$

that is, $a_k = 0$. As such, $x(t)$ is odd harmonic.