

# Maths 721 Notes

2020

## Contents

<b>1</b>	<b>Lecture 1</b>	<b>2</b>
1.1	Representations . . . . .	2
1.2	$FG$ -Modules . . . . .	4
<b>2</b>	<b>Lecture 2</b>	<b>5</b>
2.1	Module Reducibility . . . . .	7
<b>3</b>	<b>Lecture 3</b>	<b>7</b>
3.1	Module and Representation Reducibility . . . . .	7
3.2	Group Algebras . . . . .	8
3.3	The Regular $FG$ -module, $FG$ . . . . .	9

# 1 Lecture 1

In the first half of this course we will cover three main topics:

- representations;
- modules;
- characters.

We will further see that representations and modules are essentially the same, and that modules and characters are essentially the same; and hence all three are essentially the same.

From now on  $G$  is a group.

## 1.1 Representations

**Definition 1.1.** A **representation** of a group  $G$  over a field  $F$  is a group homomorphism from  $G$  to  $\mathrm{GL}(n, F)$ , where  $n$  is the **degree** of the representation.

Explicitly, a representation is a function  $\rho : G \rightarrow \mathrm{GL}(n, F)$  such that for all  $g, h \in G$ ;

- (i)  $(gh)\rho = (g\rho)(h\rho)$ ,
- (ii)  $1_G\rho = I_n$ ,
- (iii)  $g^{-1}\rho = (g\rho)^{-1}$ .

Note the use of the (incredibly shit) postfix function notation.

**Example 1.2.** Take  $D_4$ , the Dihedral group of order 8. It has the following group presentations

$$\begin{aligned} D_4 &= \langle a, b \mid a^4 = 1, b^2 = 1, a^b = a^{-1} \rangle \\ &\cong \langle (1\ 2\ 3\ 4), (1\ 4)(2\ 3) \rangle, \end{aligned}$$

where  $a^b = bab^{-1}$  is conjugation of  $a$  by  $b$ . By defining the matrix subgroup

$$H = \left\langle A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, B = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \right\rangle$$

and defining  $\rho : D_4 \rightarrow \mathrm{GL}(n, F)$  where  $F = \mathbb{R}, \mathbb{C}$ , by  $a \mapsto A$  and  $b \mapsto B$ , and  $a^i b^j \mapsto A^i B^j$  for  $0 \leq i \leq 3$ , and  $0 \leq j \leq 1$ . Hence we have  $\rho$  is a representation of  $D_4$  over  $F$ . ◀

**Example 1.3.** Take  $\mathbb{Q}_8$  the Quaternion group of order 8, which has the following group presentations

$$\begin{aligned} \mathbb{Q}_8 &= \langle a, b \mid a^4 = 1, a^2 = b^2, a^b = a^{-1} \rangle \\ &\cong \langle \bar{a} = (1\ 6\ 2\ 5)(3\ 8\ 4\ 7), \bar{b} = (1\ 4\ 2\ 3)(5\ 7\ 6\ 8) \rangle \end{aligned}$$

Define

$$H = \left\langle A = \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix}, B = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \right\rangle \subset \text{GL}(2, \mathbb{C}).$$

Then  $\rho : \mathbb{Q}_8 \rightarrow \text{GL}(2, \mathbb{C})$  defined by  $a^k b^\ell \mapsto A^k B^\ell$  is a group representation of  $\mathbb{Q}_8$  over  $\mathbb{C}$  of degree 2. ◀

**Definition 1.4.** Let  $G$  be a group and define

$$\begin{aligned} \rho : G &\rightarrow \text{GL}(n, F) \\ g\rho &= I_n \end{aligned}$$

for all  $g \in G$ . Then  $\rho$  is a representation, called the **trivial representation** of arbitrary of degree.

It follows from the trivial representation that any group  $G$  has a representation of an arbitrary degree.

Let  $\rho : G \rightarrow \text{GL}(n, F)$  be a group homomorphism, and take  $T \in \text{GL}(n, F)$ . Then

$$(T^{-1}AT)(T^{-1}BT) = T^{-1}(AB)T.$$

Thus, given  $\rho$  define  $\sigma$  such that

$$g\sigma = T^{-1}(g\rho)T$$

for all  $g \in G$ . As for all  $g, h \in G$ , one has

$$\begin{aligned} (gh)\sigma &= T^{-1}((gh)\rho)T \\ &= T^{-1}(g\rho)(h\rho)T \\ &= T^{-1}(g\rho)TT^{-1}(h\rho)T \\ &= (g\sigma)(h\sigma), \end{aligned}$$

and so  $\sigma$  is a group homomorphism; and hence a representation.

**Definition 1.5.** Define

$$\rho : G \rightarrow \text{GL}(m, F), \quad \sigma : G \rightarrow \text{GL}(n, F)$$

to both be representation of  $G$  over  $F$ . We say that  $\rho$  is **equivalent to**  $\sigma$  if  $n = m$  and there exists  $T \in \text{GL}(n, F)$  such that  $g\sigma = T^{-1}(g\rho)T$ .

**Proposition 1.6.** *Equivalence of representations is an equivalence relation.*

*Proof.* Reflexivity is clear by taking  $T = I_n$ . For symmetry, take  $T$  to be its inverse. For transitivity, if

$$g\sigma = T^{-1}(g\rho)T, \quad g\rho = S^{-1}(g\eta)S,$$

then

$$g\sigma = (ST)^{-1}(g\eta)(ST).$$

◻

**Definition 1.7.** Define the **kernel** of the representation  $\rho : G \rightarrow \text{GL}(n, F)$  as  $\ker \rho = \{g \in G \mid g\rho = I_n\}$ .

**Proposition 1.8.** The kernel of a representation of  $G$  is a normal subgroup of  $G$ ; i.e.  $\ker \rho \triangleleft G$ .

*Proof.* Suppose  $g \in \ker \rho$  and  $h \in G$  is arbitrary. Then

$$(hgh^{-1})\rho = (h\rho)(g\rho)(h^{-1}\rho) = (h\rho)I_n(h\rho)^{-1} = (h\rho)(h\rho)^{-1} = I_n,$$

and so  $hgh^{-1} \in \ker \rho$ . As  $\ker \rho$  is closed under conjugation, it is a normal subgroup of  $G$ .  $\square$

**Definition 1.9.** We say  $\rho$  is a **faithful** representation of  $G$  if  $\ker \rho = \{1_G\}$ .

**Example 1.10.** For the trivial representation  $\rho : G \rightarrow \text{GL}(n, F)$  with  $g \mapsto I_n$  for all  $g \in G$ , we have  $\ker \rho = G$ . Hence the representation is not faithful.  $\blacktriangleleft$

**Lemma 1.11.** Suppose  $G$  is a finite group, and  $\rho$  is a representation of  $G$  over  $F$ . Then  $\rho$  is faithful if, and only if,  $\text{im } \rho \cong G$ .

*Proof.* Immediate from the first isomorphism theorem.  $\square$

## 1.2 FG-Modules

Suppose  $G$  is a group, and  $F = \mathbb{R}, \mathbb{C}$ . Given  $\rho : G \rightarrow \text{GL}(n, F)$ , with  $V = F^n$ , let  $v = (\lambda_1, \dots, \lambda_n) \in V$  for  $\lambda_i \in F$  be a row vector. Moreover, note that  $g\rho$  is an  $n \times n$  matrix for all  $g \in G$ . Thus, we have  $v \cdot (g\rho) \in V$ , and satisfies the following properties:

- (i)  $v \cdot ((gh)\rho) = v \cdot (g\rho)(h\rho)$ ;
- (ii)  $v \cdot (1_G\rho) = v$ ;
- (iii)  $(\lambda v) \cdot (g\rho) = \lambda(v \cdot (g\rho))$ ;
- (iv)  $(u + v) \cdot (g\rho) = u \cdot (g\rho) + v \cdot (g\rho)$ .

We often will omit the  $\cdot$  in the operation, and write  $v(a\rho)$  for  $v \cdot (a\rho)$ .

**Example 1.12.** Recall  $D_4$  and its given presentation from a previous example. We have

$$a\rho = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad b\rho = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

If  $v = (\lambda_1, \lambda_2)$ , then we have

$$v(a\rho) = (-\lambda_2, \lambda_1), \quad v(b\rho) = (\lambda_1, -\lambda_2).$$

**Definition 1.13.** Let  $V$  a vector space over the field  $F = \mathbb{R}, \mathbb{C}$ , and let  $G$  be a group. We say  $V$  is a **FG-module** if a multiplication  $v \cdot g$  for  $v \in V$ , and  $g \in G$  is defined such that:  $\blacktriangleleft$

- (i)  $v \cdot g \in V$ ;
- (ii)  $v \cdot (gh) = (v \cdot g) \cdot h$ ;
- (iii)  $v \cdot 1_G = v$ ;
- (iv)  $(\lambda v) \cdot g = \lambda(v \cdot g)$ ;
- (v)  $(u + v) \cdot g = u \cdot g + v \cdot g$ .

This generalises the previous discussion from a matrix group to an arbitrary group.

Note that properties (i), (iv), and (v) imply that the map  $v \mapsto v \cdot g$  is an endomorphism of  $V$  (a linear map from  $V$  to itself).

**Definition 1.14.** Suppose  $V$  is an  $FG$ -module and  $B$  is a basis for  $V$ . For  $g \in G$ , let  $[g]_B$  denote the matrix of the endomorphism  $v \mapsto v \cdot g$  of  $V$  relative to the basis  $B$ .

## 2 Lecture 2

**Theorem 2.1.** Let  $\rho : G \rightarrow GL(n, F)$  be a representation of  $G$  over  $F$ .

- (I) If  $V = F^n$  is an  $FG$  module and  $G$  acts on  $V$  by  $v \cdot g = v(g\rho)$  there exists a basis  $B$  of  $V$  such that  $g\rho = [g]_B$ .
- (II) The map  $g \mapsto [g]_B$  is a representation for  $G$  over  $F$ .

*Proof.* Choose the standard basis  $B = [e_1, \dots, e_n]$ .

Since  $V$  is an  $FG$ -module we have  $v(gh) = (vg)h$  for all  $g, h \in G$  and  $v \in V$ . Thus  $[gh]_B = [g]_B[h]_B$  so the map is a homomorphism. We now check that  $[g]_B$  is invertible for all  $g \in G$ . We know  $v \cdot 1_G = (vg)g^{-1}$  so  $I_n = [g]_B[g^{-1}]_B$  and thus  $[g]_B$  has an inverse.  $\square$

**Example 2.2.** Recall the representation of  $G = D_4$  from a previous example. Define an  $FG$ -module  $V = F^2$  with the action defined by taking  $vg$  to  $v(g\rho)$ .

$$\begin{aligned} v_1 &= (1, 0), & v_1 a &= v_2, & v_1 b &= v_1, \\ v_2 &= (0, 1), & v_1 a &= -v_1, & v_1 b &= -v_2. \end{aligned}$$

In this basis we recover our representation

$$a \mapsto [a]_B = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad b \mapsto [b]_B = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$



We now provide an equivalent basis-dependent definition for an  $FG$ -module.

**Lemma 2.3.** Let  $V$  a vector space over the field  $F = \mathbb{R}, \mathbb{C}$  with basis  $B = [v_1, \dots, v_n]$ , and let  $G$  be a group. If a multiplication  $v \cdot g$  for  $v \in B$ , and  $g \in G$  is defined such that:

- (i)  $v \cdot g \in V$ ;
- (ii)  $v \cdot (gh) = (v \cdot g) \cdot h$ ;
- (iii)  $v \cdot 1_G = v$ ;
- (iv)  $(\sum_{i=1}^n \lambda_i v_i) \cdot g = \sum_{i=1}^n \lambda_i (v_i \cdot g)$  for all  $\lambda_i \in F$ ;

then  $V$  is an  $FG$ -module.

**Definition 2.4.** The trivial module of a group over  $F$  is a one dimensional vector space  $V$  over  $F$  such that  $vg = v$  for all  $v \in V$  and  $g \in G$ .

**Definition 2.5.** An  $FG$ -module is faithful if  $1_G$  is the only  $g \in G$  such that  $vg = v$  for all  $v \in V$ .

**Theorem 2.6.** Let  $V$  be an  $FG$ -module with basis  $B$  and  $\rho$  a representation of group  $G$  over  $F$  defined by taking  $g \mapsto [g]_B$ .

- (i) If  $B'$  is another basis of  $V$  then the map  $g \mapsto [g]_{B'}$  is a representation of  $G$  equivalent to  $\rho$ .
- (ii) If representation  $\sigma$  is equivalent to  $\rho$  then there exists basis  $B''$  such that  $\sigma(g) = [g]_{B''}$  for all  $g \in G$ .

*Proof.* Taking  $T$  to be the change of basis matrices, the two representations are equivalent.  $\square$

**Example 2.7.** Let  $G = C_3 = \langle a \mid a^3 = 1 \rangle$  and representation  $\rho : G \rightarrow GL(n, F)$  defined by

$$a \mapsto \begin{bmatrix} 0 & 1 \\ -1 & -1 \end{bmatrix}.$$

We attempt to construct an  $FG$ -module with group action described by  $\rho$ . Take  $V = F^2$  with basis  $B = [v_1, v_2]$ . Define the action of  $G$  on  $V$  by

$$v_1 a = v_2, \quad v_2 a = -v_1 - v_2.$$

Let us now choose alternate basis  $B' = [u_1 = v_1, u_2 = v_1 + v_2]$ . The action of  $G$  on this basis is described by

$$u_1 a = -u_1 + u_2, \quad u_2 a = -u_1.$$

This gives us a representation

$$a \mapsto [a]_{B'} = \begin{bmatrix} -1 & 1 \\ -1 & 0 \end{bmatrix}.$$

To verify this construction we write our change of basis matrix as

$$T = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$$

and verify that

$$\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ -1 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} -1 & 1 \\ -1 & 0 \end{bmatrix}.$$

**Definition 2.8.** The permutation module of a group  $G \leq S_n$  is an  $n$ -dimensional vector space  $V$  with basis  $B = [v_1, \dots, v_n]$  and action by  $G$  defined by

$$v_i g = v_{ig}$$

for all  $g \in G$  where  $ig$  is the image of  $i$  under  $g \in S_n$ .

It follows from Caley's theorem that every group has a faithful  $FG$ -module.

**Example 2.9.** Take  $G = S_4$  and pick  $g = (1\ 2)$  and  $h = (1\ 2\ 3\ 4)$ . We have representations

$$[g]_B = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \quad [h]_B = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{bmatrix}.$$

## 2.1 Module Reducibility

**Definition 2.10.** Let  $V$  be an  $FG$ -module. We call  $W$  a submodule of  $V$  if  $W$  is a vector subspace of  $V$  and  $W$  is closed under the action of  $G$ . We then write  $W < V$ .

**Example 2.11.** Let  $G = C_3 = \langle (1\ 2\ 3) \rangle$  and  $V$  the permutation module of  $G$  with basis  $B = [v_1, v_2, v_3]$ . The subspace  $W = \langle v_1 + v_2 + v_3 \rangle$  is a submodule but the subspace  $U = \langle v_1 + v_2 \rangle$  is not.

For example, consider the action of  $g = (1\ 2\ 3)$  on  $v_1 + v_2 \in U$ .

$$(v_1 + v_2)g = v_{1g} + v_{2g} = v_2 + v_3 \notin U$$

whereas  $G$  acts on  $W$  trivially.

## 3 Lecture 3

### 3.1 Module and Representation Reducibility

For any module, it is clear that we have two trivial submodules:  $0 < v$  and  $V < V$ . Where  $0 = \{0\} \subset V$ .

**Definition 3.1.** Let  $V$  be an  $FG$ -module. We say that  $V$  is irreducible if the only submodules of  $V$  are  $V$  and  $0$ . Otherwise  $V$  is reducible.

In 2.11 we showed that the permutation module of  $C_3$  is reducible.

**Definition 3.2.** Let  $\rho : G \rightarrow GL(n, F)$  be a representation. We say that  $\rho$  is irreducible if the corresponding  $FG$ -module (as constructed in 2.1) is irreducible. Otherwise  $\rho$  is reducible.

If an  $FG$ -module,  $V$  is reducible, that is,  $0 < W < V$ ,  $0 \neq W \neq V$ . Let  $B_W$  be a basis for  $W$ . If we extend  $B_W$  to  $B$  a basis of  $V$ , then we get the following representation of  $G$ :

$$g \mapsto [g]_B = \begin{bmatrix} X_g & 0 \\ Y_g & Z_g \end{bmatrix} \quad (3.1)$$

Where the matrices  $X_g, Y_g$  and  $Z_g$  are some block matrices and 0 is a block of zeros and  $X_g$  has the dimensions  $m \times m$  and, in this case,  $\dim(W) = m$ .

**Proposition 3.3.** A representation  $\rho : G \rightarrow GL(n, F)$  is reducible if and only if with respect to some basis,  $B$ , of  $F^n$ ,  $[g]_B$  has the form 3.1 for some  $0 < m < \dim(V)$  for all  $g \in G$ . Then the maps  $g \mapsto X_g$  and  $g \mapsto Z_g$  are both representations of  $G$ .

*Proof.* Suppose we have a presentation,  $\rho : G \rightarrow GL(n, F)$  and a basis  $B$  of  $V = F^n$  such that  $[g]_B$  has the form 3.1 for every  $g \in G$ . Then consider the subspace  $0 \subset W \subset V$  spanned by the first  $m$  elements of  $B$ . It is clear that  $v[g]_B \in W$  for all  $v \in W$ . Therefore the module induced by  $\rho$  is reducible, so  $\rho$  is reducible. Now, if we have a reducible representation, then the argument above this proposition shows that with respect to any basis extending  $B_W$ , the matrices  $[g]_B$  have the required form.

Now, using elementary block matrix multiplication, we get the following for  $g, h \in G$ :

$$\rho(g)\rho(h) = [g]_B[h]_B = \begin{bmatrix} X_g X_h & 0 \\ Y_g X_h + Z_g Y_h & Z_g Z_h \end{bmatrix} = [gh]_B = \rho(gh)$$

Therefore  $X_{gh} = X_g X_h$  and  $Z_{gh} = Z_g Z_h$ , so the maps  $g \mapsto X_g$  and  $g \mapsto Z_g$  are both representations of  $G$ .  $\square$

**Problem 1.** Prove that the example representation of  $D_8$  of degree 2 over  $\mathbb{R}$  or  $\mathbb{C}$  is irreducible.

### 3.2 Group Algebras

Recall that an algebra over a field  $F$  is a vector space over  $F$  equipped with a bilinear product  $A \times A \rightarrow A$  that is compatible with scalar multiplication.

**Definition 3.4.** The group algebra over a finite group  $G$  over a field  $F$  is an algebra<sup>1</sup> of dimension  $n = |G|$  over  $F = \mathbb{R}$  or  $\mathbb{C}$  called  $FG$ , with basis  $B = G = \{g_1, \dots, g_n\}$ . Where the algebra structure is given by the following for two arbitrary elements of  $FG$ ,  $u = \sum_{g \in G} \lambda_g g$ ,  $v = \sum_{g \in G} \mu_g g$ ,  $\lambda_g, \mu_g \in F$  and  $\nu \in F$ :

1.  $u + v = \sum_{i=1}^n (\lambda_i + \mu_i) g_i$

---

<sup>1</sup>See Lemma 3.6



$$2. \nu \cdot u = \sum_{i=1}^n (\nu \lambda_i) g_i$$

$$3. u \cdot v = \sum_{(h,g) \in G \times G} \lambda_g \mu_h(gh)$$

This is clearly a vector space.

**Example 3.5.** Consider  $G = C_3 = \{e, a, a^2\} = \langle a | a^3 = e \rangle$  and  $F = \mathbb{R}$  or  $\mathbb{C}$ . Then if we let  $u = e - a + 2a^2$ ,  $v = \frac{1}{2}e + 5a$ , then:

$$u + v = \frac{3}{2}e + 4a + 2a^2, \quad \frac{1}{3}u = \frac{1}{3}e - \frac{1}{3}a + \frac{2}{3}a^2, \quad uv = \frac{21}{2}e + \frac{9}{2}a - 4a^2$$

**Lemma 3.6.** Given a group algebra  $FG$ ,  $r, s, t \in FG$ ,  $\lambda \in F$ :

$$1. rs \in FG$$

$$2. (rs)t = r(st)$$

$$3. 1_G r = r 1_G = r$$

$$4. (\lambda r)s = \lambda(rs)$$

$$5. (r + s)t = rt + st$$

$$6. r(s + t) = rs + rt$$

$$7. r0 = 0r = 0$$

That is,  $FG$  is an associative algebra with unit

*Proof.* 1,3 and 7 are clear from the definition of  $FG$ , 4,5 and 6 follow from the distributive and associative laws of  $F$  and 2 follows from associativity in  $G$ .  $\square$

### 3.3 The Regular $FG$ -module, $FG$

**Problem 2.**  $V = FG$  is an  $FG$ -module with the group action defined by  $v \cdot g = vg$  for  $v \in FG$ ,  $g \in G \subset FG$ .

**Definition 3.7.** For a finite group  $G$  and  $F = \mathbb{R}$  or  $\mathbb{C}$ , the regular  $FG$ -module is  $FG$ . The associated module,  $g \mapsto [g]_B$  is called the regular representation.

**Lemma 3.8.**  $FG$  is a faithful module for  $G$  over  $F$

*Proof.* If  $vg = v$  for all  $v \in FG$ , then specifically,  $hg = h$  for all  $h \in G$ , so  $g = 1_G$ .  $\square$

**Example 3.9.** For  $C = C_3$ , over the basis  $B = G$ , we get:

$$[e]_B = I_3, \quad [a]_B = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}, \quad [a^2]_B = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$



Now, if we have an  $FG$ -module,  $V$ , then  $FG$  acts on  $V$  in the following way:

$$v \cdot r = v \cdot \left( \sum_{g \in G} \mu_g g \right) = \sum_{g \in G} \mu_g (v \cdot g)$$

**Lemma 3.10.** For  $u, v \in V$ ,  $\lambda \in F$ ,  $r, s \in FG$ :

1.  $vr \in FG$
2.  $(vr)s = v(rs)$
3.  $v1 = v$
4.  $(\lambda v)r = \lambda(vr) = v(\lambda r)$
5.  $(r + s)v = rv + sv$
6.  $r(u + v) = ru + rv$
7.  $r0 = v0 = 0$

*Proof.* 1,3 and the first part of 7 follow from  $V$  being an  $FG$ -module, the second equality of 7 follows from scalar multiplication by 0 in  $V$ . The following calculation:

$$\begin{aligned} (\lambda v)r &= \sum_{g \in G} \mu_g ((\lambda v)g) \\ &= \sum_{g \in G} \mu_g (\lambda(vg)) \\ &= \sum_{g \in G} (\lambda \mu_g)(vg) = v(\lambda r) \\ &= \lambda \sum_{g \in G} \mu_g (vg) \\ &= \lambda(vg) \end{aligned}$$

proves 4. 6 follows from the linearity of the action of  $G$  on  $V$ . 5 follows from distributivity of scalar multiplication in  $V$ . Finally, to prove 2:

$$\begin{aligned} v(rs) &= \sum_{(g,h) \in G \times G} (\mu_g \lambda_h(v(gh))) \\ &= \sum_{h \in G} \lambda_h \sum_{g \in G} \mu_g(gv)h \\ &= \sum_{h \in G} \lambda_h \left( \sum_{g \in G} \mu_g(gv) \right) h \\ &= \sum_{h \in G} \lambda_h(vr)h = (vr)s \end{aligned}$$

□