

Decomposition of Games à la Candogan

Davide Legacci

September 14, 2023

Contents

1	Combinatorial Hodge Theory	2
2	Deviation Map of Normal Form Games	10
3	Hodge Decomposition of Normal Form Games	16
A	Dual and Adjoint in Linear Algebra	24
B	Proofs	26

Introduction

Our goal is to use Hodge decomposition theorem to decompose a normal form game into some components with distinctive game-theoretical and dynamical properties.

In Section (2) we show that a normal form game can be represented as a graph, called *response graph*. Loosely speaking, this graph is the space on which the Hodge decomposition takes place.

To be more precise, a game can be represented as a *flow* living in a *chain group* of the *chain complex* built on its *response graph*; and chain groups are the spaces to which Hodge theorem applies. In Section (1) we develop some notions of simplicial homology to make sense of this statement.

In Section (3) we put these ideas together, providing an explicit algorithm to perform the decomposition of a normal form game.

1 Combinatorial Hodge Theory

Essentials 1. The key points to take home:

- A graph is a *simplicial complex* K , that is a collection of 0-dimensional faces (vertices) and 1-dimensional faces (edges)
- The *chain groups* C_k built on a simplicial complex are vector spaces spanned by the k -dimensional faces of the complex
- Elements of C_1 are called *flows*
- In Sec. (2) a game is mapped to a flow in C_1 of its response graph
- Eq. (2) gives a linear *boundary map* $\partial_k : C_k \rightarrow C_{k-1}$
- Eq. (5) gives its dual, the *coboundary map* $d_k : C_k \rightarrow C_{k+1}$
- Def. (1.13) uses these maps to define *closed*, *exact*, and *harmonic* flows
- Th. (1.14) (Hodge) states that closed flows = exact flows \oplus harmonic flows

The decomposition proposed by [Can+11] leverages the combinatorial version of Hodge theorem. This was introduced in [Eck44]; see [Fri96] for a concise overview, [Jia+11] for an operational introduction, and [Mun84] for a through treatment.

The idea is the following.

1.1 Chain complex

Chain groups A graph is an example of a *simplicial complex* K - a collection of oriented k -dimensional faces (points, segments, triangles, tetrahedrons, ...). Given a simplicial complex K one can build a family of vector spaces C_k , with k ranging from 0 to the dimension of the biggest face appearing in K . Each C_k is (formally) the space of \mathbb{R} -linear combinations of oriented k -dimensional faces, i.e. the vector space spanned by the k -dimensional faces of the complex. These vector spaces are called *chain groups*.

An oriented k -dimensional face in K contains $k + 1$ vertices v_0, \dots, v_k and is denoted by $[v_0 \dots v_k]$. A basis B_k for C_k is given by the oriented k -dimensional faces in the complex K , so the dimension of C_k is the number n_k of such k -dimensional faces. A generic vector in C_k is called *k-chain* and is then a formal¹ linear combination

$$u_k = \sum_{[e] \in B_k} \lambda_k^e [e] \in C_k \quad (1)$$

where λ_k^e are n_k real numbers. Oriented k -dimensional faces, that is basis elements in B_k , are sometimes called *elementary k-chains*.

¹We say *formal* because we have not defined what the “sum of oriented faces” is. This can be done in full rigor, paying the price of abstraction. See the very first pages of [Mun84] for details.


In the present application we need to work with C_0, C_1 , and C_2 :

- C_0 is the space of linear combinations of elementary 0-chains, i.e. vertices. Equivalently, it is the space of assignments of a number to each vertex in the complex.
- C_1 is the space of linear combinations of elementary 1-chains, i.e. edges. Equivalently, it is the space of assignments of a number to each edge in the complex.

Remark 1.1. In the language of [Can+11], an element in C_1 , that is a 1-chain, is called *flow*.

- C_2 is the space of linear combinations of elementary 2-chains (equivalently, the assignment of a number to each 2-face in the complex). Think of an elementary 2-chain as an oriented “filled” 3-vertex clique².

Example 1.2.



Here K is a simplicial complex with 4 vertices (0-faces), 5 oriented edges (1-faces) and 2 oriented 2-faces. Dropping the v from the symbol of each vertex for notational simplicity, the basis of C_0, C_1 and C_2 are respectively

$$B_0 = \{[0], [1], [2], [3]\}$$

$$B_1 = \{[01], [02], [12], [13], [23]\}$$

$$B_2 = \{[012], [123]\}$$

Generic vectors (or chains) in these spaces are

$$u_0 = \lambda_0^0 [0] + \lambda_0^1 [1] + \lambda_0^2 [2] + \lambda_0^3 [3]$$

$$u_1 = \lambda_1^{01} [01] + \lambda_1^{02} [02] + \dots + \lambda_1^{23} [23]$$

$$u_2 = \lambda_2^{012} [012] + \lambda_2^{123} [123]$$

where λ_k^e are real numbers.

²A clique is a subset of vertices of a graph such that every two vertices in the clique are adjacent.




Figure 1: Effect of boundary map, Eq. (2), on elementary chains. From [Mun84, p. 29].

Boundary operator Given the family of vector spaces C_k it is natural to look for homomorphisms (also called linear maps or linear operators) between them.

Definition 1.3. Define the linear map

$$\begin{aligned} \partial_k : C_k &\rightarrow C_{k-1} \\ [v_0 \dots v_k] &\mapsto \sum_{i=0}^k (-1)^i [v_0 \dots \hat{v}_i \dots v_k] \end{aligned} \quad (2)$$

where the symbol \hat{v}_i means that the vertex v_i is to be deleted from the array. This map is called *boundary operator*.

The boundary of a k -chain is a $(k-1)$ -chain: intuitively, the boundary of a segment is the difference between its vertices; the boundary of a triangle is a combination of its edges; and so on (see [Mun84, p. 29] for details), and Figure 1 for a geometrical intuition.

Example 1.4. Consider the same complex of Example (1.2). The boundary map acts on the elementary 2-chain $[012] \in B_2$ as

$$\begin{aligned} \partial_2 : C_2 &\rightarrow C_1 \\ [012] &\mapsto [12] - [02] + [01] \end{aligned}$$

Chain complex Crucially, the boundary map fulfills

$$\partial_k \circ \partial_{k+1} : C_{k+1} \rightarrow C_k \rightarrow C_{k-1} = 0 \quad (3)$$

that is, the boundary of a boundary vanishes identically (the proof is a straightforward direct computation). Equivalently,

$$\text{Im } \partial_{k+1} \subseteq \ker \partial_k \quad (4)$$

This is the main ingredient of a very general construction:

Definition 1.5. A *chain complex* is an algebraic structure that consists of a sequence of abelian groups and a sequence of homomorphisms between consecutive groups such that the image of each homomorphism is included in the kernel of the next.

Vector spaces are in particular abelian groups, so from a simplicial complex we can build the following chain complex:

$$\cdots \rightarrow C_{k+1} \xrightarrow{\partial_{k+1}} C_k \xrightarrow{\partial_k} C_{k-1} \rightarrow \cdots$$

In particular, we will focus on

$$C_2 \xrightarrow{\partial_2} C_1 \xrightarrow{\partial_1} C_0$$

The reason for this is that

- Hodge theorem says that each chain group C_k can be decomposed as the direct sum of three special subspaces, the objects of our interest;
- The decomposition of each C_k depends on ∂_{k+1} and ∂_k ;
- As we will see in Section (2), a game can be mapped to an object living in C_1 , that is a 1-chain, or a *flow*.

Our next goals are then to

- understand the Hodge decomposition of each chain group;
- understand how to map a normal form game to C_1 , perform the decomposition, and go back to the space of games;
- get more concrete, representing the boundary map (and the other linear maps we will introduce in the next section) as a matrix, and actually do computations.

To state Hodge theorem we need to introduce two more ingredients: the *coboundary operator* and the *Laplacian*.

1.2 Cochain complex

Coboundary operator The boundary operators allows us to go from the k -th chain group to the $(k - 1)$ -th one. Since we are in the realm linear algebra there is a natural way to obtain a map going “in the opposite direction”:

Definition 1.6. The *coboundary operator*

$$d_{k-1} := \partial_k^* : C_{k-1}^* \rightarrow C_k^* \tag{5}$$

is the dual of the boundary operator $\partial_k : C_k \rightarrow C_{k-1}$.

For the general definition of dual of a vector space and of dual of a linear map between vector spaces see Appendix (A), but for practical uses it suffices to recall that

Lemma 1.7. Let $A : V \rightarrow W$ be a linear map between vector spaces, and let $A^* : V^* \rightarrow W^*$ be its dual. The matrix representing A^* is the transpose of the matrix representing A .

Corollary 1.8. The coboundary map fulfills

$$d_k \circ d_{k-1} : C_{k-1}^* \rightarrow C_k^* \rightarrow C_{k+1}^* = 0 \quad (6)$$

or equivalently

$$\text{Im } d_{k-1} \subseteq \ker d_k \quad (7)$$

In particular, also the coboundary maps allow to build a chain complex:

$$\cdots \leftarrow C_{k+1}^* \xleftarrow{d_k} C_k^* \xleftarrow{d_{k-1}} C_{k-1}^* \leftarrow \cdots$$

Proof. Immediate from Eq. (3) and Lemma (1.7). □

The chain complex built on the dual chain groups using the coboundary maps is called *cochain complex*. In the following we will restrict our attention to the piece of the chain and cochain complex involving 0-, 1- and 2-chains:

$$\begin{array}{ccccc} & d_0 = \partial_1^* & & d_1 = \partial_0^* & \\ & \curvearrowright & & \curvearrowright & \\ C_0^* & & C_1^* & & C_2^* \\ & & & & \\ C_0 & \xleftarrow{\partial_1} & C_1 & \xleftarrow{\partial_2} & C_2 \end{array} \quad (8)$$

Constructions on inner product spaces Note that all constructions performed so far are canonical: given a simplicial complex K , the chain and cochain complexes (that is, the chain groups, the boundary operator, and their dual) are naturally defined.

The next step, on the other hand, depends on a *choice*: each chain space C_k is endowed with an inner product

$$\langle \cdot, \cdot \rangle_k : C_k \times C_k \rightarrow \mathbb{R} \quad (9)$$

that is a bilinear form symmetric and positive-definite.

This is a crucial choice, since the Hodge decomposition of each chain group will turn out to depend on these inner products. The inner product employed in [Can+11] is the Euclidean one: given two k -chains

$$\begin{aligned} u_k &= \sum_{[e] \in B_k} \alpha_k^e [e] \in C_k \\ v_k &= \sum_{[e] \in B_k} \beta_k^e [e] \in C_k \end{aligned}$$

their Euclidean inner product is

$$\langle u_k, v_k \rangle_k^{\text{eu}} = \sum_{[e] \in B_k} \alpha_k^e \beta_k^e \quad (10)$$

A vector space with an inner product is called *inner product space*.

Remark 1.9. The reader interested in concrete applications in the Euclidean setting can safely skip section (1.2.1) and jump to section (1.2.2).

1.2.1 Non-Euclidean setting

We briefly review two constructions on inner product spaces: the identification between the space and its dual; and the adjoint of a linear map.

Dual isomorphism An inner product $\langle \cdot, \cdot \rangle : V \times V \rightarrow V^*$ induces an isomorphism between an inner product space V and its dual V^* :

$$\langle \cdot, \cdot \rangle : V \xrightarrow{\sim} V^* \quad (11)$$

For details see Appendix (A); for our application it suffices to know that such isomorphism exists, and that in the case of the Euclidean inner product the matrix representing this isomorphism is the *identity matrix*, allowing for effectively identifying a vector space with its dual.

Adjoint map Given a linear map $A : V \rightarrow W$ between inner product spaces one can define the *adjoint map* $A^\dagger : W \rightarrow V$ by

$$\langle Av, w \rangle_W = \langle v, A^\dagger w \rangle_V, \quad \forall v \in V, w \in W$$

Let's remark that, given a linear map $A : V \rightarrow W$ between vector space, its dual map $A^* : W^* \rightarrow V^*$ is canonical; on the other hand its adjoint map $A^\dagger : W \rightarrow V$ depends on the inner products on V and W . Remarkably, the dual map and the adjoint map are related by the isomorphism (11):

Lemma 1.10. *Given the linear map between inner product spaces $A : V \rightarrow W$, the dual map A^* and the adjoint map A^\dagger are related by*

$$A^\dagger = \langle \cdot, \cdot \rangle_V \circ A^* \circ \langle \cdot, \cdot \rangle_W$$

that is, the following diagram commutes:

$$\begin{array}{ccc} V & \xleftarrow{A^\dagger} & W \\ \langle \cdot, \cdot \rangle_V \updownarrow & & \updownarrow \langle \cdot, \cdot \rangle_W \\ V^* & \xleftarrow{A^*} & W^* \end{array}$$

In particular, in the case of the Euclidean metric, the dual map A^* and the adjoint map A^\dagger are represented by the same matrix, that is the transpose of the matrix of A .

In the diagram above (and in the following one) black maps are canonical, blue ones represent the isomorphisms induced by inner products; and red maps depend on these inner products.

With these notions at hand we can use the inner products defined in Eq. (9) to build adjoint maps on diagram (8):

$$\begin{array}{ccccc}
 & & d_0 = \partial_1^* & & d_1 = \partial_0^* \\
 & & \curvearrowright & & \curvearrowright \\
 C_0^* & & C_1^* & & C_2^* \\
 \uparrow \langle \cdot, \cdot \rangle_0 & & \uparrow \langle \cdot, \cdot \rangle_1 & & \uparrow \langle \cdot, \cdot \rangle_2 \\
 & & d_0^\dagger & & d_1^\dagger \\
 & & \curvearrowleft & & \curvearrowleft \\
 C_0 & \xleftarrow{\partial_1} & C_1 & \xleftarrow{\partial_2} & C_2
 \end{array}$$

1.2.2 Euclidan setting

In the case of the Euclidean inner product we can forget about the difference between a space and its dual, and about the difference between an adjoint map and a dual map:

$$\begin{array}{ccccc}
 & & d_0 = \partial_1^* = \partial_1^\dagger & & d_1 = \partial_0^* = \partial_0^\dagger \\
 & & \curvearrowright & & \curvearrowright \\
 C_0 & & C_1 & & C_2 \\
 \curvearrowleft & & \curvearrowleft & & \curvearrowleft \\
 & & \partial_1 = d_0^* = d_0^\dagger & & \partial_2 = d_1^* = d_1^\dagger
 \end{array} \quad [\text{Euclidean}] \quad (12)$$

Concretely, the maps in the lower row are defined by eq. (2), and the maps in the upper row are determined by Lemma (1.7).

Laplacian The last ingredient we need for Hodge theorem is the *Laplacian operator*; we define it only on C_1 and in the Euclidean case, but it is defined analogously on chain groups of any rank and for arbitrary inner products.

Definition 1.11. The *Laplacian operator* is

$$\Delta_1 := d_0 \circ \partial_1 + \partial_2 \circ d_1 : C_1 \rightarrow C_1 \quad (13)$$

This leaves us with the following diagram:

$$\begin{array}{c}
\Delta_1 \\
\begin{array}{ccccc}
& \nearrow d_0 & & \nearrow d_1 & \\
C_0 & & C_1 & & C_2 \\
& \searrow \partial_1 & & \searrow \partial_2 & \\
& & & &
\end{array}
\end{array} \quad [\text{Euclidean}] \tag{14}$$

Remark 1.12. In the following, we consider always the Euclidean setting.

1.3 Combinatorial Hodge theorem

Recall by Eqs. (4) and (7) that

$$\begin{aligned}
\text{Im } d_0 &\subseteq \ker d_1 \\
\text{Im } \partial_2 &\subseteq \ker \partial_1
\end{aligned} \tag{15}$$

Furthermore it is a standard result that

$$\ker \Delta_1 = \ker \delta_1 \cap \ker d_1 \tag{16}$$

These vector subspaces of C_1 deserve special names:

Definition 1.13. A 1-chain in C_1 is called

- *exact* iff it belongs to $\text{Im } d_0$
- *closed* iff it belongs to $\ker d_1$
- *co-exact* iff it belongs to $\text{Im } \partial_2$
- *co-closed* iff it belongs to $\ker \partial_1$
- *harmonic* iff it belongs to $\ker \Delta_1$

Note by Eq. (15) that any (co-)exact chain is (co-)closed, and by Eq. (16) that a 1-chain is harmonic iff it is closed and co-closed. In the following we use interchangeably the terms *1-chain* and *flow* for elements of C_1 .

Theorem 1.14 (Hodge). *The chain group C_1 admits the orthogonal decomposition*

$$\begin{aligned}
C_1 &= \text{closed} \oplus \text{closed}^\perp \\
&= \text{exact} \oplus \text{harmonic} \oplus \text{co-exact}
\end{aligned} \tag{17}$$

See Appendix (B.1) for a proof sketch. The first line holds true by definition of orthogonal complement, and it is a general fact that

$$\text{closed}^\perp = (\ker d_1)^\perp = \text{Im } d_1^* = \text{Im } d_1^* \text{Im } \partial_2 = \text{co-exact}$$

The real content of Hodge theorem, and what we will actually need, is that closed flows admit the orthogonal decomposition

$$\begin{aligned} \text{closed flows} &= \text{exact flows} \oplus \text{harmonic flows} \\ \ker d_1 &= \text{Im } d_0 \oplus \ker \Delta_1 \end{aligned} \tag{18}$$

2 Deviation Map of Normal Form Games

Essentials 2. The key points to take home:

- The response graph of a normal form game skeleton $(\mathcal{N}, \mathcal{A})$ has A nodes (pure strategy profiles) and E edges (unilateral deviations)
- \mathcal{A} is the basis of C_0 and \mathcal{E} is the basis of C_1
- Sec. (2.1) gives the explicit matrix of the coboundary operator $d_0 : C_0 \rightarrow C_1$,
- Given $(\mathcal{N}, \mathcal{A})$ an utility u lives in the AN -dimensional vector space $U = C_0^N$
- Eq. (24) gives the *deviation map* $D : U \rightarrow C_1$ mapping a game to a flow (utility difference of deviating player)
- Sec. (2.3) gives the matrix of the deviation map D

The goal of this section is to represent a normal form game as a 1-chain living in C_1 on some graph - the *response graph* - , so that we can apply Hodge theorem. This will be done by a so-called *deviation map* D acting as

$$D : \text{space of games} \rightarrow C_1$$

In the next sections we build the relevant graph, define what we mean by *space of games*, and define the deviation map D in Eq. (24).

Normal form game A normal form game is a tuple $\Gamma = (\mathcal{N}, \mathcal{A}, u)$ where

- $\mathcal{N} = \{1, 2, \dots, N\}$ is the set of players
- Each player $i \in \mathcal{N}$ has a set of pure strategies

$$\mathcal{A}_i = \{1, 2, \dots, A_i\}$$

- $\mathcal{A} = \prod_{i \in \mathcal{N}} \mathcal{A}_i$ is the set of pure strategy profiles
- Each player has an individual utility function

$$u_i : \mathcal{A} \rightarrow \mathbb{R}, \quad a \mapsto u_i(a)$$




Figure 2: Response graph of a $[2, 3]$ game as in Example (2.1). Blue (vertical) edges represent deviations of the first player and red (horizontal and bent) edges represent deviations of the second player.

- The utility map of the game is

$$u : \mathcal{A} \rightarrow \mathbb{R}^N, \quad a \mapsto (u_1, \dots, u_N)(a)$$

The number of players is $N = |\mathcal{N}|$, the number of pure strategies of player $i \in \mathcal{N}$ is $A_i = |\mathcal{A}_i|$, and the number of pure strategies profiles is $A = |\mathcal{A}| = \prod_{i \in \mathcal{N}} A_i$, so the number of utilities is AN .

Game skeleton We will call the datum of a set of players and of a space of pure strategies for each player, without specifying a utility map, the *skeleton* of a normal form game, and denote it by $(\mathcal{N}, \mathcal{A})$ or by $[A_1, A_2, \dots, A_N]$. For example $[3, 4]$ is the skeleton of a normal form game with 2 players, where player 1 has 3 pure strategies and player 2 has 4 pure strategies.

Unilateral deviations Pairs of strategy profiles $a, b \in \mathcal{A}$ that differ only in the strategy of one player are called *unilateral deviations*, and their space is denoted by \mathcal{E} . One can check that given the skeleton of a normal form game $(\mathcal{N}, \mathcal{A})$ the number of unilateral deviations is

$$E := |\mathcal{E}| = \frac{A}{2} \sum_{i \in \mathcal{N}} (A_i - 1) \quad (19)$$

Response graph This allows to draw a graph from the skeleton $(\mathcal{N}, \mathcal{A})$ of a normal form game by drawing a node for each pure strategy profile in \mathcal{A} , and an edge for each unilateral deviation in \mathcal{E} . This gives a graph with A nodes and E edges. In the language of section (1.1) (in particular cf. Example (1.2))

- \mathcal{A} is the basis of the chain group C_0
- \mathcal{E} is the basis of the chain group C_1

Example 2.1. Consider a game with skeleton $[2, 3]$:

- $\mathcal{N} = \{1, 2\}$, $N = 2$
- $\mathcal{A}_1 = \{1, 2\}$, $\mathcal{A}_2 = \{1, 2, 3\}$

The space of pure strategy profiles, with cardinality $A = 6$, is

$$\text{Basis of } C_0 = \mathcal{A} = \{(1, 1), (1, 2), (1, 3), (2, 1), (2, 2), (2, 3)\} \quad (20)$$

The space of unilateral deviations, with cardinality $E = 9$, is

$$\begin{aligned} \text{Basis of } C_1 = \mathcal{E} = \{ & [(1, 1), (1, 2)], [(1, 1), (1, 3)], [(1, 1), (2, 1)], \\ & [(1, 2), (1, 3)], [(1, 2), (2, 2)], [(1, 3), (2, 3)], \\ & [(2, 1), (2, 2)], [(2, 1), (2, 3)], [(2, 2), (2, 3)] \} \end{aligned} \quad (21)$$

The corresponding response graph is shown in Figure 2.

2.1 Matrix representation of ∂_1 and d_0

Having an explicit basis for C_1 and C_0 allows for an explicit representation of the matrices of the boundary and co-boundary by Eq. (2) and Lemma (1.7).

Since we're dealing with linear maps we only need to look at how they act on basis vectors, so to write the $(A \times E)$ -dimensional matrix of $\partial_1 : C_1 \rightarrow C_0$ let's look at how it acts on \mathcal{E} .

For a game with skeleton $[2, 3]$ ordering the basis \mathcal{A} and \mathcal{E} as in Example (2.1) yields

$$\begin{aligned} \partial_1 : C_1 &\rightarrow C_0 \\ [(1, 1), (1, 2)] &\mapsto (1, 2) - (1, 1) \\ [(1, 1), (1, 3)] &\mapsto (1, 3) - (1, 1) \\ &\vdots \\ [(2, 2), (2, 3)] &\mapsto (2, 3) - (2, 2) \end{aligned}$$

corresponding to the (6×9) matrix

$$[\partial_1] = \begin{pmatrix} -1 & -1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & -1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & -1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 1 \end{pmatrix}$$

Some guesswork will show that this algorithm works for arbitrary skeletons:

```

1  def make_boundary_1_matrix(self):
2      """Matrix of partial_1: C_1 --> C_0"""
3
4      # Start with transpose
5      A = np.zeros([int(self.dim_C1), int(self.dim_C0)])
6
7      for row in range(int(self.dim_C1)):
8          basis_edge = self.edges[row]
9
10         minus_node, plus_node = basis_edge
11         minus_column = self.nodes.index( minus_node )
12         plus_column = self.nodes.index( plus_node )
13         A[row][minus_column] = -1
14         A[row][plus_column] = +1
15
16     return A.transpose()

```

Once the matrix representing $\partial_1 : C_1 \rightarrow C_0$ is known, the matrix representing $d_0 : C_0 \rightarrow C_1$ is obtained for free as its transpose by Lemma (1.7).

2.2 Vector space of games and deviation map

Vector space of games When we refer to a *game* we usually mean a utility map given a fixed skeleton. In other words, the space of games U is the space of possible utility maps u given a skeleton $(\mathcal{N}, \mathcal{A})$. This is the assignment of N utilities (one per player) to A pure strategy profiles, that is an AN -dimensional vector space.

Recall that, given a graph, C_0 is the space of possible assignments of one number to each node in the graph. Since by definition the space of games is the space of possible assignments of N numbers to each node in the response graph, the space of games is the Cartesian product of N copies of C_0 :

$$U = \underbrace{C_0 \times \cdots \times C_0}_{N \text{ times}} \quad (22)$$

A basis for U is then given by the disjoint union of N copies of \mathcal{A} .

Example 2.2. Consider a game with skeleton $[2, 3]$ as in Example (2.1), and consider the following payoff map:

$$\begin{array}{ll}
 u : \mathcal{A} \rightarrow \mathbb{R}^2 & \\
 (1,1) \mapsto (-3,3) & \\
 (1,2) \mapsto (0,-5) & \\
 (1,3) \mapsto (-3,3) & \\
 (2,1) \mapsto (3,0) & \\
 (2,2) \mapsto (-3,0) & \\
 (2,3) \mapsto (0,1) &
 \end{array}
 \quad \xleftrightarrow{\text{bimatrix notation}} \quad
 \begin{pmatrix}
 -3,3 & 0,-5 & -3,3 \\
 3,0 & -3,0 & 0,1
 \end{pmatrix}$$

The dimension of U is $AN = 12$, and a basis is given by

$$\text{Basis of } U = \mathcal{A} \sqcup \mathcal{A} = \{(1,1)_1, (1,2)_1, (1,3)_1, (2,1)_1, (2,2)_1, (2,3)_1, \\ (1,1)_2, (1,2)_2, (1,3)_2, (2,1)_2, (2,2)_2, (2,3)_2\} \quad (23)$$

The payoff map u can be seen as the vector

$$u = -3(1,1)_1 + 0(1,2)_1 + \cdots + 0(2,2)_2 + 1(2,3)_2 \in U$$

Deviation map Given a game skeleton $(\mathcal{N}, \mathcal{A})$ we are finally in the position to map a game - that is, a vector $u \in U$ - to a 1-chain, or *flow*, that is a vector in the first chain group C_1 of the response graph built from the skeleton $(\mathcal{N}, \mathcal{A})$.

Recall that a flow is the assignment of a number to each edge of the response graph. A natural way to make these assignments given a utility map u is to label each edge by the utility difference of the deviating player:

Definition 2.3.

$$\begin{array}{ll} D : U \rightarrow C_1 & \\ u \mapsto Du & \text{s.t. } \forall [a, b] \in \mathcal{E} \quad (Du)_{ab} = u_i(b) - u_i(a) \\ & \text{for } i \text{ s.t. } a_i \neq b_i \end{array} \quad (24)$$

The map D is linear, and is called *deviation map*. Given a game $u \in U$, the flow $Du \in C_1$ is called *deviation flow of the game*.

In this definition $[a, b]$ is an edge in the response graph and $(Du)_{ab}$ is the number the flow Du assigns to such edge. By convention, the edges in the response graph are oriented such that $(Du)_{ab} \geq 0$ for all $[a, b] \in \mathcal{E}$.

The deviation flow of a game captures its strategic structure: the orientation of the edges of the response graph reflects the interest of each player at each state of the game. If an arrow leaves a node, a player following the arrow does not lose.

Example 2.4. Consider the normal form game of Example (2.2), and recall that its response graph is shown in Figure 2. The deviation flow of this game is shown in Figure 3.

2.3 Matrix representation of D

Having an explicit basis for U and C_1 one can write the matrix representing the deviation map $D : U \rightarrow C_1$, of dimension $(E \times AN)$.




Figure 3: Deviation flow of the game described in Example (2.2).

Example 2.5. Consider the usual game with skeleton $[2, 3]$ as in Example (2.1); recall that in this case $N = 2$, $A = 6$, and $E = 9$. With basis for C_1 and U given respectively by Eq. (21) and Eq. (23) one gets the (9×12) matrix

$$[D] = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & -1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 1 \end{bmatrix}$$

For arbitrary skeletons, the following algorithm does the trick:

```

1
2     self.payoff_basis = [ (i.player_name-1, a) for i in self.players for
3         a in self.strategy_profiles ]
4
5     def make_pwc_matrix(self):
6         """Matrix of deviation map: U --> C_1"""
7         A = np.zeros([int(self.dim_C1), int(self.dim_CON)])
8
9         for row in range(int(self.dim_C1)):
10             edge = self.edges[row]
11             i = utils.different_index(edge)
12
13             minus_column = self.payoff_basis.index( (i, edge[0]) )
14             plus_column = self.payoff_basis.index( (i, edge[1]) )
15             A[row][minus_column] = -1
16             A[row][plus_column] = +1
17
18     return np.asmatrix(A)

```

3 Hodge Decomposition of Normal Form Games

Essentials 3. The key points to take home:

- With the explicit matrices for D and d_0 at hand we can build two orthogonal projection operators $\pi : U \rightarrow U$ and $e : C_1 \rightarrow C_1$ as in Eqs. (44) and (45) using the Moore-Penrose inverse
- Eq. (47) gives the explicit decomposition receipt based on these projections

We can finally state our decomposition theorem for normal form games, first in an abstract form (up to a normalization choice), then in a concrete choice that allows for computation (after a suitable choice of normalization).

Definition 3.1. Consider the game skeleton $(\mathcal{N}, \mathcal{A})$ and the induced deviation map $D : U \rightarrow C_1$. The space of *non-strategic games* is the kernel of the deviation map:

$$\mathcal{K} := \ker D \subseteq U \quad (25)$$

Theorem 3.2 (Abstract Decomposition Theorem). *Consider the game skeleton $(\mathcal{N}, \mathcal{A})$. The space of games U admits the orthogonal decomposition*

$$U \cong \mathcal{K} \oplus \text{exact flows} \oplus \text{harmonic flows} \quad (26)$$

Note that in this theorem the exact flows and the harmonic flows live in C_1 , while \mathcal{K} lives in the space of games U . After making a normalization choice we will state the theorem with an equality, not an isomorphism, i.e. decompose U as the orthogonal direct sum of three subspaces.

To prove this theorem we need the following lemma.

3.1 Feasible flows are precisely closed flows

Lemma 3.3. *Consider the game skeleton $(\mathcal{N}, \mathcal{A})$ and the induced deviation map $D : U \rightarrow C_1$. The image of the deviation map is precisely the space of closed flows:*

$$\text{Im } D = \ker d_1 \subseteq C_1 \quad (27)$$

Definition 3.4. The image of the deviation map is the space of *feasible flows*.

Feasible flows are the elements of C_1 that can be realized as the deviation flow of some normal form game. The content of the lemma above is that the feasible flows are precisely the *closed flows*.




Figure 4: Left: every feasible flow is a closed flow. Right: every closed flow is feasible. See Ex. (3.5) for details.

What is a closed flow? By definition a flow $X \in C_1$ is closed iff $X \in \ker d_1$, that is iff $d_1 X = 0$. One can obtain an explicit expression for the co-boundary operator $d_1 : C_1 \rightarrow C_2$ analogue to that of the boundary operator $\partial_k : C_k \rightarrow C_{k-1}$ given in Eq. (2) (see e.g. Def. 4.2 in [Jia+11]), and it turns out that a flow $X \in C_1$ is closed iff

$$X_{ab} + X_{bc} + X_{cd} = 0$$

for any 3-clique $[abc]$ in the response graph, where X_{ab} is the number the flow X assigns to the edge $[ab]$.

Sketch proof of Lemma (3.3). The fact that $\text{Im } D \subseteq \ker d_1$ is proved in [Can+11, Eq. 26] employing a relatively heavy machinery, while we developed a simple argument.

Let $u \in U$, then

$$\begin{aligned} (d_1 Du)(abc) &= (Du)_{ab} + (Du)_{bc} + (Du)_{ca} \\ &= u_i(b) - u_i(a) + u_j(c) - u_j(b) + u_h(a) - u_h(c) \\ &= 0 \text{ since } (abc) \text{ is a 3-clique} \Rightarrow i = j = h \quad \square \end{aligned}$$

The key is that all the edges in clique have the same deviating player. This means that $d_1 \circ D \equiv 0$, i.e. every feasible flow is a closed flow. Note that this is in spirit a “derivation problem”.

The proof of the fact that $\text{Im } D \supseteq \ker d_1$ is, to our knowledge, original. Given a closed flow $X \in C_1$ we need to find a utility $u \in U$ such that $Du = X$. Note that this is in spirit an “integration problem”, i.e. a differential equation, which makes it typically harder than the opposite inclusion proved above.

The idea is to look at the deviation map $D : U = C_0^N \rightarrow C_1$ as a “higher dimensional version” of the coboundary map $d_0 : C_0 \rightarrow C_1$, and to factorize the response graph into complete sub-graphs that have a unique deviating player. This allows to decouple the system of equations $Du = X$ into sub-systems relative to these sub-graphs, and with this decomposition in place the problem is reduced to showing that if X is closed then it is exact on each complete sub-graph. This is true by Poincaré lemma [DLMo5], since

each complete sub-graph is contractible (seeing the response graph as a 2-dimensional simplicial complex). \square

Example 3.5. See Figure 4.

Left : $\text{Im } D \subseteq \ker d_1$

$$\begin{aligned} (d_1 Du)(abc) &= \\ &= Du(ab) + Du(bc) + Du(ca) \\ &= b - a + c - b + a - c = 0 \end{aligned}$$

Right : $\text{Im } D \supseteq \ker d_1$

$$\begin{aligned} 0 &= (d_1 X)(abc) = \\ &= X(ab) + X(bc) + X(ca) \\ &\Rightarrow \exists u : Du = X \end{aligned}$$

3.2 Proof of the Abstract Decomposition Theorem

With this result at hand, the proof of our abstract decomposition theorem (3.2) is immediate.

Proof of Theorem (3.2). Since $D : U \rightarrow C_1$ is a linear map it holds true that

$$U \cong \ker D \oplus \text{Im } D$$

and we are done already by Lemma (3.3) and Eq. (18):

$$\text{Im } D = \text{closed flows} = \text{exact flows} \oplus \text{harmonic flows}$$

$$U \cong \mathcal{K} \oplus \text{exact flows} \oplus \text{harmonic flows}$$

\square

3.3 Normalized Decomposition Theorem

To make the decomposition concrete one can choose *any* complement $\tilde{\mathcal{K}}$ of the space of non-strategic games \mathcal{K} in U . Indeed for any choice of $\tilde{\mathcal{K}}$ such that $U = \mathcal{K} \oplus \tilde{\mathcal{K}}$ we have the following general result:

Lemma 3.6. *Consider a linear map*

$$D : U \rightarrow C_1$$

and denote $\mathcal{K} = \ker D$. Given a direct sum decomposition of the image of D

$$\text{Im } D = A \oplus B$$

then for any complement \bar{K} of K in U we have

$$\bar{K} = \left(D^{-1}(A) \cap \bar{K} \right) \oplus \left(D^{-1}(B) \cap \bar{K} \right)$$

Proof. See Appendix (B.2). □

Since by Hodge theorem and by Lemma (3.3) $\text{Im } D = \text{closed flows} = \text{exact flows} \oplus \text{harmonic flows}$ we can set $A = \text{exact}$ and $B = \text{harmonic}$ in the lemma above, so

$$\bar{K} = \left(D^{-1}(\text{exact flows}) \cap \bar{K} \right) \oplus \left(D^{-1}(\text{harmonic flows}) \cap \bar{K} \right) \quad (28)$$

Definition 3.7. Consider the game skeleton $(\mathcal{N}, \mathcal{A})$ and the induced deviation map $D : U \rightarrow C_1$.

The space of *normalized potential games* is

$$\mathcal{P} := \left(D^{-1}(\text{exact flows}) \cap \bar{K} \right) \equiv \left(D^{-1}(\text{Im } d_0) \cap \bar{K} \right) \subseteq U \quad (29)$$

The space of *normalized harmonic games* is

$$\mathcal{H} := \left(D^{-1}(\text{harmonic flows}) \cap \bar{K} \right) \equiv \left(D^{-1}(\ker \Delta_1) \cap \bar{K} \right) \subseteq U \quad (30)$$

The space of *potential games* is

$$D^{-1}(\text{exact flows}) \equiv D^{-1}(\text{Im } d_0) \subseteq U \quad (31)$$

The space of *harmonic games* is

$$D^{-1}(\text{harmonic flows}) \equiv D^{-1}(\ker \Delta_1) \subseteq U \quad (32)$$

Lemma 3.8.

$$\text{potential games} = \mathcal{P} \oplus \mathcal{K} \quad (33)$$

$$\text{harmonic games} = \mathcal{H} \oplus \mathcal{K} \quad (34)$$

Furthermore, the space of potential games $\mathcal{P} \oplus \mathcal{K}$ is precisely the space of potential games in the standard sense of Monderer and Shapley [MS96], i.e.

$$u \in \mathcal{P} \oplus \mathcal{K} \iff \exists \phi \in C_0 \text{ s.t. } (Du)_{ab} = \phi(b) - \phi(a) \quad \forall [a, b] \in \mathcal{E}$$

We refer to a choice of a complement \bar{K} for the kernel \mathcal{K} of the deviation map D as *normalization*. Given a normalization we have the Normalized Decomposition Theorem:

Theorem 3.9 (Normalized Decomposition Theorem). *Consider the game skeleton $(\mathcal{N}, \mathcal{A})$. The space of games U admits the orthogonal decomposition*

$$U = \mathcal{P} \oplus \mathcal{K} \oplus \mathcal{H} \quad (35)$$

Proof. By definition of complement $U = \mathcal{K} \oplus \bar{\mathcal{K}}$, and the conclusion follows from Eq. (28) and by definition of \mathcal{P} and \mathcal{H} . \square

Harmonic games An important consequence of Lemma (3.3) is that harmonic games just depend on ∂_1 (and not also on d_1 , as one would expect). Recall by Eq. (16) that

$$\text{harmonic flows} = \ker \Delta_1 = \ker \partial_1 \cap \ker d_1$$

Now since $\text{Im } D = \ker d_1$ we have that

$$\text{Im } D \cap \ker \Delta_1 = \text{Im } D \cap \ker \partial_1 \Rightarrow D^{-1}(\ker \Delta_1) = D^{-1}(\ker \partial_1)$$

So by definition of harmonic games we have that

Lemma 3.10. *The space of normalized harmonic games is*

$$\mathcal{H} = \left(D^{-1}(\ker \partial_1) \cap \bar{\mathcal{K}} \right) \subseteq U \quad (36)$$

This is computationally very convenient because the matrix representation of $\partial_1 : C_1 \rightarrow C_0$ is relatively inexpensive to find (c.f. Sec. (2.1)), while the matrix representation of $d_1 : C_1 \rightarrow C_2$ involves finding the 3-cliques of the response graph, which is computationally more expensive for big graphs.

3.4 Orthogonal Normalization and Moore-Penrose Inverse

In [Can+11, Def. 4.1 and Lemma 4.5] the authors choose to use the *Euclidean orthogonal* complement of \mathcal{K} as normalization space:

$$\bar{\mathcal{K}} \stackrel{!}{=} \mathcal{K}^\perp \quad (37)$$

With this choice, the decomposition of a game can be performed explicitly as follows.

By definition of $U = \mathcal{K} \oplus \mathcal{K}^\perp$ any $u \in U$ admits the unique orthogonal decomposition

$$u = u_{\mathcal{K}} + \pi u \in U \quad (38)$$

where

$$\pi : U \rightarrow U \quad (39)$$

is the orthogonal projection onto $\text{Im } \pi = \mathcal{K}^\perp$ along $\ker \pi = \mathcal{K}$.

This provides the non-strategic part of the decomposition. To get the potential and the harmonic part, map $D : u \mapsto Du \in \text{Im } D = \text{closed flows} \subseteq C_1$. By Hodge theorem Du can then be orthogonally decomposed as

$$Du = D\pi u = e Du + (1 - e) Du \in \ker d_1 \subseteq C_1 \quad (40)$$

where

$$e : C_1 \rightarrow C_1 \quad (41)$$

is the orthogonal projection onto $\text{Im } e = \text{exact flows along } \ker e = (\text{exact flows})^\perp$.

The problem is then that of writing explicitly the orthogonal projections $\pi : U \rightarrow U$ and $e : C_1 \rightarrow C_1$; and to map the exact and harmonic components of Eq. (40) from C_1 back to U .

Moore-Penrose Inverse The Moore-Penrose Inverse provides a convenient way to achieve this.

Given a linear map $A : V \rightarrow W$ between vector spaces endowed with the Euclidean inner product there exists a unique linear map $\tilde{A} : W \rightarrow V$ fulfilling a bunch of neat properties (see [KS05; RAGo8]); what we care about here is that

$$\tilde{A}A : V \rightarrow V \quad (42)$$

is the orthogonal projection

- onto $\text{Im } \tilde{A}A = \text{Im } \tilde{A} = (\ker A)^\perp$
- along $\ker \tilde{A}A = \ker A$

and analogously

$$A\tilde{A} : W \rightarrow W \quad (43)$$

is the orthogonal projection

- onto $\text{Im } A\tilde{A} = \text{Im } A$
- along $\ker A\tilde{A} = \ker \tilde{A} = (\text{Im } A)^\perp$

If A is bijective then $\tilde{A} = A^{-1}$ is just the inverse of A ; convince yourself that in this case the properties above are true.

Projection operators Recall that $D : U \rightarrow C_1$ and $d_0 : C_0 \rightarrow C_1$. With this at hand we can build the projection operator π of Eq. (39) as

$$\pi = \tilde{D}D : U \rightarrow U \quad (44)$$

that is the orthogonal projection

- onto $\text{Im } \tilde{D}D = \text{Im } \tilde{D} = (\ker D)^\perp = \mathcal{K}^\perp$

- along $\ker \tilde{D}D = \ker D = \mathcal{K}$

as we need. Analogously we can build the projection operator e of Eq. (41) as

$$e = d_0 \tilde{d}_0 : C_1 \rightarrow C_1 \quad (45)$$

that is the orthogonal projection

- onto $\text{Im } d_0 \tilde{d}_0 = \text{Im } d_0 = \text{exact flows}$
- along $\ker d_0 \tilde{d}_0 = \ker \tilde{d}_0 = (\text{Im } d_0)^\perp = (\text{exact flows})^\perp$

as we need.

The decomposition, finally This gives an explicit receipt to decompose a utility map:

Proposition 3.11. *Given a game skeleton $(\mathcal{N}, \mathcal{A})$ any utility map $u \in U$ admits the unique orthogonal decomposition*

$$u = u_{\mathcal{K}} + u_{\mathcal{P}} + u_{\mathcal{H}} \quad (46)$$

with

$$\begin{aligned} u_{\mathcal{K}} &= u - \pi u \in \mathcal{K} \\ u_{\mathcal{P}} &= \tilde{D}eDu \in \mathcal{P} \\ u_{\mathcal{H}} &= u - u_{\mathcal{N}} - u_{\mathcal{P}} \in \mathcal{H} \end{aligned} \quad (47)$$

Proof. The first line is clear by definition (44) of π . In the second line it is clear by definition (45) of e that eDu is an exact flow, so it remains to show that

$$\tilde{D}eDu \in \mathcal{P} = \left(D^{-1}(\text{exact flows}) \cap \mathcal{K}^\perp \right)$$

Once this is done, the last line follows by Theorem (3.9).

For the first part, note that $D\tilde{D} : C_1 \rightarrow C_1$ is the orthogonal projection onto $\text{Im } D$, so it acts as the identity on feasible flows. Then, since eDu is exact (hence closed hence feasible), we have that

$$D(\tilde{D}eDu) = eDu$$

is exact, i.e.

$$\tilde{D}eDu \in D^{-1}(\text{exact flows})$$

For the second part, we have that $\text{Im } \tilde{D} = (\ker D)^\perp = \mathcal{K}^\perp$, so

$$\tilde{D}eDu \in \mathcal{K}^\perp$$

and we are done. □

Decomposition implementation The `Payoff.decompose_payoff()` method implements this decomposition (the deviation map D is denoted PWC in the code).

```

1  def decompose_payoff(self):
2
3      print('start decomposition')
4
5      u = self.payoff_vector
6
7      PI = self.game.normalization_projection
8      e = self.game.exact_projection
9
10     PWC_pinv = self.game.pwc_matrix_pinv
11     PWC = self.game.pwc_matrix
12     delta_0_pinv = self.game.coboundary_0_matrix_pinv
13     delta_0 = self.game.coboundary_0_matrix
14
15     print('this seems to be the bottleneck, big matrices
multiplication')
16
17     uN = u - PI @ u
18     print('first multiplication done')
19
20     uP = PWC_pinv @ e @ PWC @ u
21     print('three more multiplications done, this is slowest step!')
22
23     uH = u - uN - uP

```

The projection operators `game.normalization_projection` and `game.exact_projection` are computed in the `Game` class:

```

1      # Pseudo-Inverse and projection block
2      # Moore-Penrose pseudo-inverse of pwc
3      print('start PINV block')
4      self.pwc_matrix_pinv = npla.pinv(self.pwc_matrix)
5
6      # PI: CON --> CON projection onto Euclidean orthogonal complement
of ker(D)
7      self.normalization_projection = np.matmul(self.pwc_matrix_pinv,
self.pwc_matrix)
8
9      # pinv(d_0): C^1 --> C^0
10     self.coboundary_0_matrix_pinv = npla.pinv(self.
coboundary_0_matrix)
11
12     # e: C1 --> C1 projection onto exact
13     self.exact_projection = np.matmul(self.coboundary_0_matrix, self.
coboundary_0_matrix_pinv)

```

Potential function The potential function relative to the potential component $u_{\mathcal{P}}$ is

$$\phi = \tilde{d}_0 Du \in C_0 \quad (48)$$

with $u \in U$, $D : U \rightarrow C_1$, and $\tilde{d}_0 : C_1 \rightarrow C_0$.

A Dual and Adjoint in Linear Algebra

A.1 Dual Homomorphism

See [RAGo8]. Let A and G be abelian groups; for definiteness think of A as a vector space and G as the real numbers.

The *dual* of A is $A^* := \text{Hom}(A, G)$. This is an abelian group if we add two homomorphism in A^* adding their value in G .

Let B be another abelian group. A homomorphism $f : A \rightarrow B$ gives rise to a *dual homomorphism*

$$\begin{aligned} f^* : B^* &\rightarrow A^* \\ \phi &\mapsto f^*\phi = \phi \circ f \end{aligned} \tag{49}$$

meaning that this diagram commutes

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ & \searrow f^*\phi & \downarrow \phi \\ & & \mathbb{R} \end{array} \tag{50}$$

This construction is canonical.

A.2 Dual Isomorphisms

Let's endow A with an inner product

$$\langle \cdot, \cdot \rangle_A : A \times A \rightarrow \mathbb{R} \tag{51}$$

This induces a non-canonical isomorphisms $A \cong A^*$ by

$$\sharp_A : A^* \xrightarrow{\sim} A \tag{52}$$

such that

$$\phi(a) = \langle \phi^{\sharp_A}, a \rangle_A \in \mathbb{R}, \quad \forall \phi \in A^*, \forall a \in A \tag{53}$$

where the notation ϕ^{\sharp_A} is used for $\sharp_A(\phi)$. The inverse of \sharp_A is denoted by

$$(\sharp_A)^{-1} = \flat_A : A \xrightarrow{\sim} A^* \tag{54}$$

hence

$$u^{\flat_A}(a) = \langle u, a \rangle_A \in \mathbb{R}, \quad \forall u, a \in A \tag{55}$$

Finally, an inner product on A induces the inner product on A^*

$$\langle \cdot, \cdot \rangle_{A^*} : A^* \times A^* \rightarrow \mathbb{R} \tag{56}$$

such that

$$\langle \phi, \psi \rangle_{A^*} = \langle \phi^{\sharp_A}, \psi^{\sharp_A} \rangle_A \quad \forall \phi, \psi \in A^* \quad (57)$$

Analogously endowing B with an inner product

$$\langle \cdot, \cdot \rangle_B : B \times B \rightarrow \mathbb{R} \quad (58)$$

one defines

$$\sharp_B : B^* \xrightarrow{\sim} B, \quad (\sharp_B)^{-1} = \flat_B : B \xrightarrow{\sim} B^*, \quad \langle \cdot, \cdot \rangle_{B^*} : B^* \times B^* \rightarrow \mathbb{R} \quad (59)$$

A.3 Adjoint Homomorphism

Given inner products on A and B as above a homomorphism $f : A \rightarrow B$ gives rise to an *adjoint homomorphism*

$$f^\dagger : B \rightarrow A \quad (60)$$

such that

$$\langle fa, b \rangle_B = \langle a, f^\dagger b \rangle_A, \quad \forall a \in A, \forall b \in B \quad (61)$$

Lemma A.1. *The dual of f and the adjoint of f are related by*

$$f^\dagger = \sharp_A \circ f^* \circ \flat_B \quad (62)$$

that is, the following diagram commutes:

$$\begin{array}{ccc} A & \xleftarrow{f^\dagger} & B \\ \sharp_A \uparrow & & \downarrow \flat_B \\ A^* & \xleftarrow{f^*} & B^* \end{array} \quad (63)$$

Proof. Take $a \in A, b \in B$. Then

$$\begin{aligned} \mathbb{R} \ni \langle f^\dagger b, a \rangle_A &= \langle b, fa \rangle_B \\ &= \flat_B (fa) = (b^{\flat_B} \circ f)(a) \\ &= (f^* b^{\flat_B})(a) \\ &= \langle (f^* b^{\flat_B})^{\sharp_A}, a \rangle_A \end{aligned}$$

and we conclude by bilinearity and non-degeneracy of the inner product. \square

B Proofs

B.1 Proof of Hodge Decomposition given Hodge Theorem

Lemma B.1.

$$A : V \rightarrow W$$

$$\frac{V}{\ker A} \cong \operatorname{Im} A$$

Lemma B.2. *For any subspace S of V*

$$V \cong S \oplus \frac{V}{S}$$

Lemma B.3.

$$A : V \rightarrow W$$

$$V \cong \ker A \oplus \frac{V}{\ker A} \cong \ker A \oplus \operatorname{Im} A$$

Lemma B.4.

$$\text{co-exact} = (\text{closed})^\perp$$

$$\operatorname{Im} (d_1^\dagger) = (\ker d_1)^\perp$$

Definition B.5. - Cohomology group

$$H_1 := \frac{\text{closed}}{\text{exact}} = \frac{\ker d_1}{\operatorname{Im} d_0}$$

- Harmonic space

$$\mathcal{H}_1 = \{c \in C^1 : \Delta_1 c = 0\}$$

Theorem B.6 (Hodge). *The space of harmonic forms is isomorphic to the cohomology group*

$$H_1 \cong \mathcal{H}_1$$

Corollary B.7 (Hodge decomposition). *The cochain group C^1 decomposes (uniquely and orthogonally) as*

$$C^1 = \text{exact} \oplus \text{co-exact} \oplus \text{harmonic}$$

$$= \operatorname{Im} d_0 \oplus \operatorname{Im} d_1^\dagger \oplus \ker \Delta_1$$

Proof.

$$C^1 = \ker d_1 \oplus (\ker d_1)^\perp = \ker d_1 \oplus \operatorname{Im} d_1^\dagger$$

First identity by fact that $V = S \oplus S^\perp$ for any subspace $S \subseteq V$ by definition of orthogonal projection; second by Lemma (4) on image of adjoint map.

So one piece is done: $\text{Im } d_1^\dagger$ are co-exact.

It is left to show that

$$\text{closed} = \ker d_1 = \text{exact} \oplus \text{harmonic} = \text{Im } d_0 \oplus \ker \Delta_1$$

Since $\text{exact} \subseteq \text{closed}$ we can quotient and by Lemma (2) above

$$\text{closed} \cong \text{exact} \oplus \frac{\text{closed}}{\text{exact}}$$

$$\ker d_1 \cong \text{Im } d_0 \oplus \frac{\ker d_1}{\text{Im } d_0}$$

So just with standard linear algebra we can get as far as

$$C^1 \cong \text{exact} \oplus \frac{\text{closed}}{\text{exact}} \oplus \text{co-exact}$$

The crucial step is now Hodge theorem: there is a unique way to choose a harmonic representative in each cohomology group. So

$$C^1 \cong \text{exact} \oplus \text{harmonic} \oplus \text{co-exact}$$

□

B.2 Proof of Lemma (3.6)

Lemma. *Given a linear map*

$$D : U \rightarrow C_1$$

denote $\mathcal{K} = \ker D$. Given a direct sum decomposition of the image of D

$$\text{Im } D = A \oplus B$$

then for any complement \bar{K} of K in U we have

$$\bar{K} = \left(D^{-1}(A) \cap \bar{K} \right) \oplus \left(D^{-1}(B) \cap \bar{K} \right)$$

Proof. By definition of complement

$$U = \mathcal{K} \oplus \bar{K}$$

The key is that the restriction of D to \bar{K} gives a non-canonical isomorphism

$$D|_{\bar{K}} : \bar{K} \xrightarrow{\sim} \text{Im } D$$

since for any $Du \in \text{Im } D$ one can decompose $u = u_{\mathcal{K}} + u_{\bar{K}}$ with $u_{\bar{K}} \in \bar{K}$ and $Du = Du_{\bar{K}}$.

[To verify: not used in the proof, but I think the inverse of this iso is realized by the Moore-Penrose pseudoinverse \tilde{D} of D .]

Now the claim is that any $u_{\tilde{\mathcal{K}}} \in \tilde{\mathcal{K}}$ can be decomposed uniquely as $u_{\tilde{\mathcal{K}}} = \alpha + \beta$ for some $\alpha \in D^{-1}(A) \cap \tilde{\mathcal{K}}$ and $\beta \in D^{-1}(B) \cap \tilde{\mathcal{K}}$. So given any $u_{\tilde{\mathcal{K}}}$, then $Du_{\tilde{\mathcal{K}}} \in \text{Im } D$ can be decomposed uniquely as $Du_{\tilde{\mathcal{K}}} = a + b$ for $a \in A$ and $b \in B$. These two components can be mapped to $\tilde{\mathcal{K}}$ via the inverse of the isomorphism above to some α and β such that $D|_{\tilde{\mathcal{K}}}\alpha = a$ and $D|_{\tilde{\mathcal{K}}}\beta = b$. So $\alpha \in D^{-1}(A) \cap \tilde{\mathcal{K}}$ and $\beta \in D^{-1}(B) \cap \tilde{\mathcal{K}}$, and $D|_{\tilde{\mathcal{K}}}u_{\tilde{\mathcal{K}}} = Du_{\tilde{\mathcal{K}}} = a + b = D|_{\tilde{\mathcal{K}}}\alpha + D|_{\tilde{\mathcal{K}}}\beta = D|_{\tilde{\mathcal{K}}}(\alpha + \beta)$, so $u_{\tilde{\mathcal{K}}} = \alpha + \beta$ since $D|_{\tilde{\mathcal{K}}}$ is an isomorphism. \square

References

- [Can+11] Ozan Candogan et al. “Flows and Decompositions of Games: Harmonic and Potential Games”. In: *Mathematics of Operations Research* 36.3 (Aug. 2011), pp. 474–503. DOI: [10.1287/moor.1110.0500](https://doi.org/10.1287/moor.1110.0500) (cit. on pp. 2, 3, 6, 17, 20).
- [DLMo5] Mathieu Desbrun, Melvin Leok, and Jerrold E. Marsden. “Discrete Poincaré Lemma”. In: *Applied Numerical Mathematics*. Tenth Seminar on Numerical Solution of Differential and Differential-Algebraic Equations (NUMDIFF-10) 53.2 (May 2005), pp. 231–248. ISSN: 0168-9274. DOI: [10.1016/j.apnum.2004.09.035](https://doi.org/10.1016/j.apnum.2004.09.035). (Visited on 01/29/2023) (cit. on p. 17).
- [Eck44] Beno Eckmann. “Harmonische Funktionen und Randwertaufgaben in einem Komplex”. In: *Commentarii Mathematici Helvetici* 17.1 (Dec. 1944), pp. 240–255. ISSN: 1420-8946. DOI: [10.1007/BF02566245](https://doi.org/10.1007/BF02566245). (Visited on 09/12/2023) (cit. on p. 2).
- [Fri96] Joel Friedman. “Computing Betti Numbers via Combinatorial Laplacians”. In: *Proceedings of the Twenty-Eighth Annual ACM Symposium on Theory of Computing - STOC '96*. Philadelphia, Pennsylvania, United States: ACM Press, 1996, pp. 386–391. ISBN: 978-0-89791-785-8. DOI: [10.1145/237814.237985](https://doi.org/10.1145/237814.237985). (Visited on 09/12/2023) (cit. on p. 2).
- [Jia+11] Xiaoye Jiang et al. “Statistical Ranking and Combinatorial Hodge Theory”. In: *Mathematical Programming* 127.1 (2011), pp. 203–244 (cit. on pp. 2, 17).
- [KS05] K. Kamaraj and K. C. Sivakumar. “Moore-Penrose Inverse in an Indefinite Inner Product Space”. In: *Journal of Applied Mathematics and Computing* 19.1 (Mar. 2005), pp. 297–310. ISSN: 1865-2085. DOI: [10.1007/BF02935806](https://doi.org/10.1007/BF02935806). (Visited on 02/13/2023) (cit. on p. 21).
- [MS96] Dov Monderer and Lloyd S. Shapley. “Potential Games”. In: *Games and Economic Behavior* 14.1 (May 1996), pp. 124–143. ISSN: 0899-8256. DOI: [10.1006/game.1996.0044](https://doi.org/10.1006/game.1996.0044). (Visited on 02/19/2023) (cit. on p. 19).
- [Mun84] James R Munkres. *Elements of Algebraic Topology*. Perseus Books, 1984 (cit. on pp. 2, 4).
- [RAGo8] Steven Roman, S Axler, and FW Gehring. *Advanced Linear Algebra*. 3rd ed. Springer, 2008 (cit. on pp. 21, 24).