

1. Recast each problem below as a problem in finding a solution  $\vec{x}$  to the vector equation  $A\vec{x} = \vec{b}$ , where  $A$  is a matrix and  $\vec{x}$  and  $\vec{b}$  are vectors. For each, define identify  $A$ ,  $\vec{x}$  and  $\vec{b}$ . Do not solve.

a. Determine if  $\begin{bmatrix} 7 \\ 5 \\ 10 \\ 6 \end{bmatrix}$  is a linear combination of  $\begin{bmatrix} 3 \\ 1 \\ 2 \\ 2 \end{bmatrix}$ ,  $\begin{bmatrix} 1 \\ 2 \\ 3 \\ 1 \end{bmatrix}$ , and  $\begin{bmatrix} 2 \\ 3 \\ 3 \\ 1 \end{bmatrix}$ .

If we label the first vector as  $\vec{w}$  and the triplet of vectors  $\vec{v}_1$ ,  $\vec{v}_2$ , and  $\vec{v}_3$ , then this problem is equivalent to determining whether there are scalar

coefficients  $x_1$ ,  $x_2$  and  $x_3$  so that  $\vec{w} = x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3 = [\vec{v}_1 | \vec{v}_2 | \vec{v}_3] \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$ .

Let  $A = [\vec{v}_1 | \vec{v}_2 | \vec{v}_3] = \begin{bmatrix} 3 & 1 & 2 \\ 1 & 2 & 3 \\ 2 & 3 & 3 \\ 2 & 1 & 1 \end{bmatrix}$ ,  $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$ , and  $\vec{b} = \vec{w} = \begin{bmatrix} 7 \\ 5 \\ 10 \\ 6 \end{bmatrix}$  and we must

determine whether  $A\vec{x} = \vec{b}$  has any solutions.

b. Find a vector in  $\mathbf{R}^3$  whose scalar products with  $\begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$ ,  $\begin{bmatrix} 2 \\ 3 \\ 1 \end{bmatrix}$ , and  $\begin{bmatrix} 4 \\ 1 \\ 2 \end{bmatrix}$

are, respectively, 3, 4, and 5.

Let  $\vec{x}$  be the vector sought, let  $\vec{b}$  be the 3-vector whose components are the specified scalar products, and let  $A$  be the matrix whose rows are the specified vectors. So, this problem is equivalent to seeking the solutions  $\vec{x}$

to  $A\vec{x} = \vec{b}$  where  $A = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 1 \\ 4 & 1 & 2 \end{bmatrix}$  and  $\vec{b} = \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix}$ .

c.  $F = \frac{1}{3} \begin{bmatrix} 2 & 1 & -2 \\ 1 & 2 & 2 \\ -2 & 2 & -1 \end{bmatrix}$  is the matrix for a reflection across a plane

through the origin in  $\mathbf{R}^3$ . Find all vectors normal (perpendicular) to this plane by using the fact that the image of a normal vector  $\vec{x}$  is  $-\vec{x}$ .

We wish to solve  $F\vec{x} = -\vec{x} = -I\vec{x}$  for  $\vec{x}$  or, equivalently, the equation

$$F\vec{x} + I\vec{x} = (F + I)\vec{x} = \vec{0}. \text{ So, we let } A = F + I = \frac{1}{3} \begin{bmatrix} 5 & 1 & -2 \\ 1 & 5 & 2 \\ -2 & 2 & 4 \end{bmatrix} \text{ and}$$

$$\vec{b} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \text{ and solve } A\vec{x} = \vec{b}.$$

2. Given  $[A|\vec{b}]_{rref}$  in each case below, solve  $A\vec{x} = \vec{b}$  completely.

$$\text{a. } [A|\vec{b}]_{rref} = \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{array} \right].$$

We have a single (unique) solution  $\vec{x} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$ .

$$\text{b. } [A|\vec{b}]_{rref} = \left[ \begin{array}{cccc|c} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 2 & 0 \\ 0 & 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right].$$

This system has no solution since the last column is a pivot column. Alternately, we note that the last row is equivalent to the inconsistent equation  $0 = 1$ .

$$\text{c. } [A|\vec{b}]_{rref} = \left[ \begin{array}{ccccc|c} 1 & 2 & 0 & 0 & 3 & 6 \\ 0 & 0 & 1 & 0 & 4 & 7 \\ 0 & 0 & 0 & 1 & 5 & 8 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right].$$

$$\vec{x} = \begin{bmatrix} 6 \\ 0 \\ 7 \\ 8 \\ 0 \end{bmatrix} + \alpha \begin{bmatrix} 2 \\ -1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + \beta \begin{bmatrix} 3 \\ 0 \\ 4 \\ 5 \\ -1 \end{bmatrix}, \text{ where } \alpha \text{ and } \beta \text{ are any reals.}$$

3. Consider the equation  $A\vec{x} = \vec{b}$  where  $A$  is an  $m \times n$  matrix,  $\vec{x}$  is in  $\mathbf{R}^n$  and  $\vec{b}$  is in  $\mathbf{R}^m$ . Show that this equation cannot have a finite number of solutions greater than 1 by proceeding as follows.

a. First, define what is meant by:  $\vec{w}$  is a solution of the equation above.  
 $A\vec{w} = \vec{b}$ .

b. Suppose that  $\vec{u}$  and  $\vec{v}$  are distinct solutions to  $A\vec{x} = \vec{b}$ . Let  $\vec{z}$  be the difference between  $\vec{v}$  and  $\vec{u}$ . Note that  $\vec{z} \neq \vec{0}$ . What is  $A(\alpha \vec{z})$  for any  $\alpha$  in  $\mathbf{R}$ ?

$$A(\alpha \vec{z}) = \alpha A\vec{z} = \alpha(A\vec{u} - A\vec{v}) = \alpha(\vec{b} - \vec{b}) = \alpha\vec{0} = \vec{0}.$$

c. Now, show that  $\{\vec{u} + \alpha \vec{z} \mid \alpha \in \mathbf{R}\}$  is a set with infinitely many (different) vectors each of which is a solution to  $A\vec{x} = \vec{b}$ .

Since  $A\vec{u} = \vec{b}$  and  $A(\alpha \vec{z}) = \vec{0}$ , we have  $A(\vec{u} + \alpha \vec{z}) = A\vec{u} + A(\alpha \vec{z}) = \vec{b} + \vec{0} = \vec{b}$  for all real  $\alpha$ . Since  $\vec{z} \neq \vec{0}$ , distinct  $\alpha$ 's give distinct solutions.

4. a. Suppose that  $f: \mathbf{R}^2 \rightarrow \mathbf{R}^2$  is linear,  $f$  triples vectors parallel to  $\begin{bmatrix} 1 \\ 2 \end{bmatrix}$  and doubles vectors parallel to  $\begin{bmatrix} 3 \\ 2 \end{bmatrix}$ . Find  $f\begin{bmatrix} x \\ y \end{bmatrix}$  for any real  $x$  and  $y$ .

Let  $A$  be the  $2 \times 2$  matrix corresponding to  $f$ . Then,  $A\begin{bmatrix} 1 & 3 \\ 2 & 2 \end{bmatrix} = \begin{bmatrix} 3 & 6 \\ 6 & 4 \end{bmatrix}$ .

$$\text{So, } A = \begin{bmatrix} 3 & 6 \\ 6 & 4 \end{bmatrix} \begin{bmatrix} 1 & 3 \\ 2 & 2 \end{bmatrix}^{-1} = \begin{bmatrix} 3 & 6 \\ 6 & 4 \end{bmatrix} \frac{1}{2-6} \begin{bmatrix} 2 & -3 \\ -2 & 1 \end{bmatrix} = \frac{-1}{4} \begin{bmatrix} -6 & -3 \\ 4 & -14 \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 6 & 3 \\ -4 & 14 \end{bmatrix}.$$

$$\text{Therefore, } f\begin{bmatrix} x \\ y \end{bmatrix} = A\begin{bmatrix} x \\ y \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 6 & 3 \\ -4 & 14 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \frac{3}{2}x + \frac{3}{4}y \\ -x + \frac{7}{2}y \end{bmatrix}.$$

b. Solve for the matrix  $X$  if  $\begin{bmatrix} 1 & 1 \\ 1 & 3 \end{bmatrix} X^{-1} \begin{bmatrix} 1 & 1 \\ 1 & 3 \end{bmatrix}^{-1} = \begin{bmatrix} 5 & 2 \\ -3 & 2 \end{bmatrix}^{-1}$ .

Inverting both sides, we have  $\begin{bmatrix} 1 & 1 \\ 1 & 3 \end{bmatrix} X \begin{bmatrix} 1 & 1 \\ 1 & 3 \end{bmatrix}^{-1} = \begin{bmatrix} 5 & 2 \\ -3 & 2 \end{bmatrix}$ . Now,

multiplying on the left by  $\begin{bmatrix} 1 & 1 \\ 1 & 3 \end{bmatrix}^{-1}$  and on the right by  $\begin{bmatrix} 1 & 1 \\ 1 & 3 \end{bmatrix}$ , we get

$$X = \begin{bmatrix} 1 & 1 \\ 1 & 3 \end{bmatrix}^{-1} \begin{bmatrix} 5 & 2 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 3 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 3 & -1 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 5 & 2 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 3 \end{bmatrix} = \begin{bmatrix} 11 & 15 \\ -4 & -4 \end{bmatrix}.$$

5. For each assertion below, indicate whether the assertion is true (**T**) or false (**F**). It is not necessary to show any work. However, a correct response earns full credit, an incorrect response earns negative half credit, and no response earns no credit.

a. A linear system of 7 equations in 8 variables always has infinitely many solutions.

**False.** It may have no solutions.

b. A  $4 \times 4$  matrix of rank 4 is always invertible.

**True.** This assertion is equivalent to the assertion that the matrix is row-reducible to the identity.

c. Given any three nonzero vectors in  $\mathbf{R}^2$ , no two of which are collinear, any one of them is always a linear combination of the other two.

**True.** This was established by a geometrical argument in class and in homework. Any pair of non-collinear vectors in  $\mathbf{R}^2$  spans  $\mathbf{R}^2$ .

d. The function  $f : \mathbf{R}^2 \rightarrow \mathbf{R}^2$  defined by  $f \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} y \\ 1 \end{bmatrix}$  is linear.

**False.** Clearly,  $f(\vec{0}) \neq \vec{0}$ . It is also easy to show that, under  $f$ , the image of a sum is not the corresponding sum of the images and the image of a multiple is not that multiple of the image.

e. There is a  $2 \times 2$  matrix  $A$ , different from the identity matrix, such that  $A^{2008}$  is the identity matrix.

**True.** There are many such matrices. For example, let  $A = -I$  or let  $A$  be a reflection across any line in  $\mathbf{R}^2$ , or let  $A$  be the matrix for rotation by  $\frac{2\pi}{2008}$ , or let  $A$  be the matrix for rotation by  $\frac{2\pi N}{2008}$  where  $N$  is any integer.

f.  $f$  and  $g$  are two linear transformations defined by  $f(\vec{x}) = A\vec{x}$  and  $g(\vec{x}) = B\vec{x}$  where  $A$  and  $B$  are  $3 \times 3$  matrices. The composite function  $f \circ g$  is also a linear transformation and its matrix is  $AB$ .

**True.** This is precisely the definition for the matrix of a composite.

g. No invertible  $10 \times 10$  matrix can have more than 90 ones among its entries.

**False.** A  $10 \times 10$  matrix with 92 or more ones would have 8 or fewer non-one entries. If the matrix had 8 or fewer non-one entries, these entries could be distributed over at most 8 rows. That would mean that, at least two rows would consist of all ones. Subtracting one of these rows of all ones

from the other would result in a row of all zeros. Therefore, the rref of such a matrix would have a row of zeros. Such a matrix would not be row-equivalent to the identity and it would therefore not be invertible. Among the  $10 \times 10$  matrices with exactly 91 ones are the following.

$$A = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 0 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 0 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 0 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 0 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 0 & 1 \end{bmatrix}$$

It is not hard to see that  $A_{rref} = I_{10}$  and so  $A$  is invertible. [Can you see how to show this? Hint: Begin by subtracting the second row from all the other rows.]