

The dynamics of neutral and trait-based multi-trophic interactions

Timothée Poisot^{1,2} and Dominique Gravel^{1,2}

¹Université du Québec à Rimouski, Rimouski, Canada.

²Quebec Center for Biodiversity Science, McGill University, Montréal, Canada.

*email: timothee.poisot@uqar.ca

Abstract

We describe *manna*, a simple modeling framework of individuals establishing trophic interactions under neutral or trait-based rules. We use this model to show that neutral and trait-based interactions differ in their temporal dynamics, and discuss how our results contribute to highlighting the need to understand spatial and temporal dynamics of species interaction networks.

1 Introduction

traits and neutral effects on networks

need to account for dynamics through space and time

simple modelling framework

main results: turnover of pred/prey, contribution to sp turnover, role of connectance

2 The model

We define an individual based model of multi-trophic interactions in a single patch. Species are defined by a vector of functional traits, them being their niche position (\mathbf{n}_i), the centroid of their feeding range (\mathbf{c}_i), the breadth of their range (\mathbf{r}_i), and their maximal population size (\mathbf{K}_i).

Some words on the logic

2.1 Generation of the species pool

The species pool is generated so as to obtain a number of species S , with an expected connectance value of Co . To generate the species pool, we draw at random in an uniform distribution ranging from 0 to 1, S values forming the vector \mathbf{n} . The vectors \mathbf{c} and \mathbf{r} are generated following the method described by Williams & Martinez [1]. **Say more.** The number of individuals in each species is \mathbf{p}_i

As a final step, the values of \mathbf{K}_i are drawn at random from the uniform distribution between 0 and $K_{\max}(i)$, where $K_{\max}(i)$ is simply $10^3 \times (1 - \mathbf{n}_i) + 100$. This way, species with

the large \mathbf{n}_i have lower carrying capacity, and species with low \mathbf{n}_i have high carrying capacity. This accounts for the well known negative relationship between trophic rank and population size. To initialize the simulation, a number of individuals \mathbf{p}_i is drawn from the uniform distribution from 10 to \mathbf{K}_i .

2.2 Simulation

Before the interactions happen and the demographic changes are calculated, there is a (possible) immigration step, to implicitly simulate some spatial context and maintain species richness. The identity of the species receiving each migrant is drawn at random. This makes it so that locally extinct species can be rescued through immigration. After the immigration is over, the probability that each individual will reproduce or die is calculated given the following set of rules. The mortality probability of individual i is

$$m_i = \mu + s_i c_\mu \quad (1)$$

, while its natality probability is

$$b_i = \nu - s_i c_\nu \quad (2)$$

, and s_i is a scaling factor related to the niche position, wherein

$$1 - (1 - \mathbf{n}_i)^k \quad (3)$$

. The exponent k regulates the strength of the scaling. The population size of species i at time $t + 1$ is calculated as

$$\min(\mathbf{K}_i, \mathbf{p}_i - \mathcal{B}(m_i, \mathbf{p}_i) + \mathcal{B}(b_i, \mathbf{p}_i)) \quad (4)$$

, where $\mathcal{B}(p, n)$ is the number of successes of a Bernoulli process with probability of success of p and n trials.

After the demographic changes are done, we pick at random 15% of the individuals, which will be seeking a prey at this timestep. Each individual has an equal probability of being picked, so that species with high abundances will be over-represented among the pools of predators. The identity of preys items is detailed in the next two sub-sections. The interactions are done on an individual basis, meaning that we pick a predator, then its prey, and reproduce this routine once we have reached enough predators.




Figure 1: Comparison of the structure of networks under the neutral (left) and trait-based (right) approaches. When the only rule for the existence of an interaction is that the two populations are large enough to interact (neutral rules), we expect more interactions than when traits are used to delineate a feeding range (trait-based rules). These trait-forbidden links are dashed in the right network. Switching from one situation to the other can lead to complex effects, both on networks dynamics and community composition.

2.3 Neutral interactions

In the neutral scenario, the only factor regulating the identity of the prey is its abundance. Once the predating individual is selected, we pick at random one individual from any species (including the possibility of cannibalism) in the pool. The predated individual is then removed.

2.4 Trait-based interactions

In the trait-based scenario, preys are selected only among the species which can be fed upon, as in the *niche* model of Williams & Martinez [1]. Preys are still selected on the basis of their abundances, *i.e.* relative to the abundance of all other preys falling within the feeding range of the predator.

2.5 Implementation and availability

A C99 implementation of the model, using the Gnu Scientific Library, is available at BitBucket - `manna`, under the conditions of the WTFPL 2.0.

3 Methods

3.1 Simulations

We used the model to generate species pools of size $S = 50$, with connectance varying from 0.01 to 0.5 by varying increments chosen to represent the changes induced by connectance. 5 individuals migrated into the system at each timestep. Each community is simulated under the neutral and trait based assumptions, and replicated 20 times. Simulations are done over 5×10^3 timesteps, as preliminary analyses showed it was long enough to reach an equilibrium in

number of persisting species, number of persisting links, and overtime dissimilarity.

3.2 Analyses

Networks were aggregated in bins of 10 timesteps for the analysis (exploratory work showed that the bin size made no quantitative difference on the results, though it did considerably speed up the calculations). Within each bin, we measure the number of species in the network (\mathcal{S}), the number of links (\mathcal{L}), the effective connectance (\mathcal{C} , expressed as $\mathcal{L}/\mathcal{S}^2$).

Within each run of the simulation, we measure components of network dissimilarity through time, using the method described in Poisot *et al.* [2]. We measure turnover of interactions and community structure between the network at time t and the network at time $t + 1$. We report the overall network dissimilarity (β_{WN}), the dissimilarity of links across shared species (β_{OS}), the species turnover (β_S), and its contribution to network turnover (β_{contrib}). In addition, we report the turnover of predators (β_U) and preys (β_L) through time.

4 Results

4.1 Community dynamics

Regardless of the rules or connectance, all simulation conditions yield a common pattern: a rapid decrease in the number of species (*i.e.* sorting) and links (Fig. 2). This results in changes in connectance, with all the simulations eventually stabilizing around a value of $\mathcal{C} \approx 0.05$. The number of species maintained is higher in the neutral conditions. When connectance increases, the number of persisting species increases as well. For a connectance of $\mathcal{C}_0 = 0.5$, there are almost no differences in species richness between the neutral and trait-based situations. The same is true of the number of interactions.

4.2 Networks dynamics

5 Discussion

1. summary of the differences
2. inferring mechanisms with temporal dynamics
3. role of connectance

References

1. Williams, R. & Martinez, N. Simple rules yield complex food webs. *Nature* **404**, 180–183 (2000).

2. Poisot, T., Canard, E., Mouillot, D., Mouquet, N. & Gravel, D. The dissimilarity of species interaction networks. *Ecology Letters* **15**, 1353–1361 (Sept. 2012).




Figure 2: Temporal dynamics of network structure, as a function of whether the system behaves neutrally (black lines) or under trait-based rules (colored lines). The first 2500 timesteps are shown, all simulations reached equilibrium at approx. $t = 1000$. **A.** Number of unique species (S). **B.** Number of unique interactions (L). **C.** Connectance of the network (C).




Figure 3: Temporal dynamics of network turnover, as a function of whether the system behaves neutrally (black lines) or under trait-based rules (colored lines). The first 2500 timesteps are shown, all simulations reached equilibrium at approx. $t = 1000$. **A.** Whole-network dissimilarity (β_{WN}) remains high under the neutral scenario. **B.** The same is true of interaction turnover across shared species, β_{OS} . **C.** Species turnover displays a hump-shaped pattern with connectance (see also [FIGURE XXX](#)). **D.** The impact of species turnover on interactions turnover is low in the neutral model, but increasingly higher when trait-based rules are used and connectance decreases.