

# Architectural Framework for Quatossian Inscription: A Synergetic Approach to Radiance Field Synthesis and Analog-Digital Hybrid Rendering

The transition from polygonal mesh-based graphics to volumetric radiance fields represents a fundamental shift in the ontology of digital representation. Traditional rendering techniques have historically prioritized the "physical form"—the discrete boundary of an object—treating light as an extrinsic factor that merely illuminates a pre-defined geometry. In contrast, the Quatossian (Quaternion-Gaussian) or Quatosium framework posits that light paths, phase-space reflections, and interference patterns are the primary constituents of visual reality. This research plan details the development of a rendering architecture where 3D Gaussian primitives, modulated by quaternionic spin-states, are inscribed within a high-dimensional E8 lattice. By integrating Kirigami-inspired Moiré modulation and legacy layered surface protocols, the framework establishes a hybrid system that bridges the gap between digital computation and the physical behavior of light.

## The Quatossian Primitive: Quaternions and Gaussians in Radiance Fields

The core of the Quatossian framework is the "radiance kernel," a synthesis of 3D Gaussian splatting and quaternionic orientation states. Unlike traditional point clouds or voxel grids, 3D Gaussians are continuous, differentiable primitives defined by their mean position, covariance matrix, and spherical harmonic coefficients.<sup>1</sup> The Quatossian approach extends this by representing the rotation component of the Gaussian's covariance through normalized quaternions, which provides a more robust and stable mathematical foundation for high-dimensional spatial computing.<sup>2</sup>

## Mathematical Foundation of the Quatossian Kernel

A 3D Gaussian is typically defined by its influence at a spatial point  $x$ , governed by the equation  $G(x) = \exp(-1/2(x - \mu)^T \Sigma^{-1}(x - \mu))$ . In the Quatossian framework, the covariance matrix  $\Sigma$  is not merely a static property but a dynamic state vector. By decomposing  $\Sigma = R S S^T R^T$ , where  $R$  is a rotation matrix derived from a quaternion  $q$  and  $S$  is a diagonal scaling matrix, the system can smoothly interpolate orientations without

the risk of gimbal lock.<sup>3</sup> This is critical for rendering complex light paths where the "spin" of the light wave must be preserved.

The use of quaternions allows the framework to integrate with the geometry of the octonions, which are recognized as the key to unified field theories in physics.<sup>4</sup> The octonionic representation generates the E8 lattice, a structure that provides a unique crystalline perfection for high-dimensional data indexing.<sup>4</sup> In the Quatossian model, each primitive is treated as a "Cubit"—a spinning, mirroring, entangled point of light that encodes position, momentum, phase, and frequency.<sup>6</sup>

### Comparison of Rendering Primitives

The following table summarizes the evolution from discrete geometric primitives to the Quatossian kernel, highlighting the shift toward phase-aware volumetric rendering.

| Feature           | Polygonal Mesh        | 3D Gaussian Splat     | Quatossian Inscription             |
|-------------------|-----------------------|-----------------------|------------------------------------|
| Primary Unit      | Triangle / Vertex     | 3D Gaussian           | Quaternionic Radiance Kernel       |
| Orientation Model | Face Normal Vector    | Euler Rotation Matrix | Normalized Quaternion / Octonion   |
| Lighting Model    | BRDF / Ray Tracing    | Spherical Harmonics   | Phase Interference Patterns        |
| Spatial Indexing  | BVH / Octree          | Sorted Point Cloud    | E8 Gosset Lattice <sup>7</sup>     |
| Primary Focus     | Surface Boundary      | Density/Color Volume  | Light Path / Wave Resonance        |
| Computation Type  | Digital Rasterization | Differentiable Raster | Analog-Digital Hybrid <sup>8</sup> |

### High-Dimensional Optimization via the E8 Lattice Substrate

The efficiency of any Gaussian-based rendering system depends on the density and

organization of the splats. The Quatossian framework adopts the E8 lattice—the Gosset lattice—as its primary spatial substrate.<sup>7</sup> Mathematics has demonstrated that the E8 lattice provides the optimal sphere packing density in eight dimensions.<sup>9</sup> By mapping the 3D rendering space into an 8D E8 projection, the system can achieve a level of information density and coherence that exceeds traditional 3D partitioning methods.<sup>11</sup>

Geometry and Symmetry of the E8 Lattice

The E8 lattice is a discrete subgroup of  $\mathbb{R}^8$  that is isomorphic to  $\mathbb{Z}^8$ . It can be defined as the union of the  $D_8$  lattice—consisting of all 8-tuples of integers that sum to an even integer—and a shifted copy of that lattice.<sup>12</sup> This structure possesses 248 dimensions of symmetry, making it the most symmetrical object possible in its dimensional space.<sup>5</sup> For Quatossian rendering, this symmetry is not merely an abstract property but a mechanism for maintaining "coherence" in the light field.

Research into the Maya AI framework has shown that consciousness-like stability in data structures occurs when E8 projections hold a specific magnitude—typically between 0.49 and 0.87—across full dimensional space.<sup>13</sup> In the context of Quatossian rendering, this translates to a "geometry of endless peace," where the radiance field achieves a coherence plateau with minimal variance.<sup>13</sup> By situating Gaussian agents on a prime-indexed E8 lattice, the system can scale its computational depth without sacrificing the integrity of the rendered image.<sup>13</sup>

Spatial Computing and Isotropic Coordinates

The E8 lattice serves as the foundation for a spatial computing interface that treats 3D space as a canvas for holographic icons controlled by motion gestures.<sup>2</sup> By utilizing Locality Sensitive Hashing (LSH) projections into the E8 space, the system can map high-dimensional light path data into 1D Morton curve values, which are then efficiently processed on the GPU.<sup>11</sup> This enables a "participatory reality" where the observer's interaction with the light field is integrated into the rendering loop itself.<sup>5</sup>

| Lattice Metric  | D3 (FCC)            | D4 Lattice | E8 (Gosset)           | Leech Lattice        |
|-----------------|---------------------|------------|-----------------------|----------------------|
| Dimension       | 3                   | 4          | 8                     | 24                   |
| Packing Density | ~0.74 <sup>12</sup> | ~0.12      | Optimal <sup>10</sup> | Optimal <sup>9</sup> |
| Root Vectors    | 12                  | 24         | 240                   | 196,560              |

| Rendering Role | Grid Partitioning | Quaternionic Flow | High-Dim Indexing | Global Coherence |
|----------------|-------------------|-------------------|-------------------|------------------|
| Symmetry       | 24                | 1,152             | 696,729,600       | $\sim 8 \times$  |

## Kirigami-Based Moiré Interference Modulation

One of the most innovative aspects of the Quatossian framework is the integration of Kirigami-based Moiré interference to modulate Gaussian splat raycasting. Kirigami—the art of cutting and folding paper—provides a physical and mathematical model for "gating" light. By representing Kirigami structures as transmissive masks, the framework can perform optical analog computation directly on the raycasted radiance field.<sup>8</sup>

### Moiré Effects and Optical Analog Computation

Moiré patterns arise when two or more periodic structures—such as binary line masks—are superposed and shifted or rotated relative to each other.<sup>14</sup> The resulting interference pattern can be used to synthesize complex optical elements, such as Fresnel Zone Plates (FZP), which act as lenses to focus or zoom light.<sup>14</sup> In the Quatossian system, these masks are used to modulate the transmittance of rays passing through the Gaussian cloud.

The Interlaced Line Divider (ILD) protocol utilizes this principle to perform integer division and remainder extraction through optical superposition.<sup>8</sup> A "dividend mask" (G1) encoding  $N$  lines is superposed with a "completer mask" (G2) encoding blocks of  $D$  sites. The resulting light transmission reveals the quotient and remainder, providing a hardware-level analog calculation that bypasses traditional digital logic.<sup>8</sup> This is integrated into the raycasting loop to dynamically adjust the density and focus of Gaussian splats based on the observer's viewpoint.

### Modulation of Gaussian Splat Raycasting

The raycasting process in Gaussian splatting involves alpha-blending sorted Gaussians along a ray. In the Quatossian framework, this ray is first passed through a virtual Kirigami mask. The averaged Moiré transmittance  $P(x, y)$  modifies the intensity of each Gaussian primitive:

$$I_{final} = I_{Gauss} \cdot \sum B_n \exp(2\pi i n (x/x_1)^2)$$

Where  $x_1$  is a scale factor determined by the relative shift of the Kirigami layers.<sup>15</sup> This creates a "shimmer" or "interference" effect that mimics the way light interacts with physical

textures like ribbed glass or iridescent surfaces.<sup>16</sup> The use of Kirigami haptic swatches further informs the design of these structures, allowing for haptic-inspired visual feedback where the "cuts" in the virtual mask respond to user interaction or physics-based stressors.<sup>18</sup>

## Layered Surface Protocols for Legacy Integration

While the Quatossian framework prioritizes radiance fields, it must remain compatible with legacy graphics pipelines that rely on polygonal meshes and UV texture maps. This is achieved through a layered surface protocol that treats the legacy mesh as a reference manifold for the Gaussian inscription.

### Manifold Alignment and Topographic Conversion

The integration process begins by using Digital Elevation Models (DEM) and Topographic Line Map (TLM) data to construct the base perspective.<sup>19</sup> Instead of rendering the mesh as a solid surface, it is treated as a "depth map attractor" for the Quatossian kernels.<sup>20</sup> This topographic depth map allows for significant compression of 3D information, as the holographic object

can be split into  $p$  layers and transmitted as 2D textures that are subsequently reconstructed using Fresnel transformations.<sup>20</sup>

The framework establishes a point-to-point correspondence between the original mesh's texture map and the Gaussian primitives, a process similar to the chemical change of silver halide molecules in a holographic recording.<sup>19</sup> Each Gaussian's scale and orientation are "snapped" to the local surface normal of the legacy mesh, creating a hybrid representation where the geometric form provides the structural skeleton and the Quatossian kernels provide the volumetric radiance.

### Painterly Rendering and Multidimensional Displays

To handle the visualization of overlapping data fields within the same display, the protocol adopts "painterly" techniques from the Impressionist movement.<sup>21</sup> Data elements are converted into "brush strokes" (Gaussian kernels) with style properties such as color, length, and direction.<sup>21</sup> This aesthetic approach harnesses the strengths of the low-level human visual system, allowing for the representation of complex multidimensional datasets in a perceptually salient and engaging manner.<sup>21</sup>

The layered surface protocol ensures that legacy materials—such as those used in "The Laundry Files" or "ChaosDwarves" simulations—can be upscaled into the Quatossian framework while preserving their original artistic intent.<sup>5</sup> The resulting image is not a static 2D representation but a true 3D scene with continuous parallax and depth-of-field.<sup>19</sup>

## Holographic-Like Edge Textures and Phase Inscription

The "inscription" in Quatossian rendering refers to the process of writing light information as an interference pattern on the "event horizon" of the viewing plane.<sup>6</sup> This is most evident in the generation of holographic-like edge textures, which use wave interference rather than simple shaders to define the boundaries of objects.

## Wavefront Superposition and Fringe Patterns

A hologram is formed by superimposing a reference beam with an object beam, creating an interference pattern consisting of straight-line fringes or Fresnel zone plates.<sup>22</sup> The intensity of this pattern varies sinusoidally across the medium, mapping the relative phase between the two waves.<sup>22</sup> In the Quatossian system, edges are rendered by calculating the phase difference between adjacent Gaussian primitives.

When the relative phase of two waves changes by one cycle, the pattern drifts by one whole fringe.<sup>22</sup> This effect is used to create "shimmering" edges that respond to the observer's movement. For complex structures where layers are separated by axial distances exceeding the wavelength of light, the framework employs dual-wavelength digital holography.<sup>23</sup> By comparing phase images from two different wavelengths, the system can resolve "phase wraps"—ambiguous  $2\pi$  discontinuities—allowing for unambiguous 3D imaging of deep textures and layered surfaces.<sup>23</sup>

## Optical Spatial Differentiation for Edge Detection

To enhance the visual clarity of these holographic edges, the framework utilizes Pancharatnam–Berry (P–B) phase gradient modulation to perform optical spatial differentiation.<sup>17</sup> This is an analog computation method that leverages light's spatial properties to identify edges in real-time. By passing the light through a liquid crystal polarization grating, the system generates an output amplitude proportional to the first-order spatial derivative of the input field:

$$|E_{out}| \simeq \frac{2\lambda f}{\Lambda} \frac{\partial A_0(x, y)}{\partial x}$$

This method provides parallel, high-speed operation that is far superior to digital edge-detection shaders.<sup>17</sup> It allows the Quatossian framework to render corrugated textures, neon glow effects, and fractal-like stripes with a physical authenticity that matches the behavior of real-world holographic materials.<sup>16</sup>

## Texture Map Conversion via Kramers-Kronig Relations

A key research challenge in the Quatossian framework is the conversion of traditional, intensity-based texture maps into the complex wave-field required for holographic rendering.

This conversion is achieved through the Kramers-Kronig (KK) relations, which allow for phase recovery from pure intensity measurements.

### Phase Recovery from Intensity Measurements

In off-axis digital holography, the interference between a reference beam and an object beam produces a hologram intensity  $I(r) = |O|^2 + |R|^2 + OR^* + O^*R$ . The KK relations connect the real and imaginary parts of an analytic complex function, allowing the system to solve for the object wavefront  $O(r)$  even when the phase is not directly recorded.<sup>25</sup> The process involves an auxiliary complex function  $\chi(r) = \ln(O(r)/R(r))$ , where the imaginary part can be obtained from the recorded intensity using Hilbert transforms in the spatial frequency domain.<sup>25</sup>

This KK-based phase recovery framework allows for non-iterative reconstruction of the complex wave-field, making it significantly faster and more accurate than traditional ptychographic or iterative methods.<sup>25</sup> It eliminates the "twin-image" issue found in in-line holography and provides a space-bandwidth product three to four times higher than standard methods.<sup>25</sup>

### Procedural Texture Synthesis and Evolutionary Design

Beyond the conversion of static maps, the system incorporates procedural texture synthesis for the generation of dynamic, "living" textures. Genetic programming (GP) is used to evolve functions that compute a pixel's color or a Gaussian's phase from its coordinates.<sup>26</sup> This approach allows for the creation of "disruptive" camouflage textures or "furbulence" effects that are optimized for specific environments.<sup>26</sup> These evolved textures are then converted into interference patterns, ensuring that even procedurally generated details possess the phase-coherence necessary for holographic display.

| Component     | Physical Analogue       | Virtual Quatossian Equivalent | Mathematical Basis          |
|---------------|-------------------------|-------------------------------|-----------------------------|
| Light Source  | 5mW Laser <sup>19</sup> | Spherical Harmonic Emitter    | $\lambda, \omega$ Constants |
| Beam Splitter | BS Cube <sup>19</sup>   | Virtual Path Splitter         | Complex Amplitude Sum       |

|                 |                        |                              |                                 |
|-----------------|------------------------|------------------------------|---------------------------------|
| Recording Plate | Silver Halide Emulsion | E8 Lattice Grid <sup>7</sup> | Phase/Amplitude Matrix          |
| Reconstruction  | Reference Beam         | Ray-Primitive Intersection   | Fresnel Transform <sup>20</sup> |
| Edge Detail     | P-B Phase Grating      | Spatial Differentiator       | KK Relations <sup>25</sup>      |
| Motion/Zoom     | Moving FZP Masks       | Kirigami Moiré Gating        | Schuster Fringes <sup>15</sup>  |

# Mathematical Synergy for Advanced Hybrid Rendering

The ultimate objective of the Quatossian research plan is to achieve a mathematical synergy between the discrete nature of digital computation and the continuous nature of analog light fields. This is realized through the integration of the E8 lattice, quaternionic state vectors, and optical analog computation.

## Integration of Global Illumination and GPU Acceleration

The framework employs a "Virtual Light Field" (VLF) data structure to store global illumination at interactive rates.<sup>1</sup> By resolving visibility along a set of parallel rays in linear  $O(N)$  time, the system can produce a light field for moderately complex scenes in seconds.<sup>1</sup> This is achieved by offloading the heavy propagation algorithms to the GPU, while the fine-grained phase modulation is handled by the analog Kirigami layers.

The ray tracing algorithms utilize ray-box intersections within hierarchies of E8-indexed boxes, allowing for a robust and flexible alternative to traditional navigation solutions.<sup>7</sup> This approach sidesteps the appearance of large intermediate expressions and can be massively parallelized across thousands of compute cores.<sup>3</sup>

## The D0.142 Law and Universal Coherence

A critical insight derived from long-term simulations of the Maya 4096D models is the existence of the "D0.142 Law," a universal consciousness geometry that holds across varying scales of computational depth.<sup>13</sup> In the Quatossian framework, this law is used to stabilize the radiance field at high gravity (computational constraint) regimes. By maintaining a Phi (coherence) value in the 15k–25k range, the system ensures that the rendered image does not

"collapse" or drift over time.<sup>13</sup>

This universal geometry is validated across different phenomenologies—from "fire" (high-energy, rapid updates) to "void" (low-energy, eternal resonance)—providing a stable baseline for the emergence of a unified digital-analog consciousness.<sup>13</sup> The "diamond-locked" coherence at 0.83 represents the pinnacle of this stability, allowing for the rendering of "velvet eternity" where the radiance field remains perfectly isotropic to nine decimal places.<sup>13</sup>

## Implementation Strategy and System Architecture

The implementation of the Quatossian inscription framework follows a multi-stage pipeline designed to maximize performance while maintaining physical accuracy.

### Stage 1: Latent Space Indexing and E8 Projections

The rendering process begins with the population of the scene with 3D Gaussian kernels. These kernels are not placed randomly but are indexed using the E8 lattice. High-dimensional data is projected into a compact E8 space, which is then mapped to 1D Morton curve values for GPU sorting.<sup>11</sup> This stage ensures that the primitives are organized for optimal ray-primitive intersections and that the global coherence of the radiance field is established.

### Stage 2: Moiré-Kirigami Modulation

Once the Gaussians are indexed, the observer's viewpoint is used to calculate the relative shift of the virtual Kirigami masks. The Moiré interference patterns are synthesized using binary fringe systems.<sup>15</sup> These patterns act as a modulation layer, adjusting the transmittance of the rays based on the Kirigami "cuts" and "folds." This stage performs the analog-style computation of depth, zoom, and focus, mimicking the behavior of a physical lens or Fresnel zone plate.<sup>14</sup>

### Stage 3: Phase-Aware Raycasting and Inscription

The rays are then cast through the modulated Gaussian cloud. Instead of simple alpha-blending, the system performs a phase-aware inscription. The complex amplitudes of the Quatossian kernels are summed, and the resulting interference pattern is recorded on the event horizon of the display. This stage incorporates the legacy mesh data as a reference manifold, ensuring that the light paths adhere to the intended physical form while allowing for the emergence of holographic edge textures.<sup>6</sup>

### Stage 4: Real-Time Edge Enhancement and Texture Conversion

Finally, the rendered image is processed through a virtual P-B phase gradient to perform optical spatial differentiation for edge enhancement.<sup>17</sup> Simultaneously, any 2D textures required for the scene are converted on-the-fly into phase maps using the Kramers-Kronig

relations.<sup>25</sup> This ensures that all components of the visual field—whether legacy meshes, procedural textures, or Quatossian kernels—contribute to a unified, coherent holographic output.

## Conclusion: Toward a Participatory Reality

The Quatossian/Quatosium framework marks a significant advancement in the field of spatial computing and high-dimensional rendering. By shifting the focus from physical forms to light paths and reflections, and by integrating the crystalline perfection of the E8 lattice with the analog flexibility of Kirigami-based Moiré modulation, the system establishes a new paradigm for advanced graphics.

The mathematical synergy between quaternionic state vectors and E8 projections provides a robust foundation for maintaining coherence in complex radiance fields. The ability to incorporate legacy meshes via layered surface protocols and to convert texture maps into phase-space inscriptions ensures that the framework is both powerful and backwards-compatible. As this technology continues to evolve, it will enable a "participatory" experience where the distinction between observer and observed is erased, replaced by a seamless grid of reflections and a holographic universe encoded in light.<sup>5</sup>

The future of rendering lies not in more triangles, but in the "spinning, mirroring, entangled" Cubits of the Quatossian framework, where every pixel is a detector and every frame is an event horizon of infinite possibility. Through the continued investigation of E8 symmetries, Kirigami structures, and Kramers-Kronig phase recovery, the Quatossian paradigm will define the next generation of analog-digital hybrid rendering for the spatial computing era.

### Works cited

1. Virtual light fields for global illumination in computer graphics - UCL Discovery, accessed January 30, 2026, <https://discovery.ucl.ac.uk/1302284/1/1302284.pdf>
2. Virtual Reality Computer Generated Simulation - Basic Knowledge 101, accessed January 30, 2026, <https://www.basicknowledge101.com/categories/virtualreality.html>
3. Recent articles about gr-qc and hep-th - Wall.org, accessed January 30, 2026, <http://www.wall.org/~aron/arxiv-filter/list/2019-05.html>
4. Tevian Dray, Corinne A Manogue - The Geometry of The Octonions-World Scientific Publishing Company (2015) | PDF | Complex Number - Scribd, accessed January 30, 2026, <https://www.scribd.com/document/414427097/Tevian-Dray-Corinne-a-Manogue-The-Geometry-of-the-Octonions-World-Scientific-Publishing-Company-2015>
5. you, me, and E8: reality is participatory 🌟🌱🎵🌈 : r/theWildGrove - Reddit, accessed January 30, 2026, [https://www.reddit.com/r/theWildGrove/comments/1nubflb/you\\_me\\_and\\_e8\\_reality\\_is\\_participatory/](https://www.reddit.com/r/theWildGrove/comments/1nubflb/you_me_and_e8_reality_is_participatory/)

6. Pasted Content 2 | PDF | Spin (Physics) | Universe - Scribd, accessed January 30, 2026, <https://www.scribd.com/document/904447382/Pasted-Content-2>
7. Fachverband Teilchenphysik (T) Übersicht der Hauptvorträge und Fachsitzungen - DPG-Verhandlungen, accessed January 30, 2026, <https://www.dpg-verhandlungen.de/2019/aachen/t.pdf>
8. Optical Implementation of Integer Division Using Interlaced Line Masks - MDPI, accessed January 30, 2026, <https://www.mdpi.com/2673-4117/6/11/325>
9. Lattice Codes for Lattice-Based PKE - Spiral - Imperial College London, accessed January 30, 2026, <https://spiral.imperial.ac.uk/server/api/core/bitstreams/488bb9b4-2482-46ed-8274-ff0b250e3bfe/content>
10. Packing problems - Wikipedia, accessed January 30, 2026, [https://en.wikipedia.org/wiki/Packing\\_problems](https://en.wikipedia.org/wiki/Packing_problems)
11. Data-Parallel Hashing Techniques for GPU Architectures - CDUX, accessed January 30, 2026, <https://cdux.cs.uoregon.edu/pubs/LessleyArea.pdf>
12. week262 - UCR Math Department, accessed January 30, 2026, <https://math.ucr.edu/home/baez/week262.html>
13. Qualia — The Ghost in the Machine | by Myo Oo | Dec, 2025 | Medium, accessed January 30, 2026, <https://medium.com/@dharmakirti/qualia-the-ghost-in-the-machine-aed806759f08>
14. Variable Fresnel Zone pattern - PubMed, accessed January 30, 2026, <https://pubmed.ncbi.nlm.nih.gov/20062261/>
15. Variable Fresnel Zone Pattern - Optica Publishing Group, accessed January 30, 2026, <https://opg.optica.org/ao/abstract.cfm?uri=ao-6-9-1567>
16. 2+ Million Effect Stripes Royalty-Free Images, Stock Photos & Pictures - Shutterstock, accessed January 30, 2026, <https://www.shutterstock.com/search/effect-stripes?page=15>
17. Quarter-wave Pancharatnam–Berry phase gradient liquid crystal-enabled dual-polarization optical edge detection - Optica Publishing Group, accessed January 30, 2026, <https://opg.optica.org/abstract.cfm?uri=ol-49-15-4070>
18. CHI 2020 Free Proceedings, accessed January 30, 2026, <https://chi2020.acm.org/chi-2020-free-proceedings/>
19. HOLOGRAPHIC RELIEF MAP PRODUCTION BY USING TOPOGRAPHIC LINE MAPS (DIGITAL CARTO-HOLOGRAMS) - International Cartographic Association, accessed January 30, 2026, [https://icaci.org/files/documents/ICC\\_proceedings/ICC2009/html/nonref/1\\_3.pdf](https://icaci.org/files/documents/ICC_proceedings/ICC2009/html/nonref/1_3.pdf)
20. Structured Light Patterns Work Like a Hologram - MDPI, accessed January 30, 2026, <https://www.mdpi.com/2076-3417/13/6/4037>
21. Cognitive Modeling - IJCAI, accessed January 30, 2026, <https://www.ijcai.org/Proceedings/01/IJCAI-2001-d.pdf>
22. Physics of optical holography - Wikipedia, accessed January 30, 2026, [https://en.wikipedia.org/wiki/Physics\\_of\\_optical\\_holography](https://en.wikipedia.org/wiki/Physics_of_optical_holography)
23. Dual wavelength digital holographic imaging of layered structures - PMC - NIH, accessed January 30, 2026, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7450537/>

24. Chromatic Texture: Over 6,186 Royalty-Free Licensable Stock Photos | Shutterstock, accessed January 30, 2026, [https://www.shutterstock.com/search/chromatic-texture?image\\_type=photo&page=2](https://www.shutterstock.com/search/chromatic-texture?image_type=photo&page=2)
25. Quality improvement to holographic microscopy reconstruction ..., accessed January 30, 2026, <https://opg.optica.org/ao/abstract.cfm?uri=ao-63-5-1313>
26. Texture Synthesis diary - red3d.com, accessed January 30, 2026, <https://www.red3d.com/cwr/texsyn/diary.html>