

*A Further Investigation of the
Symmetrical Chloride of
Paranitroorthosulphobenzoic Acid.*

Dissertation.

*Submitted to the Board of University
Studies of the Johns Hopkins University
for the degree of Doctor of Philosophy.*

by William E. Hendersen.

1897

*A Further Investigation of the
Symmetrical Chloride of
Paranitroorthosulphobenzoic Acid.*

Dissertation.

*Submitted to the Board of University
Studies of the Johns Hopkins University
for the degree of Doctor of Philosophy.*

by William E. Hendersen.

1897

The Project Gutenberg eBook of A Further Investigation of the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

Title: A Further Investigation of the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid

Author: William Edwards Henderson

Release date: April 25, 2022 [eBook #67920]

Most recently updated: October 18, 2024

Language: English

Original publication: United States: William E. Henderson, 1897

Other information and formats: www.gutenberg.org/ebooks/67920

Credits: The Online Distributed Proofreading Team at <https://www.pgdp.net>
(This file was produced from images generously made available by The Internet Archive)

*** START OF THE PROJECT GUTENBERG EBOOK A FURTHER
INVESTIGATION OF THE SYMMETRICAL CHLORIDE OF
PARANITROORTHOSULPHOBENZOIC ACID ***

A Further Investigation of the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.

Dissertation.

Submitted to the Board of University
Studies of the Johns Hopkins University
for the Degree of Doctor of Philosophy.

— by —

William E. Henderson.

1897

Acknowledgment.

The author esteems it a privilege as well as a pleasure to give expression to his sincere sense of gratitude to Prof. Remsen, under whose guidance this work was carried on not only for instruction received in the lecture room, but for his frequent suggestion, and his constant and friendly interest in the work as it progressed. These have at all times been an encouragement and an incentive.

He wishes also to express his appreciation of the instruction and kindly guidance in the laboratory, of Drs. Morse and Renouf, as well as of Dr. Ames of the Physical Laboratory.

Contents.

I.	Introduction.	<u>Page 1</u>
II.	Preparation of the Acid Potassium Salt of Paranitroorthosulphobenzoic Acid.	<u>6</u>
III.	Preparation of the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.	<u>12</u>
IV.	Properties of the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.	<u>19</u>
V.	The Action of Benzene and Aluminium Chloride on the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.	<u>22</u>
	The Barium Salts of Paranitroorthobenzoylbenzenesulphonic Acid.	<u>24</u>
VI.	The Action of Alcohols on the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.	<u>30</u>
	1. Methyl Alcohol.	<u>31</u>
	2. Ethyl Alcohol.	<u>32</u>
	Action of Ethyl Alcohol on the Unsymmetrical Chloride.	<u>36</u>
VII.	The Action of Phenols on the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.	<u>38</u>
	1. Phenol.	<u>40</u>
	2. Orthocresol.	<u>48</u>
	3. Paracresol.	<u>51</u>
	4. Hydroquinone.	<u>53</u>
	5. Resorcin.	<u>56</u>
	6. Pyrogallol.	<u>59</u>

7. β -naphthol.	<u>61</u>
VIII. The Action of Aniline on the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.	<u>62</u>
IX. The Action of Phosphorus Oxychloride on the Fusible Anilid of Paranitroorthosulphobenzoic Acid.	<u>71</u>
X. The Action of Reagents on the Dianil of Paranitroorthosulphobenzoic Acid.	<u>77</u>
1. Of Hydrochloric Acid.	<u>77</u>
2. Of Alcoholic Potash.	<u>78</u>
3. Of Glacial Acetic Acid.	<u>79</u>
XI. Conclusions.	<u>82</u>
Biographical.	<u>85</u>

I. Introduction.

The sulphobenzoic acids have been the subject of investigation in this laboratory for a number of years past. Among the many interesting facts that have been brought to light in the course of this study, perhaps no others have been attended with more interest than the discovery of well characterized isomerism in the case of the chlorides of orthosulphobenzoic acid, and its paranitro derivative; together with the preparation of a series of isomeric derivatives of these substances. The chlorides themselves have been isolated in the crystalline condition, and have been found to differ markedly, not only in chemical, but in physical properties as well.


The first evidence that such isomerism existed, was obtained by Remsen and Coates^[1] who, in the course of an investigation of the action of aniline upon the chloride of orthosulphobenzoic acid, obtained two isomeric anilids quite different in properties, which they designated as fusible and infusible respectively. The following year, Remsen and Kohler^[2] obtained one of the chlorides in crystalline form, together with an oil which they did not succeed in crystallizing.

This however was accomplished the succeeding year by Remsen and Saunders^[3], and a still more satisfactory result was obtained by Remsen and McKee^[4] in 1895. The chloride melting at 79° was found to yield only the fusible anilid, together with an anil, while from the lower melting chloride, in addition to these, the infusible anilid was also formed.


In 1895, Gray^[5] isolated the two corresponding isomeric chlorides of paranitroorthosulphobenzoic acid, the lower melting chloride being obtained in small quantity only. The succeeding year Hollis^[6] made a more careful study of this lower melting chloride, and prepared it in considerable quantity.

From evidence drawn from the action of ammonia upon these chlorides, taken in connection with a number of other facts, the higher melting chloride is identified as the one possessing a symmetrical structure, while the lower melting chloride possesses an unsymmetrical structure. The first

one, when treated with ammonia is slowly transformed into the ammonium salt of paranitrobenzoic sulphinide:



while the lower melting chloride is quickly transformed into the ammonium salt of paranitroorthocyanbenzenesulphonic acid:



Gray's study of the symmetrical chloride was confined for the most part to the preparation of a series of salts of this latter acid, and to an investigation of the action of aniline upon the chloride itself. It was thought to be of interest to extend this study to a wider range of reactions, as well as to improve, if possible, the method of preparing the chloride in pure condition. At the suggestion of Prof. Remsen this work was accordingly undertaken.

II. Preparation of Material.

The method employed in the preparation of paranitroorthosulphobenzoic acid was essentially that described by Hart,^[7] Kastle,^[8] Gray^[9] and Hollis.^[10] The details of it are repeated here for the purpose of calling attention to certain facts that came under the author's notice.

100 grams of paranitrotoluene are added to 400 grams of fuming sulfuric acid, and the mixture heated in a balloon flask at 100° on a water bath. The heating is continued until a few drops of the mixture, added to cold water, dissolves completely to a clear solution. The time required for this operation varies much with the conditions. Continued stirring very considerably hastens the reaction, as paranitrotoluene forms a layer on the acid, which presents a small surface to its action. With constant stirring the reaction is complete in a few hours, whereas if no stirring is resorted to, as much as several days may be required, especially when large quantities are employed at one time.


When the reaction is complete, the mixture is poured into a large volume of water, and neutralized with calcium carbonate. In the filtrate from calcium sulphate, the calcium salt of paranitroorthotoluene sulphonic acid is found, and this is converted into the potassium salt in the usual way.

The oxidation of the potassium salt is effected as follows. 50 grams of the salt are dissolved in 2½ litres of water, and to this is added a solution of 15 grams of potassium hydroxide. The mixture is heated to 100° on a water-bath, and when this temperature is reached, 110 grams of potassium permanganate are added. Heating is continued until the solution is decolorized, care being taken to prevent the evolution of free oxygen.


The oxides of manganese are then filtered off, the filtrate neutralized with hydrochloric acid, and evaporated to about one fifth of its original volume. Strong hydrochloric acid is then added in excess, and on cooling the acid potassium salt of paranitroorthosulphobenzoic acid separates in very slender colorless needles completely filling the liquid.

For the success of this operation it is important that the potassium salt of paranitroorthotoluenesulphonic acid and the potassium hydroxide should both be perfectly dissolved before they are heated together. If the two substances lie together in solid form at the bottom of the flask, a very slight elevation of temperature leads to the formation of an extremely troublesome red substance, which is very difficult to remove. It is almost impossible to remove it from the oxidation product by recrystallization, since any considerable amount of it has a marked influence on the solubility of the salt, rendering it much more soluble. It persists throughout all subsequent transformations of paranitroorthosulphobenzoic acid, and should therefore be carefully avoided.

Otto Fischer^[11] has shown that in concentrated solution, potassium hydroxide acts on nitro derivatives of toluene, with the formation of various colored substances derived from stilbene. In the case of paranitroorthotoluenesulphonic acid, he describes the substance formed as possessing a cherry red color. The reactions involved in its formation are:



By oxidation this passes to a nitro compound of the composition



It was no doubt the formation of substances of this nature that occasioned the color observed in some of the oxidations.

The only effective method of separating this colored substance was found to be to pass to the neutral salt of paranitroorthosulphobenzoic acid, by making the solution slightly alkaline. The salt of this colored substance is also formed and the two can be separated by a few recrystallizations in a fairly satisfactory manner.

The yield in both of the transformations involved in the preparation of paranitroorthosulphobenzoic acid does not fall far short of the theoretical.

III. Preparation of the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.

This chloride was first separated from its unsymmetrical isomer by Gray^[12]. It was obtained by allowing a chloroform solution of the mixed chlorides to evaporate until the chloroform had almost entirely disappeared. In the thick liquid so obtained, crystals of the symmetrical chloride were formed. It was also obtained by applying the method devised by Bucher in connection with the corresponding chloride of orthosulphobenzoic acid—i.e. by the action of dilute ammonia on the mixed chlorides. Gray also found that the best conditions for securing a relatively large proportion of the symmetrical chloride were, the employment of as low a temperature as possible in the formation of the chlorides, and of as small an excess of phosphorus pentachloride as would suffice for the reaction.


After many experiments, under widely differing conditions, the following method of procedure, embodying the results of Gray's work, was adopted.

Dehydrated acid potassium salt of paranitroorthosulphobenzoic acid, and phosphorus pentachloride, in the ratio of 40: 55 grams, are brought together in a mortar and intimately mixed. The mixture is put into an evaporating dish, and placed on a sulphuric acid bath, previously heated to 150°. As soon as the action has been well started, the dish is removed, and the reaction allowed to proceed without further heating. When it is complete, and the contents of the dish has cooled down to the temperature of the room, the oily product is poured slowly into a salts bottle containing ice water, the bottle being frequently shaken during the process. The shaking is continued with renewed portions of water, as long as the wash water is cloudy. The water is then poured off, the brownish gummy chloride dissolved in chloroform, and the solution placed in a good-sized separating funnel. Ice water is then added, and the contents of the funnel treated with successive portions of ammonia (desk ammonia diluted one half). Shaking


is continued after each addition until the odor of ammonia has disappeared, and ice is added from time to time as may be required.

When it is found that the odor of ammonia persists after several minutes' shaking, the chloroform layer, which is usually filled with a solid substance that has separated during the process, is drawn off, filtered, and dried with calcium chloride.

By this process all of the unsymmetrical chloride is converted into the ammonium salt of paranitroorthocyanbenzenesulphonic acid, according to the equation:



while the symmetrical chloride remains for the most part unchanged, though some of it is converted into the ammonium salt of paranitrobenzoic sulphinide:



It was found that working in this way the symmetrical chloride could be prepared in pure condition, free from its isomer. The chloroform completely evaporates in a short time leaving fine crystals of the symmetrical chloride. In case the evaporation is slow and incomplete, it may be concluded that not all of the unsymmetrical chloride has been removed. The yield was uniformly about 40 per cent of the theoretical.

From the water used to wash the chlorides a considerable amount of the original salt can be recovered, as the reaction under the conditions employed, is never complete.

An examination was made of the substance mentioned as separating in the chloroform solution of the chlorides, during the treatment with ammonia, and it was found to possess the following properties. It is insoluble in benzene, chloroform, acetone, ether and ligroin; soluble in glacial acetic acid, from which it separates on cooling in colorless, crystalline condition; insoluble in the cold in water, alcohol and ammonia, but by boiling with these reagents, or by long standing in the cold, it is dissolved with decomposition. It was dissolved in hot water and the solution, which was acid in reaction, was neutralized with potassium carbonate. On adding an excess of hydrochloric acid to the solution, and allowing it to cool, characteristic crystals of acid potassium salt of paranitroorthosulphobenzoic acid separated. These properties identify the substance as the anhydride of this acid.

The formation of the corresponding anhydride of orthosulphobenzoic acid by the action of phosphorus pentachloride upon its acid potassium salt was observed by Sohoni^[13], who made use of the reaction to prepare this anhydride in quantity.

IV. Properties of the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.


As first obtained, the crystals of the symmetrical chloride resemble irregularly shaped pieces of amber, both in color, and in lustre. On recrystallization from chloroform or ether, they may be obtained perfectly colorless, and are often of very simple crystallographic form. The chloride crystallizes in the monoclinic system, and possesses a very remarkable crystallizing power, in which respect it differs noticeably from its isomer. Even in chloroform solution that is far from dry, crystals appear with the greatest ease.

The habit of the crystals differs very much according to the conditions of crystallization. Not infrequently almost perfectly formed crystals of the simplest form—the oblique octahedron—were obtained though for the most part the form was much more complicated, pinacoid and dome faces, together with basal planes being prominent. As a rule, the crystals were not suitable for crystallographic work, as the faces are usually uneven and the edges rounded. By proper precautions however, good ones were obtained, and measurements of these will be found in this dissertation when it appears in print.

The size of some of the crystals obtained was unusual for substances of this class. One crystal obtained with no special precautions, save letting a solution of the chloride stand undisturbed for several days, in a rather cool place measured $3 \times 2.5 \times 1.5$ cm., and weighed 11.2 grams. The crystals are quite compact, their density being about 1.85. They melt at 98° (uncorr.) The chloride is quite stable in crystalline condition. Even in moist air the crystals were unchanged, and retained their lustre as long as they were in my possession.

An analysis for chlorine gave the following results.

.2200 gram gave .2212 gram AgCl.



V. The Action of Benzene and Aluminium Chloride on the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.

Hollis^[14] in his study of the action of these reagents upon the unsymmetrical chloride, tested their action upon one portion of the symmetrical chloride, and found the products to be identical in the two cases. A few experiments were made in confirmation of these results, and the same products, in general, were obtained. It was observed however that the reactions differ in the relative ease with which they are brought about. In the case of the symmetrical chloride, the reaction is a much more vigorous one. On adding aluminium chloride to a solution of the symmetrical chloride, in benzene, action begins at once the temperature of the hand, and very little external heat, and that only in the latter stages of the operation, is needful for the completion of the reaction. The application of much heat converts all of the product into thick tarry substances from which nothing satisfactory could be obtained.

When the reaction was complete, the resulting product was isolated and purified in accordance with the directions given by Hollis. Repeated trials showed that, as in the case of the unsymmetrical chloride, only one phenyl group could be introduced by this method. The resulting compound, paranitroorthobenzoylbenzenesulphon chloride, was identical with that derived from the unsymmetrical chloride. Owing, however, to the fact that so much more decomposition occurs in the reaction with the symmetrical chloride, in paranitroorthobenzoylbenzene sulphon chloride could not be obtained in perfectly pure condition. In appearance it agreed closely with that described by Hollis, forming very characteristic greenish, rhombic crystals. These melted, not very sharply, at 174° instead of 177° as observed by Hollis.

Accordingly, to establish the identity of the two compounds beyond any doubt, the material on hand was converted into the barium salt of

paranitroorthobenzoylbenzene sulphonic acid. This was done by boiling the sulphon chloride with dilute hydrochloric acid until complete solution had been effected; evaporating to dryness on a water-bath; dissolving the residue in hot water, and neutralizing with barium carbonate. On filtering the hot solution from the excess of carbonate, and allowing it to cool, the barium salt separated.

The solution was somewhat colored by impurities, and the long needles in which the salt crystallized were also somewhat colored. They were analysed with the expectation that they would prove to be specimens of the salt described by Hollis as having three, or three and a half molecules of water of crystallization, in as much as the conditions under which they were formed were favorable to the formation of salts with these ratios of water of crystallization. Hollis found that this salt could be obtained with at least four different ratios of water of crystallization viz. three, three and a half, six and seven molecules respectively. The analysis was as follows, the amount of barium being calculated on the basis of the anhydrous salt.

0.3087 gram lost 0.064 gram at 210°,

and gave 0.0759 gram BaSO₄.

Cal. for	Found.
(C₁₃H₈O₆NS)₂Ba + 11H₂O	
H ₂ O = 20.90	20.73
Ba = 18.29	18.23

The mother-liquor, in which the crystals remaining from analysis were redissolved, was warmed, but not boiled, with boneblack, to remove impurities. When filtered, the solution was perfectly colorless, and on standing for some time, well formed colorless, rhombic crystals appeared. On analysis they gave results as follows.

0.2804 gram lost 0.0405 gram at 210°,

and gave 0.0759 gram BaSO₄.

Cal. for	Found.
(C₁₃H₈O₆NS)₂Ba + 7H₂O.	

H ₂ O = 14.40	14.44.
Ba = 18.29	18.03.

In making a further supply of the salt it was found that if the solution, after filtering from the barium carbonate, was diluted to such an extent that no crystals separated on cooling, then on slow evaporation under a bell-jar the first crystals to appear were very long slender needles. As evaporation proceeded, these needles became much thicker assuming prismatic proportions, and corresponded in appearance to the salt described by Hollis as having six molecules of crystal water.

As growth proceeded, the crystals became dark in color, and the mother-liquor correspondingly clearer, the crystals evidently absorbing the impurity in their growth.

When the solution had become quite colorless, rhombic crystals of the salt containing seven molecules of water of crystallization appeared. The larger prismatic crystals were carefully removed, and redissolved in water in order to see if the same phenomena would repeat themselves. This in fact was the case, crystals of both types appearing in the same way as described. Without separating the crystals in this second experiment, water was added, and the crystals dissolved. The solution was then warmed briskly with boneblack, and filtered. From the filtrate, which was colorless, nothing but rhombic crystals having seven molecules of water of crystallization could be obtained, although a great many variations in the conditions were tried. Analysis of these last crystals was as follows:

0.2400 gram lost 0.035 gram at 210°,
and gave 0.0637 gram BaSO₄.




Cal. for	Found.
(C₁₃H₈O₆NS)₂Ba + 7H₂O.	
H ₂ O = 14.40	14.58
Ba = 18.29	18.27

Hollis states that treatment with boneblack decomposes this salt, and hence he did not purify it prior to crystallization. From the experiments just described it seems probable that the impurities present affect the crystalline

habit, and the degree of hydration of this salt in a very striking manner. By careful warming with boneblack no decomposition was observed, and the crystals so obtained have constantly seven molecules of crystal water.

VI. The Action of Alcohols upon the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.

Kastle^[15] found that when the chlorides of paranitroorthosulphobenzoic acid (which he supposed to be an individual) were dissolved in alcohol, and the solution boiled for some time, the acid ethereal salt of paranitroorthosulphobenzoic acid was the final product. The reactions were shown to be:



Kastle, it will be observed, gave the symmetrical formula to this mixture of chlorides. Several acid ethereal salts were made, and a series of the neutral salts of various metals described by him.


The action of pure symmetrical chloride was studied in the same general manner to see if the resulting products would be the same as those formed from the mixed chlorides.

1. Action of Methyl Alcohol upon the Symmetrical chloride.

A portion of the chloride was dissolved in methyl alcohol, and the solution boiled until a drop added to cold water gave no precipitate, of unchanged chloride. The alcohol was then distilled off, and the thick syrup remaining, diluted with water. This solution was neutralized with barium carbonate and filtered. On cooling, the barium salt crystallized in shining mica-like plates, or in yellowish needles corresponding accurately with those described by Kastle. They gave the following analytical results.

0.2664 gram lost 0.0211 gram at 150°,

and gave 0.0870 gram BaSO₄.



H ₂ O = 7.79	7.88
[anhydrous salt] Ba = 20.85	20.85

2. In like manner the barium ethyl salt was made. It also agreed perfectly with Kastle's description, crystallizing in fine, colorless needles, forming in tufts from a not too concentrated solution. In case it is necessary to concentrate these solutions, it is of advantage to add a small quantity of alcohol to the solution as this prevents any great amount of saponification, which otherwise takes place to a noticeable extent.

Analysis.

I. 0.2824 gram lost 0.0276 gram at 180°,
and gave 0.0860 gram BaSO₄.

II. 0.2655 gram lost 0.0262 gram at 190°,
and gave 0.0815 gram BaSO₄.

$\begin{array}{c} \text{COOC}_2\text{H}_5 \\ / \\ \text{Cal. for } \left[\text{C}_6\text{H}_3\text{—SO}_2\text{O} \right] \text{Ba} + 4\text{H}_2\text{O}. \\ \backslash \\ \text{NO}_2 \end{array} \quad \quad \quad \mathbf{2} \quad \quad \quad \mathbf{Found.}$			
		I	II
H ₂ O =	9.51	9.77	9.86
Ba =	20.00	19.84	20.02

Kastle also found that by dissolving the mixed chlorides in alcohol in the cold, and allowing the solution to evaporate, there separated after a time, crystals of the chloride of the acid etherial salt of paranitroorthosulphobenzoic acid whose formation and composition are represented in equation I.

This same product was sought for when pure symmetrical chloride was employed, but without success. In every case, crystals of unchanged chloride separated, or else it was found that it had been completely converted into the acid etherial salt. In another trial cold water was carefully added in small portions, since Kastle found that such treatment facilitated the separation of the substance; the chloride alone appeared. Still other attempts were made to obtain the substance by adding a large amount of water to the solution of the chloride in alcohol, after it had stood for some time. In this way, quite a precipitate was thrown down, and this was filtered off and crystallized from ether. It always proved to be the symmetrical chloride, and none of the other substance was obtained.

Karslake^[16] in working with the symmetrical chloride of orthosulphobenzoic acid, was unable to isolate the analogous compound, although from the mixed chlorides, by the action of alcohols, Remsen and Dohme^[17] had obtained chloro-etherial salts.

In as much as the pure symmetrical chloride is relatively stable in cold alcohol (it can be crystallized from warm alcohol with very little loss), it is possible that it is more stable than the chloro etherial salt, and that in consequence the latter, when formed, yields more readily to the further action of alcohol than does the unacted on chloride. Hence when the action begins, it at once proceeds to the limit. The fact that the symmetrical chloride is rather sparingly soluble in cold alcohol, making the use of concentrated solutions impossible, may also be a factor in the case. Whatever may be the cause, this substance could not be obtained under any conditions that were devised.

Having in my possession a very small specimen of crystallized unsymmetrical chloride, it was submitted to the action of ethyl alcohol, under as nearly as possible the conditions employed by Kastle. Crystals of a colorless substance were obtained, which in every respect agreed with Kastle's description of the chloride of the acid ethyl etherial salt of paranitroorthosulphobenzoic acid. Crystallized from ether they melted at 68°.

The conditions employed by Kastle in preparing the chloride would undoubtedly lead to a relatively large proportion of unsymmetrical chloride, and it is to this chloride that the formation of the chloro etherial salt is apparently due.

VII. The Action of Phenols upon the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.

Remsen and Saunders^[18] in their investigation of the chlorides of orthosulphobenzoic acid, studied the action of phenol upon these substances, and from both the symmetrical chloride and the mixed chlorides, they obtained a normal diphenyl ether together with a red substance which was not further studied. It was formed in small quantity and was probably the corresponding sulphonphthalein. Later McKee^[19] obtained these same substances from both the symmetrical and the unsymmetrical chlorides. R. Meyer^[20] obtained analogous substances by the action of various phenols upon phthalyl chloride. It seemed probable, therefore, that the chlorides of paranitroorthosulphobenzoic acid would yield similar derivatives, and a study was accordingly made of the reaction of the symmetrical chloride with a series of phenols. The products in some instances were exceedingly difficult to deal with, possessing properties that made it impossible to prepare them for analysis, but even in such cases there could be little doubt as to the general nature of the reactions which had occurred.

1. The Action of Phenol upon the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.


A portion of the symmetrical chloride was brought together with somewhat more than double the molecular amount of phenol. The mixture was placed in a good-sized test-tube and the temperature gradually raised by means of a sulphuric acid bath.

As soon as the phenol melts, some slight action occurs, as is indicated by the fact that the mixture assumes a bright red color. No appreciable amount of hydrochloric acid gas is evolved however, until the liquid mixture has reached a temperature of about 115°. At this point the gas is freely evolved, and the action is complete at a temperature of 125°. The temperature observations were made by means of a thermometer used as a stirring rod in the mixture. During the heating, the color of the liquid becomes a much more intense red, growing darker in shade, and the liquid itself becomes somewhat viscous but does not solidify while hot.

When cool, the melt was repeatedly extracted with boiling water, the aqueous solution being very deep purple in color. The colored matter was removed very slowly in this manner, and so the process was continued with dilute alkali. A solid

insoluble residue was thus obtained, of a light-brownish color. This was dissolved in alcohol, boiled with boneblack and filtered. On cooling, needles of a straw yellow color were deposited from the alcoholic solution.

This proved to be the normal diphenyl etherial salt of paranitroorthosulphobenzoic acid, the formation of this substance being expressed by the equation:



Analysis of the substance gave the following results:

- I. 0.1627 gram gave 0.3398 gram CO₂ and 0.0510 gram H₂O.
- II. 0.1999 gram gave 0.4180 gram CO₂ and 0.0600 gram H₂O.
- III. 0.2649 gram gave 0.1561 gram BaSO₄.

	$ \begin{array}{c} \text{COOC}_6\text{H}_5 \\ / \\ \text{C}_6\text{H}_3\text{—SO}_2\text{OC}_6\text{H}_5 \\ \backslash \\ \text{NO}_2 \end{array} $			
Cal. for			Found.	
		I	II	III
C = 57.14		56.97	57.03	—
H = 3.26		3.47	3.33	—
S = 8.02		—	—	8.09

This substance melts at 119° (uncorr).

It possesses properties similar to those of the diphenyl etherial salt of orthosulphobenzoic acid described by Saunders. It is insoluble in water, and is unaffected by hydrochloric acid or aqueous alkali. On heating for a short time with alcoholic potash, the needles were transformed into a voluminous precipitate. This


was filtered off, dissolved in water, and hydrochloric acid was added. On cooling, characteristic crystals of the acid potassium salt of paranitroorthosulphobenzoic separated.

Analysis.

0.1392 gram lost 0.009 gram at 150° and gave 0.0385 gram K₂SO₄.

$\begin{array}{c} \text{COOH} \\ / \\ \text{Cal. for } \text{C}_6\text{H}_3\text{—SO}_2\text{OK} + \text{H}_2\text{O} \\ \backslash \\ \text{NO}_2 \end{array}$	
H ₂ O = 5.95	Found. H ₂ O = 6.51
K = 13.65	K = 13.35

No attempt was made to isolate the corresponding intermediate chlor-etherial salt of the composition





or its acid as was done by McKee^[21] in his work on the analogous etherial salt of orthosulphobenzoic acid.

On evaporating the aqueous extract from the original melt almost to dryness on the water-bath, there was a deposit on the sides of the dish of scales possessing a beautiful bronze-green metallic lustre. They formed a deep purple solution in alkalis, or magenta, if the solution was very dilute, and orange-yellow in acids. On acidifying the alkaline extract with hydrochloric acid, this same substance was precipitated as a brownish flocculent precipitate. It was, however, found to be impossible to obtain this substance in pure condition. The amount formed in the reaction is small, and its properties were such as to render work with it very difficult. The method of precipitation is not satisfactory because, owing to the fact that the substance is soluble in acid solutions to an unusual extent for substances of this class, the solution had to be concentrated to such a degree as to render the precipitated substance very impure from acids and alkali salts. These could not be removed by washing, obviously, without again dissolving the substance. From its properties however, and its color reactions, there can be little doubt that the


substance is a sulphonphthaleïn, and that it is always formed in considerable quantities in the reaction of phenol upon the symmetrical chloride.

It was noticed that the aqueous extract of the mass left after fusion was almost always decidedly acid in reaction, and it was thought that this might be due to the formation of an acid etherial salt, whose formation would be expressed by the equations:



Accordingly, the solution was saturated with barium carbonate, the excess of carbonate removed by filtration, the filtrate concentrated, and allowed to cool. Crystals in the form of pearly scales separated, which upon analysis proved to be the neutral barium salt of paranitroorthosulphobenzoic acid.

0.2291 gram anhydrous salt gave 0.1386 gram BaSO₄.



This would seem to indicate that the reaction is an incomplete one even in the presence of excess of phenol. No indications of the formation of an acid etherial

salt was observed.

2. The Action of Orthocresol upon the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.

With orthocresol the reaction proceeds with more difficulty. A higher temperature was required (135° - 145°), and quite an amount of tarry material was obtained from which very little could be extracted. The product was warmed repeatedly with dilute alkali, the solution so obtained neutralized with hydrochloric acid, and distilled with steam for several hours to free it from cresol. The resulting solution was then evaporated to small volume, and acidified with hydrochloric acid. A considerable precipitate was thrown down, which was easily filtered off and dried. In this condition it is a dark purple-red powder, lumps of which possessed a yellowish-bronze metallic lustre. In dilute alkaline solution it forms a deep-bluish purple solution, while in acids it is crimson, or light yellow if the solution is dilute. It is an excellent indicator, especially with ammonia.

In the insoluble tarry substance the etherial salt was sought for and obtained in small quantity only. As this substance is soluble in alcohol, and separates again on cooling in much the same condition, the etherial salt could not be isolated by crystallization from this solvent. By boiling the substance with benzene, purifying the filtrate with boneblack, and allowing the benzene to evaporate, an almost colorless gummy substance was obtained, which when dissolved in alcohol, crystallizes in small colorless needles which melt at 89° - 90° . They were not obtained in quantity sufficient for analysis, but there was little doubt that they were crystals of the diorthocresol etherial salt.

Apparently much more decomposition occurred in this reaction than when paracresol was employed, probably in consequence of the higher temperature required for the reaction.

3. The Action of Paracresol upon the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.

This reaction was conducted in the same manner as with phenol. No hydrochloric acid was evolved until a temperature of about 110° was reached, although after melting, the solution had steadily darkened to a deep reddish-brown color. At 130° , after heating for several hours, hydrochloric acid ceased to be evolved. The product was treated as in the last experiment. The alkaline extract did not exhibit any marked color reactions, such as were observed in most of these experiments, being dull reddish-brown in both acid and alkaline solution.

The insoluble residue crystallized from alcohol in light brown transparent crystals, which did not lose their color by repeated crystallization, and boiling with

boneblack, and melted sharply at 117°. From benzene they crystallized in colorless needles or flat, narrow plates. These become opaque on exposure to the air, apparently through loss of benzene of crystallization.

Analysis of the needles from alcohol gave the following results:

I. 0.2372 gram of substance gave 0.5137 gram CO₂ and 0.0965 gram H₂O.

II. 0.2223 gram gave 0.1203 gram BaSO₄.

	$\text{COOC}_6\text{H}_4\cdot\text{CH}_3$		
	/		
Cal. for	$\text{C}_6\text{H}_3\text{—SO}_2\text{OC}_6\text{H}_4\cdot\text{CH}_3$		
	\		
	NO_2		
		Found.	
		I	II
C =	59.08	59.06	
H =	3.98	4.52	
S =	7.49		7.43


4. The Action of Hydroquinone upon the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.

Action with hydroquinone occurs at 120°-135°, the mixture at the same time becoming dark colored and viscous.

On cooling, the product was powdered and treated with dilute alkali. It readily dissolved, without residue, forming a dark red solution. In concentrated solution the addition of acid produces a voluminous precipitate, dark brown in color, which when washed, and dried in paper forms an almost black powder. A dilute solution of this powder is dark red when alkaline, orange-yellow when acid.


From the way in which this powder was obtained, and owing to the fact that its solubility prevented repeated washing, it was evident that it would not give close analytical results for a calculated formula. It was thought, however, that analysis would give a general idea of the composition.

Analysis of different specimens gave results for sulphur which averaged about 5.5%. The percentage required for the formula



which represents the simplest sulphonfluoresceïn, is 7.43.

The compound could hardly have been so far from pure as to occasion such a discrepancy in results as this. It would appear, therefore, that more than two molecules of hydroquinone enter into the reaction with one molecule of the chloride. Should four molecules be involved in the reaction, leading to a compound of some such formula as



the theoretical percentage of sulphur would be 6.00 which corresponds much more closely with the results obtained.


This is in accord with the observations of a number of workers in this laboratory—Lyman, Gilpin, Linn and others—who have worked on various sulphonfluoresceïns, and have found that in many cases four, six and even eight phenol residues condense with one molecule of the anhydrous acid. Lyman^[22] especially describes a tetra hydroquinone sulphonfluoresceïn derived from orthosulphoparatoluic acid. No etherial salt was observed.

5. The Action of Resorcin upon the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.

The reaction of resorcin with the chloride is a much cleaner one and proceeds more easily than in the case just described, leading apparently to an individual compound which is well characterized.


During the reaction, which is complete at 125°, the mixture becomes almost perfectly solid, and when cool, it is quite brittle. It was reduced to a reddish powder in a mortar and dissolved in sodium hydroxide, there being no insoluble residue. By the addition of hydrochloric acid, the sulphonfluorescein was thrown down as a chocolate-brown precipitate, which was filtered off, washed to neutral reaction on a filter, and dried on paper. In this condition it is a light chocolate-brown powder. In dilute alkaline solution it possesses a slight fluorescence being pink by transmitted and yellow by reflected light, suggesting eosin in a general way. It is interesting to note that the sulphonfluorescein of orthosulphobenzoic acid possesses a fluorescence that can hardly be distinguished from ordinary fluorescein and that the introduction of a nitro group into the acid residue produces much of the same effect as do the four bromine atoms in eosin. In acid solution the color is reddish-orange.


Analysis of the compound, prepared as above described, gave the following results.



Analysis of the substance gave the following results:

- I. 0.1745 gram gave 0.3339 gram of CO₂ and 0.059 gram H₂O.
- II. 0.1467 gram gave 0.2820 gram CO₂ and 0.0432 gram H₂O.
- III. 0.1732 gram gave 0.3345 gram CO₂ and 0.0571 gram H₂O.
- IV. 0.2000 gram gave 0.1104 gram BaSO₄.
- V. 0.1505 gram gave 0.0820 gram BaSO₄.





	I	II	III	IV	V
C = 52.66	52.18	52.42	52.67	—	—
H = 3.46	3.76	3.27	3.66	—	—
S = 7.39	—	—	—	7.57	7.48

An effort to obtain the anhydride was unsuccessful. Some loss of weight was observed, but the compound underwent decomposition before this loss amounted to much.

6. The Action of Pyrogallol upon the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.

The product of this action dissolves readily in dilute sodium hydroxide without residue, producing a very deep purple-black color when concentrated, passing to grayish-violet as the solution is diluted. On adding hydrochloric acid, precipitation occurs, as in most of these reactions. On attempting to filter off this precipitate, it forms a sticky, black mass on the filter with which little can be done. It is best to evaporate to dryness before filtration and powder the residue. This powder can then be washed fairly clean from alkali salts and acid.

Nothing to suggest the formation of an etherial salt was observed.

Analysis of this product for sulphur showed that in this gallein, as in the case of the hydroquinone phthalein more than two pyrogallol residues had entered the acid residue. The indications were that six had entered into one of the chloride. This also agrees with the observation of Lyman^[23], who describes a hexapyrogallol gallein of orthosulphoparatoluic acid.

Probably a mixture of varying composition was obtained, and little importance was attached to the results save as they showed that no etherial salt is formed in the reaction.

7. The Action of β -Naphthol upon the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.

It was hoped that here, as in the case of the monohydroxy phenols an etherial salt would be obtained. It was found, however, that very little action occurred, save such as was indicated by the development of a bright carmine color in the melted mixture, until a temperature of about 160° was reached. At this point hydrochloric acid was evolved, but the chloride itself undergoes decomposition. Nothing definite could be isolated among the reaction products, save unchanged β -Naphthol.

VIII. The Action of Aniline upon the Symmetrical Chloride of Paranitroorthosulphobenzoic Acid.

As has been pointed out in the Introduction, it was in connection with the aniline derivatives of orthosulphobenzoic acid, that the isomerism of the chlorides was first noticed, two anilids being obtained. Accordingly, when Gray began his study of the chlorides of paranitroorthosulphobenzoic acid, his first effort was to obtain evidence of the existence of two anilids. These were not obtained, however, until after the chlorides themselves had been isolated, as their properties made their isolation and preparation a matter of difficulty.



Some points still remained in doubt after Gray's study, and a further investigation was thought to be desirable to clear these up.

Some time was spent in an endeavor to obtain a method by which a good yield of fusible, or symmetrical, anilid could be obtained. The yield in all cases tried, is not a good one. The presence of the nitro group appears to complicate the reaction, leading to secondary reactions whose course could not be followed. Upon bringing aniline and the chloride together, a very vivid red color was always observed, and the same was true when it was necessary to employ alkali. The fact that such colors develop when nitro compounds are treated with alkali has been noticed in many instances and some progress has been made in the study of these compounds. Jackson and Ittner^[24] have lately reviewed this subject.

If a solution of the symmetrical chloride in ether is slowly added to a similar solution of aniline, no appreciable amount of heat is evolved. If the resulting solution is allowed to stand at ordinary temperatures, action proceeds very slowly, aniline hydrochloride being precipitated as the reaction proceeds. This can be filtered off from time to time and the rate of action so observed. In such a way it was found that five grams of chloride required about fifty hours time to react completely with an excess of aniline. Similar results were obtained with chloroform as the solvent. By boiling the solution for an hour or more the reaction is complete.


The method employed was to bring the chloride and an excess of aniline—somewhat more than four molecules—together in chloroform solution. The flask was then boiled for about an hour, when the chloroform was distilled off. During the boiling as well as the distillation more or less bumping occurs in consequence of the aniline hydrochloride which separates, and constant shaking of the flask is sometimes necessary. The residue which is in a thick, gummy condition in consequence of the presence of an excess of aniline, was digested with water acidulated with hydrochloric acid. The excess of aniline is thus removed, and the reaction product obtained as a reddish-brown solid substance. This was treated with dilute sodium hydroxide, all lumps being broken up with a stirring rod. The undissolved substance is largely anil, which was filtered off. The anilid was then regained by acidifying the alkaline solution, in which it was dissolved. It separates immediately as a curdy colorless precipitate, though it is frequently colored pink by impurity. It was found that this color could be removed, in case not much was present, by redissolving the anilid in alkali, and slowly pouring the solution into an excess of dilute acid.

In all cases a considerable amount of anil was obtained, even when the substances were employed in the molecular ratios of 1:10. The reactions involved, so far as the formation of anilid and anil are concerned are,



On the whole the reaction seemed to be the most satisfactory in chloroform solution, the main objection being, that, owing to the simultaneous presence of chloroform, alkali, and a trace of aniline, phenyl isocyanide is always formed, and renders the work more or less unpleasant.

A number of experiments were also made to see if the yield could be increased by employing a modification of the “Schotten-Baumann Reaction”^[25] for the formation of anilids. For this purpose an ethereal solution of the chloride was added to a like solution of aniline in which was suspended finely powdered anhydrous potassium carbonate. The proportions of the substances were those demanded by the equation




Very little anilid was, however obtained, but in its place a substance soluble in water, of acid reaction capable of forming salts and yielding several well characterized derivatives. I hope to investigate this reaction more fully at some future time.

The anilid is rather sparingly soluble in alcohol, from which it is deposited on cooling in very small needles. These melt, as stated by Gray, at 222°. It is also soluble in chloroform and glacial

acetic acid, but does not form well defined crystals from any solvent. It dissolves in dilute alkali from which solution acids precipitate it unchanged.

The anil is also soluble in alcohol, glacial acetic acid etc. It crystallizes in much better-formed crystals than does the anilid. These melt at 188°.


On boiling the anil with aniline for a time, it is converted into the anilid



In none of these reactions was any infusible anilid observed.

IX. The Action of Phosphorus Oxychloride upon the Fusible Anilid.

Hunter^[26] found that when either of the anilids of orthosulphobenzoic acid were treated with phosphorus oxychloride, or similar dehydrating agents, a molecule of water was abstracted with the formation of a new substance. A careful study of the compound led to the belief that it was a dianil, and that its formation and structure could be represented by the equation



A corresponding study of the fusible anilid of paranitroorthosulphobenzoic acid was undertaken.

The method employed in this study was as follows. A tubulated retort of convenient size was fused onto the inner tube of a small condenser. This was done to avoid connections, which are nearly always attacked by the oxychloride. Another satisfactory plan is to have the neck of the retort of the same size as the inner tube of the condenser. The ends are placed in contact, and the tubes bound in position by wrapping with asbestos paper. Over the joint so made, a tight rubber tube is drawn.

A convenient amount of phosphorus oxychloride (50 c.c.) was placed in the retort and the anilid (5 gr.) added through the tubulus. On boiling, with the condenser inverted, the anilid soon dissolved, with evolution of hydrochloric acid gas, and the solution became bright yellow in color,

sometimes inclining to orange. The boiling was continued as long as hydrochloric acid was given off. The oxychloride was then distilled off under diminished pressure, care being taken to shake the retort constantly during the distillation as violent bumping is almost sure to occur especially towards the end of the operation. The product remaining, spattered over the walls of the retort, was a greenish yellow solid.

Water was then added, and the whole allowed to stand for an hour or so to thoroughly dissolve the phosphoric acid formed in the reaction.


In case the anilid is not perfectly dry, a much more energetic reaction occurs, and on distilling off the oxychloride, the product remains as a dark, gummy mass. This should be spread out on the sides of the retort while still liquid. On cooling and adding water, this gum gradually disappears, and in its place is found the yellow solid product just described. The gum appears to be a solution of this substance in phosphoric acid.

After the substance is filtered off and dried, it can be crystallized from acetone, benzene, glacial acetic acid or alcohol. From these solvents it crystallizes in small yellow needles resembling quinone in appearance.

The crystals obtained from acetone are rather larger than those from the other solvents, and are more nearly orange in color, apparently because of their greater compactness. When glacial acetic acid is used, care must be taken to avoid any unnecessary heating, as continued heating produces a change that will presently be described. The substance melts at 208° .

Analysis of the substance resulted as follows:

- I. 0.3822 gram gave 0.8334 gram CO_2 and 0.1272 gram H_2O .
- II. 0.2645 gram gave 0.5812 gram CO_2 and 0.0910 gram H_2O .
- III. 0.2023 gram gave 0.1283 gram BaSO_4 .
- IV. 0.2061 gram gave 0.1280 gram BaSO_4 .
- V. 0.1853 gram gave 16.73 C.C.N (Standard).



Found.


	I	II	III	IV	V
C = 60.11	59.47	59.93	—	—	—
H = 3.44	3.69	3.82	—	—	—
S = 8.45	—	—	8.70	8.52	—
N = 11.08	—	—	—	—	11.35

For analyses I & II I am indebted to Mr. Nakaseko, who kindly made them for me.

X. The Action of Reagents upon the Dianil of Paranitroorthosulphobenzoic Acid.

1. The Action of Hydrochloric Acid on the Dianil

When the dianil is boiled for some time with concentrated hydrochloric acid, the yellow color of the substance disappears, and the dianil is converted into a colorless substance without, however, passing into solution. The substance so obtained was filtered off, and crystallized from alcohol. It crystallized in small colorless needles, which melted at 183°, and possessed all the properties of the anil, which, in fact, it proved to be. The reaction was therefore



This reaction also explains the fact that some anil was always obtained in making the dianil from the anilid. Hydrochloric acid is formed in the reaction, and in turn acts on the dianil in the sense of the equation just given.

2. The Action of Alcoholic Potash on the Dianil.

On boiling the dianil with alcoholic potash for a time, the solution turned red, and nothing but tarry products were obtained. In this respect the dianil differs from the dianil of orthosulphobenzoic acid, which under similar conditions, is transformed into infusible anilid. This observation is, however, in keeping with the fact that the nitro derivative, is in general much less stable in the presence of alkali.

3. The Action of Glacial Acetic Acid on the Dianil.

When the dianil is boiled with glacial acetic acid for some time, the color of the solution changes to a much lighter shade of yellow, or becomes colorless. On evaporating the solution to small volume, and allowing it to cool, a colorless substance separates. This is infusible anilid. It could not be obtained in crystals from any solvent, but always separated in flakes. It does not melt or undergo change at 340°.


Like the fusible anilid it dissolves in dilute alkali, but on acidifying the solution it does not immediately reappear. After standing for some time, however, it gradually separates in perfectly pure form. In this particular my observation differs from that of Gray,^[27] who states that this anilid is decomposed by solution in alkali.

A specimen that had been repeatedly precipitated gave the following results on analysis.

- I. 0.1607 gram gave 13.88 C.C.N. (standard).
- II. 0.2195 gram gave 0.1285 gram BaSO₄.
- III. 0.1357 gram gave 0.0807 gram BaSO₄.

	$ \begin{array}{c} \text{C[NH.C}_6\text{H}_5\text{]}_2 \\ / \quad \backslash \\ / \quad \text{O} \\ / \quad / \\ \text{Cal. for C}_6\text{H}_3\text{—SO}_2 \\ \backslash \\ \text{NO}_2 \end{array} $		
	Found.		
	I.	II.	III.
N = 10.58	10.85	——	——
S = 8.06	——	8.00	8.16

By this series of transformations it is possible to pass from one anilid to the other, the steps being:



This is of special interest as affording a means of passing from a derivative of one of the chlorides, to a substance derived from the other, by steps that can be clearly followed.

Conclusions.

In the course of this investigation several facts have been established.

1. By the methods described, the symmetrical chloride of paranitroorthosulphobenzoic acid can be obtained in fine crystalline form, perfectly free from its isomer, with an average yield of forty percent.

2. By treatment of the chloride with benzene and aluminium chloride, only one chlorine atom can be replaced by a phenyl group.

3. The barium salt of paranitroorthobenzoyl benzenesulphonic acid, when perfectly pure, crystallizes constantly with seven molecules of water of crystallization.

4. With alcohols, the symmetrical chloride yields directly the acid etherial salt of paranitroorthosulphobenzoic acid, no evidence having been obtained of an intermediate chloro-etherial salt. The unsymmetrical chloride on the other hand yields the intermediate product.

5. With phenols, two series of derivatives are obtained.

(1) With monohydroxy phenols, both etherial salts and sulphonphthaleïns are formed, the former predominating.

(2) With polyhydroxy phenols no etherial salts were obtained, but compounds of the unsymmetrical type, usually containing more than two phenol residues.

6. With aniline an anil and an anilid of symmetrical constitution are formed.

7. With phosphorus oxychloride, the anilid, by loss of water, forms a dianil.

8. This dianil undergoes transformation with

- (1) Glacial acetic acid, forming an anilid of unsymmetrical constitution.
- (2) Hydrochloric acid forming the anil.
- (3) Alcoholic potash, with the formation of colored decomposition products.

Biographical.

The author of the foregoing dissertation was born at Wilkinsburg, Pa., Jan. 29., 1870. Owing to prolonged sickness in childhood his education, prior to entering college, was much interrupted, and was largely confined to instruction received at home.

In the fall of 1887 he entered Wooster University (Ohio), from which institution he received the degree of Bachelor of Arts in 1891. The two following years were spent as a teacher of Sciences in the College of Emporia (Kansas). In 1893 he entered the Johns Hopkins University where he has since been a student of chemistry, with physics and mathematics as subordinate studies.

In 1895 he was appointed University Scholar in Chemistry. During 1895-6 he served as lecture assistant to Prof. Remsen and Dr. Renouf in the undergraduate courses. In the spring of 1896 he was appointed Fellow for the present year.

Footnotes:

- [1] Am. Chem. Journ. XVII, 311.
- [2] Ibid XVII, 230.
- [3] Ibid XVII, 354.
- [4] Ibid XVIII, 794.
- [5] Inaug. Diss. J. H. Univ. 1895.
- [6] Inaug. Diss. J. H. Univ. 1896.
- [7] Am. Chem. Journ. I, 350.
- [8] Ibid XI, 177.
- [9] Inaug. Diss. J. H. Univ. 1845.
- [10] Inaug. Diss. J. H. Univ. 1896.
- [11] Ber. XXVI-2231; XXVIII-2281
- [12] Inaug. Diss. J. H. Univ. 1895.
- [13] Inaug. Diss. J. H. Univ. 1896.
- [14] Inaug. Diss. J. H. Univ. 1896.

- [\[15\]](#) Am. Ch. Journ. XI—281.
- [\[16\]](#) Inaug. Diss. J. H. Univ. 1895.
- [\[17\]](#) Am. Ch. Journ. XI, 341.
- [\[18\]](#) Am. Chem. Journ. XVII, 347.
- [\[19\]](#) Ibid. XVIII, 798.
- [\[20\]](#) Ber. XXVI, 204.
- [\[21\]](#) Am. Ch. Journ. XVIII-799
- [\[22\]](#) Am. Chem. Journ. XVI-525
- [\[23\]](#) Am. Ch. Journ. XVI-527.
- [\[24\]](#) Am. Chem. Journ. XIX-199
- [\[25\]](#) Ber. XVII-2545; XXIII, 3430.
- [\[26\]](#) Am. Ch. Journ. XVIII-810.
- [\[27\]](#) Inaug. Diss. J. H. Univ. 1895.

Transcriber's Notes:

The cover image was created by the transcriber, and is in the public domain.
Typographical and punctuation errors have been silently corrected.

*** END OF THE PROJECT GUTENBERG EBOOK A FURTHER
INVESTIGATION OF THE SYMMETRICAL CHLORIDE OF
PARANITROORTHOSULPHOBENZOIC ACID ***

Updated editions will replace the previous one—the old editions will be renamed.

Creating the works from print editions not protected by U.S. copyright law means that no one owns a United States copyright in these works, so the Foundation (and you!) can copy and distribute it in the United States without permission and without paying copyright royalties. Special rules, set forth in the General Terms of Use part of this license, apply to copying and distributing Project Gutenberg™ electronic works to protect the PROJECT GUTENBERG™ concept and trademark. Project Gutenberg is a registered trademark, and may not be used if you charge for an eBook, except by following the terms of the trademark license, including paying royalties for use of the Project Gutenberg trademark. If you do not charge anything for copies of this eBook, complying with the trademark license is very easy. You may use this eBook for nearly any purpose such as creation of derivative works, reports, performances and research. Project Gutenberg eBooks may be modified and printed and given away—you may do practically ANYTHING in the United States with eBooks not protected by U.S. copyright law. Redistribution is subject to the trademark license, especially commercial redistribution.

START: FULL LICENSE

THE FULL PROJECT GUTENBERG™ LICENSE

PLEASE READ THIS BEFORE YOU DISTRIBUTE OR USE THIS WORK

To protect the Project Gutenberg™ mission of promoting the free distribution of electronic works, by using or distributing this work (or any other work associated in any way with the phrase “Project Gutenberg”), you agree to comply with all the terms of the Full Project Gutenberg License available with this file or online at www.gutenberg.org/license.

Section 1. General Terms of Use and Redistributing Project Gutenberg electronic works

1.A. By reading or using any part of this Project Gutenberg electronic work, you indicate that you have read, understand, agree to and accept all the terms of this license and intellectual property (trademark/copyright) agreement. If you do not agree to abide by all the terms of this agreement, you must cease using and return or destroy all copies of Project Gutenberg electronic works in your possession. If you paid a fee for obtaining a copy of or access to a Project Gutenberg electronic work and you do not agree to be bound by the terms of this agreement, you may obtain a refund from the person or entity to whom you paid the fee as set forth in paragraph 1.E.8.

1.B. “Project Gutenberg” is a registered trademark. It may only be used on or associated in any way with an electronic work by people who agree to be bound by the terms of this agreement. There are a few things that you can do with most Project Gutenberg electronic works even without complying with the full terms of this agreement. See paragraph 1.C below. There are a lot of things you can do with Project Gutenberg electronic works if you follow the terms of this agreement and help preserve free future access to Project Gutenberg electronic works. See paragraph 1.E below.

1.C. The Project Gutenberg Literary Archive Foundation (“the Foundation” or PGLAF), owns a compilation copyright in the collection of Project Gutenberg electronic works. Nearly all the individual works in the collection are in the public domain in the United States. If an individual work is unprotected by copyright law in the United States and you are

located in the United States, we do not claim a right to prevent you from copying, distributing, performing, displaying or creating derivative works based on the work as long as all references to Project Gutenberg are removed. Of course, we hope that you will support the Project Gutenberg mission of promoting free access to electronic works by freely sharing Project Gutenberg works in compliance with the terms of this agreement for keeping the Project Gutenberg name associated with the work. You can easily comply with the terms of this agreement by keeping this work in the same format with its attached full Project Gutenberg License when you share it without charge with others.

1.D. The copyright laws of the place where you are located also govern what you can do with this work. Copyright laws in most countries are in a constant state of change. If you are outside the United States, check the laws of your country in addition to the terms of this agreement before downloading, copying, displaying, performing, distributing or creating derivative works based on this work or any other Project Gutenberg work. The Foundation makes no representations concerning the copyright status of any work in any country other than the United States.

1.E. Unless you have removed all references to Project Gutenberg:

1.E.1. The following sentence, with active links to, or other immediate access to, the full Project Gutenberg License must appear prominently whenever any copy of a Project Gutenberg work (any work on which the phrase “Project Gutenberg” appears, or with which the phrase “Project Gutenberg” is associated) is accessed, displayed, performed, viewed, copied or distributed:

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg™ License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

1.E.2. If an individual Project Gutenberg electronic work is derived from texts not protected by U.S. copyright law (does not contain a notice indicating that it is posted with permission of the copyright holder), the work can be copied and distributed to anyone in the United States without paying any fees or charges. If you are redistributing or providing access to a work with the phrase “Project Gutenberg” associated with or appearing on the work, you must comply either with the requirements of paragraphs 1.E.1 through 1.E.7 or obtain permission for the use of the work and the Project Gutenberg trademark as set forth in paragraphs 1.E.8 or 1.E.9.

1.E.3. If an individual Project Gutenberg electronic work is posted with the permission of the copyright holder, your use and distribution must comply with both paragraphs 1.E.1 through 1.E.7 and any additional terms imposed by the copyright holder. Additional terms will be linked to the Project Gutenberg License for all works posted with the permission of the copyright holder found at the beginning of this work.

1.E.4. Do not unlink or detach or remove the full Project Gutenberg License terms from this work, or any files containing a part of this work or any other work associated with Project Gutenberg.

1.E.5. Do not copy, display, perform, distribute or redistribute this electronic work, or any part of this electronic work, without prominently displaying the sentence set forth in paragraph 1.E.1 with active links or immediate access to the full terms of the Project Gutenberg License.

1.E.6. You may convert to and distribute this work in any binary, compressed, marked up, nonproprietary or proprietary form, including any word processing or hypertext form. However, if you provide access to or distribute copies of a Project Gutenberg work in a format other than “Plain Vanilla ASCII” or other format used in the official version posted on the official Project Gutenberg website (www.gutenberg.org), you must, at no additional cost, fee or expense to the user, provide a copy, a means of exporting a copy, or a means of obtaining a copy upon request, of the work in its original “Plain Vanilla ASCII” or other form. Any alternate format must include the full Project Gutenberg License as specified in paragraph 1.E.1.

1.E.7. Do not charge a fee for access to, viewing, displaying, performing, copying or distributing any Project Gutenberg works unless you comply with paragraph 1.E.8 or 1.E.9.

1.E.8. You may charge a reasonable fee for copies of or providing access to or distributing Project Gutenberg electronic works provided that:

- You pay a royalty fee of 20% of the gross profits you derive from the use of Project Gutenberg works calculated using the method you already use to calculate your applicable taxes. The fee is owed to the owner of the Project Gutenberg trademark, but he has agreed to donate royalties under this paragraph to the Project Gutenberg Literary Archive Foundation. Royalty payments must be paid within 60 days following each date on which you prepare (or are legally required to prepare) your periodic tax returns. Royalty payments should be clearly marked as such and sent to the Project Gutenberg Literary Archive Foundation at the address specified in Section 4, "Information about donations to the Project Gutenberg Literary Archive Foundation."
- You provide a full refund of any money paid by a user who notifies you in writing (or by e-mail) within 30 days of receipt that s/he does not agree to the terms of the full Project Gutenberg™ License. You must require such a user to return or destroy all copies of the works possessed in a physical medium and discontinue all use of and all access to other copies of Project Gutenberg™ works.
- You provide, in accordance with paragraph 1.F.3, a full refund of any money paid for a work or a replacement copy, if a defect in the electronic work is discovered and reported to you within 90 days of receipt of the work.
- You comply with all other terms of this agreement for free distribution of Project Gutenberg™ works.

1.E.9. If you wish to charge a fee or distribute a Project Gutenberg™ electronic work or group of works on different terms than are set forth in this agreement, you must obtain permission in writing from the Project Gutenberg Literary Archive Foundation, the manager of the Project

Gutenberg™ trademark. Contact the Foundation as set forth in Section 3 below.

1.F.

1.F.1. Project Gutenberg volunteers and employees expend considerable effort to identify, do copyright research on, transcribe and proofread works not protected by U.S. copyright law in creating the Project Gutenberg™ collection. Despite these efforts, Project Gutenberg™ electronic works, and the medium on which they may be stored, may contain “Defects,” such as, but not limited to, incomplete, inaccurate or corrupt data, transcription errors, a copyright or other intellectual property infringement, a defective or damaged disk or other medium, a computer virus, or computer codes that damage or cannot be read by your equipment.

1.F.2. LIMITED WARRANTY, DISCLAIMER OF DAMAGES - Except for the “Right of Replacement or Refund” described in paragraph 1.F.3, the Project Gutenberg Literary Archive Foundation, the owner of the Project Gutenberg™ trademark, and any other party distributing a Project Gutenberg™ electronic work under this agreement, disclaim all liability to you for damages, costs and expenses, including legal fees. YOU AGREE THAT YOU HAVE NO REMEDIES FOR NEGLIGENCE, STRICT LIABILITY, BREACH OF WARRANTY OR BREACH OF CONTRACT EXCEPT THOSE PROVIDED IN PARAGRAPH 1.F.3. YOU AGREE THAT THE FOUNDATION, THE TRADEMARK OWNER, AND ANY DISTRIBUTOR UNDER THIS AGREEMENT WILL NOT BE LIABLE TO YOU FOR ACTUAL, DIRECT, INDIRECT, CONSEQUENTIAL, PUNITIVE OR INCIDENTAL DAMAGES EVEN IF YOU GIVE NOTICE OF THE POSSIBILITY OF SUCH DAMAGE.

1.F.3. LIMITED RIGHT OF REPLACEMENT OR REFUND - If you discover a defect in this electronic work within 90 days of receiving it, you can receive a refund of the money (if any) you paid for it by sending a written explanation to the person you received the work from. If you received the work on a physical medium, you must return the medium with your written explanation. The person or entity that provided you with the defective work may elect to provide a replacement copy in lieu of a refund. If you received the work electronically, the person or entity providing it to

you may choose to give you a second opportunity to receive the work electronically in lieu of a refund. If the second copy is also defective, you may demand a refund in writing without further opportunities to fix the problem.

1.F.4. Except for the limited right of replacement or refund set forth in paragraph 1.F.3, this work is provided to you 'AS-IS', WITH NO OTHER WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR ANY PURPOSE.

1.F.5. Some states do not allow disclaimers of certain implied warranties or the exclusion or limitation of certain types of damages. If any disclaimer or limitation set forth in this agreement violates the law of the state applicable to this agreement, the agreement shall be interpreted to make the maximum disclaimer or limitation permitted by the applicable state law. The invalidity or unenforceability of any provision of this agreement shall not void the remaining provisions.

1.F.6. INDEMNITY - You agree to indemnify and hold the Foundation, the trademark owner, any agent or employee of the Foundation, anyone providing copies of Project Gutenberg™ electronic works in accordance with this agreement, and any volunteers associated with the production, promotion and distribution of Project Gutenberg™ electronic works, harmless from all liability, costs and expenses, including legal fees, that arise directly or indirectly from any of the following which you do or cause to occur: (a) distribution of this or any Project Gutenberg work, (b) alteration, modification, or additions or deletions to any Project Gutenberg work, and (c) any Defect you cause.

Section 2. Information about the Mission of Project Gutenberg

Project Gutenberg is synonymous with the free distribution of electronic works in formats readable by the widest variety of computers including obsolete, old, middle-aged and new computers. It exists because of the

efforts of hundreds of volunteers and donations from people in all walks of life.

Volunteers and financial support to provide volunteers with the assistance they need are critical to reaching Project Gutenberg's goals and ensuring that the Project Gutenberg collection will remain freely available for generations to come. In 2001, the Project Gutenberg Literary Archive Foundation was created to provide a secure and permanent future for Project Gutenberg and future generations. To learn more about the Project Gutenberg Literary Archive Foundation and how your efforts and donations can help, see Sections 3 and 4 and the Foundation information page at www.gutenberg.org.

Section 3. Information about the Project Gutenberg Literary Archive Foundation

The Project Gutenberg Literary Archive Foundation is a non-profit 501(c)(3) educational corporation organized under the laws of the state of Mississippi and granted tax exempt status by the Internal Revenue Service. The Foundation's EIN or federal tax identification number is 64-6221541. Contributions to the Project Gutenberg Literary Archive Foundation are tax deductible to the full extent permitted by U.S. federal laws and your state's laws.

The Foundation's business office is located at 41 Watchung Plaza #516, Montclair NJ 07042, USA, +1 (862) 621-9288. Email contact links and up to date contact information can be found at the Foundation's website and official page at www.gutenberg.org/contact

Section 4. Information about Donations to the Project Gutenberg Literary Archive Foundation

Project Gutenberg™ depends upon and cannot survive without widespread public support and donations to carry out its mission of increasing the number of public domain and licensed works that can be freely distributed in machine-readable form accessible by the widest array of equipment

including outdated equipment. Many small donations (\$1 to \$5,000) are particularly important to maintaining tax exempt status with the IRS.

The Foundation is committed to complying with the laws regulating charities and charitable donations in all 50 states of the United States. Compliance requirements are not uniform and it takes a considerable effort, much paperwork and many fees to meet and keep up with these requirements. We do not solicit donations in locations where we have not received written confirmation of compliance. To SEND DONATIONS or determine the status of compliance for any particular state visit www.gutenberg.org/donate.

While we cannot and do not solicit contributions from states where we have not met the solicitation requirements, we know of no prohibition against accepting unsolicited donations from donors in such states who approach us with offers to donate.

International donations are gratefully accepted, but we cannot make any statements concerning tax treatment of donations received from outside the United States. U.S. laws alone swamp our small staff.

Please check the Project Gutenberg web pages for current donation methods and addresses. Donations are accepted in a number of other ways including checks, online payments and credit card donations. To donate, please visit: www.gutenberg.org/donate.

Section 5. General Information About Project Gutenberg electronic works

Professor Michael S. Hart was the originator of the Project Gutenberg concept of a library of electronic works that could be freely shared with anyone. For forty years, he produced and distributed Project Gutenberg eBooks with only a loose network of volunteer support.

Project Gutenberg eBooks are often created from several printed editions, all of which are confirmed as not protected by copyright in the U.S. unless a

copyright notice is included. Thus, we do not necessarily keep eBooks in compliance with any particular paper edition.

Most people start at our website which has the main PG search facility:
www.gutenberg.org.

This website includes information about Project Gutenberg, including how to make donations to the Project Gutenberg Literary Archive Foundation, how to help produce our new eBooks, and how to subscribe to our email newsletter to hear about new eBooks.