

THE
HEATHERY;
OR A
MONOGRAPH
OF
THE GENUS ERICA:
VOL. 1.



THE
HEATHERY;
OR A
MONOGRAPH
OF
THE GENUS ERICA:
VOL. I.



The Project Gutenberg eBook of The heathery; or, A monograph of the genus Erica. vol. 1

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

Title: The heathery; or, A monograph of the genus Erica. vol. 1

Author: active 1799-1828 Henry Cranke Andrews

Release date: December 26, 2022 [eBook #69639]

Language: English

Original publication: United Kingdom: Henry G. Bohn, 1845

Other information and formats: www.gutenberg.org/ebooks/69639

Credits: Chuck Greif and the Online Distributed Proofreading Team at <http://www.pgdp.net> (This file was produced from images available at The Internet Archive)

*** START OF THE PROJECT GUTENBERG EBOOK THE HEATHERY; OR, A
MONOGRAPH OF THE GENUS ERICA. VOL. 1 ***

INTRODUCTION A SHORT DISSERTATION MODE OF CULTURE ALPHABETICAL INDEX

THE
HEATHERY;
OR A
MONOGRAPH
OF
THE GENUS ERICA:

CONTAINING
COLOURED ENGRAVINGS,
WITH
LATIN AND ENGLISH DESCRIPTIONS, DISSECTIONS, ETC.
OF ALL
THE KNOWN SPECIES OF THAT EXTENSIVE AND DISTINGUISHED
TRIBE OF PLANTS.

By H. C. ANDREWS.

IN SIX VOLUMES.

VOL. I.

SECOND EDITION, CORRECTED AND ENLARGED.

LONDON:
HENRY G. BOHN, YORK STREET, COVENT GARDEN.
1845.

INTRODUCTION.

As an apology for the present undertaking, the Author has only to observe, that the very great encouragement he has received in his folio edition of the ERICAS, joined with the repeated application of many of its subscribers, for a small, concise, and pleasant reference to that beautiful and extensive tribe, has induced him to offer to the amateurs of plants in general, the present Monograph; in which will be given all the known species, and likewise all those successively that may hereafter be introduced; with Latin and English descriptions, dissections, &c. on the same plan as the folio work.

The Author therefore hopes it will be regarded as an agreeable green-house companion, calculated to preserve the folio edition for the use of the library, to which on account of its size it is best adapted.

When one volume is completed, every necessary requisite for binding will be given.

A SHORT DISSERTATION.

THIS beautiful and extended tribe of plants, at present so much admired and cultivated in our British Gardens, is but of recent introduction, at least the major part of them, as antecedent to the year 1772 the few species then known were the *E. vulgaris*, *E. Tetralix*, *E. cinerea*, and *E. vagans*, natives; the *E. Dabœcii*, from Ireland; the *E. arborea* from Madeira in 1748; the *E. herbacea* or *carnea* in 1763 from Switzerland; the *E. mediterranea* in 1765 from Minorca; the *E. scoparia*, *E. viridipurpurea*, *E. australis*, *E. ciliaris*, and *E. umbellata*, from Portugal, between the years 1768 and 1707. The two other European species we possess, the *E. stricta*, and *E. multiflora*, natives of Spain, have been but 14 years in cultivation with us; and the African species found within the district of the Cape of Good Hope and the adjacent territory, which have swelled the Genus to so great an extent, and by the extreme brilliance of their flowers have contributed so much to the present splendour of our green-houses, were unknown to our English botanists, but by name, till the above æra. In the year 1772 seeds of two species were sent from the Cape.—Both vegetated. The first was the *E. tubiflora* of the *Sp. Plant.* of Linnæus, the other the *E. concinna*. In 1774 the superb collection at Kew was enriched by nearly 20 species sent by Mr. F. Masson, His Majesty's collector at the Cape, for which we refer to the 2nd vol. of the Catalogue of that garden. From this period, till within these few years, the accession has been so rapid, so many different collectors producing new species, that it would be only a list of names to enumerate them, and no way illustrate the present subject; nor would it be any way relevant to the subject, to know how many were enumerated by Petiver, Plukenet, Hermann, Oldenland, Ray, &c. &c. as, before Linnæus had by his mode of classification determined the precise limits of the Genus, the confusion that then pervaded all the elder botanists is such, that any comment from them would rather perplex than elucidate. Therefore, beginning with the *Systema Naturæ* of Linnæus, vol. ii. of 1767, including the European species, he there enumerates but 42; and Dahlgren, in 1770, edited a dissertation (under his inspection) on the Genus, containing a catalogue of 58 names from Bergius, the *Mantissa*, &c. Thunberg, on his return from Africa, added 13 to the number, all of which were inserted in the *Supplementum Plantarum* of 1781. From this and some other sources Murray has in his *Syst. Veg.* of 1784 made a list of 74 names, and Martyn in his edition of Miller's Dictionary, 1795, enumerates but 84. Willdenow, collating from all the foregoing, &c., has mounted the list to 137 in his *Species Plantarum*. This is certainly far short of the number at present cultivated in Britain; and from the variety of beautiful new specimens and seeds lately received by G. Hibbert, Esq. from his collector at the Cape, many of which have vegetated and are in high perfection at the Clapham Gardens, we may fairly conjecture that the Genus is by no means bounded by the species we at present possess.

MODE OF CULTURE.

VERY few, if any, of the species, but have been increased by cuttings. The only method we shall therefore treat of is this, and by seeds. The process for making and planting the cuttings is as follows: Let a middle-sized pot be prepared by filling it within three inches of the margin with small broken potsherds, or some such matter, that there may be a constant and ready drain for the water, which it will be necessary to give the cuttings at any time when the earth appears dry. Then, consulting the Index, take a sufficient quantity of sandy loam, sandy peat, or common sand, or a mixture of any of them, finely sifted, to fill the pot up to the margin, and press it lightly with the hand. Thus prepared, it is fit to receive the cuttings. These must be taken about an inch long, or shorter, from the smallest and tenderest shoots, cutting off the leaves two-thirds of the length, close to the shoot. When a few are thus made ready, take a small pointed piece of wood and dibble them into the pot; fixing the lower end of the cutting as firm as possible. When the pot is filled with them at about half an inch distance each way, cover them with a small bell-glass, to fit as near as possible. Then place it under a large hand-glass, or frame, where it must be kept from the stronger rays of the sun till the cuttings are rooted. Soon after the small bell-glass must be first removed, and in a week they may be exposed to the open air in the shade of a north wall, or, in winter, in the shady part of the green-house, to protect them from the sun and winds till they are removed into separate pots. The seeds of some species which ripen in this country, as well as those procured from the Cape, may be sown about the middle of March, in pots prepared the same way as for the cuttings; with this exception, that the earth be invariably sandy peat finely sifted. The seeds should be very slightly covered, and watered with a pot whose rose or head should only admit the water like a light dew, lest the seeds be all washed together. The smallest pots that can be procured are the best for transplanting. The earth in which most Heaths thrive best is a soft, loose, sandy peat.

ERICA.

CLASSIS VIII. ORDO I.

OCTANDRIA MONOGYNIA.

Character Genericus Essentialis.

CALYX. Perianthium tetraphyllum.

HEATH.

CLASS VIII. ORDER I.

EIGHT CHIVES. ONE POINTAL.

Essential Generic Character.

EMPALEMENT. Cup four-leaved.

COROLLA monopetala, quadrifida.

STAMINA. Filamenta octo, capillaria, receptaculo inserta. Antheræ apice bifidæ.

PISTILLUM. Germen superum. Stylus filiformis. Stigma coronulatum. Capsula quadrilocularis. Dissepimenta valvularum margine.

SEMINA numerosa, minima.

Obs. *Antheræ in aliis emarginatæ, in aliis bicornutæ, cristatæ, aristatæ, muticæ, vel penicillatæ; inclusæ vel exsertæ.*

BLOSSOM one petal, four-cleft.

CHIVES. Threads eight, hair-like, inserted into the receptacle. Tips two-cleft at the end.

POINTAL. Seed-bud above. Shaft, thread-shaped. Summit crowned. Capsule four-celled. Partitions from the edge of the valves.

SEEDS numerous, very small.

Obs. *The tips in some are notched at the end, in others two-horned, crested, bearded, beardless, or pencilled; within the blossom, or without.*

ERICA acuta.

DESCRIPTIO.

ANTHERÆ cristatæ, inclusæ. Flores in apice omnium ramulorum, terni, cernui. Corolla urceolata, purpurea, nitida. Folia quaterna, subulata, mucronata, glabra, erecta. Caulis erectus, pedalis, gracilis. Rami pauci; ramuli numerosi, brevi, erecti.

DESCRIPTION.


TIPS crested, within the blossom. Flowers grow by threes at the ends of all the small branches, nodding. Blossom pitcher-shaped, purple, and shining. Leaves by fours, awl-shaped, sharp-pointed, smooth, and upright. Stem upright, a foot high, slender. Branches few; the small branches numerous, short, and upright.

Native of the Cape of Good Hope.

In bloom from May till July.

REFERENCE.

1. A leaf magnified.
2. The empalement magnified.
3. A blossom.
4. The chives and pointal, one tip magnified.
5. Seed-bud and pointal, summit magnified.



E. acuta

ERICA albens.

DESCRIPTIO.

ANTHERÆ muticæ, inclusæ. Flores in axillis foliorum, solitarii, cernui. Corolla ventricosa, ore arctata, sub-albicans. Folia terna, trigona, linearia, acuta, glabra. Caulis fruticosus, pedalis, erectus, ramosus, gracilis. Rami simplices.

DESCRIPTION.

TIPS beardless, within the blossom. The flowers grow from the base of the foot-stalks of the leaves, singly, and are bent downwards. Blossom swelled at the base, straightened at the mouth, and whitish. Leaves by threes, which are three-sided, linear, sharp-pointed, and smooth. Stem shrubby, a foot high, upright, branching and slender. Branches simple.





Native of the Cape of Good Hope.

In bloom from April till July.

REFERENCE.

1. The empalement and floral leaves.
2. The chives and pointal, one tip magnified.
3. Seed-bud and pointal, summit magnified.



E. albens

Chen. Act. Pinetale 1845.

14. 36. Andrews. 5.

ERICA Archeria.

DESCRIPTIO.

ANTHERÆ muticæ, inclusæ, stylo exserto. Corolla tubuloso-clavata, pollicaris, villosa, rubro-purpurea, viscosa. Folia sena, linearia, rigida, patentia, margine serrato-ciliata. Caulis fruticosus, erectus, sesquipedalis. Rami erecti, simplices.

DESCRIPTION.

TIPS beardless, within the blossom, shaft without. Blossoms tubularly club-shaped, an inch long, hairy, of a red purple colour, and clammy. Leaves by sixes, linear, harsh, spreading, and sawed like a fringe at the margin. Stem shrubby, upright, a foot and a half high. Branches upright and simple.



Native of the Cape of Good Hope.

In bloom from August till November.

REFERENCE.

1. The empalement.
2. The chives and pointal.
3. Seed-bud and pointal, summit magnified.
4. The seed-bud magnified.



E. Archeria

at Dore's Port, 1804

by J. B. Andrews & Houghlin

ERICA barbata, *major*.

DESCRIPTIO.

ANTHERÆ cristatæ, inclusæ. Flores in ultimis ramulis terminales, umbellati, cernui. Pedunculi longi, pilosi. Corolla urceolata, alba, pilosa, viscosa. Perianthium tetraphyllum, foliolis spathulatis, barbatis, glanduloso-pilosis. Folia quaterna, ciliata, barbata, ovata. Caulis fruticosus, pedalis. Ramuli villosiusculi.

DESCRIPTION.

TIPS crested, within the blossom. Flowers terminate the small branches in umbels, nodding. The peduncles are long and hairy. Blossom pitcher-shaped, white, hairy and viscous. Cup four-leaved: leaflets spathula-shaped, bearded and glandularly hairy. Leaves by fours, fringed, bearded, and oval. Stem shrubby, a foot high. The smaller branches a little hairy.




Native of the Cape of Good Hope.

In bloom from May till August.

REFERENCE.

1. A leaf magnified.
2. The empalement magnified.
3. The chives detached, one tip magnified.
4. The pointal magnified.



E. barbata. major

from the ...

by the ...

ERICA barbata minor.

DESCRIPTIO.

ANTHERÆ cristatæ, subexsertæ. Flores terminales, umbellati, cernui. Corolla urceolata, alba, pilosa, viscosa, calyce spathulato, glanduloso-piloso. Pedunculi colorati. Folia quaterna, barbata, ovata, patentia. Caulis fruticosus, bipedalis. Rami filiformes, laxi, patentes.

DESCRIPTION.

TIPS crested, just without the blossom. Flowers terminate the branches in umbels, nodding. Blossom pitcher-shaped, white, hairy and viscous, with the cup spathula-shaped, and glandularly hairy. Footstalks coloured. Leaves by fours, bearded, ovate and spreading. Stem shrubby, two feet high. Branches thread-shaped, limber and spreading.

Native of the Cape of Good Hope.

In bloom from August till March.

REFERENCE.

1. The under side of a leaf magnified.
2. The empalement magnified.
3. The chives detached, one tip magnified.
4. The seed-bud magnified.



E. barbata minor

Wm. Pursh, Jun. Sculp.

Wm. Woodhouse, Sculp.

ERICA Bruinades.

DESCRIPTIO.

ANTHERÆ muticæ, exsertæ. Flores urceolati, pubescentes, albi, in toto tecti calyce lanato; calycis foliolis ovatis, carneis. Folia terna, linearia, obtusa, margine pilis longis hirta. Caulis filiformis, flexuosus. Ramuli capillares, frequentes, tomentosi.

DESCRIPTION.

TIPS beardless, without the blossom. Flowers pitcher-shaped, downy and white, and covered totally by a woolly cup; the leaflets of the cup are egg-shaped, and flesh-coloured. Leaves by threes, linear and blunt, the edges covered with long hairs. Stem thread-shaped, very flexible. The smaller branches hair-like, numerous and downy.



Native of the Cape of Good Hope.

In bloom from February till June.

REFERENCE.

1. The empalement.
2. The blossom.
3. The chives detached from the pointal, one tip magnified.
4. The seed-bud and pointal magnified.



E. Bruniades

The Act Directa Feb. 1805

by H. Andrews

ERICA caffra.

DESCRIPTIO.

ANTHERÆ aristatæ, inclusæ, stylo exserto. Corolla campanulata, minuta, cernua, odoratissima, alba. Caulis fruticosus, pedalis. Rami erecti. Folia quaterna, linearia, obtusa, sub-ciliata, patentia.

DESCRIPTION.

TIPS bearded, within the blossom, shaft without. Blossoms bell-shaped, small, nodding, sweet-scented, and white. Stem shrubby, grows a foot high. The branches are upright. Leaves by fours, linear, obtuse, a little hairy, and spreading.



Native of the Cape of Good Hope.

In bloom from October till February.

REFERENCE.

1. A leaf magnified.
2. The empalement and floral leaves magnified.
3. The chives and pointal magnified.
4. The seed-bud magnified.



ERICA calycina.

DESCRIPTIO.

ANTHERÆ cristatæ, inclusæ. Flores plures, in ultimis ramulis terminales, umbellati. Calyx tetraphyllus, foliolis lato-ovatis, carneis, corolla longiora, pedunculis longissimis. Corolla minuta, urceoliformis, apice dilute carnea, ad basin virescens; oris laciniis reflexis, saturate carneis. Folia terna, lanceolata, supra concava, adpressa, glabra. Caulis filiformis, spithameus, erectus.

DESCRIPTION.

TIPS crested, within the blossom. Flowers numerous, terminating the small branches in bunches. Cup four-leaved, of a broad oval form, flesh-coloured, and longer than the blossom. The foot-stalks very long. Blossom small, of a pitcher-shape, light flesh-colour at the end, and greenish at the base. The segments of the mouth are bent back, and deep flesh-coloured. Leaves by threes, lance-shaped, concave on the upper surface, pressed to the stem, and smooth. Stem thread-shaped, half a foot high, and upright.

Native of the Cape of Good Hope.

In bloom from August till December.

REFERENCE.

1. The empalement and floral leaves.
2. The blossom.
3. The chives detached from the pointal, one tip magnified.
4. The seed-bud and pointal; summit magnified.



E. calycina

Det. M. B. S.

by H. Andromed. & R. H. S.

ERICA cernua.

DESCRIPTIO.

ANTHERÆ cristatæ, inclusæ. Flores umbellati, cernui, secundi, subovati, carnei; laciniis limbi acuminatis, sub-erectis. Folia quaterna, linearia, obtusa; petiolis brevissimis. Caulis fruticosus, erectus, pedalis. Rami sub-simplices, erecto-patentes.

DESCRIPTION.

TIPS crested, within the blossom. The flowers grow in bunches, nodding, all pointing one way; nearly egg-shaped, of a pale flesh colour; the segments of the border tapered, and nearly upright. Leaves grow by fours, are linear and blunt, having very short footstalks. Stem shrubby, upright, a foot high. The branches are almost simple, upright, and spreading.

Native of the Cape of Good Hope.

In bloom from August till December.

REFERENCE.

1. The empalement and floral leaves magnified.
2. The chives detached from the pointal, one tip magnified.
3. The seed-bud and pointal magnified.



E. cernua

by Th. Anderson & J. D. Hooker del.

ERICA comosa.

DESCRIPTIO.

ANTHERÆ muticæ, inclusæ. Flores in ramulis terminales, quasi spicam comosam densam formantes. Pedunculi brevissimi. Corolla ventricosa, alba, basi inflata, apice arctata. Folia quaterna, obtusa, patentia, parva, glabra. Caulis fruticosus, subpedalis, ramosus. Rami virgati, penduli.

DESCRIPTION.



TIPS beardless, within the blossom. Flowers terminate the small branches, forming as it were a thick tufted spike. Footstalks very short. Blossom bellied, white, swelled at the base, and narrowed at the mouth. Leaves by fours, blunt, spreading, small, and smooth. Stem shrubby, near a foot high, and branching. The branches are twiggy, and hang down.



Native of the Cape of Good Hope.
In bloom from June till September.

REFERENCE.

1. The empalement magnified.
2. The chives and pointal.
3. Seed-bud and pointal, summit magnified.
4. The seed-bud magnified.



E. comosa

at Dorset House

by H. Andrews

ERICA comosa, rubra.

DESCRIPTIO.

ANTHERÆ muticæ, inclusæ. Flores in ramulis terminales. Pedunculi brevissimi. Corolla ventricosa, carnea, basi inflata, apice arctata. Folia quaterna, obtusa, parva, glabra. Caulis fruticosus, pedalis, ramosus. Rami erecti, virgati, numerosi.

DESCRIPTION.

TIPS beardless, within the blossom. Flowers terminate the small branches. Foot-stalks very short. Blossom bellied, flesh-coloured, swelled at the base, and narrowed at the mouth. Leaves by fours, blunt, small, and smooth. Stem shrubby, a foot high, branching. The branches are numerous, upright, and twiggy.

Native of the Cape of Good Hope.

In bloom from August till December.

REFERENCE.

1. The empalement and floral leaves magnified.
2. The chives and pointal.
3. A chive magnified.
4. The seed-bud and pointal, summit magnified.



E. Comosa rubra

ERICA conspicua.

DESCRIPTIO.

ANTHERÆ basi bicornes, exsertæ. Corolla clavata, sesquipollicaris, lutea, basi attenuata; oris laciniis magnis, revolutis. Folia quaterna, obtusa, linearia, crassa. Flores subterni. Caulis fruticosus, erectus, tripedalis, ramosus.

DESCRIPTION.

TIPS two-horned at their base, and without the blossom. The blossoms are club-shaped, an inch and a half long, yellow, and tapering to the base: the segments of the mouth are large, and rolled back. Leaves grow by fours, are blunt, linear, and thick. Flowers grow mostly by threes. Stem shrubby, upright, grows three feet high, and branching.

Native of the Cape of Good Hope.

In bloom from June till August.

REFERENCE.

1. The empalement with its three floral leaves.
2. The chives detached from the pointal, one tip magnified.
3. The seed-bud and pointal, summit magnified.



Castroville, Cal. 1891

E. conspicua

by H. G. Anderson

ERICA costata.

DESCRIPTIO.

ANTHERÆ muticæ, inclusæ, stylo exserto. Corolla sub-cylindræa, incarnata, costata, oris laciniis erectis albidis. Folia caulina terna, linearia, obtusa, pubescentia; folia ramentacea erectiora, subserrata. Caulis fruticosus, bipedalis, erectus, ramosus. Ramuli frequentissimi.

DESCRIPTION.



TIPS beardless, and within the blossom; pointal without. Blossoms nearly cylindrical, of a flesh colour, and ribbed; the segments of the mouth upright and white. The leaves of the stem grow by threes, linear, blunt, and downy; those on the smaller branches more upright, and a little sawed at the edges. Stem shrubby, two feet high, upright and branching. The smaller branches numerous.

Native of the Cape of Good Hope.

In bloom from August till February.

REFERENCE.

1. The empalement.
2. The chives and pointal.
3. A chive magnified.
4. Seed-bud and pointal, summit magnified.



E. costata

ERICA cubica.

DESCRIPTIO.

ANTHERÆ basi bicornutæ, sub-inclusæ. Flores subterminales, cernui; pedunculis longissimis. Corolla globoso-campanulata, purpurea; limbo maximo, expanso. Folia quaterna, sæpe quina; nitida, obtusa, reflexa, apice incurvata. Caulis erectus, fruticosus, pedalis. Rami et ramuli filiformes.

DESCRIPTION.

TIPS two-horned at the base, nearly within the blossom. Flowers near the summits of the branches, hanging down; with very long foot-stalks. Blossom globularly bell-shaped and purple; the border very large and spreading. Leaves by fours, and often fives; shining, blunt, bent back, and turned inward at the point. Stem upright, shrubby, a foot high. The larger and smaller branches thread-shaped.




Native of the Cape of Good Hope.

In bloom from April till July.

REFERENCE.

1. The empalement and floral leaves magnified.
2. The chives and pointal, one tip magnified.
3. The seed-bud and pointal, summit magnified.



ERICA cubica, minor.

DESCRIPTIO.

ANTHERÆ muticæ, sub-inclusæ. Flores sub-terminales, umbellati; pedunculis longissimis. Corolla sub-campanulata, purpurea; limbo maximo. Folia quaterna, obtusa, linearia, nitida. Caulis erectus, ramosus, palmaris. Rami et ramuli verticillati.

DESCRIPTION.

TIPS beardless, just within the blossom. Flowers near the ends of the branches in umbels. Foot-stalks very long. Blossom nearly bell-shaped, and purple; border very large. Leaves by fours, blunt, linear, and shining. Stem upright, branching, a span high. The larger and smaller branches grow in whorls.



Native of the Cape of Good Hope.

In bloom from August till November.

REFERENCE.

1. The empalement and floral leaves magnified.
2. The chives and pointal.
3. The same magnified.



E. cubica minor

ERICA curviflora.

DESCRIPTIO.

ANTHERÆ muticæ, inclusæ. Corolla curvata, clavato-cylindræa, pubescens, terminalis, luteo-aurantia. Folia quaterna, linearia, glabra. Caulis fruticosus, bipedalis, laxis, superne villosus. Rami laxi, numerosi; ramulis brevissimis, frequentissimis, sparsis.

DESCRIPTION.

TIPS beardless, within the blossom, which is curved, cylindrically club-shaped, downy, terminating the branches, and of a yellow gold colour. The leaves grow by fours, are linear and smooth. Stem shrubby, grows two feet high, weak, and hairy at the top. Branches weak, and numerous; the smaller branches are very short, numerous, and scattered.

Native of the Cape of Good Hope.

In bloom from July till November.

REFERENCE.

1. The empalement and floral leaves.
2. The chives detached from the pointal, summit magnified.
3. Seed-bud and shaft, summit magnified.



E. curviflora

ERICA depressa.

DESCRIPTIO.

ANTHERÆ aristatæ, inclusæ. Flores in ramulis terminales, subterni, sub-sessiles; bracteis adpressis, imbricatis. Corolla sub-cylindrica, sub-pollicaris, flava. Folia quaterna, crassiuscula, obtusa. Caulis fruticosus, robustus. Rami divaricati, flexuosi, depressi.

DESCRIPTION.


TIPS bearded, within the blossom. Flowers terminate the small branches, in general sitting close to the stem. Floral leaves tiled, and pressed to the cup. Blossom almost cylindrical, near an inch long, and yellow. Leaves by fours, thickish and blunt. Stem shrubby and stout. The branches are straggling, zigzag, and pressed downwards.

Native of the Cape of Good Hope.

In bloom from May till August.

REFERENCE.

1. The empalement and floral leaves.
2. The chives detached from the pointal, one tip magnified.
3. The seed-bud and pointal magnified.



ERICA droseroides.

DESCRIPTIO.

ANTHERÆ cristatæ, inclusæ. Corolla ventricosa, læte purpurea, costata, viscosa, pubescens; ore obliqua, arctata, profunde sanguinea; pedunculi longissimi. Folia alterna, sparsa, obtusa, apice reflexa, pilis glandulosis micantibus hirta. Caulis erectus, spithamæus. Rami filiformes, sparsi, erecto-patentes, virgati.

DESCRIPTION.

TIPS crested, within the blossom, which is bellied, bright purple, ribbed, clammy, and downy; with the mouth oblique, narrowed, and of a deep blood colour; foot-stalks very long. Leaves alternate, scattered and blunt, the end reflexed, and thick with shining glandular hairs. Stem upright, a span high. Branches thread-shaped, scattered, upright, spreading, and twiggy.



Native of the Cape of Good Hope.

In bloom from July till October.

REFERENCE.

1. The empalement and floral leaves magnified.
2. The chives detached from the pointal, one tip magnified.
3. Seed-bud and pointal, summit magnified.



E. droseroides

ERICA empetroides.

DESCRIPTIO.

ANTHERÆ cristatæ, inclusæ, stylo exserto. Flores in medio ramorum densissimè spicati, horizontaliter siti. Pedunculi brevissimi. Corolla urceolata, carnea, odorata. Folia sena, linearia, villosa, obtusa, horizontalia. Caulis debilis, pedalis. Rami filiformes, laxi.

DESCRIPTION.

TIPS crested, within the blossom, shaft without. Flowers in close spikes in the centre of the branches, standing out horizontally, with very short footstalks. Blossom pitcher-shaped, flesh-coloured, and sweet-scented. Leaves by sixes, linear, hairy, blunt and horizontal. Stem weak, a foot high. The branches are thread-shaped and loose.

Native of the Cape of Good Hope.

In bloom from May till August.

REFERENCE.

1. A leaf magnified.
2. Empalement and blossom.
3. Empalement magnified.
4. The chives and pointal.
5. A chive magnified.
6. Seed-bud and pointal, summit magnified.
7. The seed-bud magnified.



E. empetroides

Walt. Jon. 2. 1805.

by J. G. Andreux & H. G. Andreux

ERICA exsurgens.

DESCRIPTIO.

ANTHERÆ muticæ, exsertæ. Corolla clavato-cylindræa, pollicaris, aurantia. Flores verticillati; verticillis alter altero exsurgentibus. Folia quaterna, linearia, glabra, apice reflexo, mucronato. Caulis fruticosus, pedalis et ultra, erectus. Rami verticillati.

DESCRIPTION.

TIPS beardless, without the blossom. Blossoms cylindrically club-shaped, an inch long, and orange-coloured; grow in whorls, the whorls rising out of each other in succession. Leaves grow by fours, are linear, smooth, and bent downward at the point, which is sharp. Stem shrubby, rather more than a foot high, upright. The branches grow in whorls.

Native of the Cape of Good Hope.

In bloom from September till March.

REFERENCE.

1. The empalement and floral leaves.
2. The chives detached from the pointal.
3. The seed-bud and pointal, summit magnified.



Not direct. ex. linn.

E. exurgens.

by J. C. Burmann. Sculp. by J. G. de Vries.

ERICA exsurgens, *minor*.

DESCRIPTIO.

ANTHERÆ muticæ, sub-exertæ. Flores in medio ramorum verticillati, recti, verticilli alter alteri exsurgentes. Corolla clavato-cylindrica, saturate aurantia, semi-uncialis. Folia sena, linearia, glabra, apice mucronata. Caulis fruticosus, pedalis. Rami erecti, simplices.

DESCRIPTION.


TIPS beardless, just without the blossom. Flowers grow in whorls near the middle of the branches, straight out; the whorls rise out of each other in succession. Blossom cylindrically club-shaped, of a deep orange-colour, half an inch long. Leaves by sixes, linear, smooth, and sharp at the point. Stem shrubby, a foot high. The branches are upright and simple.

Native of the Cape of Good Hope.

In bloom from June till October.

REFERENCE.

1. The empalement and blossom.
2. Flower of a yellow variety.
3. The chives spread open, one tip magnified.
4. The seed-bud and pointal, summit magnified.
5. The seed-bud magnified.



E. exurgens. minor

ERICA filamentosa.

DESCRIPTIO.

ANTHERÆ muticæ, inclusæ. Flores axillares sub apice ramorum, verticillati, numerosi. Pedunculi longissimi. Corolla tubuloso-campanulata, purpurea. Folia sena, tremulantia, linearia, tenuia. Caulis erectus, sesquipedalis. Rami simplices, verticillati.

DESCRIPTION.

TIPS beardless, within the blossom. Flowers grow from the insertion of the leaves near the ends of the branches in whorls, and numerous. The footstalks very long. Blossom tubularly bell-shaped and purple. Leaves by sixes, tremulous, linear and slender. Stem upright, a foot and a half high. Branches simple and in whorls.

Native of the Cape of Good Hope.

In bloom from November till April.

REFERENCE.

1. The empalement and floral leaves.
2. The chives detached, one tip magnified.
3. The seed-bud and pointal, summit magnified.
4. The seed-bud magnified.



E. filamentosa

Feb. 1. 1825

ERICA flammea.

DESCRIPTIO.

ANTHERÆ muticæ, inclusæ. Flores in ramulis terminales, racemum formantes prope caulis summitatem. Corolla tubiformis, sub-pollicaris, pallide lutea; apice sub-albidâ. Folia terna, sæpe quaterna, obtusa, linearia. Caulis erectus, bipedalis. Ramuli filiformes, virgati.

DESCRIPTION.


TIPS beardless, within the blossom. Flowers terminating the small branches, making a long spike near the summit of the stem. Blossom trumpet-shaped, near an inch long, of a pale yellow; the end whitish. Leaves by threes, often in fours, blunt, and linear. Stem upright, two feet high. Branches thread-shaped, and twiggy.

Native of the Cape of Good Hope.

In bloom from October till February.

REFERENCE.

1. The empalement and floral leaves magnified.
2. The chives and pointal, one tip magnified.
3. Seed-bud and pointal, summit magnified.



E. flammea

ERICA gelida.

DESCRIPTIO.

ANTHERÆ aristatæ, inclusæ. Floras in ultimis ramis axillares, spicati, imbricati, dependentes. Corolla cylindrica, apice arctata, pollicaris, albicante-virescens, apice atro-virens. Folia ramulorum, subquaterna; ramorum sena, curvata, acuta, patentia. Caulis strictissimus, fruticosus, tripedalis. Rami pauci. Ramuli verticillati.

DESCRIPTION.



TIPS bearded, within the blossom. Flowers grow near the top of the branches, close to the stem, in spikes, imbricated hanging down. Blossom cylindrical, narrowed at the top, an inch long, of a whitish green, and dark green at the end. Leaves on the small branches mostly by fours; those on the larger by sixes, curved, pointed, and spreading. Stem very upright, shrubby, three feet high. Few large branches. The small branches grow in whorls.

Native of the Cape of Good Hope.

In bloom from April till July.

REFERENCE.

1. The empalement and floral leaves.
2. The chives detached, one tip magnified.
3. Seed-bud and pointal, summit magnified.



E. gelida

by H. Andrews & Knight

ERICA glauca.

DESCRIPTIO.

ANTHERÆ cristatæ, inclusæ. Flores terminales, umbellati, nutantes. Pedunculi longissimi, colorati. Corolla sub-conica, atro-purpurea, ore arctata. Folia terna, acuta, patentia, glabra, carnosæ, glauca. Caulis erectus, fragilis, bipedalis. Rami erecti, longi, sub-simplices.

DESCRIPTION.

TIPS crested, within the blossom. Flowers terminate the branches in bunches, nodding. The footstalks are very long, and coloured. Blossom nearly conical, dark purple, and narrowed at the mouth. Leaves by threes, linear-pointed, and spreading, smooth, fleshy, and of a sea-green colour. Stem upright, brittle, two feet high. The branches are upright and long, with few small branches.



Native of the Cape of Good Hope.

In bloom from May till July.

REFERENCE.

1. The empalement and floral leaves.
2. The chives detached from the pointal.
3. A chive magnified.
4. Seed-bud and pointal, summit magnified.



E. glauca

W. A. N. S. 1871

ERICA ignescens.

DESCRIPTIO.

ANTHERÆ muticæ, sub-exsertæ. Flores terminales, solitarii; pedunculis fere nullis. Corolla clavata, tomentosa, ad basin gibbosa, rubro-coccinea. Folia quaterna, linearia, glabra. Caulis erectus, sesquipedalis. Ramuli virgati, filiformes, numerosi.

DESCRIPTION.

TIPS beardless, within the blossom. Flowers terminal, solitary; scarcely any foot-stalks. Blossom club-shaped, downy, gouty at the base, and of a red-scarlet colour. Leaves by fours, linear, and smooth. Stem upright, a foot and a half high. The branches are twiggy, thread-shaped, and numerous.

Native of the Cape of Good Hope.

In bloom from March till June.

REFERENCE.

1. The empalement and floral leaves magnified.
2. The chives and pointal, one tip magnified.
3. Seed-bud and pointal, summit magnified.
4. The seed-bud magnified.

ERICA jasminiflora.

DESCRIPTIO.

ANTHERÆ basi bicornes, inclusæ; stylo exserto. Corolla viscosa, ampullacea, sesquipollicaris, summa cylindracea, ad basin inflata, ore arctata; laciniis cordatis, patentibus. Folia terna, trigona, subulata, erecta, patentia. Caulis filiformis, erectus. Rami simplicissimi, filiformes, longi, patentes.

DESCRIPTION.

TIPS two-horned at the base, and within the blossom; the shaft without. Blossoms clammy, flask-shaped, an inch and a half long, the upper part cylindrical, swelled at the base, and pinched in at the mouth; the segments heart-shaped and spreading. Leaves grow by threes, are three-sided, awl-shaped, upright and spreading. Stem thread-shaped and upright. The branches are quite simple, thread-shaped, long, and spreading.



Native of the Cape of Good Hope.

In bloom from July till November.

REFERENCE.

1. The empalement and floral leaves.
2. The chives detached from the pointal, one tip magnified.
3. The seed-bud and pointal, summit magnified.



E. jasminiflora.

by H. C. Andrews & Knightbridge.

ERICA leucanthera.

DESCRIPTIO.

ANTHERÆ cristatæ, inclusæ. Flores in ramulis terminales, conferti, racemosi. Pedunculi breves. Corolla urceolata, alba. Folia terna, trigona, glabra, obtusa, adpressa. Caulis fruticosus, pedalis. Rami et ramuli filiformes.

DESCRIPTION.

TIPS crested, within the blossom. Flowers terminate the small branches, clustered together, and forming long bunches. Footstalks short. Blossom pitcher-shaped and white. Leaves by threes, three-sided, smooth, blunt, and pressed to the stem. Stem shrubby, a foot high. The larger and smaller branches are thread-shaped.



Native of the Cape of Good Hope.

In bloom from January till March.

REFERENCE.


1. A blossom with its empalement.
2. The empalement magnified.
3. The chives and pointal magnified.



1



2



3

E. leucanthera

Currier, April 1851

by J. B. Andrews & Horner

ERICA lutea.

DESCRIPTIO.

ANTHERÆ muticæ, inclusæ. Corolla lutea, ovato-acuminata, ore arctata. Flores plures, conglomerati. Folia opposita, linearia, adpressa, triquetra, nitida. Caulis laxus, filiformis, gracilis, ad basin ramosus. Ramuli conferti.

DESCRIPTION.



TIPS beardless, within the blossom, which is yellow, of a pointed oval shape, and compressed at the mouth. The flowers are numerous, and close together. The leaves grow opposite in pairs, are linear, three-sided, pressed to the stem, and shining. Stem flexible, thread-shaped, slender, branching to the bottom. The branches crowded together.

Native of the Cape of Good Hope.

In bloom from September till May.

REFERENCE.

1. The empalement and floral leaves.
2. The chives detached from the pointal, one tip magnified.
3. The seed-bud and pointal, summit magnified.



E. lutea.

Wm. B. Woodcock & Son

ERICA Melastoma.

DESCRIPTIO.

ANTHERÆ muticæ, penicillatæ, exsertæ, longitudine corollæ, attenuatæ in filamenta plana. Flores flavi, conici, parum curvati, laciniis longissimis adpressis, nigris. Folia quaterna, linearia, sub-scabrida, rigida. Caulis laxis, erectus, bipedalis. Ramuli brevissimi, foliosi.

DESCRIPTION.

TIPS beardless, pencil-like, and without the blossom, tapering into threads, which are flat. Flowers yellow, conical, slightly curved, the segments of the border very long, pressed to the chives, and black. Leaves by fours, linear, roughish, and stiff. Stem weak, upright, grows two feet high. The small branches very short, and covered with leaves.




Native of the Cape of Good Hope.

In bloom from February till July.

REFERENCE.

1. The empalement.
2. A flower divested of its empalement.
3. The chives detached from the pointal.
4. The seed-bud and pointal, summit magnified.



E. melastoma

by H. Andrews & Son

ERICA nigrita.

DESCRIPTIO.

ANTHERÆ muticæ, exertæ, atræ. Flores in apicibus ramulorum terminales, sub-terni. Corolla campanulata, albida; laciniis oris revolutis, et tubo majoribus. Folia terna, glabra, nitida, sub-triquetra, obtusa, crassa. Caulis fruticosus, erectus, pedalis. Rami et ramuli frequentissimi, sub-erecti, virgati.

DESCRIPTION.

TIPS beardless, without the blossom, and black. Flowers grow from the ends of the small branches, mostly by threes. Blossom bell-shaped and white; the segments of the mouth are rolled back, and larger than the tube. Leaves by threes, smooth, shining, nearly three-sided, blunt, and thick. Stem shrubby, upright, a foot high. The larger and smaller branches are numerous, nearly upright, and twiggy.




Native of the Cape of Good Hope.

In bloom from April till July.

REFERENCE.

1. The empalement and floral leaves magnified.
2. The chives spread open, one tip magnified.
3. Seed-bud and pointal magnified.



E. nigrita

14 1/2 1/2

ERICA obbata.

DESCRIPTIO.

ANTHERÆ basi bicornes, inclusæ. Flores terminales, quaterni, erecti, inflati ore arctati, leviter striati, nitidi. Limbus maximus, glandulæ octo fauci circumpositæ. Folia quaterna, reflexa, rigida, ciliata. Caulis fruticosus, pedalis, ramosus.

DESCRIPTION.

TIPS two-horned at the base, within the blossom. Flowers grow by fours at the ends of the branches, upright, swelled out and contracted at the mouth, slightly striped and shining. Border very large, with eight glands surrounding the mouth. Leaves by fours, reflexed, harsh, and fringed. Stem shrubby, a foot high and branching.

Native of the Cape of Good Hope.

In bloom from April till June.

REFERENCE.

1. The empalement.
2. A segment of the border magnified.
3. The chives, one tip magnified.
4. The pointal, summit magnified.




1



2



3



4

E. obbata

ERICA obliqua.

DESCRIPTIO.

ANTHERÆ ad basin bicornutæ, inclusæ. Flores numerosi, ovati, purpurei, ad apices ramorum umbellati; pedunculis flore triplo longioribus, coloratis, viscidis. Folia sparsa, obliqua, truncata; petiolis tenuissimis. Caulis fruticosus, pedalis. Rami simplices, longi, filiformes.

DESCRIPTION.


TIPS two-horned at the base, and within the blossom. Flowers numerous, egg-shaped, purple, growing in umbels at the ends of the branches. Foot-stalks thrice the length of the flowers, coloured and clammy. Leaves obliquely scattered, seeming cut off at the ends, and with very slender foot-stalks. Stem shrubby, a foot high. The branches simple, long, and thread-shaped.

Native of the Cape of Good Hope.

In bloom from August till December.

REFERENCE.

1. The empalement magnified.
2. The seed-bud, chives and pointal.
3. A chive magnified.
4. The seed-bud and pointal magnified.



E. obliqua.

ERICA physodes.

DESCRIPTIO.

ANTHERÆ cristatæ, inclusæ. Corolla ovata, inflata, glabra, viscosa, crystallina, alba, magnitudine pisi; laciniis acuminatis erectis. Folia quaterna, obtusa, margine glandulosa, viscosa, patentia. Caulis fruticosus, bipedalis, erectus, ramulosus.

DESCRIPTION.

TIPS crested, within the blossoms, which are egg-shaped, swelled out, smooth, clammy, like crystal, and white, the size of a pea; segments of the border tapering and upright. Leaves by fours, are blunt, the edges glandular, clammy, and spreading. Stem shrubby, grows two feet high, is upright, and has numerous little branches.



Native of the Cape of Good Hope.

In bloom from February till July.

REFERENCE.

1. The empalement and floral leaves magnified.
2. The chives and pointal.
3. A chive magnified.
4. The seed-bud and pointal magnified.



Del. 1. 1804.

E. physodes

by J. L. Smith & J. H. Smith

ERICA pinea.

DESCRIPTIO.

ANTHERÆ muticæ, inclusæ. Corolla tubuloso-clavata, sub-albida. Flores in medio ramorum verticillati, horizontales, pollicares. Folia sena, linearia, glabra, patentia, longissima. Caulis erectus, bipedalis, robustus. Rami verticillati, erecto-patentes.

DESCRIPTION.

TIPS beardless, within the blossom, which is tubularly club-shaped, and nearly white. Flowers grow in whorls about the middle of the branches, horizontal, an inch in length. Leaves by sixes, linear, smooth, spreading outward, very long. Stem upright, two feet high, very stout. Branches in whorls, spreading, upright.

Native of the Cape of Good Hope.

In bloom from August till December.

REFERENCE.

1. The empalement and floral leaves.
2. The chives detached from the pointal, a tip magnified.
3. Seed-bud and pointal, summit magnified.
4. The seed-bud magnified.



E. pinsea

ERICA plumosa.

DESCRIPTIO.

ANTHERÆ aristatæ, inclusæ. Flores axillares, verticillati, sub-solitarii, cernui. Corolla urceolata, glabra, purpurea. Folia quaterna, obtusa, villosa, patentia, linearia. Caulis fruticosus, flexuosus, pedalis. Rami et ramuli filiformes.

DESCRIPTION.

TIPS bearded, within the blossom. Flowers grow from the insertion of the leaves, whorled, mostly solitary, and nodding. Blossom pitcher-shaped, smooth, and purple. Leaves by fours, blunt, hairy, linear, and spreading. Stem shrubby, and taking various directions, a foot high. The large and small branches are thread-shaped.



Native of the Cape of Good Hope.

In bloom from March till July.

REFERENCE.

1. A leaf magnified.
2. The empalement and floral leaves magnified.
3. The chives and pointal, one tip magnified.
4. The seed-bud and pointal, summit magnified.
5. The seed-bud magnified.



E. plumosa

by H. Andrews & Hays

ERICA pubescens.

DESCRIPTIO.

ANTHERÆ aristatæ, inclusæ. Flores in ramulis terminales, umbellati, bini, terni, vel plures, copiosi. Corolla ovata, obtusa, villosa, purpurea. Folia quaterna, obtusa, villosa, incurva, arcuata. Caulis fruticosus, sesquipedalis, ramosus, pilosus. Ramuli pilosi, filiformes, virgati.

DESCRIPTION.

TIPS bearded, within the blossom. Flowers terminate the branches in umbels of two, three, or more, and are very abundant. Blossom egg-shaped, blunt, covered with soft hairs, and purple. Leaves by fours, blunt, hairy, turned inwards, and bowed. Stem shrubby, a foot and half high, branching, and hairy. The branches are hairy, thread-shaped, and twiggy.



Native of the Cape of Good Hope.

In bloom from August till February.

REFERENCE.

1. The empalement and floral leaves magnified.
2. The chives and pointal, one tip magnified.
3. The seed-bud and pointal, summit magnified.
4. The seed-bud magnified.



E. pubescens

by H. Andrews. s. p.

ERICA pubescens, *minor*.

DESCRIPTIO.

ANTHERÆ aristatæ, inclusæ. Flores in ramulis terminales, umbellati, numerosi. Corolla ovata, obtusa, villosa; laciniis minutis, erectis. Folia quaterna, obtusa, hirta, erecta. Caulis sesquipedalis, ramosus, pilosus. Rami et ramuli pilosi, virgati, erecti.

DESCRIPTION.

TIPS bearded, within the blossom. Flowers terminate the small branches in numerous umbels. Blossom egg-shaped, obtuse, covered with soft hairs; segments of the border very small and upright. Leaves by fours, blunt, hairy and upright. Stem a foot and a half high, branching and hairy. The large and small branches are hairy, twiggy, and upright.

Native of the Cape of Good Hope.

In bloom from October till March.

REFERENCE.

1. The empalement and floral leaves magnified.
2. The chives and pointal.
3. A chive magnified.
4. Seed-bud and pointal magnified.



E. pubescens. minor.

Mon. 1808

by H. Andrews

ERICA pubescens, *minima*.

DESCRIPTIO.

ANTHERÆ aristatæ, inclusæ. Flores in ultimis ramulis terminales, umbellati, cernui. Corolla minuta, urceolata, purpurea, villosa. Folia quaterna, obtusa, villosa, incurva, arcuata. Caulis fruticosus, pedalis, ramosus, pilosus. Ramuli pilosi, virgati, filiformes.

DESCRIPTION.

TIPS bearded, within the blossom. Flowers terminate the small branches in umbels, nodding. Blossom very small, pitcher-shaped, purple, and hairy. Leaves by fours, blunt, covered with soft hairs, turned inwards, and bowed. Stem shrubby, a foot high, branching, and hairy. The branches are hairy, twiggy, and thread-shaped.

Native of the Cape of Good Hope.

In bloom from November till April.

REFERENCE.

1. The empalement magnified.
2. A blossom magnified.
3. The chives and pointal magnified.
4. The seed-bud magnified.



E. pubescens minima

ERICA pulchella.

DESCRIPTIO.

ANTHERÆ muticæ, inclusæ. Flores in summis ramulorum, paniculati, axillares, sub-globosi, incarnati. Folia terna, glabra, obtusa, trigona, sexfariam, sub-erecta; petiolis brevissimis. Caulis flexuosus, pedalis, glaber. Rami flexuosi, sub-simplices, filiformes, longi.

DESCRIPTION.

TIPS beardless, within the blossom. Flowers grow in loose spikes, near the ends of the branches, from the base of the leaves, are nearly globular, and flesh-coloured. Leaves by threes, smooth, blunt, three-sided, forming six edges, and almost upright; with very short foot-stalks. Stem grows zigzag, a foot high, and smooth. Branches like the stem, grow nearly simple, thread-shaped, and long.

Native of the Cape of Good Hope.

In bloom from August till February.

REFERENCE.

1. The empalement and floral leaves magnified.
2. The chives and pointal.
3. A chive magnified.
4. The seed-bud and pointal magnified.



E. pulchella

by H. Andrews & Knight

ERICA quadriflora.

DESCRIPTIO.

ANTHERÆ cristatæ, inclusæ, stylo incluso. Corolla globosa, purpurea, cernua, glabra. Flores ramulorum quaterni, calyce reflexo. Folia quaterna, linearia, ciliata, flaccida. Caulis debilis, sesquipedalis. Rami plurimi, virgati.

DESCRIPTION.

TIPS crested, within the blossom, shaft included. Flowers globular, purple, nodding, and smooth. Flowers grow by fours, at the ends of the small branches. Cup reflexed. Leaves by fours, linear, fringed and flaccid. Stem weak, a foot and a half high. Branches numerous and twiggy.



Native of the Cape of Good Hope.

In bloom from June till August.

REFERENCE.

1. The empalement magnified.
2. The chives and pointal, summit magnified.
3. A chive magnified.
4. The seed-bud magnified.



ERICA radiata.

DESCRIPTIO.

ANTHERÆ muticæ, inclusæ, stylo exserto. Flores in ramulorum apice verticillati, horizontaliter siti. Corolla sub-cylindræa, oris laciniis revolutis. Folia quaterna, linearia, patentia. Caulis fruticosus, erectus, subpedalis, ramosus. Rami sub-simplices.

DESCRIPTION.

TIPS beardless, within the blossom, shaft without. Flowers grow in whorls at the ends of the small branches, standing horizontally. Blossoms almost cylinder-shaped, the segments of the mouth rolled back. Leaves by fours, linear and spreading. Stem shrubby, upright, near a foot high, and branching. The branches nearly simple.




Native of the Cape of Good Hope.

In bloom from August till November.

REFERENCE.

1. The empalement.
2. The chives and pointal.
3. The seed-bud and pointal, summit magnified.



E. radiata

ERICA rubens.

DESCRIPTIO.

ANTHERÆ cristatæ, inclusæ. Flores terminales, in umbellis nutantibus vel cernuis 3-8-floris. Pedunculi longiusculi. Corolla late ovata, rubro-purpurea. Folia quaterna, erecta, linearia, glabra. Caulis fruticosus, sub-erectus, ramis patentibus, vel subinde pendulis.

DESCRIPTION.


TIPS crested, within the blossom. Flowers terminal, in nodding or cernuous umbels of from 3 to 8 flowers. The peduncles longish. Blossom broad egg-shaped, and of a red purple. Leaves by fours, upright, linear, smooth. Stem shrubby, nearly upright. Branches spreading, or sometimes nodding.

Native of the Cape of Good Hope.

In bloom from June till September.

REFERENCE.

1. The empalement magnified.
2. The chives and pointal, one tip magnified.
3. Seed-bud and pointal, summit magnified.
4. The seed-bud magnified.



E. rubens

W. D. Dore

by F. Anderson

ERICA serratifolia.

DESCRIPTIO.

ANTHERÆ muticæ, sub-exsertæ. Flores terminales, in apicibus ramulorum bini vel terni, pollicares, sub-cylindrici, luteo aurantii. Folia quaterna, rigida, acuta, serrata. Caulis erectus, rigidus, pedalis, ramosus. Ramuli frequentissimi, breves.

DESCRIPTION.

TIPS beardless, just out of the blossom. Flowers terminal, two or three together at the end of the smaller branches, an inch long, almost cylindrical, and of an orange yellow. Leaves by fours, harsh, sharp, and sawed at the edges. Stem upright, harsh, a foot high and much branched. The smaller branches are numerous and short.

Native of the Cape of Good Hope.

In bloom from August till December.

REFERENCE.

1. The empalement and floral leaves magnified.
2. The chives detached from the pointal, summit magnified.
3. Seed-bud and pointal, summit magnified.



ERICA spicata.

DESCRIPTIO.

ANTHERÆ aristatæ, inclusæ. Flores plures, sub-terminales, arcte stipati in spicam duram, ima parte luteoli, summa virescentes, teretiusculi, basi attenuati; ore obtuso, quadrifido, æquali, connivente. Perianthium persistens. Folia subsena, mucronata, patentia. Caulis fruticosus, pedalis, erectus; ramis pluribus adscendentibus.

DESCRIPTION.

TIPS bearded, within the blossom. Flowers numerous, nearly at the ends of the branches, in hard close set spikes, the lower part light yellow, and the extremity of a light green; somewhat cylindrical, with a tapered base; the mouth blunt, and divided into a four-cleft equal border; the segments tend to each other. Cup permanent. Leaves mostly by sixes, sharp-pointed, and spreading. Stem shrubby, a foot high, upright, with numerous ascending branches.

Native of the Cape of Good Hope.

In bloom from November till April.

REFERENCE.

1. The empalement and blossom.
2. The empalement and floral leaves magnified.
3. The chives detached from the pointal, one tip magnified.
4. The seed-bud and pointal, summit magnified.



E. spicata

Nov. 7. 1895

by H. Andrews & K. A. Schott

ERICA tubiflora.

DESCRIPTIO.

ANTHERÆ muticæ, sub-inclusæ. Flores sub-solitarii, sessiles, terminales, clavato-cylindracei, curvati, pubescentes. Folia quaterna, tenuia, obtusa, ciliata, subtus sulcata. Caulis flexibilis, erectus. Rami sparsi, filiformes, frequentes, villosi.

DESCRIPTION.

TIPS beardless, just within the blossom. Flowers grow singly, sessile, at the ends of the branches; between club and cylinder-shaped, curved, downy. Leaves by fours, thin, blunt, fringed with hair at the edges, and furrowed beneath. Stem flexible, upright. Branches scattered, thread-shaped, numerous, hairy.



Native of the Cape of Good Hope.

In bloom from April till July.

REFERENCE.

1. The empalement and floral leaves.
2. The chives detached from the pointal, one tip magnified.
3. The seed-bud and pointal, summit magnified.
4. The seed-bud magnified.



E. tubiflora

ERICA versicolor.

DESCRIPTIO.

ANTHERÆ muticæ, subinclusæ. Corolla sub-cylindræa, medio paululum arctata, ima parte aurantia, apice virescens. Folia terna, linearia, subulata, subtus sulcata, glabra, saturate viridia. Caulis fruticosus, bipedalis, erectus, attenuatus in apicem. Rami subsimplices.

DESCRIPTION.

TIPS beardless, just within the blossom, which is nearly cylindrical, a little narrowed at the middle, the lower part of a deep orange colour, and the mouth greenish. Leaves by threes, linear, awl-shaped, furrowed on the under surface, smooth, and of a deep green. Stem shrubby, two feet high, erect, tapering to the top. The branches nearly simple.

Native of the Cape of Good Hope.

In bloom from October till April.

REFERENCE.

1. The empalement magnified.
2. The chives and pointal.
3. A chive magnified.
4. The seed-bud and pointal, summit magnified.



E. versicolor

ERICA verticillata.

DESCRIPTIO.

ANTHERÆ aristatæ, inclusæ. Flores in apicibus ramorum, verticillati, nutantes, conferti, ventricosi-cylindrici, pollicares, ad basin tetragoni, profunde costati, ore arctato, saturate aurantio. Folia quaterna, glabra, linearia, acuta. Caulis erectus, bipedalis. Rami verticillati.

DESCRIPTION.

TIPS bearded, within the blossom. Flowers grow in whorls at the upper part of the branches, hanging downwards and close together, of a swelled cylinder-shape, an inch long, four-cornered at the base and deeply ribbed. The mouth is narrowed, and of a deep orange colour. Leaves by fours, smooth, linear, and sharp. Stem upright, two feet high. The branches grow in whorls.




Native of the Cape of Good Hope.

In bloom from July till November.

REFERENCE.

1. The empalement and floral leaves.
2. The chives detached from the pointal.
3. Seed-bud and pointal, summit magnified.



E. verticillatus

ERICA viscaria.

DESCRIPTIO.

ANTHERÆ muticæ, inclusæ. Corolla campanulata, valde mucosa, purpurascens. Folia quaterna, linearia, acuta, glabra; juniora in apicibus ramulorum viscosa. Caulis fruticosus, bipedalis, erectus. Rami simplices, longi, superne viscosi.

DESCRIPTION.

TIPS beardless, within the blossoms, which are bell-shaped, very clammy, and purple. Leaves grow by fours, are linear, pointed, and smooth: the younger ones are clammy at the ends of the branches. Stem shrubby, two feet high, and upright. Branches simple, long, and clammy at the ends.

Native of the Cape of Good Hope.

In bloom from April till July.

REFERENCE.

1. The empalement and floral leaves magnified.
2. The blossom.
3. The chives and pointal.
4. A chive magnified.
5. The seed-bud and pointal, summit magnified.



Edm. M.

E. viscaria

ERICA Walkeria.

DESCRIPTIO.

ANTHERÆ muticæ, sub-inclusæ. Flores sessiles, erecti, fastigiati, quaterni, terminales, fere pedunculati. Corolla ventricosa, pellucida, ore arcuato, extus pallide sanguineo, intus albo, laciniis expansis maximis. Folia quaterna, nitida, patentia. Caulis fruticosus, spithameus, erectus, ramulis numerosis.

DESCRIPTION.


TIPS beardless, within the blossom. Flowers sit close in upright bunches by fours, terminal, with scarcely any footstalks. Blossom swelled out and pellucid; the mouth narrowed, of a pale red without and white within; the segments spreading and large. Leaves by fours, shining, and spreading. Stem shrubby, a span high, upright, with numerous little branches.

Native of the Cape of Good Hope.

In bloom from February till June.

REFERENCE.

1. The empalement magnified.
2. A blossom.
3. The chives and pointal, one tip magnified.
4. Seed-bud and pointal magnified.



E. Walkeria

ALPHABETICAL INDEX

TO THE

FIGURES IN VOL. I.

01. ERICA [acuta.](#)
02. [albens.](#)
03. [Archeria.](#)
04. [barbata, major.](#)
05. [β minor.](#)
06. [Bruinades.](#)
07. [caffra.](#)
08. [calycina.](#)
09. [cernua.](#)
10. [comosa.](#)
11. [β rubra.](#)
12. [conspicua.](#)
13. [costata.](#)
14. [cubica.](#)
15. [α minor.](#)
16. [curviflora.](#)
17. [depressa.](#)
18. [droseroides.](#)
19. [empetroides.](#)
20. [exsurgens.](#)
21. [α minor.](#)
22. [filamentosa.](#)
23. [flammea.](#)
24. [gelida.](#)
25. [glauca.](#)

- 26. ignescens.
 - 27. jasminiflora.
 - 28. leucanthera.
 - 29. lutea.
 - 30. Melastoma.
 - 31. nigrita.
 - 32. obbata.
 - 33. obliqua.
 - 34. physodes.
 - 35. pinea.
 - 36. plumosa.
 - 37. pubescens.
 - 38. α minor.
 - 39. β minima.
 - 40. pulchella.
 - 41. quadriflora.
 - 42. radiata.
 - 43. rubens.
 - 44. serratifolia.
 - 45. spicata.
 - 46. tubiflora.
 - 47. versicolor.
 - 48. verticillata.
 - 49. viscaria.
 - 50. Walkeria.
-
-

*** END OF THE PROJECT GUTENBERG EBOOK THE HEATHERY;
OR, A MONOGRAPH OF THE GENUS ERICA. VOL. 1 ***

Updated editions will replace the previous one—the old editions will be renamed.

Creating the works from print editions not protected by U.S. copyright law means that no one owns a United States copyright in these works, so the Foundation (and you!) can copy and distribute it in the United States without permission and without paying copyright royalties. Special rules, set forth in the General Terms of Use part of this license, apply to copying and distributing Project Gutenberg™ electronic works to protect the PROJECT GUTENBERG™ concept and trademark. Project Gutenberg is a registered trademark, and may not be used if you charge for an eBook, except by following the terms of the trademark license, including paying royalties for use of the Project Gutenberg trademark. If you do not charge anything for copies of this eBook, complying with the trademark license is very easy. You may use this eBook for nearly any purpose such as creation of derivative works, reports, performances and research. Project Gutenberg eBooks may be modified and printed and given away—you may do practically ANYTHING in the United States with eBooks not protected by U.S. copyright law. Redistribution is subject to the trademark license, especially commercial redistribution.

START: FULL LICENSE

THE FULL PROJECT GUTENBERG™ LICENSE

PLEASE READ THIS BEFORE YOU DISTRIBUTE OR USE THIS WORK

To protect the Project Gutenberg™ mission of promoting the free distribution of electronic works, by using or distributing this work (or any other work associated in any way with the phrase “Project Gutenberg”), you agree to comply with all the terms of the Full Project Gutenberg License available with this file or online at www.gutenberg.org/license.

Section 1. General Terms of Use and Redistributing Project Gutenberg electronic works

1.A. By reading or using any part of this Project Gutenberg electronic work, you indicate that you have read, understand, agree to and accept all the terms of this license and intellectual property (trademark/copyright) agreement. If you do not agree to abide by all the terms of this agreement, you must cease using and return or destroy all copies of Project Gutenberg electronic works in your possession. If you paid a fee for obtaining a copy of or access to a Project Gutenberg electronic work and you do not agree to be bound by the terms of this agreement, you may obtain a refund from the person or entity to whom you paid the fee as set forth in paragraph 1.E.8.

1.B. “Project Gutenberg” is a registered trademark. It may only be used on or associated in any way with an electronic work by people who agree to be bound by the terms of this agreement. There are a few things that you can do with most Project Gutenberg electronic works even without complying with the full terms of this agreement. See paragraph 1.C below. There are a lot of things you can do with Project Gutenberg electronic works if you follow the terms of this agreement and help preserve free future access to Project Gutenberg electronic works. See paragraph 1.E below.

1.C. The Project Gutenberg Literary Archive Foundation (“the Foundation” or PGLAF), owns a compilation copyright in the collection of Project Gutenberg electronic works. Nearly all the individual works in the collection are in the public domain in the United States. If an individual work is unprotected by copyright law in the United States and you are

located in the United States, we do not claim a right to prevent you from copying, distributing, performing, displaying or creating derivative works based on the work as long as all references to Project Gutenberg are removed. Of course, we hope that you will support the Project Gutenberg mission of promoting free access to electronic works by freely sharing Project Gutenberg works in compliance with the terms of this agreement for keeping the Project Gutenberg name associated with the work. You can easily comply with the terms of this agreement by keeping this work in the same format with its attached full Project Gutenberg License when you share it without charge with others.

1.D. The copyright laws of the place where you are located also govern what you can do with this work. Copyright laws in most countries are in a constant state of change. If you are outside the United States, check the laws of your country in addition to the terms of this agreement before downloading, copying, displaying, performing, distributing or creating derivative works based on this work or any other Project Gutenberg work. The Foundation makes no representations concerning the copyright status of any work in any country other than the United States.

1.E. Unless you have removed all references to Project Gutenberg:

1.E.1. The following sentence, with active links to, or other immediate access to, the full Project Gutenberg License must appear prominently whenever any copy of a Project Gutenberg work (any work on which the phrase “Project Gutenberg” appears, or with which the phrase “Project Gutenberg” is associated) is accessed, displayed, performed, viewed, copied or distributed:

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg™ License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

1.E.2. If an individual Project Gutenberg electronic work is derived from texts not protected by U.S. copyright law (does not contain a notice indicating that it is posted with permission of the copyright holder), the work can be copied and distributed to anyone in the United States without paying any fees or charges. If you are redistributing or providing access to a work with the phrase “Project Gutenberg” associated with or appearing on the work, you must comply either with the requirements of paragraphs 1.E.1 through 1.E.7 or obtain permission for the use of the work and the Project Gutenberg trademark as set forth in paragraphs 1.E.8 or 1.E.9.

1.E.3. If an individual Project Gutenberg electronic work is posted with the permission of the copyright holder, your use and distribution must comply with both paragraphs 1.E.1 through 1.E.7 and any additional terms imposed by the copyright holder. Additional terms will be linked to the Project Gutenberg License for all works posted with the permission of the copyright holder found at the beginning of this work.

1.E.4. Do not unlink or detach or remove the full Project Gutenberg License terms from this work, or any files containing a part of this work or any other work associated with Project Gutenberg.

1.E.5. Do not copy, display, perform, distribute or redistribute this electronic work, or any part of this electronic work, without prominently displaying the sentence set forth in paragraph 1.E.1 with active links or immediate access to the full terms of the Project Gutenberg License.

1.E.6. You may convert to and distribute this work in any binary, compressed, marked up, nonproprietary or proprietary form, including any word processing or hypertext form. However, if you provide access to or distribute copies of a Project Gutenberg work in a format other than “Plain Vanilla ASCII” or other format used in the official version posted on the official Project Gutenberg website (www.gutenberg.org), you must, at no additional cost, fee or expense to the user, provide a copy, a means of exporting a copy, or a means of obtaining a copy upon request, of the work in its original “Plain Vanilla ASCII” or other form. Any alternate format must include the full Project Gutenberg License as specified in paragraph 1.E.1.

1.E.7. Do not charge a fee for access to, viewing, displaying, performing, copying or distributing any Project Gutenberg works unless you comply with paragraph 1.E.8 or 1.E.9.

1.E.8. You may charge a reasonable fee for copies of or providing access to or distributing Project Gutenberg electronic works provided that:

- You pay a royalty fee of 20% of the gross profits you derive from the use of Project Gutenberg works calculated using the method you already use to calculate your applicable taxes. The fee is owed to the owner of the Project Gutenberg trademark, but he has agreed to donate royalties under this paragraph to the Project Gutenberg Literary Archive Foundation. Royalty payments must be paid within 60 days following each date on which you prepare (or are legally required to prepare) your periodic tax returns. Royalty payments should be clearly marked as such and sent to the Project Gutenberg Literary Archive Foundation at the address specified in Section 4, "Information about donations to the Project Gutenberg Literary Archive Foundation."
- You provide a full refund of any money paid by a user who notifies you in writing (or by e-mail) within 30 days of receipt that s/he does not agree to the terms of the full Project Gutenberg™ License. You must require such a user to return or destroy all copies of the works possessed in a physical medium and discontinue all use of and all access to other copies of Project Gutenberg™ works.
- You provide, in accordance with paragraph 1.F.3, a full refund of any money paid for a work or a replacement copy, if a defect in the electronic work is discovered and reported to you within 90 days of receipt of the work.
- You comply with all other terms of this agreement for free distribution of Project Gutenberg™ works.

1.E.9. If you wish to charge a fee or distribute a Project Gutenberg™ electronic work or group of works on different terms than are set forth in this agreement, you must obtain permission in writing from the Project Gutenberg Literary Archive Foundation, the manager of the Project

Gutenberg™ trademark. Contact the Foundation as set forth in Section 3 below.

1.F.

1.F.1. Project Gutenberg volunteers and employees expend considerable effort to identify, do copyright research on, transcribe and proofread works not protected by U.S. copyright law in creating the Project Gutenberg™ collection. Despite these efforts, Project Gutenberg™ electronic works, and the medium on which they may be stored, may contain “Defects,” such as, but not limited to, incomplete, inaccurate or corrupt data, transcription errors, a copyright or other intellectual property infringement, a defective or damaged disk or other medium, a computer virus, or computer codes that damage or cannot be read by your equipment.

1.F.2. LIMITED WARRANTY, DISCLAIMER OF DAMAGES - Except for the “Right of Replacement or Refund” described in paragraph 1.F.3, the Project Gutenberg Literary Archive Foundation, the owner of the Project Gutenberg™ trademark, and any other party distributing a Project Gutenberg™ electronic work under this agreement, disclaim all liability to you for damages, costs and expenses, including legal fees. YOU AGREE THAT YOU HAVE NO REMEDIES FOR NEGLIGENCE, STRICT LIABILITY, BREACH OF WARRANTY OR BREACH OF CONTRACT EXCEPT THOSE PROVIDED IN PARAGRAPH 1.F.3. YOU AGREE THAT THE FOUNDATION, THE TRADEMARK OWNER, AND ANY DISTRIBUTOR UNDER THIS AGREEMENT WILL NOT BE LIABLE TO YOU FOR ACTUAL, DIRECT, INDIRECT, CONSEQUENTIAL, PUNITIVE OR INCIDENTAL DAMAGES EVEN IF YOU GIVE NOTICE OF THE POSSIBILITY OF SUCH DAMAGE.

1.F.3. LIMITED RIGHT OF REPLACEMENT OR REFUND - If you discover a defect in this electronic work within 90 days of receiving it, you can receive a refund of the money (if any) you paid for it by sending a written explanation to the person you received the work from. If you received the work on a physical medium, you must return the medium with your written explanation. The person or entity that provided you with the defective work may elect to provide a replacement copy in lieu of a refund. If you received the work electronically, the person or entity providing it to

you may choose to give you a second opportunity to receive the work electronically in lieu of a refund. If the second copy is also defective, you may demand a refund in writing without further opportunities to fix the problem.

1.F.4. Except for the limited right of replacement or refund set forth in paragraph 1.F.3, this work is provided to you 'AS-IS', WITH NO OTHER WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR ANY PURPOSE.

1.F.5. Some states do not allow disclaimers of certain implied warranties or the exclusion or limitation of certain types of damages. If any disclaimer or limitation set forth in this agreement violates the law of the state applicable to this agreement, the agreement shall be interpreted to make the maximum disclaimer or limitation permitted by the applicable state law. The invalidity or unenforceability of any provision of this agreement shall not void the remaining provisions.

1.F.6. INDEMNITY - You agree to indemnify and hold the Foundation, the trademark owner, any agent or employee of the Foundation, anyone providing copies of Project Gutenberg™ electronic works in accordance with this agreement, and any volunteers associated with the production, promotion and distribution of Project Gutenberg™ electronic works, harmless from all liability, costs and expenses, including legal fees, that arise directly or indirectly from any of the following which you do or cause to occur: (a) distribution of this or any Project Gutenberg work, (b) alteration, modification, or additions or deletions to any Project Gutenberg work, and (c) any Defect you cause.

Section 2. Information about the Mission of Project Gutenberg

Project Gutenberg is synonymous with the free distribution of electronic works in formats readable by the widest variety of computers including obsolete, old, middle-aged and new computers. It exists because of the

efforts of hundreds of volunteers and donations from people in all walks of life.

Volunteers and financial support to provide volunteers with the assistance they need are critical to reaching Project Gutenberg's goals and ensuring that the Project Gutenberg collection will remain freely available for generations to come. In 2001, the Project Gutenberg Literary Archive Foundation was created to provide a secure and permanent future for Project Gutenberg and future generations. To learn more about the Project Gutenberg Literary Archive Foundation and how your efforts and donations can help, see Sections 3 and 4 and the Foundation information page at www.gutenberg.org.

Section 3. Information about the Project Gutenberg Literary Archive Foundation

The Project Gutenberg Literary Archive Foundation is a non-profit 501(c)(3) educational corporation organized under the laws of the state of Mississippi and granted tax exempt status by the Internal Revenue Service. The Foundation's EIN or federal tax identification number is 64-6221541. Contributions to the Project Gutenberg Literary Archive Foundation are tax deductible to the full extent permitted by U.S. federal laws and your state's laws.

The Foundation's business office is located at 41 Watchung Plaza #516, Montclair NJ 07042, USA, +1 (862) 621-9288. Email contact links and up to date contact information can be found at the Foundation's website and official page at www.gutenberg.org/contact

Section 4. Information about Donations to the Project Gutenberg Literary Archive Foundation

Project Gutenberg™ depends upon and cannot survive without widespread public support and donations to carry out its mission of increasing the number of public domain and licensed works that can be freely distributed in machine-readable form accessible by the widest array of equipment

including outdated equipment. Many small donations (\$1 to \$5,000) are particularly important to maintaining tax exempt status with the IRS.

The Foundation is committed to complying with the laws regulating charities and charitable donations in all 50 states of the United States. Compliance requirements are not uniform and it takes a considerable effort, much paperwork and many fees to meet and keep up with these requirements. We do not solicit donations in locations where we have not received written confirmation of compliance. To SEND DONATIONS or determine the status of compliance for any particular state visit www.gutenberg.org/donate.

While we cannot and do not solicit contributions from states where we have not met the solicitation requirements, we know of no prohibition against accepting unsolicited donations from donors in such states who approach us with offers to donate.

International donations are gratefully accepted, but we cannot make any statements concerning tax treatment of donations received from outside the United States. U.S. laws alone swamp our small staff.

Please check the Project Gutenberg web pages for current donation methods and addresses. Donations are accepted in a number of other ways including checks, online payments and credit card donations. To donate, please visit: www.gutenberg.org/donate.

Section 5. General Information About Project Gutenberg electronic works

Professor Michael S. Hart was the originator of the Project Gutenberg concept of a library of electronic works that could be freely shared with anyone. For forty years, he produced and distributed Project Gutenberg eBooks with only a loose network of volunteer support.

Project Gutenberg eBooks are often created from several printed editions, all of which are confirmed as not protected by copyright in the U.S. unless a

copyright notice is included. Thus, we do not necessarily keep eBooks in compliance with any particular paper edition.

Most people start at our website which has the main PG search facility:
www.gutenberg.org.

This website includes information about Project Gutenberg, including how to make donations to the Project Gutenberg Literary Archive Foundation, how to help produce our new eBooks, and how to subscribe to our email newsletter to hear about new eBooks.

THE
HEATHERY;
OR A
MONOGRAPH
OF
THE GENUS ERICA:
VOL. I.



[back](#)



[back](#)




E. albens

Chapm. Bot. Diarctia 1845.

14. E. albens 5. Knightbridge

[back](#)

[back](#)



1



2



3



4

E. barbata. major

1820. E. barbata. major

1820. E. barbata. major

[back](#)



E. barbata minor

Wm. Pursh, Jun. Sculp.

Wm. Woodcock Sculp.

[back](#)



E. Bruniades

The New York Botanical Garden

by H. Andrews & J. D. Hooker

[back](#)



E. castra

[back](#)



E. calycina

Bot. Novartis

ex. Hb. Andromed. & Knightbridge

[back](#)




det. D. B. S. 1844

. *E. cernua*

by H. Andrews & J. Hughes 1844

[back](#)



St. Dorothea 1784.

by H. Andrews & Knightbridge

[back](#)



E. Comosa rubra

[back](#)



as the old records have it

E. conspicua.

by H.G. Andrews, 1847

[back](#)




1



2



3



4

E. costata

[back](#)




E. cubica

at Dordrecht April 1793

by J. G. Lindley del.

[back](#)



E. culicis minor

[back](#)



E. curviflora

[back](#)



1



2




3

[back](#)



E. droseroides

[back](#)



E. empetroides

[back](#)




Not direct. v. l. l. l.

E. exurgens.

by F. C. Hermann. T. J. G. v. d. G.

[back](#)



E. exurgens. minor

[back](#)





[back](#)



E. flammea

[back](#)



E. gelida

by H. Andrews & Knight

[back](#)



E. glauca

W. A. Anderson

[back](#)



E. ignescens

Wm. B. Wood, 1848

by H. R. Anderson & W. B. Wood

[back](#)




at Dorset Oct. 1, 1844

E. jasmineflora.

by H. C. Andrews & Knightbridge

[back](#)



E. leucanthera


Linnaea, Nov. 1753

by J. B. deCandolle & J. G. deCandolle

[back](#)



[back](#)



E. melastoma

by H. Anderson & Son

[back](#)



E. nigrita

Prodr. 1845


1845

[back](#)



E. obtata

[back](#)



E. obliqua.

[back](#)



E. physodes

[back](#)



E. pinea

[back](#)



[back](#)



E. pubescens

by H. Andrews, sculp.

[back](#)




E. pubescens minor.

Nov. 1868

by H. Andrews

[back](#)



E. pubescens minima

[back](#)



E. pulchella

by H. Andrews & Houghton

[back](#)




[back](#)



E. radiata

[back](#)




E. rubens

Ad. Directa

by T. Anderson, sc.

[back](#)



E. serratifolia

Feb. 1805

by J. B. Andrews & H. Knight

[back](#)



E. spicata

1897. 7. 1907



by H. Andromeda. H. Andromeda.

[back](#)



E. tubiflora


[back](#)



E. versicolor

by H. Andrews's Florist

[back](#)



E. verticillata


[back](#)



Ed. n. 10

E. viscaria

[back](#)



E. Walkeria

[back](#)