

ON
SULPHONFLUORESCEIN
AND SOME OF ITS DERIVATIVES.

DISSERTATION
Presented for the Degree
of
Doctor of Philosophy.
at the
Johns Hopkins University

by
C. Willard Hayes.

1887.

ON
SULPHONFLUORESCEIN
AND SOME OF ITS DERIVATIVES.

DISSERTATION
Presented for the Degree
of
Doctor of Philosophy.
at the
Johns Hopkins University

by
C. Willard Hayes.

1887.

The Project Gutenberg eBook of On Sulphonfluoresceïn and Some of Its Derivatives

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

Title: On Sulphonfluoresceïn and Some of Its Derivatives

Author: C. W. Hayes

Release date: November 22, 2015 [eBook #50531]

Most recently updated: October 22, 2024

Language: English

Other information and formats: www.gutenberg.org/ebooks/50531

Credits: Produced by Paul Marshall and the Online Distributed Proofreading Team at <http://www.pgdp.net> (This file was produced from images generously made available by The Internet Archive)

*** START OF THE PROJECT GUTENBERG EBOOK ON
SULPHONFLUORESCẺİN AND SOME OF ITS DERIVATIVES ***

ON SULPHONFLUORESCEIN AND SOME OF ITS DERIVATIVES.

DISSERTATION

Presented for the Degree
of
Doctor of Philosophy
at the
Johns Hopkins University

by
C. Willard Hayes,

1887.

CONTENTS

	Introduction	p. 3.
I	Ortho-sulpho-benzoic acid.	
	Preparation from toluene and H_2SO_4	7.
	" " para-nitro-toluene and H_2SO_4	8.
	Oxidation of toluene - o - sodium sulphonate	12.
	Preparation of Sulphinide from " "	14.
	" " " by chlorsulphonic acid.	17.
	Conversion of sulphinide into o-sulphobenzoic acid.	25.
	Analysis of and crystallography of " " "	28.

II	Sulphonfluorescein—previous attempts	<u>29.</u>
	Preparation and purification of s-fluorescein	<u>30.</u>
	Analyses of	<u>34.</u>
	Properties of	<u>35.</u>
	Salts. Ba. and Ca. of	<u>37.</u>
	Acetyl derivative of	<u>41.</u>
	Br substitution products of	<u>42.</u>
	Action of H ₂ SO ₄ on	<u>44.</u>
	Action of HCl on	<u>45.</u>
	Reduction of	<u>46.</u>
	Conclusion.	<u>47.</u>

On Sulphonfluorescein and some of its Derivatives.

Introduction.

The close analogy in composition and structure between phthalic acid and ortho-sulpho-benzoic acid suggests the possibility of obtaining from the latter, by its action on phenols, substances analogous to the phthaleïns. If such compounds could be made they would afford a favorable opportunity of studying the effects produced in the properties of a compound by the replacement of CO by SO₂. It was with a view to such study that the following discussed work was undertaken at the suggestion of Prof. Remsen and carried on under his constant guidance.

Some experiments previously performed by Remsen and Palmer (A.G.) indicated the possibility of the formation of a fluorescent substance by the action of ortho-sulpho-benzoic acid on resorcin but they did not succeed in obtaining any definite crystallized compound from the reaction.

The chief obstacle to be overcome in the work is the difficulty in obtaining the o-sulpho-benzoic acid and a large proportion of the work here described was applied in that direction.

Ortho-sulpho-benzoic acid: Methods for its preparation.

1. From toluene and H_2SO_4 .

A method employed by Remsen and Fahlberg (Am. Ch. Jour. Vol. 1. p ___) for getting the sulphonic acid group in the ortho position to methyl was; (a) treat toluene with fuming H_2SO_4 forming thus ortho- and para-toluene sulphonic acids, (b) make the calcium salt of the sulphonic acids thus formed and from this the potassium salt. (c) treat this mixture of potassium toluene sulphonates with phosphorous penta-chloride forming the corresponding sulphonchlorides. One of these (para) being a solid and the other (ortho) an oily liquid a nearly complete separation could be effected. The difficulty with this method is however that the larger part of the product is the para and only a comparatively small proportion of the ortho compound is formed.

2. From p-nitro-toluene and H_2SO_4 .

A second method employed consists in starting with p-nitro-toluene. This when treated with H_2SO_4 forms toluene p-nitro o-sulphonic acid. If now a method could be obtained for removing the nitro group the desired result would be attained.


The attempt was made by Remsen and Palmer (A.G.) to accomplish this by (a) reducing the nitro compound to the amide, (b) making the diazo compound and (c) boiling this with absolute alcohol. According to generally accepted views this should effect the removal of the diazo group and its replacement by hydrogen.

Experiments however showed that the replacement was made not by hydrogen but by the ethoxy group $-\text{OC}_2\text{H}_5$. This method was therefore impracticable.

A modification of this method was suggested by an observation of Baeyer and Liebermann that if phenyl hydrazine be boiled with a dilute solution of copper sulphate the hydrazine group is replaced by hydrogen and benzene thus formed. Hence it was believed that if the hydrazine compound should be made from diazo compound mentioned above, the corresponding hydrocarbon, i.e. toluene o-sulphonic acid could be obtained. The results of experiments showed that this afforded a practicable method of preparing toluene ortho-sulphonic acid.

After experimenting with various modifications of the method the following was found to be the best adapted to the purpose.


The potassium salt of toluene p-nitro-o-sulphonic acid is easily obtained, as already stated, by heating p-nitro toluene on the water bath with three times its weight of fuming H_2SO_4 , neutralizing with chalk and to the solution of calcium salt thus obtained adding a slight excess of K_2CO_3 . On filtering from the precipitated CaCO_3 and evaporating slightly, the salt is obtained in long needle shaped crystals of a pale straw yellow color. This is



The reduction of the nitro group is best effected by means of tin and HCl, in the proportion, salt 5 parts, tin 6 parts and concentrated HCl 30 parts.

The amido acid forms a compound with tin which crystallizes from the HCl together with stannous chloride. This compound may be broken up and the tin removed by continued boiling with water.


A better method of removing the tin is by dissolving the compound in Na_2CO_3 . This forms a salt with the amido acid and throws down the tin as $\text{Sn}(\text{OH})_2$, a white flocculent precipitate. On filtering and adding to the solution conc. HCl, the free amido acid is deposited in characteristic colorless, rhombic crystals, having the formula







The method at first employed for preparing the hydrazine compound consisted in treating the amido acid, suspended in HCl, with potassium nitrite and then with stannous chloride. The tin was then removed from the solution by the addition of sodium carbonate and the hydrazine compound thrown down with HCl. This method however gave poor results the yield being only about 50% of the theoretical.



Another method was accordingly substituted for the above, namely that of Strecker and Römer (Ber. IV. s 784.) By this the diazo compound is made first and isolated. This is done by suspending the finely powdered acid in absolute alcohol, cooling and passing a current of the oxides of nitrogen through in the ordinary way. The acid changes in appearance, becoming more crystalline and slightly darker and settles quickly on being shaken. The reaction here may be expressed thus—



When the reaction is completed as shown by the appearance of the suspended powder it is filtered and while still fresh is added to a solution of acid sodium sulphite as long as it continues to dissolve readily.


To this solution there is added a quantity of solution of acid sodium sulphite equivalent to that already used and the solution is then boiled. It has at first a deep red color but in a few moments becomes light reddish yellow. The reaction of HNaSO_3 on the diazo compound may be represented in two stages, the first portion forming an addition product and the second acting as a reducing agent. Thus,






To the hot solution an excess of conc. HCl is added when the hydrazine compound separates in a few moments in lustrous yellow scales which completely fill the solution. On the addition of the HCl a large amount of SO_2 is given off from the excess of HNaSO_3 and the solution becomes deep red. When the hydrazine has separated the mother liquor is again yellow.

The reaction is represented as follows:



The yield of hydrazine when both the diazo and the NaHSO_3 are freshly prepared is practically quantitative.


The hydrazine thus prepared was treated with a hot 10% solution of copper sulphate till a permanent blue color was obtained in the solution. Nitrogen is evolved and the copper sulphate is reduced to cuprous oxide which is precipitated as a red powder. The reaction is as follows.



Chalk was added to the solution to precipitate the H_2SO_4 and form a calcium salt of toluene-o-sulphonic acid. From this the sodium salt was made by adding a slight excess of Na_2SO_3 and evaporating to dryness. The salt is very soluble being deliquescent in the air while the corresponding potassium salt is not. From 1538 gr. of para-nitro-toluene, 655 gr. of toluene ortho-sodium sulphonate were obtained.


Having thus obtained the toluene ortho-sulphonic acid the next step in the problem was to find a convenient method for converting this into ortho-sulph-benzoic acid. Two ways present themselves for accomplishing this end. (1) direct oxidation of this salt and (2) conversion into benzoic sulphinide from which the acid may be obtained. Both of these methods were tried.

Oxidation of toluene-o-sodium sulphonate.



The sodium salt of toluene-o-sulphonic acid is oxidized to ortho-sulphobenzoic acid with considerable difficulty by KMnO_4 in neutral solution.

Thus two experiments showed that the oxidation was not complete after 24 hours boiling with excess of permanganate. If the solution be made alkaline however, the oxidation is completed in a few hours, yet the greatest difficulty still remains in the separation of the free acid from the products of oxidation in the solution. If HCl be added to the solution the acid salt




is formed and this has nearly the same solubility as the KCl also present. A better method therefore is to add a slight excess of H_2SO_4 and evaporate nearly to dryness. In this way are formed sulphates and the free acid presumably. The mixture is heated with alcohol (95%) which extracts the acid leaving the greater part of the manganese salts. This extract is evaporated and reextracted with alcohol. To this solution BaCO_3 is added to precipitate the H_2SO_4 and form the Barium salt of the o-sulphobenzoic acid. The solution is filtered from the BaSO_4 and just enough H_2SO_4 is added to exactly precipitate the barium. The solution should thus contain only the free acid sought, which crystallizes out on evaporating to a small volume. While the method is theoretically possible it presents so many difficulties that it is practically useless. The yield is extremely small; only enough acid being obtained in this way to show that it was possible.


Formation of Sulphinide from toluene-o-sodium sulphonate.

The second method for obtaining free o-sulphobenzoic acid from toluene-ortho-sulphonic acid is by the conversion of the latter first into benzoic sulphinide and then into the free acid. The sulphinide was made essentially as described by Remsen (Am. Ch. Jour. Vol. I. p. 428) with a few changes in the details as follows.


The salt



finely pulverized and in portions of from 10 to 50 gr. was placed in a Florence flask; an equivalent quantity of PCl_5 added; An inverted condenser was then attached and the flask shaken. The action takes place at once and involves sufficient heat to distill off the oxychloride formed in the reaction. This being returned to the flask by the condenser furnishes a liquid medium in which the reaction takes place more readily and completely than when it is not present. It is best to cool the flask at first and afterwards heat gently on the water bath. The reaction which takes place may be represented as follows.



On the addition of water the chloride separates as a light yellow oil. This is washed with water and concentrated aqueous ammonia added, which forms toluene-o-sulphonamide thus.



The reaction is accompanied by a slight evolution of heat and the formation, apparently, of an intermediate product having a yellowish color, which passes over on longer standing into the white amide. After standing several hours the excess of ammonia was driven off by very gentle heating

on the water bath. If the heat is too high a large amount of a tarry product is formed and the yield of amide is correspondingly small. In any case some of this tarry product is formed. When nearly all the ammonia had been driven off the mass was boiled with water which dissolves everything except the tar. The hot solution was filtered through charcoal and on cooling the amide separated in white feathery crystals which melt at 155°-156°.

The amide thus obtained was oxidized as described by Remsen (loc. cit.) with potassium permanganate in neutral solution. The proportions are 10 gr amide. 40 gr KMnO_4 and 1 L. water. The oxidation was usually effected in from four to six hours.

To obtain the sulphinide from this solution after oxidation, the latter, after filtration from the precipitated oxides of manganese, was slightly acidified with HCl and evaporated to about one fourth its original volume. On the addition of concentrated HCl to this solution, the sulphinide separated out in white or slightly yellowish feather shaped crystals melting at 212° and having the characteristic intensely sweet taste.

Formation of Sulphinide from Toluene by means of the chlorsulphonic acid reaction.

Before passing on to the methods used for converting the sulphinide into free acid another method should be described by which the former was obtained in larger quantities and much more easily than by the one above described.

Beckurts and Otto (Ber. XI. 2061) found that by treatment of toluene with sulphuryl hydroxy-chloride or chlorsulphonic acid, ClSO_2OH , both o- and p- and as they supposed also m-toluene sulphonchlorides were formed together with the corresponding sulphonic acid.

Claesson and Wallin (Ber. XII. p. 1848) repeated the work reaching practically the same results and finally Noyes (Am. Ch. Jour. Vol. VIII. p. 176) employed the reaction as a convenient method for obtaining toluene o-sulphon-chloride.

Chlorsulphonic acid is made by passing dry HCl over solid sulphuric acid so long as it continues to be absorbed. Since no solid sulphuric acid

was at hand, ordinary fuming Nordhausen acid was taken and from one of two equal portions the SO_3 was driven over into the other. HCl was passed into the latter and the resulting chlorsulphonic acid distilled off at about 156° .


This was placed in a flask, provided with a drop funnel and exit tube, in portions of 150 gr. and to each portion 60 gr. of toluene was added, very slowly, with constant shaking, the temperature being kept near 10° . The action is violent and if any toluene is allowed to collect on the surface of the liquid it is apt to produce disastrous results. Large quantities of HCl are given off and the liquid in the flask assumes a brown color. When all the toluene has been added, it is poured into a large quantity of ice water, when the sulphon-chlorides separate out, the ortho- as a heavy oil and the para- as a white crystalline solid. After allowing to stand some time in order that as much of the para-chloride might crystallize as possible the ortho- was drawn off and subjected to a freezing temperature for several hours. By this means more of the p-chloride was removed and the operation was replicated as long as any crystals continued to form, generally two or three times. In this way the greater part of the para- may be removed, though some still remains dissolved in the liquid chloride, which cannot be removed by repeated freezings.

The chloride thus obtained was treated with strong aqueous ammonia. The conversion to the sulphamide does not take place so readily as in case of the pure o-chloride obtained from the sulphonic acid and phosphorus pentachloride.


After standing about two days the whole of the oily chloride had solidified to a yellowish brown mass. The excess of ammonia was driven off by gentle heating on the water bath and the mass then boiled with water. Not enough water was added at first for complete solution but when the first portion was saturated it was poured off through a filter and from it the amide separated in yellowish feathery crystals which melted at 105° - 125° and consisted therefore as shown by Fahlberg (Am. Ch. Jour. Vol. I. p. 170) of a mixture of o- and p-sulphonamides. It was recrystallized and from it was obtained a portion melting at 153° - 5° and one at 108° - 20° .

Since this mixture cannot be completely separated by recrystallization another method was suggested. Remsen has shown that $\text{K}_2\text{Cr}_2\text{O}_7$ in acid

solution does not oxidize the methyl group in



but does oxidize that in



It was thought that in a mixture of the two the former might be left unchanged while the latter was oxidized to p-sulphamine benzoic acid.

To test this 15 gr of the mixture, melting at 105°-125°, was heated with 40 gr K₂Cr₂O₇, 55 gr H₂SO₄ and 2 vols. of water for about two hours. It was then tested and shown to be still a mixture of p- & o-amides, since it was again heated for several hours with half the original quantity of oxidizing mixture, then diluted, filtered and washed. The white crystalline residue was treated with sodium carbonate to dissolve the benzoic para-sulphamide and the residue was found to be pure toluene o-sulphonamide melting at 153°-155°. The small quantity remaining, 3 gr., indicated that part of the o-amide had been completely broken down by the strong oxidizing agent, though the proportion of o- & p-amides in the original mixtures was known only approximately. The evolution of gas during the oxidation would point to the same conclusion.

Although this effects a complete separation it is hardly economical since it will be shown later that a separation can be conveniently effected after the oxidation with KMnO₄ so that the o-amide contained in the mixture need not be lost.

The original mass was treated with successive portions of water till nothing remained but a black tarry substance. The amide which separated from these extracts was perfectly white and melted at 153-5°. It was

therefore regarded as practically pure o-amide. The yield in amide melting above 153° was a little over one sixth the weight of toluene used.

The amide obtained in this way was oxidized in the manner already described. It was found however that there was always some benzoic p-sulphamide in the solution of the oxidation, due to the slight admixture of p- with the o-amide used. This is thrown down with the sulphinide on acidifying the solution and may be removed by re-crystallization since it is somewhat less soluble in hot and cold water than sulphinide.

A better way to effect the separation, however, was found to be the following. After having evaporated the solution containing the products of oxidation, nearly neutralized with HCl, to about one fifth its original volume, it is made very slightly acid and allowed to cool. In this way very nearly all the benzoic p-sulphamide is separated from the solution and none of the sulphinide. After filtering, strong HCl is added and the sulphinide then separates in its characteristic form. This indicates that sulphinide forms an alkaline salt which is not decomposed by diluted HCl while the p-sulphamide does not.

The mixture of amides meeting at 105° - 120° was oxidized and the products separated in this way gave about equal quantities of sulphinide and benzoic p-sulphamide.


When toluene is treated with chlorsulphonic acid there are formed besides the ortho- and para- chlorides also ortho and para sulphonic acids. These of course are in solution in the water from which the chlorides separated. In order to recover the ortho-acid, the solution was neutralized with chalk forming the calcium salt: this converted into the potassium salt which by evaporating the solution to dryness was obtained as a white crystalline powder. When treated with PCl_5 in the manner already described this gave a mixture of ortho and para sulphonchlorides consisting of about $\frac{1}{3}$ ortho and $\frac{2}{3}$ para.

Formation of Orthosulphobenzoic acid from Sulphinide.

Benzoic sulphinide may be converted into a sulpho-benzoic acid (1) by boiling with $\text{Ba}(\text{OH})_2$, (2) by heating in a closed tube with conc. HCl or (3) by evaporating on the water bath with HCl.

1. Three gramms of sulphinide were boiled in a flask connected with an inverted condenser for about two days with an excess of Ba(OH)_2 . There was formed in the flask a hard mineral-like mass which was insoluble in water and cold diluted HCl but dissolved in hot HCl with effervescence. This was a Barium salt, probably basic (?) of ortho sulphobenzoic acid. There was also formed an easily soluble barium salt of that acid. The former was dissolved in H_2SO_4 and treated with BaCO_3 ; the filtrate from the BaSO_4 which contained a soluble barium salt was added to that above mentioned and the barium exactly precipitated with H_2SO_4 and the filtrate evaporated to dryness giving the free acid but not in a perfectly pure condition.

2. 2.75 gr. of sulphinide was sealed up in a tube with pure conc. HCl and heated two hours to 150° . On cooling nothing separated; the liquid was evaporated to dryness giving 3.2 gr of acid and ammonium chloride. The reaction taking place here may be represented thus:





3. A more convenient method for obtaining the acid than either of the above, is to heat the sulphinide with conc. HCl on the water bath for two days. Then evaporate to dryness and dissolve the residue in a small quantity of water. If the sulphinide contained any para-sulphamide, as is usually the case, this will be left undissolved and most of the NH_4Cl will crystallize on standing. This solution by slow evaporation deposits large colorless crystals of the free acid.

This acid is soluble in about two parts of cold water, very difficultly soluble in absolute alcohol and almost completely insoluble in ether. It does not melt under 250° but considerably above that it melts, at first apparently without change and then with slight sublimation of a very deliquescent substance, probably the anhydride.

Two determinations of the S. made by Mr. A. F. Linn, gave the following results.

I	·1358	gr substance	gave	·1855	gr BaSO ₄	representing	15·72%
II	”	”	”	”	”	”	”

Calculated for the formula



A crystallographic examination of the acid showed it to belong to the orthorhombic system. Axial ratio: $a : b : c = \cdot 8507 : 1 : \cdot 8121$. Planes. P and $\alpha P \dot{\alpha}$.

Angles measured	{	$P \wedge P$	Edge X =
			$131^\circ 8'$
			” Y =
			$82^\circ 18'$
	{	$P \wedge \alpha P \dot{\alpha}$	” Z =
			$118^\circ 40'$
			=
			$114^\circ 38'$


The crystals are up to 8 mm in length.
The pyramidal faces are generally etched so that the image is poor.

Sulphonfluoresceïn.

Several attempts had already been made to obtain from the action of o-sulphobenzoic acid and resorcin a substance analogous to the fluoresceïn obtained by Baeyer^[1] from phthalic anhydride and resorcin but while a strongly fluorescent substance was easily obtained, no definite compound could be separated from it. Thus Palmer obtained, by heating together the above named substances to 160°(?) a solid mass, part soluble in water and part insoluble as a dark brown amorphous powder. Both parts gave a strong fluorescence with alkalis. He was unable however to obtain the substance itself or any derivative in a crystalline form.

The first experiments in this series gave the same negative results. The mixture of acid and resorcin was heated in a sulphuric acid both for several hours to 150°-170°, and as it showed no sign of solidification the temperature was raised to 200° and then to 235° where it was kept several hours longer. The black viscous mass obtained in this way became vitreous on cooling, and in all respects resembled that described by Palmer. This however is not the normal course of the reaction as shown later but is probably due to a decomposition of the normal product produced by too high heating.

An experiment was made with the acid salt of ortho-sulphobenzoic acid.



resorcin and H₂SO₄ heating the mixture to 150°-170°. A solid black mass was obtained strongly fluorescent in alkaline solution and in all other respects like the substance obtained above.

Preparation of Sulphonfluoresceïn.

As the result of a number of experiments the following method of preparing and purifying the sulphonfluorescein was found to give the best results.

The free acid is mixed with resorcin in the proportion of 1 part acid to 1.2 parts resorcin, or a slight excess over two molecules of the latter to one of the former. The mixture is placed in a deep vessel, a test tube or beaker, which is placed in a sulphuric acid bath, a thermometer being suspended in the mixture. The bath is heated and at about 100° the resorcin melts and the acid slowly dissolves in the liquid. No action appears to take place till the temperature reaches 175° where water begins to be given off and the liquid slowly assumes a darker color. White crystals of resorcin collect on the sides of the vessel. After the heating has been continued for about seven hours at 178° - 185° the liquid has a clear deep red color but shows no signs of becoming viscous. At length lustrous yellow crystalline plates appear in the liquid and soon the whole mass becomes a thick nearly solid yellow paste. Continuous heating at the same temperature causes no further apparent change. This mass, which on cooling is made up of yellow crystals imbedded in a red vitreous matrix, is then dissolved in hot water forming a clear red solution or at most one containing but a small quantity of a brown flocculent precipitate. This solution is filtered if necessary and evaporated to a small volume from which the substance separates on cooling in reddish yellow radial crystals. These are filtered and washed with ether till the washings are perfectly colorless. The substance on repeated crystallization from water has a pale straw yellow color and when deposited slowly forms transparent crystal blocks from 2·6 mm long, arranged in radial groups.

A considerable amount of resorcin is lost by sublimation during the reaction especially if the operation is carried on in a beaker so that some excess should be added. But even when the resorcin is present in excess at the end of the reaction some free acid is always left which may be obtained from the mother liquid in the characteristic colorless orthorhombic crystals.

The water of crystallization and sulphur were determined in the new compound. In estimating the water, the substance was heated to 106° - 123° for about ten hours till it attained a constant weight. On standing in the air it quickly regains its original weight. When heated to 130° - 140° for some time it turns slightly reddish and loses over 10% of its weight which is not regained by standing in the air.

Estimation No. I. was made by fusion with KOH and KNO₃. Nos. II and III were made by Mr. Mindileff by Morse's method, oxidizing with KMnO₄ in KOH solution.


·3882 gr. heated to 106°-123° lost ·0302 gr. = 8·5%
 Calculation for 2H₂O - C₁₉H₁₂O₆S + 2H₂O = 8·9% water.

I. ·2007 gr sub. gave ·1286 gr BaSO₄ = 8·77% S.
 II. gr " " " gr " = 7·91" "
 III. gr " " " gr " = 7·89" "


Calculation for C₁₉H₁₂O₆S + 2H₂O = 7·92% S.

These analyses show with but little doubt that the substance has the composition indicated above i.e. C₁₉H₁₂O₆S + 2H₂O. The reaction therefore which takes place between ortho-sulpho benzoic acid and resorcin from its analogy to that taking place between phthalic anhydride and resorcin may be represented thus, as shown by Baeyer in his second paper (Ann. 202. S. 43)


Representing the formation of the anhydride as the first action.



and the action of resorcin on this anhydride thus.



The substance thus formed would naturally receive the name Sulphonfluoresceïn from its analogy with Fluoresceïn.



Properties of S.fluoresceïn.

This compound shows a marked similarity to the fluoresceïn described by Baeyer as would naturally be expected from its great similarity of composition and constitution, but it also shows decided differences which may be attributed to the replacement of CO by SO₂.

Dissolved in water it shows a weak green fluorescence which in alkaline solution becomes much deeper but not by any means so strong as that of fluoresceïn. The dilute alkaline solution by transmitted light is almost perfectly colorless and by reflected light a clear green. Unlike fluoresceïn it is extremely soluble in water, about one part in two or three of hot and five or six of cold water. It is also soluble in absolute alcohol forming a yellow solution with weak fluorescence. It is soluble with difficulty in ether but when in solution is deposited only on evaporating to a small volume.

It does not melt at 250° but if held at a lower temperature for a long time becomes red undergoing some decomposition. If quickly heated somewhat above 300° it melts to a deep red liquid and then solidifies. If the mass is treated with water it partly dissolves leaving a dark brown flocculent precipitate which dissolves on the addition of an alkali, the

solution having an intense fluorescence, nearly if not quite equaling that of fluoresceïn. This change produced by heating was not further studied.

The crystals are very thin blades, apparently monoclinic, showing the clinopinacoid $\alpha P\alpha$ and a very narrow prism αP and clinodome $P\alpha$. The angle $\beta = 75^\circ$ and the extinction angle against the c axis = 20° . The axial ratio could not be accurately determined.

Salts of sulphonfluoresceïn.

The influence of the SO_2 group is shown by the fact that the substance acts as an acid decomposing carbonates and forming salts which is not the case with fluoresceïn.

Barium salt.

The substance was boiled with an excess of carefully purified $BaCO_3$ for two hours. The filtrate from the $BaCO_3$ evaporated to a small volume deposited yellow crystals resembling the original substance but in shorter and thicker prisms. These were twice recrystallized and had then a light straw yellow color.

A determination of the Ba gave the following results. The salt was dried in the air.

Transcriber's Note:

The following table was crossed out on the original. A note on the previous page beside the table was:

“All these calculations are wrong. J.R.”

I	·1078	gr salt	gave	·0304	gr BaSO ₄	= 15·73%	Ba.
II	·1641	" "	" "	·0457	" "	= 15·53"	" "
III	·2425	" "	" "	·0680	" "	= 15·65"	" "
IV	·2860	" "	" "	·0798	" "	= 15·54"	" "
V	·1843	" "	" "	·0498	" "	= 15·08"	" "
VI	·2620	" "	" "	·0708	" "	= 15·08"	" "
VII	·3230	" "	" "	·0906	" "	= 15·65"	" "
VIII	·2875	" "	" "	·0807	" "	= 15·66"	" "



Calculated for C₁₉H₁₃O₇SBa = 15·10% Ba.


In the above determinations the salt analysed was taken from specimens made at three different times and purified in slightly different ways, Nos 1, 2, & 3 being washed with absolute alcohol. Nos V and V were made by precipitating the Ba with H₂SO₄ from a solution of the salt.

The water was determined by heating at 110° till constant weight was reached. Part only of the weight lost was regained on standing in the air.

·3943 gr salt lost at 110° ·0286 gr = 7.25%
 Water calculated for C₁₉H₁₃O₇SBa+2H₂O = 7.35%

Although these analyses show a per cent. of Ba somewhat above that required by a compound having the formula C₁₉H₁₃O₇SBa still this appears to be the most probable formula which can be assigned to the substance. If this is the true composition of the salt, then in sulphonfluoresceïn the anhydride condition must be broken up by boiling with BaCO₃ forming the salt thus.





By treating the salt with H_2SO_4 the original substance is reformed.

Calcium Salt.

Attempts were made to prepare the calcium salt but without success. The S-fluoresceïn was boiled several hours with very finely powdered calcite, and some salt was formed as shown by the CO_2 evolved but on evaporating the solution and recrystallizing the substance deposited it was found to be the unchanged S-fluoresceïn. Some Ca. salt was in the mother liquors but its extreme solubility prevented a separation being made.

Acetyl derivative of S.fluoresceïn.

S.fluoresceïn was boiled with an excess of acetic anhydride for about three hours. The solution became quite dark and when evaporated on the water bath left a black tarry residue. This was treated with water which dissolved a part leaving a dark flocculent precipitate. The solution was

boiled with animal charcoal and evaporated nearly to dryness. On cooling there separated a light yellow flocculent precipitate very soluble in hot water and but slightly less so in cold. This was dissolved in a small quantity of alcohol from which it separated on evaporation in small radial crystals having a light lavender color & satiny luster. They also have a peculiar odor resembling slippery elm which is not removed by recrystallization. They show a tendency to decompose, becoming yellow on exposure to the air. The substance does not melt or change in appearance under 245° . With alkalis it gives a slight greenish fluorescence. From the method of its formation this was taken to be an acetyl derivative of S.fluoresceïn but whether the mono-or di-acetyl could not be determined without analysis for which the substance did not suffice.

Bromine substitution products of S-fluoresceïn.

It was especially interesting to see what influence the SO_2 group would exert upon the introduction of Bromine into the compound. In the case of fluoresceïn four Bromine atoms enter easily and special precautions are necessary to obtain a product containing a smaller number. The case however is different with S.fluoresceïn.

The latter was dissolved in glacial acetic acid in which it is soluble with some difficulty and to the solution was added a 20% solution of bromine in acetic acid, in sufficient quantity to make eight atoms of bromine to one molecule of S.fluoresceïn. This solution was evaporated on the water bath and while still having a considerable volume, small, red, sharply defined crystals began to separate. The solution was evaporated to a small volume and allowed to cool but nothing further separated. These crystals are difficultly soluble in water, alcohol and ether. The alkaline solution shows a green fluorescence and slight red color by transmitted light. These crystals were dissolved in a large quantity of alcohol which on evaporation gradually deposited very small yellow crystals, which were dried in the air and taken for analysis. The Br. was determined by Carius method.

- I. ·2345 gr sub. gave ·1718 gr AgBr = 31·17% Br.
- II. ·2786 " " " ·1815 " " " = 27·72% Br.

Calculated for $C_{19}H_{10}Br_2O_6S$ = 30.42% Br.

These results, though not conclusive, indicate that under the given conditions it is the di-bromsulphonfluoresceïn which is formed. Whether this is due to the presence in the compound of the SO_2 group or simply to the greater insolubility of the di- than of the tetra-brom product cannot be definitely stated. When the original acetic acid mother liquor was evaporated to dryness, a red non-crystalline substance remained which more closely resembled rosin than the crystals. The concentrated alkaline solution had a deep red color without fluorescence and acted as a red dye stuff. The dilute alkaline solution showed the characteristic delicate pink of rosin.

Action of H_2SO_4 on S.fluoresceïn.

A test tube in which S.fluoresceïn was being made just at the end of the reaction broke and allowed the contents to run out into the sulphuric acid bath, which had a temperature of 175° . On standing several days the solution deposited a heavy precipitate which was separated by filtering through glass wool. When dry it formed a light yellow powder extremely soluble in water, alcohol and ether.

The alkaline solution had an intense green fluorescence with delicate shades of pink by transmitted light. On account of its great solubility it was impossible to purify it by crystallization, hence the Ba salt was made. The substance decomposed $BaCO_3$ with great ease forming an easily soluble salt. When it was attempted to evaporate the solution of this salt to crystallization the latter came out in a hard insoluble granular form and on continuous boiling of the solution turned brown. To avoid this undesirable form it was converted into the Ca. salt by treatment with H_2SO_4 and then $CaCO_3$. This also formed a hard granular insoluble mass on boiling but did not change in color. As there was no guarantee as to its purity and only a small quantity was obtained it was not analyzed.

Action of HCl on S.fluoresceïn.

Hydrochloric acid does not dissolve S.fluoresceïn but converts it into a light yellow granular powder. When recrystallized from water in which it is quite easily soluble it melts partially at 130° apparently with some decomposition. This compound was not further studied.


Reduction of S.fluoresceïn.

When treated with zinc dust in a strong alkaline solution sulphonfluoresceïn is reduced to a colorless substance probably analogous to fluoresceïn which is formed in the same manner. On account of its great solubility it could not be obtained in the free state. It is quickly oxidized to s.fluoresceïn by oxidizing agents as KMnO_4 and HNO_3 and passes back spontaneously on standing in the air. The latter action is however much slower than in case of fluoresceïn.

Conclusion.

The principal results relating to s.fluoresceïn which have been reached in this work may be briefly summarized as follows. Orthosulphobenzoic acid acts on resorcin at a temperature of about 180° giving off water and forming a substance analogous to fluoresceïn but having the CO group replaced by SO_2 . This substance sulphonfluoresceïn crystallizes from water in light yellow monoclinic crystals having the composition $\text{C}_{19}\text{H}_{12}\text{O}_6\text{S} + 2\text{H}_2\text{O}$. It is very soluble in alcohol and water and with difficulty in ether. It does not melt under 250° but above 300° melts with decomposition. It shows in alkaline solution a clear green fluorescence. It acts as an acid, decomposing carbonates and forming salts, the Ba salt having the composition $\text{C}_{19}\text{H}_{13}\text{O}_7\text{SBa}$. It forms an acetyl compound when boiled with acetic anhydride. It forms substitution products with Br, probably the dibrom-product most easily. It forms a compound with H_2SO_4 , probably a substitution product, whose composition was not determined. It is reduced by zinc dust and KOH to a colorless substance analogous to fluoresceïn.

Finally in terms of the prevalent theory the substance itself may be represented thus—



\ /
C
H

[1]

Annalin. No. 183. S. 1. No. 202. S. 36 & S. 153.
Berichte. No. IV. S. 457. 555. 658. 662.
" " VIII. S. 66. 146.

Transcriber's Notes:

Illustration has been moved so it does not break up the paragraph.

As this was a hand-written thesis, the spelling, punctuation and hyphenation is very inconsistent.

The original spelling, hyphenation and punctuation have been left unchanged.

*** END OF THE PROJECT GUTENBERG EBOOK ON
SULPHONFLUORESCENCE AND SOME OF ITS DERIVATIVES ***

Updated editions will replace the previous one—the old editions will be renamed.

Creating the works from print editions not protected by U.S. copyright law means that no one owns a United States copyright in these works, so the Foundation (and you!) can copy and distribute it in the United States without permission and without paying copyright royalties. Special rules, set forth in the General Terms of Use part of this license, apply to copying and distributing Project Gutenberg™ electronic works to protect the PROJECT GUTENBERG™ concept and trademark. Project Gutenberg is a registered trademark, and may not be used if you charge for an eBook, except by following the terms of the trademark license, including paying royalties for use of the Project Gutenberg trademark. If you do not charge anything for copies of this eBook, complying with the trademark license is very easy. You may use this eBook for nearly any purpose such as creation of derivative works, reports, performances and research. Project Gutenberg eBooks may be modified and printed and given away—you may do practically ANYTHING in the United States with eBooks not protected by U.S. copyright law. Redistribution is subject to the trademark license, especially commercial redistribution.

START: FULL LICENSE

THE FULL PROJECT GUTENBERG™ LICENSE

PLEASE READ THIS BEFORE YOU DISTRIBUTE OR USE THIS WORK

To protect the Project Gutenberg™ mission of promoting the free distribution of electronic works, by using or distributing this work (or any other work associated in any way with the phrase “Project Gutenberg”), you agree to comply with all the terms of the Full Project Gutenberg License available with this file or online at www.gutenberg.org/license.

Section 1. General Terms of Use and Redistributing Project Gutenberg electronic works

1.A. By reading or using any part of this Project Gutenberg electronic work, you indicate that you have read, understand, agree to and accept all the terms of this license and intellectual property (trademark/copyright) agreement. If you do not agree to abide by all the terms of this agreement, you must cease using and return or destroy all copies of Project Gutenberg electronic works in your possession. If you paid a fee for obtaining a copy of or access to a Project Gutenberg electronic work and you do not agree to be bound by the terms of this agreement, you may obtain a refund from the person or entity to whom you paid the fee as set forth in paragraph 1.E.8.

1.B. “Project Gutenberg” is a registered trademark. It may only be used on or associated in any way with an electronic work by people who agree to be bound by the terms of this agreement. There are a few things that you can do with most Project Gutenberg electronic works even without complying with the full terms of this agreement. See paragraph 1.C below. There are a lot of things you can do with Project Gutenberg electronic works if you follow the terms of this agreement and help preserve free future access to Project Gutenberg electronic works. See paragraph 1.E below.

1.C. The Project Gutenberg Literary Archive Foundation (“the Foundation” or PGLAF), owns a compilation copyright in the collection of Project Gutenberg electronic works. Nearly all the individual works in the collection are in the public domain in the United States. If an individual work is unprotected by copyright law in the United States and you are

located in the United States, we do not claim a right to prevent you from copying, distributing, performing, displaying or creating derivative works based on the work as long as all references to Project Gutenberg are removed. Of course, we hope that you will support the Project Gutenberg mission of promoting free access to electronic works by freely sharing Project Gutenberg works in compliance with the terms of this agreement for keeping the Project Gutenberg name associated with the work. You can easily comply with the terms of this agreement by keeping this work in the same format with its attached full Project Gutenberg License when you share it without charge with others.

1.D. The copyright laws of the place where you are located also govern what you can do with this work. Copyright laws in most countries are in a constant state of change. If you are outside the United States, check the laws of your country in addition to the terms of this agreement before downloading, copying, displaying, performing, distributing or creating derivative works based on this work or any other Project Gutenberg work. The Foundation makes no representations concerning the copyright status of any work in any country other than the United States.

1.E. Unless you have removed all references to Project Gutenberg:

1.E.1. The following sentence, with active links to, or other immediate access to, the full Project Gutenberg License must appear prominently whenever any copy of a Project Gutenberg work (any work on which the phrase “Project Gutenberg” appears, or with which the phrase “Project Gutenberg” is associated) is accessed, displayed, performed, viewed, copied or distributed:

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg™ License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

1.E.2. If an individual Project Gutenberg electronic work is derived from texts not protected by U.S. copyright law (does not contain a notice indicating that it is posted with permission of the copyright holder), the work can be copied and distributed to anyone in the United States without paying any fees or charges. If you are redistributing or providing access to a work with the phrase “Project Gutenberg” associated with or appearing on the work, you must comply either with the requirements of paragraphs 1.E.1 through 1.E.7 or obtain permission for the use of the work and the Project Gutenberg trademark as set forth in paragraphs 1.E.8 or 1.E.9.

1.E.3. If an individual Project Gutenberg electronic work is posted with the permission of the copyright holder, your use and distribution must comply with both paragraphs 1.E.1 through 1.E.7 and any additional terms imposed by the copyright holder. Additional terms will be linked to the Project Gutenberg License for all works posted with the permission of the copyright holder found at the beginning of this work.

1.E.4. Do not unlink or detach or remove the full Project Gutenberg License terms from this work, or any files containing a part of this work or any other work associated with Project Gutenberg.

1.E.5. Do not copy, display, perform, distribute or redistribute this electronic work, or any part of this electronic work, without prominently displaying the sentence set forth in paragraph 1.E.1 with active links or immediate access to the full terms of the Project Gutenberg License.

1.E.6. You may convert to and distribute this work in any binary, compressed, marked up, nonproprietary or proprietary form, including any word processing or hypertext form. However, if you provide access to or distribute copies of a Project Gutenberg work in a format other than “Plain Vanilla ASCII” or other format used in the official version posted on the official Project Gutenberg website (www.gutenberg.org), you must, at no additional cost, fee or expense to the user, provide a copy, a means of exporting a copy, or a means of obtaining a copy upon request, of the work in its original “Plain Vanilla ASCII” or other form. Any alternate format must include the full Project Gutenberg License as specified in paragraph 1.E.1.

1.E.7. Do not charge a fee for access to, viewing, displaying, performing, copying or distributing any Project Gutenberg works unless you comply with paragraph 1.E.8 or 1.E.9.

1.E.8. You may charge a reasonable fee for copies of or providing access to or distributing Project Gutenberg electronic works provided that:

- You pay a royalty fee of 20% of the gross profits you derive from the use of Project Gutenberg works calculated using the method you already use to calculate your applicable taxes. The fee is owed to the owner of the Project Gutenberg trademark, but he has agreed to donate royalties under this paragraph to the Project Gutenberg Literary Archive Foundation. Royalty payments must be paid within 60 days following each date on which you prepare (or are legally required to prepare) your periodic tax returns. Royalty payments should be clearly marked as such and sent to the Project Gutenberg Literary Archive Foundation at the address specified in Section 4, "Information about donations to the Project Gutenberg Literary Archive Foundation."
- You provide a full refund of any money paid by a user who notifies you in writing (or by e-mail) within 30 days of receipt that s/he does not agree to the terms of the full Project Gutenberg™ License. You must require such a user to return or destroy all copies of the works possessed in a physical medium and discontinue all use of and all access to other copies of Project Gutenberg™ works.
- You provide, in accordance with paragraph 1.F.3, a full refund of any money paid for a work or a replacement copy, if a defect in the electronic work is discovered and reported to you within 90 days of receipt of the work.
- You comply with all other terms of this agreement for free distribution of Project Gutenberg™ works.

1.E.9. If you wish to charge a fee or distribute a Project Gutenberg™ electronic work or group of works on different terms than are set forth in this agreement, you must obtain permission in writing from the Project Gutenberg Literary Archive Foundation, the manager of the Project

Gutenberg™ trademark. Contact the Foundation as set forth in Section 3 below.

1.F.

1.F.1. Project Gutenberg volunteers and employees expend considerable effort to identify, do copyright research on, transcribe and proofread works not protected by U.S. copyright law in creating the Project Gutenberg™ collection. Despite these efforts, Project Gutenberg™ electronic works, and the medium on which they may be stored, may contain “Defects,” such as, but not limited to, incomplete, inaccurate or corrupt data, transcription errors, a copyright or other intellectual property infringement, a defective or damaged disk or other medium, a computer virus, or computer codes that damage or cannot be read by your equipment.

1.F.2. LIMITED WARRANTY, DISCLAIMER OF DAMAGES - Except for the “Right of Replacement or Refund” described in paragraph 1.F.3, the Project Gutenberg Literary Archive Foundation, the owner of the Project Gutenberg™ trademark, and any other party distributing a Project Gutenberg™ electronic work under this agreement, disclaim all liability to you for damages, costs and expenses, including legal fees. YOU AGREE THAT YOU HAVE NO REMEDIES FOR NEGLIGENCE, STRICT LIABILITY, BREACH OF WARRANTY OR BREACH OF CONTRACT EXCEPT THOSE PROVIDED IN PARAGRAPH 1.F.3. YOU AGREE THAT THE FOUNDATION, THE TRADEMARK OWNER, AND ANY DISTRIBUTOR UNDER THIS AGREEMENT WILL NOT BE LIABLE TO YOU FOR ACTUAL, DIRECT, INDIRECT, CONSEQUENTIAL, PUNITIVE OR INCIDENTAL DAMAGES EVEN IF YOU GIVE NOTICE OF THE POSSIBILITY OF SUCH DAMAGE.

1.F.3. LIMITED RIGHT OF REPLACEMENT OR REFUND - If you discover a defect in this electronic work within 90 days of receiving it, you can receive a refund of the money (if any) you paid for it by sending a written explanation to the person you received the work from. If you received the work on a physical medium, you must return the medium with your written explanation. The person or entity that provided you with the defective work may elect to provide a replacement copy in lieu of a refund. If you received the work electronically, the person or entity providing it to

you may choose to give you a second opportunity to receive the work electronically in lieu of a refund. If the second copy is also defective, you may demand a refund in writing without further opportunities to fix the problem.

1.F.4. Except for the limited right of replacement or refund set forth in paragraph 1.F.3, this work is provided to you 'AS-IS', WITH NO OTHER WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR ANY PURPOSE.

1.F.5. Some states do not allow disclaimers of certain implied warranties or the exclusion or limitation of certain types of damages. If any disclaimer or limitation set forth in this agreement violates the law of the state applicable to this agreement, the agreement shall be interpreted to make the maximum disclaimer or limitation permitted by the applicable state law. The invalidity or unenforceability of any provision of this agreement shall not void the remaining provisions.

1.F.6. INDEMNITY - You agree to indemnify and hold the Foundation, the trademark owner, any agent or employee of the Foundation, anyone providing copies of Project Gutenberg™ electronic works in accordance with this agreement, and any volunteers associated with the production, promotion and distribution of Project Gutenberg™ electronic works, harmless from all liability, costs and expenses, including legal fees, that arise directly or indirectly from any of the following which you do or cause to occur: (a) distribution of this or any Project Gutenberg work, (b) alteration, modification, or additions or deletions to any Project Gutenberg work, and (c) any Defect you cause.

Section 2. Information about the Mission of Project Gutenberg

Project Gutenberg is synonymous with the free distribution of electronic works in formats readable by the widest variety of computers including obsolete, old, middle-aged and new computers. It exists because of the

efforts of hundreds of volunteers and donations from people in all walks of life.

Volunteers and financial support to provide volunteers with the assistance they need are critical to reaching Project Gutenberg's goals and ensuring that the Project Gutenberg collection will remain freely available for generations to come. In 2001, the Project Gutenberg Literary Archive Foundation was created to provide a secure and permanent future for Project Gutenberg and future generations. To learn more about the Project Gutenberg Literary Archive Foundation and how your efforts and donations can help, see Sections 3 and 4 and the Foundation information page at www.gutenberg.org.

Section 3. Information about the Project Gutenberg Literary Archive Foundation

The Project Gutenberg Literary Archive Foundation is a non-profit 501(c)(3) educational corporation organized under the laws of the state of Mississippi and granted tax exempt status by the Internal Revenue Service. The Foundation's EIN or federal tax identification number is 64-6221541. Contributions to the Project Gutenberg Literary Archive Foundation are tax deductible to the full extent permitted by U.S. federal laws and your state's laws.

The Foundation's business office is located at 41 Watchung Plaza #516, Montclair NJ 07042, USA, +1 (862) 621-9288. Email contact links and up to date contact information can be found at the Foundation's website and official page at www.gutenberg.org/contact

Section 4. Information about Donations to the Project Gutenberg Literary Archive Foundation

Project Gutenberg™ depends upon and cannot survive without widespread public support and donations to carry out its mission of increasing the number of public domain and licensed works that can be freely distributed in machine-readable form accessible by the widest array of equipment

including outdated equipment. Many small donations (\$1 to \$5,000) are particularly important to maintaining tax exempt status with the IRS.

The Foundation is committed to complying with the laws regulating charities and charitable donations in all 50 states of the United States. Compliance requirements are not uniform and it takes a considerable effort, much paperwork and many fees to meet and keep up with these requirements. We do not solicit donations in locations where we have not received written confirmation of compliance. To SEND DONATIONS or determine the status of compliance for any particular state visit www.gutenberg.org/donate.

While we cannot and do not solicit contributions from states where we have not met the solicitation requirements, we know of no prohibition against accepting unsolicited donations from donors in such states who approach us with offers to donate.

International donations are gratefully accepted, but we cannot make any statements concerning tax treatment of donations received from outside the United States. U.S. laws alone swamp our small staff.

Please check the Project Gutenberg web pages for current donation methods and addresses. Donations are accepted in a number of other ways including checks, online payments and credit card donations. To donate, please visit: www.gutenberg.org/donate.

Section 5. General Information About Project Gutenberg electronic works

Professor Michael S. Hart was the originator of the Project Gutenberg concept of a library of electronic works that could be freely shared with anyone. For forty years, he produced and distributed Project Gutenberg eBooks with only a loose network of volunteer support.

Project Gutenberg eBooks are often created from several printed editions, all of which are confirmed as not protected by copyright in the U.S. unless a

copyright notice is included. Thus, we do not necessarily keep eBooks in compliance with any particular paper edition.

Most people start at our website which has the main PG search facility:
www.gutenberg.org.

This website includes information about Project Gutenberg, including how to make donations to the Project Gutenberg Literary Archive Foundation, how to help produce our new eBooks, and how to subscribe to our email newsletter to hear about new eBooks.