


THE
HEATHERY;
OR A
MONOGRAPH
OF
THE GENUS ERICA:
VOL. IV.



THE
HEATHERY;
OR A
MONOGRAPH
OF
THE GENUS ERICA:
VOL. IV.



The Project Gutenberg eBook of The heathery; or, A monograph of the genus Erica. vol. 4

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

Title: The heathery; or, A monograph of the genus Erica. vol. 4

Author: active 1799-1828 Henry Cranke Andrews

Release date: February 20, 2023 [eBook #70085]

Language: English

Original publication: United Kingdom: Henry G. Bohn, 1845

Other information and formats: www.gutenberg.org/ebooks/70085

Credits: Chuck Greif and the Online Distributed Proofreading Team at <https://www.pgdp.net> (This file was produced from images generously made available by Biodiversity Heritage Library.)

*** START OF THE PROJECT GUTENBERG EBOOK THE
HEATHERY; OR, A MONOGRAPH OF THE GENUS ERICA. VOL. 4

ALPHABETICAL INDEX
TO THE
FIGURES IN VOL. IV.

THE
HEATHERY;

OR A

MONOGRAPH

OF

THE GENUS ERICA:

CONTAINING

COLOURED ENGRAVINGS,

WITH

LATIN AND ENGLISH DESCRIPTIONS, DISSECTIONS, ETC.

OF ALL

THE KNOWN SPECIES OF THAT EXTENSIVE AND DISTINGUISHED TRIBE OF PLANTS.

By H. C. ANDREWS.

IN SIX VOLUMES.

VOL. IV.

SECOND EDITION, CORRECTED AND ENLARGED.

LONDON:
HENRY G. BOHN, YORK STREET, COVENT GARDEN.
1845.

ERICA Andromedæflora.

DESCRIPTIO.

Antheræ cristatæ, inclusæ. Flores axillares, sub-terminales, declinati, plerumque terni. Calyx magnus, lato-ovatus, carneus: corollis ovatis, magnis, incarnatis. Folia ternata, subulata, incurvata, acuta, petiolis longis adpressis. Caulis erectus, pedalis et ultra. Rami pauci. Ramuli verticillati, ternati.

DESCRIPTION.

Tips crested, and within the blossom. Flowers grow from the axillæ of the leaves, are nearly terminal, and bend downwards, mostly in threes. The empalement is large, of a broad ovate form, and flesh-coloured. Blossom egg-shaped, large, and flesh-coloured. Leaves by threes, awl-shaped, incurved, sharp-pointed, with long footstalks pressed to the branches. Stem upright, a foot or more high. Large branches few. The smaller branches in whorls, by threes.



Native of the Cape of Good Hope.
In bloom from March till May.

REFERENCE.

1. The empalement.
2. A blossom.
3. The same spread open.
4. Chives and pointal, one tip magnified.
5. Seed-bud and pointal, summit magnified.



E. Andromedaeflora

ERICA aristata.

DESCRIPTIO.

Antheræ inclusæ, basi bicornes. Flores in ramis terminales, fasciculati, plerumque quaterni. Pedunculi longi, colorati. Corolla inflato-tubulosa, ore arctata, saturate purpurea, pollicaris, lucida; laciniis limbi rotundatis, expansis, albicantibus. Folia quaterna et quina, revoluta, lucida, apice aristata. Caulis fruticosus, vix pedalis, ramosus. Rami et ramuli flexuosi, filiformes.

DESCRIPTION.

Tips within the blossoms, and two-horned at their base. Flowers terminate the branches in bunches, mostly in fours. Footstalks long, and coloured. Blossom of an inflated tubular form, narrowed at the mouth, which is of a deep purple, an inch long, and shining. The segments of the border are rounded, spreading, and white. Leaves by fours and fives, rolled back, and shining, with a bristle at the point. Stem shrubby, scarce a foot high, branching. The small and large branches are flexuose, and thread-shaped.



Native of the Cape of Good Hope.
In bloom in spring, and again late in autumn.

REFERENCE.

1. A leaf magnified.
2. The empalement.
3. Chives and pointal, one tip magnified.
4. Seed-bud and pointal, summit magnified.



E. aristata

ERICA aurea.

DESCRIPTIO.

Antheræ muticæ, sub-inclusæ. Flores horizontaliter verticillati, prope ramorum summitatem. Pedunculi brevissimi, bracteis tribus instructi. Corolla cylindræa, glabra, aurantia: laciniis oris parum reflexis. Folia sena, linearia, glabra, apice incurvata, patentia. Caulis erectus, fruticosus, pedalis. Rami verticillati, patentes.

DESCRIPTION.


Tips beardless, just within the blossom. Flowers grow horizontally in whorls near the summit of the branches. Footstalks very short, and furnished with three floral leaves. Blossom cylinder-shaped, smooth, and of a gold colour: segments of the mouth a little reflexed. Leaves by sixes, linear, and smooth, turned inward at the point, and spreading. Stem upright, shrubby, a foot high. The branches grow in whorls, and are spreading.



Native of the Cape of Good Hope.
In bloom from July till September.

REFERENCE.

1. The empalement magnified.
2. The chives and pointal, one tip magnified.
3. Seed-bud and pointal, summit magnified.



E. aurea.

ERICA Blandfordia.

DESCRIPTIO.

Antheræ aristatæ, inclusæ. Flores plures, sessiles, et horizontaliter siti, ramulos plerumque binatim terminantes: sub apicibus ramorum conferti in ramulos brevissimos, spicam formantes. Corolla urceolata, glabra, lutea, ore arctata: laciniis reflexis. Folia quaterna, linearia, obtusa, glabra. Caulis fruticosus, sesquipedalis, erectus.

DESCRIPTION.

Tips bearded, and within the blossom. Flowers numerous, sessile, in a horizontal direction, and terminate the smaller branches mostly in twos: near the upper part of the larger branches they are crowded together upon very short branches, and form a spike. Blossom jar-shaped, smooth and yellow, narrowed at the mouth, with the segments reflexed. Leaves by fours, linear, obtuse, and smooth. Stem shrubby, a foot and a half high, and upright.



Native of the Cape of Good Hope.
In bloom from March till June.

REFERENCE.

1. The empalement magnified.
2. The seed-bud, chives, and pointal.
3. A chive magnified.
4. Seed-bud and pointal, summit magnified.



ERICA calycina, *major*.

DESCRIPTIO.

Antheræ cristatæ, inclusæ. Flores in ultimis ramulis terminales, umbellati. Pedunculi carnei. Perianthium tetraphyllum; foliolis spathulatis, acuminatis, carneis, ad basin virescentibus, corolla longioribus. Corolla urceolata, saturate carnea, ad basin pallida; oris laciniis patentibus, acuminatis. Folia terna, lanceolata, trigona, acuminata, adpressa. Caulis virgatus, pedalis. Rami et ramuli flexuosissimi.

DESCRIPTION.

Tips crested, and within the blossom. Flowers terminate the smaller branches in umbels. Footstalks flesh-coloured. Cup four-leaved. Leaflets spathula-shaped, sharp-pointed, flesh-coloured, greenish at the base, and longer than the flower. Blossom pitcher-shaped, of a deep flesh-colour, pale at the base: segments of the border spreading, and sharp-pointed. Leaves by threes, lance-shaped, three-sided, sharp-pointed, and pressed to the branches. Stem twiggy, a foot high. The small and large branches are very flexuose.



Native of the Cape of Good Hope.
In bloom from June till September.

REFERENCE.

1. The empalement.
2. A blossom.
3. A chive magnified.
4. The seed-bud and pointal, summit magnified.
5. Seed-bud magnified.



E. calycina major

Wm. Andrews del. H. G. S.

ERICA canaliculata.

DESCRIPTIO.

Antheræ muticæ, sub-exertæ, nigræ. Flores in ultimis ramulis terminales, plerumque terni, cernui. Pedunculi longi, colorati. Corolla campanulata, pallide purpurea; oris laciniis ovatis, acutis. Folia terna, erecta, obtusa, supra plana, subtus canaliculata. Caulis fruticosus, bipedalis, erectus, ferrugineus. Rami et ramuli verticillati, numerosi.

DESCRIPTION.

Tips beardless, just without the blossom, and black. Flowers terminate the ends of the smaller branches, mostly by threes, hanging down. Footstalks long and coloured. Blossom bell-shaped, and of a pale purple: segments of the border are egg-shaped, and sharp-pointed. Leaves by threes, upright, and blunt, flat on the upper surface, and channelled beneath. Stem shrubby, two feet high, upright, and rusty. The large and smaller branches are whorled and numerous.



Native of the Cape of Good Hope.
In bloom from March till May.

REFERENCE.

1. A leaf magnified.
2. The empalement magnified.
3. A blossom.
4. Chives and pointal, one tip magnified.
5. The seed-bud and pointal, summit magnified.



E. canaliculata

ERICA canaliculata, *minor*.

DESCRIPTIO.

Antheræ muticæ, sub-inclusæ. Flores in ramulis terminales, plerumque terni, cernui. Pedunculi colorati. Calyx glandulosus. Corolla parva, campanulata, pallide purpurea: oris laciniis ovatis, acutis. Folia terna, erecta, obtusa, supra plana, subtus canaliculata. Caulis fruticosus, bipedalis, erectus. Rami et ramuli verticillati, numerosi.

DESCRIPTION.


Tips beardless, and just within the blossom. Flowers terminate the ends of the smaller branches mostly by threes, hanging down. Footstalks coloured. Empalement glandular. Blossom small, bell-shaped, and of a pale purple: the segments are ovate, and sharp-pointed. Leaves by threes, upright and obtuse, flat on the upper surface, and channelled beneath. Stem shrubby, two feet high, and upright. The larger and smaller branches are whorled, and numerous.



Native of the Cape of Good Hope.
In bloom from March till June.

REFERENCE.

1. A leaf magnified.
2. The empalement magnified.
3. A chive magnified.
4. Seed-bud and pointal, summit magnified.



E. canaliculata minor

ERICA cordata.

DESCRIPTIO.

Antheræ muticæ, sub-exertæ. Floras terminales, in apicibus ramulorum confertissimi, capitati, erecti. Pedunculi breves. Corolla globosa, glabra, albida; laciniis limbi rotundatis, patentibus. Folia terna, cordata, patentia, marginibus revolutis, supra villosa, subtus lanata. Caulis humilis, spithamæus. Rami et ramuli numerosi, conferti, flexuosi, patentes.

DESCRIPTION.


Tips beardless, just without the blossom. Flowers terminal, crowded very close together at the ends of the branches in upright heads. Footstalks short. Blossom globular, smooth, and white. Segments of the border round and spreading. Leaves by threes, heart-shaped, spreading, and rolled back at the edges, hairy on the upper surface, and woolly beneath. Stem low, a span high. The large and small branches are numerous, crowded together, flexuose, and spreading.



Native of the Cape of Good Hope.
In bloom from April till June.

REFERENCE.

1. A leaf magnified.
2. The empalement and blossom.
3. The chives and pointal, one tip magnified.
4. Seed-bud and pointal, summit magnified.
5. Seed-bud magnified.



E. cordata

W. B. Andrews. England

ERICA decora.

DESCRIPTIO.

Antheræ muticæ, inclusæ. Flores spicati, laxi, in medio ramulorum, axillares, dependentes. Pedunculi longi, colorati. Corolla campanulata, costata, saturate carnea: oris laciniis reflexis. Folia sena, linearia, obtusa, arcuata, petiolis longis adpressis. Caulis fruticosus, sesquipedalis, sub-erectus, gracilis. Rami et ramuli verticillati.

DESCRIPTION.

Tips beardless, and within the blossom. Flowers grow in large spikes, about the middle of the branches, from the footstalks of the leaves, and hang down. Flower-stalk long, and coloured. Blossom bell-shaped, ribbed, and of a deep flesh-colour: the segments of the border are reflexed. Leaves by sixes, linear, obtuse, and bowed, with long footstalks pressed to the branches. Stem shrubby, a foot and a half high, nearly upright, and slender. The small and large branches grow in whorls.



Native of the Cape of Good Hope.
In bloom from November till February.

REFERENCE.

1. The empalement magnified.
2. The chives magnified.
3. Seed-bud and pointal, summit magnified.
4. Seed-bud magnified.



ERICA discolor.

DESCRIPTIO.

Antheræ aristatæ, inclusæ. Flores plerumque terni, sub-cernui, ramulis terminalibus. Calyx duplex. Corolla tubulosa, sub-cylindræa, ima parte carnea, summa flavo-albicante. Folia terna, linearia, nitida, erecto-patentia. Caulis fruticosus, tripedalis, erectus, ramosus.

DESCRIPTION.


Tips bearded, and within the blossom. Flowers mostly by threes, bending a little downwards, and terminating the smaller branches. Empalement double. Blossom tubular, nearly cylindrical, flesh-coloured at the base, and of a yellowish white at the mouth. Leaves by threes, linear, shining, and between erect and spreading. Stem shrubby, three feet high, upright, and branching.



Native of the Cape of Good Hope.
In bloom from November till April.

REFERENCE.

1. The empalement magnified.
2. The chives and pointal, one tip magnified.
3. Seed-bud and pointal, summit magnified.



E. Discolor

ERICA Echiiflora.

DESCRIPTIO.

Antheræ muticæ, inclusæ. Flores fere terminales, in spicam dense verticillati, horizontales. Corolla tubulosa, brevis, inflata, instar Echii, saturate carnea, costata, viscosa; laciniis patentibus, rotundatis. Folia sena vel septena, irregulariter verticillati. Caulis fruticosus, bipedalis.

DESCRIPTION.


Tips beardless, within the blossom. Flowers almost terminal, in a spike, densely whorled, and horizontal. Blossom tubular, short, inflated, and resembling an Echium, of a deep flesh-colour, ribbed, and viscous; segments spreading, and rounded. Leaves by sixes or sevens, and irregularly whorled. Stem shrubby, and two feet high.



Native of the Cape of Good Hope.
In bloom from February till May.

REFERENCE.

1. The empalement magnified.
2. A blossom.
3. Chives and pointal, one tip magnified.
4. Seed-bud and pointal, summit magnified.



E. Echiiflora

43. Andrews & Long

ERICA ferruginea.

DESCRIPTIO.

Antheræ muticæ, inclusæ. Flores in ramulis terminales, simpliciter verticillati, horizontales. Pedunculi colorati, hispiduli, bracteis tribus hirsutis instructi. Corolla basi inflata, apice attenuata; ima pars carnea, summa saturate purpurea, ore arctata: laciniis acutis, patentibus. Folia subquaterna, obtusa, pilis longis ferrugineis circumscissa. Caulis humilis, fruticosus, erectus, sub-pedalis. Rami et ramuli flexuosi, filiformes, foliis subtecti.

DESCRIPTION.

Tips beardless, and within the blossom. Flowers terminate the smaller branches in single horizontal whorls. Footstalks coloured, slightly hispid, and furnished with three hairy floral leaves. Blossom inflated at the base, tapering to the point, the lower part of a flesh-colour, the upper part of a deep purple, straightened at the mouth: the segments are pointed, and spreading. Leaves mostly by fours, blunt-ended, and beset with long rusty-coloured hairs. Stem low, shrubby, upright, and scarce a foot high. The large and small branches are flexuose, thread-shaped, and nearly covered by the leaves.

—

Native of the Cape of Good Hope.
In bloom from April till July.

REFERENCE.

1. A leaf magnified.
2. The empalement magnified.
3. Chives and pointal, one tip magnified.
4. Seed-bud and pointal, summit magnified.



ERICA fragrans.

DESCRIPTIO.

Antheræ basi bicornes, exertæ. Flores plerumque terni, laterales et terminales, parvi: pedunculis longiusculis. Corolla campanulata, fragrans, pallide purpurea: laciniis majusculis revolutis. Folia ternata, subulata, erecta, glauca, acuta, supra plana, subtus sulcata. Caulis fruticosus, semipedalis, valde ramosus: ramis patentibus.

DESCRIPTION.


Tips two-horned at the base, and without the blossom. Flowers grow mostly by threes, lateral, and terminal, small, and with longish peduncles. Blossom bell-shaped, sweet-scented, pale purple, with largish revolute segments. Leaves in threes, awl-shaped, upright, of a glaucous green, and sharp-pointed, flat on their upper surface, and furrowed beneath. Stem shrubby, half a foot high, very much branched, with the branches spreading.



Native of the Cape of Good Hope.
In bloom from March till May.

REFERENCE.

1. A leaf magnified.
2. The empalement magnified.
3. Seed-bud, chives, and pointal, magnified.



E. fragrans

St. Andrews & Hayfield reges

ERICA halicacaba.

DESCRIPTIO.

Antheræ muticæ, inclusæ. Flores sub-terni, cernui, terminales. Pedunculi breves. Perianthium duplex, corollis adpressum. Corolla ovata, inflata, maxima, pallide lutea: oris laciniis erectis, acuminatis, conniventibus, obtusis. Folia terna, subulata, recurvata, trigona, glabra. Caulis robustus, pedalis, valde ramosus. Ramuli patentes, plurimi, oppositi.

DESCRIPTION.


Tips beardless, within the blossom. Flowers grow mostly by threes, nodding, and terminating the branches. Footstalks short. Cup double, and pressed to the blossom. Blossom egg-shaped, inflated, large, and of a pale yellow colour: the segments of the border are erect, tapered, approaching, and blunt-ended. Leaves by threes, awl-shaped, recurved, three-sided, and smooth. Stem sturdy, a foot high, and very branching. The smaller branches are spreading, numerous, and opposite.



Native of the Cape of Good Hope.
In bloom from May till July.

REFERENCE.

1. The empalement.
2. Chives and pointal, one tip magnified.
3. Seed-bud and pointal, summit magnified.



E. halicacaba

ERICA hirta.

DESCRIPTIO.

Antheræ aristatæ, inclusæ. Flores terminales, terni, cernui. Pedunculi breves. Corolla clavata, pollicaris, ad basin profunde rubro-purpurea, apice virescens, ore parum arctata; laciniis obtusis, erectis. Folia terna, linearia, obtusa, hirta. Caulis erectus, ramosus, bipedalis.

DESCRIPTION.

Tips bearded, and within the blossom. Flowers terminate the branches by threes, hanging down. Footstalks short. Blossom club-shaped, an inch long, of a deep red purple at the base, greenish at the end, a little narrowed at the mouth: segments blunt, and upright. Leaves by threes, linear, blunt, and hairy. Stem upright, branching, and two feet high.



Native of the Cape of Good Hope.
In bloom from January till April.

REFERENCE.

1. The empalement.
2. The chives and pointal spread.
3. Seed-bud and pointal, summit magnified.



ERICA hirta; *Var. viridiflora*.

DESCRIPTIO.

Antheræ aristatæ, inclusæ. Flores in ramulis terminales, plerumque terni, patente-cernui. Pedunculi brevissimi. Corolla cylindræa, pollicaris, viridis: limbi laciniis obtusis, erectis. Folia terna et quaterna, linearia, recta, obtusa, hirta, supra plana, subtus sulcata. Caulis erectus, sesquipedalis, villosus. Ramuli terni, verticillati.

DESCRIPTION.


Tips bearded, and within the blossom. Flowers terminate the smaller branches mostly by threes, between spreading and hanging down. Footstalks very short. Blossom cylindrical, an inch long, and green. The segments of the border are obtuse, and upright. Leaves by threes and fours, linear, straight, blunt, and hairy, flat on their upper surface, and furrowed beneath. Stem upright, a foot and a half high, and hairy. The small branches grow by threes, in whorls.



Native of the Cape of Good Hope.
In bloom from October till January.

REFERENCE.

1. A leaf magnified.
2. The empalement.
3. Chives and pointal, one tip magnified.
4. Seed-bud and pointal, summit magnified.



E. hirta var. *viridiflora*

ERICA Hyacinthoides.

DESCRIPTIO.

Antheræ muticæ, inclusæ. Flores sessiles, erecti, fastigiati, quaterni, conferti, terminales, fere pedunculati. Corolla ventricosa, incarnata, ore arctata; laciniis sub-cordatis, undulatis, expansis, maximis. Folia quaterna, linearia, glabra, crassiuscula, nitida. Caulis fruticosus, spithameus, erectus, ramosus. Ramuli frequentissimi, erecto-patentes.

DESCRIPTION.


Tips beardless, within the blossom. Flowers sit close in upright bunches by fours, on the ends of the branches with scarcely any footstalks. Blossom swelled out, flesh-coloured, and narrowed at the mouth; the segments are nearly heart-shaped, waved, spreading, and very large. Leaves by fours, linear, smooth, thickish, and shining. Stem shrubby, a span high, upright, and branching. The smaller branches are numerous, and between erect and spreading.



Native of the Cape of Good Hope.
In bloom from June till August.

REFERENCE.

1. The empalement magnified.
2. The chives and pointal, one tip magnified.
3. Seed-bud and pointal, summit magnified.
4. Seed-bud magnified.



E. Hyacinthoides

Ed. Harvey, & Th. G. Smith

ERICA incarnata.

DESCRIPTIO.

Antheræ cristatæ, sub-inclusæ. Flores in ramulis terminales, in racemum densum coalescentes. Pedunculi longi, purpurei. Corolla sub-ovata, cernua, carnea; ore contracto, lævissime quadrifido, laciniis erectis. Folia quaterna, obtusa, linearia, glabra. Caulis sub-erectus, pedalis, ramosissimus; rami et ramuli flexuosi, divaricatopatentes.

DESCRIPTION.


Tips crested, and just within the blossom. The flowers grow at the ends of the smaller branches, forming a close bunch. Footstalks are long, and purple. Blossom nearly egg-shaped, bending down, and flesh-coloured; the mouth contracted, and slightly cut into four upright segments. Leaves by fours, obtuse, linear, and smooth. Stem nearly upright, a foot high, and very much branched; the large and smaller branches are flexuose, and spread outwards.



Native of the Cape of Good Hope.
In bloom from July till October.

REFERENCE.

1. The empalement magnified.
2. The chives spread open, one tip magnified.
3. Seed-bud and pointal, summit magnified.



E. incarnata

ERICA incurva.

DESCRIPTIO.

Antheræ aristatæ, sub-inclusæ: stylo sub-exerto. Flores in ramulis terminales, umbellati, terni. Pedunculi aculeati. Folia terna, farinose viridia, arcuata, incurva, aculeata, subtus exarata. Corolla urceolata, purpurea, odorata: oris laciniis rotundatis, sub-erectis. Caulis fruticosus, ramosus. Rami et ramuli hirsuti, erecto-patentes.

DESCRIPTION.

Tips bearded, and just within the blossom. Shaft just without. Flowers terminate the small branches in umbels by threes. Footstalks prickly. Leaves by threes, of a mealy green, bowed, turned inwards, prickly, and deeply furrowed beneath. Blossom pitcher-shaped, purple, and sweet-scented: the segments of the border are roundish, and nearly upright. Stem shrubby, and branching. The large and small branches are roughly haired, and between erect and spreading.



Native of the Cape of Good Hope.
In bloom from March till May.

REFERENCE.

1. A leaf magnified.
2. The empalement magnified.
3. The chives and pointal, one tip magnified.
4. The seed-bud and pointal, summit magnified.
5. A flower and leaf of a small variety.



E. incurva

1827

by H. Andrews & R. Brown

ERICA lachnæa, *purpurea*.

DESCRIPTIO.

Antheræ cristatæ, inclusæ. Flores terni, in apice ramulorum, cernui. Perianthium duplex; pedunculis brevissimis. Corolla cylindraceo-urceolata, purpureo-pulverulenta; laciniis limbi obtusis, sub-erectis. Folia terna, obtusa, conferta, tomentosa, farinaria, viridia. Caulis erectus, pedalis, fruticosus. Rami numerosi, ramulis brevissimis tecti.

DESCRIPTION.

Tips crested, and within the blossom. Flowers grow by threes, at the ends of the small branches, nodding. Cup double; footstalks very short. Blossom cylindrically pitcher-shaped, and of a powdery purple; segments of the border are obtuse, and nearly upright. Leaves by threes, blunt, crowded, downy, and of a mealy green. Stem upright, a foot high, and shrubby. Branches numerous, and covered by very short smaller ones.



Native of the Cape of Good Hope.
In bloom from July till October.

REFERENCE.

1. A leaf magnified.
2. The empalement magnified.
3. Chives and pointal, one tip magnified.
4. The seed-bud and pointal, summit magnified.
5. Seed-bud magnified.



E. lachnaea purpurea

ERICA Lambertia.

DESCRIPTIO.

Antheræ cristatæ, inclusæ. Flores in medio ramulorum, solitarii, vel terni, cernui. Pedunculi et calyx colorati. Corolla sub-globosa, inflata, alba, glabra, magnitudine pisi magni; laciniis ovatis, ad basin gibbis, erectis. Folia quaterna, acuminata, glabra; juniora erecto-patentia, seniora reflexa. Caulis fruticosus, sub-pedalis, ramosissimus.

DESCRIPTION.



Tips crested, and within the blossom. Flowers grow in the middle of the small branches, solitary, or by threes, nodding. The footstalks and empalement coloured. Blossom nearly globular, swelled out, white, smooth, and of the size of a large pea; the segments egg-shaped, bunched at the base, and upright. Leaves by fours, sharp-pointed, and smooth; the younger ones erect and spreading, the older ones reflexed. Stem shrubby, near a foot high, and very much branched.



Native of the Cape of Good Hope.
In bloom from May till July.

REFERENCE.

1. The empalement magnified.
2. The chives and pointal.
3. A chive magnified.
4. The seed-bud and pointal, summit magnified.
5. Seed-bud magnified.



E. Lambertia

by H. Andrews & Th. Knightbridge

ERICA lucida.

DESCRIPTIO.

Antheræ cristatæ, sub-inclusæ. Flores terminates, in umbellis trifloris. Pedunculi breves. Perianthium tetraphyllum, foliolis lato-ovatis, acutis, ciliatisque. Corolla breviter campanulata, laciniis patulis, pallide carneis et lucidis. Folia ternata, subulata, erecta, lucida. Caulis fruticosus, strictus, gracilis, bipedalis. Rami plerumque ternati, erecto-patentes.

DESCRIPTION.


Tips crested, just within the blossom. Flowers terminal, in three-flowered umbels. Footstalks short. Cup four-leaved: leaflets broadly egg-shaped, sharp-pointed, and ciliated. Blossom shortly bell-shaped, with spreading segments of a pale shining flesh-colour. Leaves by threes, awl-shaped, upright, and shining. Stem shrubby, upright, slender, and two feet high. Branches mostly by threes, between erect and spreading.



Native of the Cape of Good Hope.
In bloom from April till June.

REFERENCE.

1. The empalement magnified.
2. A blossom magnified.
3. The chives and pointal, magnified.
4. The seed-bud and pointal, magnified.



E. lucida

D. Moore

by F. Anderson & T. Knight

ERICA Monsoniana.

DESCRIPTIO.

Antheræ aristatæ, inclusæ. Flores in ramulorum apice terni, nutantes, spicam quasi formantes prope caulis summitatem. Pedunculi fere nulli. Corolla tubulosa, fauce arctata, candidissima; laciniis sub-erectis. Folia terna, nitida, linearia, erecto-patentia. Caulis tripedalis, teres, erectus.

DESCRIPTION.

Tips bearded, and within the blossom. Flowers grow by threes at the ends of the smaller branches, hanging down, and forming a sort of spike near the end of the stem; with hardly any footstalks. Blossom tubular, narrowed at the mouth, and of a very clear white. Segment of the border nearly upright. Leaves by threes, shining, linear, and between erect and spreading. Stem three feet high, round and upright.



Native of the Cape of Good Hope.
In bloom from May till July.

REFERENCE.

1. A flower-branch with its involucrum.
2. The empalement.
3. The chives and pointal spread.
4. The seed-bud and pointal, summit magnified.



E. Monsoniana

W. & A. N. Y.

by J. B. Sowerby & H. J. Sowerby

ERICA mucosa.

DESCRIPTIO.

Antheræ aristatæ, inclusæ. Flores plures, terminales, fasciculati, cernui, purpurei, fulgentes, mucosi; pedunculis longis, purpureis. Corolla globosa; laciniis semi-orbiculatis, æqualibus, erectis. Folia quaterna, sub-trigona, obtusa, glaberrima. Caulis fruticosus, tripedalis, ramosus; ramis longis, virgatis, flexuosis.

DESCRIPTION.

Tips bearded, and within the blossom. Flowers many, terminating the branches in bunches, hanging down, of a brilliant purple colour, and clammy. Footstalks long and purple. Blossom globular; the segments of the border are half round, equal, and upright. Leaves by fours, nearly three-sided, blunt, and smooth. Stem shrubby, three feet high, and branching; the branches are long, twiggy, and flexuose.



Native of the Cape of Good Hope.
In bloom from June till September.

REFERENCE.

1. The empalement magnified.
2. The chives and pointal, summit magnified.
3. Seed-bud and pointal, summit magnified.



E. mucosa

W. G. Anderson

ERICA multiflora.

DESCRIPTIO.

Antheræ muticæ, exertæ. Flores axillares, sub-terminales, sub-spicati. Pedunculi capillares, longissimi, bracteis tribus minutis instructi. Corolla cylindraceo-urceolata, costata, sub-carnea; laciniis minutis, reflexis. Folia quaterna seu quina, obtusa, glabra, linearia, recta. Caulis fruticosus, erectus, ramosus. Rami et ramuli rigidi, erecti.

DESCRIPTION.

Tips beardless, and without the blossom. Flowers grow from the insertion of the leaves, nearly terminal, and almost spiked. Footstalks hair-like, very long, and furnished with three small floral leaves. Blossom cylindrically pitcher-shaped, ribbed, and slightly flesh-coloured: segments small and reflexed. Leaves by fours or fives, blunt, smooth, linear, and straight. Stem shrubby, upright, and branching. The small and larger branches are harsh, and upright.



Native of Portugal.
In bloom from October till December.

REFERENCE.

1. An umbel of flowers.
2. The empalement magnified.
3. The chives and pointal, one tip magnified.
4. Seed-bud and pointal, magnified.



E. multiflora

ERICA mutabilis.

DESCRIPTIO.

Antheræ sub-exertæ, basi bicornes. Flores terminales, in umbellis irregularibus, plerumque horizontaliter siti. Pedunculi longissimi, capillares, colorati, glandulis tecti. Calyx lanceolatus, coloratus, glandulosus, adpressus. Corolla tubulosa, costata, leviter pilosa, imprimis saturate purpurea, denique pallidior. Folia terna, quaterna, ad basin plantæ quina, supra convexa, subtus sulcata, margine pilis longis obsita. Caulis fruticosus. Rami numerosi, erecto-patentes.

DESCRIPTION.

Tips just without the blossom, with two horns at their base. Flowers terminate the branches in irregular umbels, mostly in a horizontal direction. Footstalks very long, thread-shaped, coloured, and covered with glands. The empalement is lance-shaped, coloured, glandular, and pressed to the flower. Blossom tubular, ribbed, and slightly hairy, of a deep purple at first, and afterwards paler. Leaves by threes and fours, and towards the base of the plant they grow in fives, convex on their upper surface, and furrowed beneath, with long hairs on their margins. Stem shrubby. The branches are numerous, and between erect and spreading.



Native of the Cape of Good Hope.
In bloom from February till October.

REFERENCE.

1. A leaf magnified.
2. Chives and pointal, one tip magnified.
3. Seed-bud and pointal, summit magnified.
4. The seed-bud magnified.



ERICA odorata.

DESCRIPTIO.

Antheræ muticæ, inclusæ. Flores numerosi, sub apices ramorum umbellati. Pedunculi flore triple longiores, viscosi, bracteis tribus adpressis. Corolla campanulata, magna, alba, viscosa, lucida, melliodora: laciniis rotundatis, reflexis. Folia sparsa, sæpe vero quaterna, linearia, margine glanduloso-denticulata, truncata: petiolis longis. Caulis fruticosus, pedalis. Rami simplices, longi.

DESCRIPTION.


Tips beardless, and within the blossom. Flowers numerous, growing in umbels at the ends of the branches. The footstalks thrice the length of the flowers, clammy, with three floral leaves pressed to them. Blossom bell-shaped, large, white, viscous, shining, and honey-scented: the segments of the border are rounded, and reflexed. Leaves scattered, but sometimes by fours, linear, appearing toothed by small glands at the edge, and seeming cut off at the ends. Footstalks long. Stem shrubby, a foot high. The branches are simple, and long.



Native of the Cape of Good Hope.
In bloom from April till July.

REFERENCE.

1. A leaf shown from the under side, magnified.
2. The empalement.
3. Chives and pointal, summit magnified.
4. The seed-bud magnified.



ERICA oppositifolia.

DESCRIPTIO.

Antheræ muticæ, inclusæ. Flores in ramulis terminales, plerumque bini. Pedunculi breves, bracteis duabus minutis adpressis. Calyx magnus, ovatus. Corolla urceolata, alba: laciniis reflexis. Folia binatim opposita, triquetra, nitida, adpressa. Caulis semipedalis, filiformis, ad basin ramosissimus, flexuosus.

DESCRIPTION.

Tips beardless, and within the blossom. Flowers terminate the branches mostly by twos. Footstalks short, having two small floral leaves pressed to them. The empalement is large, and egg-shaped. Blossom pitcher-shaped, and white. The segments of the border are reflexed. Leaves grow opposite in pairs, three-sided, shining, and pressed to the stem. Stem half a foot high, and thread-shaped: the lower part of the plant very much branched and flexuose.



Native of the Cape of Good Hope.
In bloom from April till July.

REFERENCE.

1. The empalement.
2. A blossom.
3. The chives and pointal, one tip magnified.
4. Seed-bud and pointal, summit magnified.
5. Seed-bud magnified.



E. oppositifolia

ERICA oppositifolia, *major*.

DESCRIPTIO.

Antheræ muticæ, inclusæ. Flores in ramulis terminales, plerumque bini. Pedunculi breves, bracteis duabus minutis adpressis. Calyx magnus, ovatus, acutus. Corolla urceolata, alba: laciniis suberectis. Folia binatim opposita, triquetra, nitida, adpressa, magna. Caulis pedalis, filiformis. Rami et ramuli numerosi, erecti.

DESCRIPTION.

Tips beardless, and within the blossom. Flowers terminate the branches mostly by twos. Footstalks short, having two small floral leaves pressed to them. The empalement is large, egg-shaped, and pointed. Blossom pitcher-shaped, and white. The segments of the border are nearly upright. Leaves grow opposite in pairs, three-sided, shining, pressed to the stem, and large. Stem a foot high, and thread-shaped: the large and smaller branches are numerous and upright



Native of the Cape of Good Hope.
In bloom from March till May.

REFERENCE.

1. The empalement.
2. A blossom.
3. The chives and pointal, one tip magnified.
4. Seed-bud and pointal, summit magnified.



E. oppositifolia major

ERICA oppositifolia, *rubra*.

DESCRIPTIO.

Antheræ muticæ, inclusæ. Flores terminales, plerumque bini. Pedunculi breves, bracteis duabus. Perianthium magnum, ovatum, acutum. Corolla urceolata, rubra: limbo quadrilobo, sub-reflexo. Folia binatim opposita, triquetra, nitida, adpressa. Caulis semi-pedalis, filiformis, gracilis, ad basin ramosus. Ramuli conferti, flexuosi.

DESCRIPTION.


Tips beardless, and within the blossom. Flowers terminate the branches mostly in pairs. Footstalks short, having two floral leaves. Cup large, oval, and sharp-pointed. Blossom pitcher-shaped, and red, having a slightly reflexed four-lobed border. Leaves grow opposite in pairs, three-sided, shining, and pressed to the stem. Stem half a foot high, thread-shaped, and slender, branching to the bottom. The branches crowded together, and flexuose.



Native of the Cape of Good Hope.
In bloom from May till August.

REFERENCE.

1. The empalement.
2. A blossom.
3. The chives and pointal, one tip magnified.
4. Seed-bud and pointal, summit magnified.
5. Seed-bud magnified.
6. A branch with the involucrum of the flowers.



E. oppositifolia rubra.

ERICA Patersonia.

DESCRIPTIO.

Antheræ aristatæ, inclusæ. Flores plures, axillares et in medio caulis, e foliis recte emergentes; pedunculis fere nullis. Perianthium duplex. Corolla cylindrica, sub-pollicaris, glabra, lutea, ore arctata; limbo reflexo. Folia quaterna, linearia, incurvata, acuta, glabra. Caulis fruticosus, bipedalis, erectus; basi ramosus, dein simplicissime pyramidatus, ramulis brevissimis vestitus.

DESCRIPTION.

Tips bearded, within the blossom. Flowers numerous, growing close to and about the middle of the stem, peeping straight out from amongst the leaves, with scarce any footstalks. Cup double. Blossom cylindrical, near an inch long, smooth, yellow, and narrow at the mouth; border reflexed. Leaves by fours, linear, and incurved, sharp-pointed, and smooth. Stem shrubby, two feet high, and upright; branching at the base, then forming a simple pyramidal spike covered with very short branches.



Native of the Cape of Good Hope.
In bloom from March till August.

REFERENCE.

1. The empalement magnified.
2. The chives spread open, one tip magnified.
3. The shaft and its summit magnified.



E. Patersonia

ERICA pallens.

DESCRIPTIO.

Antheræ muticæ, inclusæ. Flores terminales, in umbellis ternis vel senis, sub-sessiles. Corolla tubulosa, luteo-alba, horizontaliter sita: oris laciniis rotundatis, erecto-patentibus. Folia terna, subulata, tomentosa, albo-virescentia. Caulis humilis, fruticosus, semi-pedalis. Rami et ramuli numerosi, patentes.

DESCRIPTION.


Tips beardless, and within the blossom. Flowers terminate the branches in umbels of from three to six, nearly sessile. Blossom tubular, of a yellowish white colour, and standing in a horizontal direction. Segments of the border roundish, and between erect and spreading. Leaves by threes, awl-shaped, downy, and of a whitish green. Stem low, shrubby, half a foot high. The small and large branches are numerous, and spreading.



Native of the Cape of Good Hope.
In bloom from April till July.

REFERENCE.

1. The empalement magnified.
2. The chives and pointal.
3. Seed-bud and pointal, summit magnified.
4. Seed-bud magnified.



E. pallens

ERICA pellucida.

DESCRIPTIO.

Antheræ muticæ, inclusæ. Flores in ramulis terminales, plerumque quaterni, cernui. Pedunculi longi, glandulosi, bracteis tribus lanceolatis, glandulosis, instructi. Corolla tubulato-ventricosa, pilosa, costata, alba, et pellucida; limbi laciniis semi-orbicularibus, sub-erectis. Folia quaterna, lanceolata, lucida, aculeata, sulco exarata. Caulis gracilis, sesquipedalis, hirsutus. Rami et ramuli hirsuti, flexuosi.

DESCRIPTION.

Tips beardless, and within the blossom. Flowers terminate the small branches mostly by fours, nodding. Footstalks long, glandular, and furnished with three lance-shaped, glandular floral leaves. Blossom of an inflated tubular form, hairy, ribbed, white, and transparent. The segments of the border are half round, and nearly upright. Leaves by fours, lance-shaped, shining, prickly, and deeply furrowed beneath. Stem slender, a foot and a half high, harshly haired. The large and small branches are hirsute and flexuose.



Native of the Cape of Good Hope.
In bloom from November till June.

REFERENCE.

1. The empalement magnified.
2. A chive magnified.
3. Seed-bud and pointal, summit magnified.
4. Seed-bud magnified.



ERICA pinifolia.

DESCRIPTIO.

Antheræ muticæ, inclusæ. Flores sub apices ramorum verticillatim conferti, horizontaliter siti. Pedunculi colorati. Corolla clavata, pollicaris, alba, pilis tomentosis densissime circumsessæ: oris laciniis revolutis, obsolete crenulatis. Folia sena vel octona, linearia, tremula, glabra, subtus sulcata, attenuata in petiolos capillares. Caulis erectus, sesquipedalis, parum ramosus. Ramuli simplices, foliis tecti.

DESCRIPTION.


Tips beardless, and within the blossom. Flowers grow near the ends of the branches in crowded whorls, and in an horizontal direction. Footstalks coloured. Blossom club-shaped, an inch long, white, and closely beset with downy hairs: the segments of the border rolled back, and slightly notched. Leaves by sixes or eights, linear, tremulous, and smooth, furrowed beneath, and tapering into hair-like footstalks. Stem upright, a foot and a half high, branching but little. The small branches are simple, and covered with leaves.



Native of the Cape of Good Hope.
In bloom from February till July.

REFERENCE.

1. The empalement magnified.
2. The chives and pointal, one tip magnified.
3. Seed-bud and pointal, summit magnified.
4. Seed-bud magnified.



E. pinifolia

By Andrew C. Bingham

ERICA planifolia.

DESCRIPTIO.

Antheræ exsertæ, basi bicornes. Flores sub apices ramorum, simplices, axillares. Pedunculi longi, colorati, glanduloso-pilosi. Corolla urceolata, purpurea: oris laciniis obtusis, erectis. Folia terna, spathulata, supra plana, subtus carinata: marginibus glandulosis, revolutis. Caulis fruticosus, sesquipedalis. Rami et ramuli numerosi, erecto-patentes.

DESCRIPTION.

Tips without the blossom, and two-horned at their base. Flowers grow near the ends of the branches, singly, from the insertion of the leaves. Footstalks long, coloured, and glandularly hairy. Blossom pitcher-shaped, and purple: the segments obtuse, and upright. Leaves by threes, spathula-shaped, flat on their upper surface, and keeled beneath: the margins are glandular, and rolled back. Stem shrubby, a foot and a half high. The large and small branches are numerous, and between erect and spreading.



Native of the Cape of Good Hope.
In bloom from August till November.

REFERENCE.

1. A leaf shown from the under side, magnified.
2. The empalement magnified.
3. Chives and pointal, one tip magnified.
4. Seed-bud and pointal, summit magnified.



E. planifolia

ERICA Plukenetia.

DESCRIPTIO.

Antheræ muticæ, exertæ, longissimæ. Flores sub-solitarii, pendentes a ramulis, in medio ramorum spicam laxam formantes. Pedunculi longi, colorati: bracteis tribus infra minutis. Corolla conica, purpurea: laciniis exiguis, erectis, luteo-virentibus. Folia terna, linearia, arcuata, acuta, glabra, fasciculata. Caulis erectus, bipedalis, fruticosus. Rami sub-simplices, ramulis brevissimis, confertis, tecti.

DESCRIPTION.


Tips beardless, without the blossom, and very long. Flowers nearly solitary, hanging down from the smaller branches, and forming a loose spike about the middle of the branches. Footstalks long, coloured: with three minute floral leaves on the lower part. Blossom conical, and purple. The segments of the border are small, upright, and of a yellowish green. Leaves by threes, linear, bowed, pointed, smooth, and bundled together. Stem upright, two feet high, and shrubby. The large branches are nearly simple, long, and covered with very short branches crowded together.



Native of the Cape of Good Hope.
In bloom from May till August.

REFERENCE.

1. The empalement magnified.
2. Chives and pointal, one tip magnified.
3. Seed-bud and pointal, summit magnified.
4. The seed-bud magnified.



ERICA Plukenetia, *albens*.

DESCRIPTIO.

Antheræ muticæ, exertæ, longissimæ. Flores sub-solitarii, pendentes a ramulis, in medio ramorum crescentes, et spicam laxam formantes. Pedunculi longi: bracteis tribus minutis ad basin instructis. Corolla conica, exalbida: laciniis erectis. Folia terna, fasciculata, linearia, arcuata, acuta, glabra. Caulis erectus, sesquipedalis, fruticosus. Rami sub-simplices, ramulis numerosis brevissimis tecti.

DESCRIPTION.


Tips beardless, without the blossom, and very long. Flowers nearly solitary, hanging down from the small branches, growing near the middle of the branches, and forming a loose spike. Footstalks long, and furnished at the base with three small floral leaves. Blossom conical and whitish. Segments of the border are upright. Leaves by threes, bundled together, linear, bowed, sharp-pointed, and smooth. Stem upright, a foot and a half high, and shrubby. The large branches nearly simple, and covered with numerous very short branches.



Native of the Cape of Good Hope.
In bloom from May till September.

REFERENCE.

1. The empalement.
2. The chives and pointal.
3. Seed-bud and pointal, summit magnified.
4. The seed-bud magnified.
5. Flower of a large variety.



ERICA racemifera.

DESCRIPTIO.

Antheræ aristatæ, inclusæ. Flores sub-terminales, racemosi, cernui, plerumque duo; pedunculis longissimis. Corolla sub-globosa, purpurea; oris laciniis erecto-patentibus. Folia sena, linearia, conferta, oblique verticillata, tremula. Caulis fruticosus, sub-pedalis. Rami virgati, sub-simplices.

DESCRIPTION.

Tips bearded, within the blossom. Flowers nearly terminate the branches in clusters, hanging down, mostly two together, with very long footstalks. Blossom nearly globular, and purple; the segments of the border are between erect and spreading. Leaves by sixes, linear, crowded, irregularly whorled, and tremulous. Stem shrubby, near a foot high. Branches twiggy, and nearly simple.

—

Native of the Cape of Good Hope.
In bloom from April till June.

REFERENCE.

1. An umbel of flowers.
2. The empalement magnified.
3. Chives and pointal.
4. A chive magnified.
5. Seed-bud and pointal, summit magnified.
6. Seed-bud magnified.



E. racemifera

by L. Anderson - Glasgow

ERICA Sebana, *nana*.

DESCRIPTIO.

Antheræ muticæ, exertæ, longissimæ. Flores duo, in apicibus ramulorum, dependentes. Pedunculi breves, nudi. Perianthium duplex. Corolla clavata, incurvata, rugosa, saturate aurantia; laciniis limbi erectis, incurvatis. Folia terna, linearia, arcuata, conferta, sub-trigona, glabra, apice acuta. Caulis erectus, robustus, humilis, ramosissimus.

DESCRIPTION.

Tips beardless, without the blossom, and very long. Flowers grow by twos, at the ends of the small branches, hanging down. The footstalks are short, and naked. Cup double. Blossom club-shaped, turned inwards, wrinkled, and of a deep orange colour; the segments of the border are upright, and turned inwards. Leaves by threes, linear, bowed, and crowded, nearly three-sided, smooth, and sharp-pointed. Stem upright, sturdy, low, and very much branched.



Native of the Cape of Good Hope.
In bloom from July till January.

REFERENCE.

1. The empalement magnified.
2. A blossom.
3. The chives spread open.
4. Seed-bud and pointal, summit magnified.



E. Sebana nana

ERICA Sebana, *spicata*.

DESCRIPTIO.

Antheræ muticæ, exertæ, longissimæ. Flores terni, in apicibus ramulorum, penduli; pedunculis brevissimis. Corolla clavata, incurvata, saturate coccinea, sulcata: laciniis limbi obtusis, incurvis. Folia terna, linearia, arcuata, conferta, reflexa, apice acuta. Caulis erectus, bipedalis. Rami pauci, ramulis brevissimis confertis tecti.

DESCRIPTION.

Tips beardless, without the blossom, and very long. Flowers grow by threes at the ends of the small branches, hanging down, with very short footstalks. Blossom club-shaped, turned inwards, of a deep scarlet colour, and furrowed. The segments of the mouth are blunt, and turned inwards. Leaves by threes, linear, curved, crowded, reflexed, and sharp-pointed. Stem upright, two feet high. The larger branches few, covered with small branches crowded together.



Native of the Cape of Good Hope.
In bloom from September till December.

REFERENCE.

1. The empalement magnified.
2. The chives and pointal, one tip magnified.
3. Seed-bud and pointal, summit magnified.
4. Seed-bud magnified.



E. Lebrana spicata

Ed. Anderson del. H. B. K. sculp.

ERICA sordida.

DESCRIPTIO.

Antheræ muticæ, exertæ. Flores sub-sessiles, in apicibus ramulorum, nunc solitarii, nunc bini, vel terni. Corolla tubulosa, clavata, curvata, pollicaris, sordide lutea, pilosa, oris laciniis recurvatis. Folia quaterna, linearia, pilosa, obtusa. Caulis fruticosus, sesqui-pedalis, erectus, apice pilosus.

DESCRIPTION.

Tips beardless, and without the blossom. Flowers nearly sessile, upon the ends of the smaller branches, growing solitarily, or by twos and threes. Blossom tubularly club-shaped, curved, an inch long, of a dirty yellow colour, and hairy: the segments of the mouth rolled back. Leaves grow by fours, linear, hairy, and blunt. Stem shrubby, a foot and a half high, upright, and hairy at the top.



Native of the Cape of Good Hope.
In bloom from August till December.

REFERENCE.

1. A leaf magnified.
2. The empalement magnified.
3. Chives and pointal, summit magnified.



ERICA speciosa.

DESCRIPTIO.

Antheræ aristatæ, inclusæ. Flores 2-4-nati, sed plerumque ternati, cernui, ramulos laterales terminantes. Pedunculi colorati. Corolla cylindrica, uncialis, curvata, viscosa, lucida, saturate et læte rubra, sed apicem versus viridis: laciniis superne patentibus. Folia terna, linearia, obtusa, pubescentia. Caulis fruticosus, bipedalis, erectus, ramosus.

DESCRIPTION.

Tips bearded, and within the blossom. Flowers in twos and fours, but mostly by threes, bowing, terminating the lateral branches. Footstalks coloured. Blossom cylindrical, an inch long, curved, viscous, and shining, of a deep bright red colour, but towards the apex green; with the upper part of the segments spreading. Leaves by threes, linear, blunt, and downy. Stem shrubby, two feet high, upright, and branching.



Native of the Cape of Good Hope.
In bloom from June till September.

REFERENCE.

1. The empalement.
2. The chives and pointal, one tip magnified.
3. The seed-bud and pointal, summit magnified.



E. speciosa

ERICA staminea.

DESCRIPTIO.

Antheræ muticæ. Filamenta capillaria, prælonga, extra flores pendula. Flores solitarii, terminales, in ramulis pernumerosis lateralibus ternatis. Corolla parva, luteo-albida: laciniis patentibus, rotundatis. Folia ternata, erecta, parva, carnosæ, lucida. Caulis fruticosus, bipedalis: ramis numerosis, verticillatis, patulis.

DESCRIPTION.


Tips beardless. Threads hair-like, very long, and hanging out of the flower. Flowers solitary, and terminal, on very numerous lateral ternate branches. Blossom small, of a yellow white colour, with segments spreading and rounded. Leaves by threes, erect, small, fleshy, and shining. Stem shrubby, two feet high, with numerous whorled branches slightly spreading.



Native of the Cape of Good Hope.
In bloom from June till September.

REFERENCE.

1. The empalement magnified.
2. A blossom.
3. Blossom of a white variety.
4. The chives and pointal, magnified.
5. Seed-bud and pointal, summit magnified.



E. staminea

ERICA tenuiflora, *alba*.

DESCRIPTIO.

Antheræ muticæ, inclusæ. Flores ad apices ramulorum quaterni, pedunculis brevibus. Corolla tubæformis, tenuissima: oris laciniis acuminatis, patentibus, albicantibus: tubo pellucido. Folia quaterna, linearia, obtusa, glabra, erecto-patentia. Caulis fruticosus, debilis, pedalis et ultra. Rami et ramuli filiformes, virgati.

DESCRIPTION.

Tips beardless, within the blossom. Flowers grow by fours at the ends of the small branches. Footstalks short. Blossom trumpet-shaped, and very slender. The segments of the border are sharp-pointed, spreading, and very white. The tube is of a transparent white. Leaves by fours, linear, blunt, smooth, and between erect and spreading. Stem shrubby, weak, a foot or more high. The large and small branches are thread-shaped, and twiggy.



Native of the Cape of Good Hope.
In bloom from July till December.

REFERENCE.

1. The empalement magnified.
2. Chives and pointal, summit magnified.
3. Seed-bud and pointal, summit magnified.
4. The seed-bud magnified.



ERICA thymifolia.

DESCRIPTIO.

Antheræ basi bicornes, sub-inclusæ. Flores axillares, solitarii; pedunculis basi bracteis duabus instructis. Corolla parva, urceolata, saturate rosea; laciniis expansis, subtus concavis. Folia ternata, cordata, ciliata, subtus glauca. Caulis fruticosus, spithamæus, dumosus, perflexus; ramulis filiformibus, expansis, decumbentibus.

DESCRIPTION.

Tips two-horned at the base, and just within the blossom. Flowers axillary and solitary, with the peduncles furnished with two floral leaves at the base. Blossom small, pitcher-shaped, of a deep rose colour, with the segments expanded, and concave beneath. Leaves in threes, heart-shaped, hairy, and glaucous beneath. Stem shrubby, a span high, bushy, very much bent; with the branches thread-shaped, expanded, and decumbent.



Native of the Cape of Good Hope.
In bloom from March till July.

REFERENCE.

1. A leaf magnified.
2. The empalement magnified.
3. A chive magnified.
4. Seed-bud and pointal, summit magnified.
5. Seed-bud magnified.



1867

ERICA tiaræflora.

DESCRIPTIO.

Antheræ muticæ, exertæ. Flores in ramulis terminales, terni, cernui. Pedunculi longi. Corolla tiaræformis, apice incarnata, ad basin albida; laciniis minutis, obtusis, erectis. Folia terna, obtusa, linearia, rigida, erecto-patentia. Caulis fruticosus, erectus, sub-sesquipedalis, ramosus. Rami terni, patentes, erecti.

DESCRIPTION.


Tips beardless, and without the blossom. Flowers terminate the branches by threes, nodding. Footstalks long. Blossom turban-shaped, flesh-coloured at the ends, and whitish at the base; the segments of the border are very small, obtuse, and upright. Leaves by threes, blunt, linear, harsh, and between erect and spreading. Stem shrubby, upright, near a foot and a half high, and branching. Branches by threes, spreading, and upright.



Native of the Cape of Good Hope.
In bloom from May till July.

REFERENCE.

1. Empalement and blossom, magnified.
2. A blossom without its empalement, magnified.
3. A chive magnified.
4. The seed-bud and pointal, summit magnified.



E. tiariflora

By the Hon. Mr. J. E. Gray

ERICA ventricosa.

DESCRIPTIO.

Antheræ basi bicornes, inclusæ. Flores terminales, numerosi, fastigiati, erecti, fasciculati. Pedunculi purpurei, basi bracteis binis parvis oppositis instructi. Corolla oblongo-ovata, apice arctata, alba, glabra, nitida: limbo æquali, sub-reflexo, in centro rubro. Folia quaterna, linearia, acuta, basi recurvata, apice adscendentia, ciliata, nitida. Caulis fruticosus, semi-pedalis, teres, dumosus.

DESCRIPTION.

Tips two-horned at their base, and within the blossom. Flowers terminal, numerous, level, upright, and in bunches. The footstalks are purple, and furnished with two small opposite floral leaves at their base. Blossom of an oblong ovate form, and narrowed towards the end, white, smooth, and shining: segments of the border equal, slightly reflexed, and red in the centre. Leaves by fours, linear, sharp-pointed, with a recurved base and an ascending point, ciliated, and shining. Stem shrubby, half a foot high, round, and bushy.



Native of the Cape of Good Hope.
In bloom from April till September.

REFERENCE.

1. The empalement.
2. Chives and pointal, one tip magnified.
3. Seed-bud and pointal, summit magnified.



E. ventricosa

By J. H. Maiden, F.R.S. (1914)

ERICA vestita, *coccinea*.

DESCRIPTIO.

Antheræ muticæ, sub-exertæ. Flores in ramulis mediis conferti, erecto-patentes. Pedunculi brevissimi. Corolla clavata, pollicaris, saturate coccinea; oris laciniis revolutis. Folia sena, conferta, linearia, tremula, glabra, attenuata in petiolos capillares. Caulis erectus, bipedalis, parum ramosus; ramuli simplices, foliis tecti.

DESCRIPTION.

Tips beardless, just without the blossom. Flowers are clustered together about the middle of the branches, between erect and spreading. Footstalks very short. Blossom club-shaped, an inch long, and of a deep scarlet colour; the segments of the mouth are rolled back. Leaves by sixes, crowded together, linear, tremulous, smooth, and tapering into hair-like footstalks. Stem upright, two feet high, seldom branching; the smaller branches are simple, and covered with leaves.




Native of the Cape of Good Hope.
In bloom from August till November.

REFERENCE.


1. The empalement magnified.
2. The chives spread open, one tip magnified.
3. Seed-bud and pointal, summit magnified.



1



2



3

E. vestita coccinea

in D. v. 116

in H. v. 116

ERICA vestita, *purpurea*.

DESCRIPTIO.

Antheræ muticæ, sub-exertæ. Flores in ramulorum mediis verticillati, recti; corollis clavatis, rectis, pollicaribus, purpureis. Folia sena, linearia, tremulenta, acuta, glabra, attenuata in petiolos capillares semi-unguiculares. Caulis fruticosus, bipedalis, erectus, basi simplex, dein verticillatim ramosus, ramis simplicibus, adscendentibus.

DESCRIPTION.

Tips beardless, just without the blossom. Flowers grow in whorls about the middle of the branches, straight out. Blossom club-shaped, straight out, an inch long, and purple. Leaves by sixes, linear, tremulous, sharp-pointed, smooth, and tapering into hair-like footstalks a quarter of an inch in length. Stem shrubby, two feet high, and upright, simple at the base, then branching out from a whorl into simple ascending branches.



Native of the Cape of Good Hope.
In bloom from July till January.

REFERENCE.

1. A leaf.
2. The empalement magnified.
3. The chives and pointal, one tip magnified.
4. Seed-bud and pointal, summit magnified.



E. vestita purpurea

by J. Anderson & C. H. S. S. S. S. S.

ERICA villosa.

DESCRIPTIO.

Antheræ muticæ, exertæ. Flores in extremis ramulis terminales, sub terni, penduli. Pedunculi longi, rubri. Perianthium lato-ovatum; margine calycis læte viridi. Corolla urceolata, villosa, alba; oris laciniis obtusis, erectis. Folia terna, linearia, obtusa; margine pilis longis hirta. Caulis fruticosus, flexuosus, purpurascens. Rami et ramuli filiformes, villosi.

DESCRIPTION.

Tips beardless, and without the blossom. Flowers grow at the extremity of the smaller branches, mostly by threes, hanging down. The footstalks are long, and of a red colour. Cup broadly egg-shaped; the edges of the cup are of a bright green colour. Blossom pitcher-shaped, villous, and white; the segments of the border are obtuse, and upright. Leaves by threes, linear, and blunt; the edges rough, with long hairs. Stem shrubby, flexuose, and of a purplish colour. The small and large branches are thread-shaped, and hairy.



Native of the Cape of Good Hope.
In bloom from February till June.

REFERENCE.

1. A leaf magnified.
2. The empalement magnified.
3. Chives and pointal, magnified.



E. villosa

ALPHABETICAL INDEX

TO THE

FIGURES IN VOL. IV.

- 151. Erica andromedæflora.
- 152. aristata.
- 153. aurea.
- 154. Blandfordia.
- 155. calycina *major*.
- 156. canaliculata.
- 157. canaliculata, *minor*.
- 158. cordata.
- 159. decora.
- 160. discolor.
- 161. echiflora.
- 162. ferruginea.
- 163. fragrans.
- 164. halicacaba.
- 165. hirta.
- 166. hirta *viridiflora*.
- 167. hyacinthoides.
- 168. incarnata.
- 169. incurva.
- 170. lachnæa *purpurea*.
- 171. Lambertia.
- 172. lucida.
- 173. Monsoniana.
- 174. mucosa.
- 175. multiflora.
- 176. mutabilis.
- 177. odorata.

- 178. oppositifolia.
 - 179. oppositifolia, *major*.
 - 180. oppositifolia *rubra*.
 - 181. Patersonia.
 - 182. pallens.
 - 183. pellucida.
 - 184. pinifolia.
 - 185. planifolia.
 - 186. Pluckenetia.
 - 187. Pluckenetia *albens*.
 - 188. racemifera.
 - 189. Sebana nana.
 - 190. Sebana spicata.
 - 191. sordida.
 - 192. speciosa.
 - 193. staminea.
 - 194. tenuiflora, *alba*.
 - 195. thymifolia.
 - 196. tiaræflora.
 - 197. ventricosa.
 - 198. vestita, *purpurea*.
 - 199. vestita, *coccinea*.
 - 200. villosa.
-
-

*** END OF THE PROJECT GUTENBERG EBOOK THE HEATHERY;
OR, A MONOGRAPH OF THE GENUS ERICA. VOL. 4 ***

Updated editions will replace the previous one—the old editions will be renamed.

Creating the works from print editions not protected by U.S. copyright law means that no one owns a United States copyright in these works, so the Foundation (and you!) can copy and distribute it in the United States without permission and without paying copyright royalties. Special rules, set forth in the General Terms of Use part of this license, apply to copying and distributing Project Gutenberg™ electronic works to protect the PROJECT GUTENBERG™ concept and trademark. Project Gutenberg is a registered trademark, and may not be used if you charge for an eBook, except by following the terms of the trademark license, including paying royalties for use of the Project Gutenberg trademark. If you do not charge anything for copies of this eBook, complying with the trademark license is very easy. You may use this eBook for nearly any purpose such as creation of derivative works, reports, performances and research. Project Gutenberg eBooks may be modified and printed and given away—you may do practically ANYTHING in the United States with eBooks not protected by U.S. copyright law. Redistribution is subject to the trademark license, especially commercial redistribution.

START: FULL LICENSE

THE FULL PROJECT GUTENBERG™ LICENSE

PLEASE READ THIS BEFORE YOU DISTRIBUTE OR USE THIS WORK

To protect the Project Gutenberg™ mission of promoting the free distribution of electronic works, by using or distributing this work (or any other work associated in any way with the phrase “Project Gutenberg”), you agree to comply with all the terms of the Full Project Gutenberg License available with this file or online at www.gutenberg.org/license.

Section 1. General Terms of Use and Redistributing Project Gutenberg electronic works

1.A. By reading or using any part of this Project Gutenberg electronic work, you indicate that you have read, understand, agree to and accept all the terms of this license and intellectual property (trademark/copyright) agreement. If you do not agree to abide by all the terms of this agreement, you must cease using and return or destroy all copies of Project Gutenberg electronic works in your possession. If you paid a fee for obtaining a copy of or access to a Project Gutenberg electronic work and you do not agree to be bound by the terms of this agreement, you may obtain a refund from the person or entity to whom you paid the fee as set forth in paragraph 1.E.8.

1.B. “Project Gutenberg” is a registered trademark. It may only be used on or associated in any way with an electronic work by people who agree to be bound by the terms of this agreement. There are a few things that you can do with most Project Gutenberg electronic works even without complying with the full terms of this agreement. See paragraph 1.C below. There are a lot of things you can do with Project Gutenberg electronic works if you follow the terms of this agreement and help preserve free future access to Project Gutenberg electronic works. See paragraph 1.E below.

1.C. The Project Gutenberg Literary Archive Foundation (“the Foundation” or PGLAF), owns a compilation copyright in the collection of Project Gutenberg electronic works. Nearly all the individual works in the collection are in the public domain in the United States. If an individual work is unprotected by copyright law in the United States and you are

located in the United States, we do not claim a right to prevent you from copying, distributing, performing, displaying or creating derivative works based on the work as long as all references to Project Gutenberg are removed. Of course, we hope that you will support the Project Gutenberg mission of promoting free access to electronic works by freely sharing Project Gutenberg works in compliance with the terms of this agreement for keeping the Project Gutenberg name associated with the work. You can easily comply with the terms of this agreement by keeping this work in the same format with its attached full Project Gutenberg License when you share it without charge with others.

1.D. The copyright laws of the place where you are located also govern what you can do with this work. Copyright laws in most countries are in a constant state of change. If you are outside the United States, check the laws of your country in addition to the terms of this agreement before downloading, copying, displaying, performing, distributing or creating derivative works based on this work or any other Project Gutenberg work. The Foundation makes no representations concerning the copyright status of any work in any country other than the United States.

1.E. Unless you have removed all references to Project Gutenberg:

1.E.1. The following sentence, with active links to, or other immediate access to, the full Project Gutenberg License must appear prominently whenever any copy of a Project Gutenberg work (any work on which the phrase “Project Gutenberg” appears, or with which the phrase “Project Gutenberg” is associated) is accessed, displayed, performed, viewed, copied or distributed:

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg™ License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

1.E.2. If an individual Project Gutenberg electronic work is derived from texts not protected by U.S. copyright law (does not contain a notice indicating that it is posted with permission of the copyright holder), the work can be copied and distributed to anyone in the United States without paying any fees or charges. If you are redistributing or providing access to a work with the phrase “Project Gutenberg” associated with or appearing on the work, you must comply either with the requirements of paragraphs 1.E.1 through 1.E.7 or obtain permission for the use of the work and the Project Gutenberg trademark as set forth in paragraphs 1.E.8 or 1.E.9.

1.E.3. If an individual Project Gutenberg electronic work is posted with the permission of the copyright holder, your use and distribution must comply with both paragraphs 1.E.1 through 1.E.7 and any additional terms imposed by the copyright holder. Additional terms will be linked to the Project Gutenberg License for all works posted with the permission of the copyright holder found at the beginning of this work.

1.E.4. Do not unlink or detach or remove the full Project Gutenberg License terms from this work, or any files containing a part of this work or any other work associated with Project Gutenberg.

1.E.5. Do not copy, display, perform, distribute or redistribute this electronic work, or any part of this electronic work, without prominently displaying the sentence set forth in paragraph 1.E.1 with active links or immediate access to the full terms of the Project Gutenberg License.

1.E.6. You may convert to and distribute this work in any binary, compressed, marked up, nonproprietary or proprietary form, including any word processing or hypertext form. However, if you provide access to or distribute copies of a Project Gutenberg work in a format other than “Plain Vanilla ASCII” or other format used in the official version posted on the official Project Gutenberg website (www.gutenberg.org), you must, at no additional cost, fee or expense to the user, provide a copy, a means of exporting a copy, or a means of obtaining a copy upon request, of the work in its original “Plain Vanilla ASCII” or other form. Any alternate format must include the full Project Gutenberg License as specified in paragraph 1.E.1.

1.E.7. Do not charge a fee for access to, viewing, displaying, performing, copying or distributing any Project Gutenberg works unless you comply with paragraph 1.E.8 or 1.E.9.

1.E.8. You may charge a reasonable fee for copies of or providing access to or distributing Project Gutenberg electronic works provided that:

- You pay a royalty fee of 20% of the gross profits you derive from the use of Project Gutenberg works calculated using the method you already use to calculate your applicable taxes. The fee is owed to the owner of the Project Gutenberg trademark, but he has agreed to donate royalties under this paragraph to the Project Gutenberg Literary Archive Foundation. Royalty payments must be paid within 60 days following each date on which you prepare (or are legally required to prepare) your periodic tax returns. Royalty payments should be clearly marked as such and sent to the Project Gutenberg Literary Archive Foundation at the address specified in Section 4, "Information about donations to the Project Gutenberg Literary Archive Foundation."
- You provide a full refund of any money paid by a user who notifies you in writing (or by e-mail) within 30 days of receipt that s/he does not agree to the terms of the full Project Gutenberg™ License. You must require such a user to return or destroy all copies of the works possessed in a physical medium and discontinue all use of and all access to other copies of Project Gutenberg™ works.
- You provide, in accordance with paragraph 1.F.3, a full refund of any money paid for a work or a replacement copy, if a defect in the electronic work is discovered and reported to you within 90 days of receipt of the work.
- You comply with all other terms of this agreement for free distribution of Project Gutenberg™ works.

1.E.9. If you wish to charge a fee or distribute a Project Gutenberg™ electronic work or group of works on different terms than are set forth in this agreement, you must obtain permission in writing from the Project Gutenberg Literary Archive Foundation, the manager of the Project

Gutenberg™ trademark. Contact the Foundation as set forth in Section 3 below.

1.F.

1.F.1. Project Gutenberg volunteers and employees expend considerable effort to identify, do copyright research on, transcribe and proofread works not protected by U.S. copyright law in creating the Project Gutenberg™ collection. Despite these efforts, Project Gutenberg™ electronic works, and the medium on which they may be stored, may contain “Defects,” such as, but not limited to, incomplete, inaccurate or corrupt data, transcription errors, a copyright or other intellectual property infringement, a defective or damaged disk or other medium, a computer virus, or computer codes that damage or cannot be read by your equipment.

1.F.2. LIMITED WARRANTY, DISCLAIMER OF DAMAGES - Except for the “Right of Replacement or Refund” described in paragraph 1.F.3, the Project Gutenberg Literary Archive Foundation, the owner of the Project Gutenberg™ trademark, and any other party distributing a Project Gutenberg™ electronic work under this agreement, disclaim all liability to you for damages, costs and expenses, including legal fees. YOU AGREE THAT YOU HAVE NO REMEDIES FOR NEGLIGENCE, STRICT LIABILITY, BREACH OF WARRANTY OR BREACH OF CONTRACT EXCEPT THOSE PROVIDED IN PARAGRAPH 1.F.3. YOU AGREE THAT THE FOUNDATION, THE TRADEMARK OWNER, AND ANY DISTRIBUTOR UNDER THIS AGREEMENT WILL NOT BE LIABLE TO YOU FOR ACTUAL, DIRECT, INDIRECT, CONSEQUENTIAL, PUNITIVE OR INCIDENTAL DAMAGES EVEN IF YOU GIVE NOTICE OF THE POSSIBILITY OF SUCH DAMAGE.

1.F.3. LIMITED RIGHT OF REPLACEMENT OR REFUND - If you discover a defect in this electronic work within 90 days of receiving it, you can receive a refund of the money (if any) you paid for it by sending a written explanation to the person you received the work from. If you received the work on a physical medium, you must return the medium with your written explanation. The person or entity that provided you with the defective work may elect to provide a replacement copy in lieu of a refund. If you received the work electronically, the person or entity providing it to

you may choose to give you a second opportunity to receive the work electronically in lieu of a refund. If the second copy is also defective, you may demand a refund in writing without further opportunities to fix the problem.

1.F.4. Except for the limited right of replacement or refund set forth in paragraph 1.F.3, this work is provided to you 'AS-IS', WITH NO OTHER WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR ANY PURPOSE.

1.F.5. Some states do not allow disclaimers of certain implied warranties or the exclusion or limitation of certain types of damages. If any disclaimer or limitation set forth in this agreement violates the law of the state applicable to this agreement, the agreement shall be interpreted to make the maximum disclaimer or limitation permitted by the applicable state law. The invalidity or unenforceability of any provision of this agreement shall not void the remaining provisions.

1.F.6. INDEMNITY - You agree to indemnify and hold the Foundation, the trademark owner, any agent or employee of the Foundation, anyone providing copies of Project Gutenberg™ electronic works in accordance with this agreement, and any volunteers associated with the production, promotion and distribution of Project Gutenberg™ electronic works, harmless from all liability, costs and expenses, including legal fees, that arise directly or indirectly from any of the following which you do or cause to occur: (a) distribution of this or any Project Gutenberg work, (b) alteration, modification, or additions or deletions to any Project Gutenberg work, and (c) any Defect you cause.

Section 2. Information about the Mission of Project Gutenberg

Project Gutenberg is synonymous with the free distribution of electronic works in formats readable by the widest variety of computers including obsolete, old, middle-aged and new computers. It exists because of the

efforts of hundreds of volunteers and donations from people in all walks of life.

Volunteers and financial support to provide volunteers with the assistance they need are critical to reaching Project Gutenberg's goals and ensuring that the Project Gutenberg collection will remain freely available for generations to come. In 2001, the Project Gutenberg Literary Archive Foundation was created to provide a secure and permanent future for Project Gutenberg and future generations. To learn more about the Project Gutenberg Literary Archive Foundation and how your efforts and donations can help, see Sections 3 and 4 and the Foundation information page at www.gutenberg.org.

Section 3. Information about the Project Gutenberg Literary Archive Foundation

The Project Gutenberg Literary Archive Foundation is a non-profit 501(c)(3) educational corporation organized under the laws of the state of Mississippi and granted tax exempt status by the Internal Revenue Service. The Foundation's EIN or federal tax identification number is 64-6221541. Contributions to the Project Gutenberg Literary Archive Foundation are tax deductible to the full extent permitted by U.S. federal laws and your state's laws.

The Foundation's business office is located at 41 Watchung Plaza #516, Montclair NJ 07042, USA, +1 (862) 621-9288. Email contact links and up to date contact information can be found at the Foundation's website and official page at www.gutenberg.org/contact

Section 4. Information about Donations to the Project Gutenberg Literary Archive Foundation

Project Gutenberg™ depends upon and cannot survive without widespread public support and donations to carry out its mission of increasing the number of public domain and licensed works that can be freely distributed in machine-readable form accessible by the widest array of equipment

including outdated equipment. Many small donations (\$1 to \$5,000) are particularly important to maintaining tax exempt status with the IRS.

The Foundation is committed to complying with the laws regulating charities and charitable donations in all 50 states of the United States. Compliance requirements are not uniform and it takes a considerable effort, much paperwork and many fees to meet and keep up with these requirements. We do not solicit donations in locations where we have not received written confirmation of compliance. To SEND DONATIONS or determine the status of compliance for any particular state visit www.gutenberg.org/donate.

While we cannot and do not solicit contributions from states where we have not met the solicitation requirements, we know of no prohibition against accepting unsolicited donations from donors in such states who approach us with offers to donate.

International donations are gratefully accepted, but we cannot make any statements concerning tax treatment of donations received from outside the United States. U.S. laws alone swamp our small staff.

Please check the Project Gutenberg web pages for current donation methods and addresses. Donations are accepted in a number of other ways including checks, online payments and credit card donations. To donate, please visit: www.gutenberg.org/donate.

Section 5. General Information About Project Gutenberg electronic works

Professor Michael S. Hart was the originator of the Project Gutenberg concept of a library of electronic works that could be freely shared with anyone. For forty years, he produced and distributed Project Gutenberg eBooks with only a loose network of volunteer support.

Project Gutenberg eBooks are often created from several printed editions, all of which are confirmed as not protected by copyright in the U.S. unless a

copyright notice is included. Thus, we do not necessarily keep eBooks in compliance with any particular paper edition.

Most people start at our website which has the main PG search facility:
www.gutenberg.org.

This website includes information about Project Gutenberg, including how to make donations to the Project Gutenberg Literary Archive Foundation, how to help produce our new eBooks, and how to subscribe to our email newsletter to hear about new eBooks.




E. Andromedaeflora

[back](#)



E. aristata

[back](#)



E. aurea.

[back](#)



[back](#)



E. calycina major

[back](#)



E. canaliculata


Ed. Leach & Thos. G. Smith

[back](#)



E. canaliculata minor

[back](#)




E. cordata

W. B. Andrews. England

[back](#)




[back](#)



E. Discolor

[back](#)




E. Echiiflora

W. B. Woodhouse & Son

[back](#)





[back](#)



E. fragrans

St. Andrews & Hayfield reg.

[back](#)




E. halicacaba

[back](#)



[back](#)



E. hirta var. *viridiflora*


[back](#)



E. Hyacinthoides

24. 18. 1840. 1840. 1840.

[back](#)



E. incarnata

[back](#)



1



2



3



4



5

E. incurva

1827

by H. Andrews & R. Brown



[back](#)



E. lachnaea purpurea

by R. Anderson & Th. G. G. G.


[back](#)



E. Lambertia

by H. Andrews & Threlkeld

[back](#)



E. lucida

D. Don

by F. Linderoth & J. Knight

[back](#)



E. Monsoniana

W. H. H. 117

by J. B. Sowerby & H. S. G. Sowerby

[back](#)



E. mucosa

W. G. Anderson

[back](#)




E. multiflora

[back](#)



[back](#)



[back](#)




E. oppositifolia

[back](#)



E. oppositifolia major

[back](#)



E. oppositifolia rubra.

[back](#)



E. Patersonia

[back](#)




E. pallens

[back](#)



[back](#)



E. pinifolia


By Andrew C. Bingham

[back](#)




E. planifolia

[back](#)



[back](#)



[back](#)



E. racemifera

by S. Anderson - Glasgow

[back](#)



E. Sebana nana

[back](#)



[back](#)



[back](#)



E. speciosa


[back](#)



2



3



4



5

E. staminea

[back](#)




E. tenuiflora alba.

[back](#)



E. thymifolia

[back](#)



E. tiariflora



By the Rev. J. D. Hooker, Esq. & J. D. Hooker, Esq.

[back](#)



E. ventricosa

[back](#)



E. vestita coccinea

in D. v. 116

in H. v. 116

[back](#)



E. vestita purpurea

by J. Anderson & C. H. S. S. S. S. S.

[back](#)



E. villosa

[back](#)