


The Botanical Magazine, Vol. 09


William Curtis



Project Gutenberg

The Botanical Magazine, Vol. 09

William Curtis



Project Gutenberg

The Project Gutenberg eBook of The Botanical Magazine, Vol. 09

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

Title: The Botanical Magazine, Vol. 09

Author: William Curtis

Release date: December 22, 2011 [eBook #38382]

Language: English

Other information and formats: www.gutenberg.org/ebooks/38382

Credits: Produced by Jason Isbell and the Online Distributed Proofreading Team at <http://www.pgdp.net>

*** START OF THE PROJECT GUTENBERG EBOOK THE
BOTANICAL MAGAZINE, VOL. 09 ***

THE BOTANICAL MAGAZINE;

OR,

Flower-Garden Displayed:

IN WHICH

The most Ornamental FOREIGN PLANTS, cultivated in the Open Ground, the Greenhouse, and the Stove, are accurately represented in their natural Colours.

TO WHICH ARE ADDED,

Their Names, Class, Order, Generic and Specific Characters, according to the celebrated LINNÆUS; their Places of Growth, and Times of Flowering:

TOGETHER WITH

THE MOST APPROVED METHODS OF CULTURE.

A WORK

Intended for the Use of such LADIES, GENTLEMEN, and GARDENERS, as wish to become scientifically acquainted with the Plants they cultivate.

By **WILLIAM CURTIS**,

Author of the *FLORA LONDINENSIS*.

VOL. IX.

"But softer tasks divide Florella's hours;
"To watch the buds just op'ning on the day;
"With welcome shade to screen the languid flowers,
"That sicken in the summer's parching ray.

"Oft will she stoop amidst her evening walk,
"With tender hand each bruised plant to rear;
"To bind the drooping lily's broken stalk,
"And nurse the blossoms of the infant year."

Mrs. BARBAULD.

LONDON:

PRINTED BY STEPHEN COUCHMAN,

For W. CURTIS, N^o 3, *St. George's Crescent*, Black-Friars-Road;
And Sold by the principal Booksellers in Great-Britain and Ireland.

M DCC XCV.

**CONVOLVULUS LINEARIS. NARROW-LEAVED
CONVOLVULUS.**

Class and Order.

PENTANDRIA MONOGYNIA.

Generic Character.

Corolla campanulata, plicata. *Stigmata* 2. *Caps.* 2-locularis:
loculis dispermis.

Specific Character.

CONVOLVULUS *linearis* caulibus erectis fruticosis, foliis
linearibus acutis piloso-sericeis, floribus terminalibus
umbellato-paniculatis, calycibus pilosis.



The plant here represented has long been cultivated as a greenhouse plant in this country under the name of *Convolvulus Cantabrica*, but it differs so essentially from that plant, as figured and described by Prof. JACQUIN in his *Flora Austr.* and accords so little with the other species described by LINNÆUS, that we have been induced to regard it as a perfectly distinct

species; in most points it agrees with *Convolvulus Cneorum*, but differs in having leaves much narrower, more pointed, and less silky.

It strikes most readily from cuttings, is a hardy greenhouse plant, and flowers during most of the Summer, qualities which many of the modern and more shewy greenhouse plants cannot boast.

The precise time of its introduction here, together with its particular place of growth, we have not as yet been able satisfactorily to ascertain.

[290]

AMARYLLIS LUTEA. YELLOW AMARYLLIS.

Class and Order.

HEXANDRIA MONOGYNIA.

Generic Character.

Corolla hexapetaloidea, irregularis. *Filamenta* fauci tubi inserta, declinata, inæqualia proportione vel directione. *Linn. Fil.*

Specific Character and Synonyms.

AMARYLLIS *lutea* spatha indivisa obtusa, flore sessili, corolla campanulata erecta basi breve tubulosa, staminibus erectis, alternis brevioribus. *Linn. Fil. Ait. Kew. v. 21. p. 415.*

COLCHICUM luteum majus. *Bauh. Pin. p. 69.*

NARCISSUS autumnalis major. The greater Autumne or Winter Daffodill. *Park. Parad. p. 77. 75. f. 7.*



The *Amaryllis lutea* is a hardy perennial bulbous plant, a native of Spain, and other of the more Southern parts of Europe, and was cultivated in our Gardens in the time of GERARD, and PARKINSON.

Flora, who commences her revolutionary reign, by enlivening the flower border with the Spring Crocus, and its numerous varieties, terminates it with flowers equally pleasing, and of similar hues; thus we have the present

plant, the Saffron Crocus, and the Colchicum, flowering nearly at the same time, from the end of September, through October, and sometimes part of November.

Similar as the Amaryllis is to the yellow Spring Crocus, in the colour, and form of its flowers, it differs obviously in the number of its stamina, the breadth of its leaves, and the size and colour of its root.

Authors describe it as varying in size, in the breadth of its leaves, the height of its flowers, and multiplication of the Corolla.

The Dutch Florists export it under the title of yellow Colchicum, following the name of some of the old writers.

It succeeds best in a soil moderately moist, in which it increases considerably by offsets, and flowers to the most advantage when the roots have remained for some few years undisturbed in the same spot.

[291]

CAPPARIS SPINOSA. THE CAPER SHRUB.

Class and Order.

POLYANDRIA MONOGYNIA.

Generic Character.

Cal. 4-phyllus, coriaceus. *Petala* 4. *Stamina* longa. *Bacca* corticosa, unilocularis, pedunculata.

Specific Character and Synonyms.

CAPPARIS *spinosa* pedunculis unifloris solitariis, stipulis spinosis, foliis annuis, capsulis ovalibus. *Linn. Syst. Vegetab. ed. 14. Murr. p. 487. Ait. Kew. v. 2. p. 221.*

CAPPARIS *spinosa* fructu minore, folio rotundo. *Bauh. Pin. p. 480.*



We are happy in having it in our power to lay before our readers a representation of the Caper shrub, whose blossoms are rarely seen in this country, though its flower-buds are in very general use as a pickle; indeed, so great is their consumption, that they form a very considerable article of commerce.

The plant grows spontaneously in the more southern parts of Europe, especially in Italy and the Levant; in its wild state it forms a shrub of low growth, having numerous, spreading, spinous branches, somewhat thickly beset with smooth roundish leaves; the blossoms grow alternately on the branches, and when the plant begins to flower, one opens generally every other morning, but so delicate are its parts, that on a hot summer's day it fades before noon: the petals are white; the filaments, which are extremely numerous, are white below, and of a rich purple above; in these the beauty of the flower chiefly consists, as in the pistillum or pointal does its great singularity; at first view, one would be led to conclude, that the part so conspicuous in the centre of the flower was the style terminated by the stigma in the usual way; but if we trace this part of the flower to a more advanced state, we shall perceive, that what we took for the style, was merely an elongation of the flower-stalk, and what we took for the stigma, was in reality the germen placed on it, crowned with a minute stigma, without any intervening style; this germen swells, turns downward, and ultimately becomes the seed-vessel, rarely ripening in this country.

MILLER observes, that these plants are with difficulty preserved in England, for they delight to grow in crevices of rocks, and the joints of old walls and ruins, and always thrive best in an horizontal position; so that when they are planted either in pots or the full ground, they rarely thrive, though they may be kept alive for many years.

It flowers in May and June, and is usually raised from seeds.

Mr. AITON regards it as a greenhouse plant, and informs us that it was cultivated by GERARD in 1596.

[292]

**PASSERINA GRANDIFLORA. GREAT-FLOWERED
PASSERINA.**

Class and Order.

OCTANDRIA MONOGYNIA.

Generic Character.

Cal. 0. *Cor.* 4-fida. *Stamina* tubo imposita. *Sem.* 1. corticatum.

Specific Character and Synonyms.

PASSERINA *grandiflora* glaberrima, foliis oblongis acutis
concavis extrinsecus rugosis, floribus terminalibus sessilibus
solitariis. *Linn. Suppl. Pl. p.* 226.



The *Passerina* here figured, distinguished from all the other known species by the largeness of its flowers, is described in the *Suppl. Pl.* of the younger LINNÆUS, but not enumerated in the *Hortus Kewensis* of Mr. AITON: it is indeed a plant recently introduced to this country from the Cape; we saw it last Summer in great perfection, at Messrs. LEE and KENNEDY'S, Hammersmith; it forms a small neat shrub, somewhat like the *Phyllica*

ericoides, is a hardy greenhouse plant, flowering in May and June, and increased without difficulty from cuttings.

[293]

CATANANCHE CÆRULEA. BLUE CATANANCHE.

Class and Order.

SYNGENESIA POLYGAMIA ÆQUALIS.

Generic Character.

Recept. paleaceum. *Cal.* imbricatus. *Pappus* aristatus, caliculo 5 seto.

Specific Character and Synonyms.

CATANANCHE cærulea squamis calicis inferioribus ovatis. *Linn.*
Syst. Vegetab. ed. 14. Murr. p. 722. Ait. Kew. v. 3. p. 134.

CHONDRILLA cærulea cyani capitulo. *Bauh. Pin. 130.*



The *Catananche caerulea* is a native of the South of France, where it grows in hilly situations that are stony: it is a perennial herbaceous plant, moderately hardy, and has long been cultivated in our gardens, Mr. AITON says, by PARKINSON in 1640: MILLER, who treats of it in his Dictionary, describes it as a pretty ornament to a garden, and one that is easily kept within bounds; there is certainly much about it to excite our admiration, more especially in the structure of the calyx, and the florets: the flowers,

which are of a pale blue colour with a dark eye, make their appearance from July to October.

It is propagated by seeds, which MILLER recommends to be sown in the Spring; the seedlings should be transplanted in the Autumn, into the borders where they are to remain; it may also be increased by slips: the plant requires a situation moderately dry, and is most productive of flowers and seeds when it stands long in one spot.

In the 14th *edit.* of the *Systema Vegetab.* of Prof. MURRAY, mention is made of a variety with double flowers, which we believe has not been seen in this country.

[294]

AMARYLLIS SARNIENSIS. GUERNSEY AMARYLLIS.

Class and Order.

HEXANDRIA MONOGYNIA.

Generic Character.

Cor. hexapetaloidea irregularis. *Filamenta* fauci tubi inserta declinata inæqualia proportionem vel directionem. *Linn. fil. Ait. Kew. p. 415.*

Specific Character and Synonyms.

AMARYLLIS *sarniensis*, petalis linearibus planis, staminibus pistilloque rectiusculis corolla longioribus, stigmatibus partitis revolutis. *Linn. fil. Ait. Kew. v. 1. p. 420. Thunb. Jap. p. 131.*

LILIUM *sarniense*. *Dougl. Monogr. t. 1, 2.*

NARCISSUS *japonicus* rutilo flore. *Corn. Canad. Kæmpf. Amæn. p. 872.*



The Guernsey Lily, as it is most commonly called, is originally a native of Japan; where it is described to grow by KÆMPFER and THUNBERG, who visited that island, the latter says on the hills about Nagasaki, from thence roots are said to have been introduced to the garden of JOHANNES MORINUS at Paris, in which it flowered, October 1634: its introduction to this country, which was subsequent to that date, as Dr. DOUGLASS relates in his *Monographia* on this plant, "happened by a very singular melancholy

accident, of which Dr. MORISON, who no doubt had it from some persons then residing in Guernsey, gives us the following account: A Dutch or English ship, it is uncertain which, coming from Japan, with some of the roots of this flower on board, was cast away on the island of Guernsey; the roots were thrown upon a sandy shore, and so by the force of the winds and waves, were soon buried in sand; there they remained for some years, and afterwards, to the great surprise and admiration of the inhabitants, the flowers appeared in all their pomp and beauty." Some of these soon made their appearance in this country: Mr. AITON relates, that the plant was cultivated here in 1659, by General LAMBERT, at Wimbledon.

Fatal as Guernsey proved to the unfortunate mariners, it afforded the roots of our plant a soil and situation apparently congenial to their own; in that island they have flourished ever since, there they are propagated in the open borders of the flower-garden with the least possible trouble, flowering most readily, but we believe never producing any ripe seeds; from thence most of the roots which flower with the curious here, are yearly imported in the Autumn.

In Guernsey, the cold of the Winter is far less intense than with us; many of those plants which we keep in our greenhouses, stand with them in the open ground; the superior mildness of the climate enables them to cultivate this plant with more success than we can do, even perhaps with all the expence and trouble to which we might subject ourselves; to such, however, whose situations may be favourable, and who may be fond of making experiments, we recommend the perusal of FAIRCHILD'S Directions, a practical Gardener of great ingenuity, and who appears to have had much experience in the culture of this plant^[A].

It is usual to plant the imported bulbs in pots of sand, or light loam, as soon as they arrive, and place them in the parlour window, or greenhouse; they blossom in September and October; the flowers, which continue about a month in perfection, are inodorous, but make up for that deficiency by the superior splendour of their colours: Dr. DOUGLASS thus describes them, each flower when in its prime looks like a fine gold tissue wrought on a rose-coloured ground, but when it begins to fade and decay, it looks more like a silver tissue, or what they call a pink colour: when we look upon the flower in full sun-shine, each leaf appears to be studded with thousands of little

diamonds, sparkling and glittering with a most surprising and agreeable lustre; but if we view the same by candle-light, these numerous specks or spangles look more like fine gold dust.

Both KÆMPFER and THUNBERG agree, that the Japanese regard the root as poisonous.

[A] "They love a light earth, made with dung and sand, and a little lime rubbish with it does very well, it keeps the roots sound; for if the earth be too stiff or wet, you may keep them for many years before they blow. If they are in pots, they should be put in the house in Winter, to keep them from the severe frosts, which are apt to rot the roots. The time of moving them is when they have no leaves on the root, that is from June to August: those that come with six leaves this year, seldom fail blowing the next year: they need not be put in fresh earth above once in two or three years: by this method of management I have had the same roots blow again in four years time. The many miscarriages that happen to the Guernsey Lily, are by letting the leaves be killed by the fierceness of the frost in Winter, or by cutting them off, as some people do, when they are green, which will so much weaken the plants, that they may keep them twenty years and not have them blow; by the above management, where there is a stock, there will be continually some blowing.

"MILLER recommends for these roots the following compost: Take a third-part of fresh virgin earth from a pasture-ground which is light, then put near an equal part of sea-sand, to which you should add rotten dung and sifted lime rubbish, of each an equal quantity."

The great business in the culture of this flower, next to a proper soil and situation, seems to consist in giving the plant as much air as possible, and in preserving the foliage in the Winter from the injury of frost.

[295]

AGROSTEMMA CÆLI ROSA. SMOOTH-LEAV'D COCKLE, or ROSE CAMPION.

Class and Order.

DECANDRIA PENTAGYNIA.

Generic Character.

Cal. 1-phyllus, coriaceus. *Petala* 5 unguiculata: limbo obtuso indiviso. *Caps.* 1 locularis.

Specific Character and Synonyms.

AGROSTEMMA *C[oe]lli* rosa glabra, foliis lineari-lanceolatis, petalis emarginatis coronatis. *Linn. Syst. Vegetab. ed. 14. p. 435.* *Ait. Kew. v. 2. p. 116.*

LYCHNIS segetum, Nigellastrum minus glabrum dicta, flore eleganter rubello. *Moris. Hist. 2. p. 543. s. 5. t. 22. f. 32.*

LYCHNIS pseudomelanthis similis africana glabra angustifolia. *Herm. Leyd. 391. t. 393.*



Mr. AITON informs us in his *Hortus Kewensis*, that the charming annual here figured, the liveliness of whose colours no paint can express, was cultivated by MILLER in 1739; seeing it is a plant of such beauty, and honoured with so distinguished an appellation, it is singular that it should not by this time have made its way more generally into our gardens.

The Cockle of our corn-fields is an ornamental plant, the present species resembles it; but while the plant itself is much smaller, its flowers are proportionably larger, and their colours more vivid.

It is an annual of ready growth, a native of Sicily and the Levant, flowering in July and August, and ripening its seeds in September and October.

It appears to most advantage when several plants of it grow together; the best mode, therefore, is to sow about a dozen seeds early in April on the several spots of the flower-border where you intend they shall remain; no other care is necessary than to keep the plants free from weeds and vermin.

The *Agr. Cæli rosa* of *Miller's Gard. Dict. ed. 6 4to.* is the *coronaria*.

[296]

SEMPERVIVUM TORTUOSUM. GOUTY HOUSELEEK.

Class and Order.

DODECANDRIA DODECAGYNIA.

Generic Character.

Cal. 12-partitus. *Petala* 12. *Capsulæ* 12 polyspermæ.

Specific Character.

SEMPERVIVUM *tortuosum* foliis obovatis subtus gibbis villosis,
nectariis bilobis. *Ait. Kew. v. 2. p. 148.*



We know of no figure of this plant, the first and only account of which is to be found in the *Hort. Kew.* of Mr. AITON; there it is specifically described, and from thence we discover that it was introduced from the Canary Islands, where it is a native, by Mr. MASSON, in 1779.

It is a shrubby plant of low growth, producing numerous fleshy leaves growing thickly together, which being evergreen, and making a pretty appearance the year through, render the plant worthy a place in all general

collections at least, of greenhouse plants; and though it cannot vie with many of the more showy high-priced novelties, it is an abiding plant, not subject to casualties, while many of those are here to day and gone to morrow.

It throws up its flowering stems, supporting numerous, starry, stonecrop-like flowers, in July and August, and is most readily propagated by cuttings.

It is one of those species of Houseleek which connect the genera *Sedum* & *Sempervivum*.

[297]

DIANTHUS SUPERBUS. SUPERB PINK.

Class and Order.

DECANDRIA DIGYNIA.

Generic Character.

Cal. cylindricus, 1-phyllus: basi squamis 4. *Petala* 5 unguiculata.
Caps. cylindrica, 1-locularis.

Specific Character and Synonyms.

DIANTHUS *superbus* floribus paniculatis: squamis calycinis brevibus acuminatis, corollis multifido-capillaribus, caule erecto. *Linn. Syst. Vegetab. ed. 14. Murr. p. 418.*

CARYOPHYLLUS simplex alter, flore laciniato odoratissimo. *Bauh. Pin. 210.*

CARYOPHYLLUS plumarius Austriacus sive superba Austriaca. The feathered Pinke of Austria. *Park. Parad. p. 316. 4.*



Many of the plants of this genus diffuse an agreeable odour, which renders them most desirable objects for the flower-garden: this quality the present species possesses in a superior degree to most others; a few of its flowers communicate to a nosegay a delicate and most delicious smell, or placed in a vial of water they will even scent a small apartment^[B]: it is to be regretted, however, that the blossoms, unless placed in water, from their extreme delicacy, flag soon after they are gathered.

It may be doubted whether the *Dianthus superbus* of Miller's *Dict. ed. 6. 4to.* be our plant; if it be, the description is not drawn up with that accuracy which distinguishes his descriptions in general; the mode of culture, however, which he recommends is strictly applicable to it, as the plant rarely continues in vigour more than two years, and as it is in its greatest beauty the first year of its flowering, he recommends that young plants should be annually raised for succession from seeds, which are plentifully produced; the seeds of this plant ought therefore to be kept in the shops with annuals and biennials.

The *Dianthus superbus* is a native of Germany, Switzerland, France, and Denmark: CLUSIUS found it growing in the moist meadows about Vienna, and on the borders of woods adjoining to such, with some of its flowers white, others purplish; PARKINSON describes them of these two colours, but says the most ordinary with us are pure white, which is contrary to what we now find them: they are rarely produced before August, from which period they will continue frequently to blossom till October.

The Spring is the best time for sowing its seeds; the plants require no very nice or particular treatment.

[B] This fragrance has been noticed by all the old authors who have treated of the plant: CLUSIUS describes the flowers as *suavissimi odoris et è longinquo nares ferientis*, of which words PARKINSON'S are almost a literal translation "of a most fragrant sent, comforting the spirits and senses afarre off."

[298]

ORIGANUM DICTAMNUS. DITTANY OF CRETE.

Class and Order.

DIDYNAMIA GYMNOSPERMIA.

Generic Character.

Strobilus tetragonus, spicatus, calyces colligens.

Specific Character and Synonyms.

ORIGANUM *Dictamnus* foliis inferioribus tomentosis, spicis
nutantibus. *Linn. Syst. Vegetab. ed. 14. Murr. p. 541. Ait.*
Kew. v. 2 p. 311. Dittany of Candia.

DICTAMNUS creticus. *Bauh. Pin. p. 222.*



By the name of Dittany of Crete, the species of *Origanum* here figured, has long been known in this country as a medicinal plant; to the purposes of physic it still indeed continues to be applied, as imported in a dried state from the Levant: when bruised, the whole plant gives forth an aromatic fragrance, highly grateful; as an ornamental plant, it has also been long, and is now, very generally cultivated in this country. TURNER, whose Herbal was printed in 1568, writes thus concerning it, "I have sene it growynge in

England in Maister Riches garden naturally, but it groweth no where ellis that I know of, saving only in Candy." As at this period no idea was entertained of a greenhouse, the plant must have been cultivated in the open ground, where it would doubtless grow readily, if secured from the severity of the weather, it being more hardy than many plants usually kept in greenhouses.

This plant is at all times ornamental, but more particularly so when in flower, in which state it appears during most of the summer and autumnal months.

It is usually increased by cuttings, which strike readily.

[299]

HERMANNIA ALNIFOLIA. ALDER-LEAVED HERMANNIA.

Class and Order.

MONADELPHIA PENTANDRIA.

Generic Character.

Pentagyna. Caps. 5-locularis. Petala basi semitubulata, obliqua.

Specific Character and Synonyms.

HERMANNIA *alnifolia* foliis cuneiformibus lineatis plicatis crenato-emarginatis. *Linn. Syst. Vegetab. ed. 14. Murr. p. 610. Ait. Kew. v. 2. p. 412.*

ARBUSCULA africana tricapsularis ononidis vernæ singulari folio. *Pluk. Mant. 14. t. 239. f. 1.*



Hermannia is a genus of plants named in honour of Dr. PAUL HERMAN, a Dutch Botanist of great celebrity, author of the *Paradisus Batavus*, and other valuable works: twenty-six species are enumerated in the 13th edition of the *Syst. Naturæ* of LINNÆUS by Prof. GMELIN, and eight in the *Hortus Kewensis* of Mr. AITON; most of those in the latter work are cultivated in the nurseries near town: they form a set of the more hardy greenhouse plants,

grow readily, and flower freely; their blossoms are for the most part yellow, and have a considerable affinity with those of the *Mahernia*.

The present species flowers very early in the spring, from February to May, producing a great profusion of bloom during that period; is a native of the Cape, and was cultivated by Mr. MILLER, in 1728.

It rarely ripens its seeds with us, but is readily increased by cuttings.

The nurserymen near town regard this plant as the *grossularifolia* of LINNÆUS, calling another, equally common species, with longer and narrower leaves, *alnifolia*, and which does not appear to be described by LINNÆUS or mentioned by Mr. AITON; our plant accords exactly with the Linnæan description of *alnifolia*, and there is we think no doubt of its being the *alnifolia* of the *Hortus Kewensis*, and Mr. MILLER'S *Dictionary*.

[300]

GNAPHALIUM EXIMIUM. GIANT CUDWEED.

Class and Order.

SYNGENESIA POLYGAMIA SUPERFLUA.

Generic Character.

Recept. nudum. *Pappus* plumosus vel capillaris. *Cal.* imbricatus, squamis marginalibus rotundatis, scariosis, coloratis.

Specific Character and Synonyms.

GNAPHALIUM *eximium* foliis sessilibus ovatis confertis erectis tomentosis, corymbo sessili. *Linn. Mant. Pl. p. 573. Syst. Nat. ed. 13. Gmel.*

ELYCHRYSUM *africanum* foliis lanceolatis integris tomentosis decurrentibus, capitulus congestis ex rubello aureis. *Edw. Av. t. 183.*



In the summer of 1794, towards the end of July, the *Gnaphalium* here figured, the most magnificent and shewy of all the species hitherto introduced to this country, flowered in great perfection at Messrs. LEE and KENNEDY'S, Hammersmith: Mr. LEE informs me, that he raised it from seeds given him by Capt. WILLIAM PATERSON, author of a Narrative of four journeys into the country of the Hottentots, and Caffraria, and who has most laudably exerted himself in introducing many new and interesting plants to this country; this gentleman assured Mr. LEE, that the plant was found in a

wild state, five hundred miles from the Cape, on the borders of the Caffre country, from whence the natives bring bundles of the dried plant to the Cape as presents; in the state the plant has long since been imported from that fertile coast: if we mistake not, a specimen of this sort is figured in PETIVER'S works, and a coloured representation is given of it in EDWARDS'S History of Birds, taken from a dried plant, brought from the Cape, by Capt. ISAAC WORTH, in 1749.

The plants we saw were about a foot and a half high, the stalks shrubby, and but little branched; the foliage and flowers as represented on the plate.

Several of the Gnaphaliums it is well known are liable to be killed by moisture, especially in the winter season; during that time, this plant in particular, should be kept as dry as possible, and, if convenient, on a shelf, separate from the other plants of the greenhouse; when it is necessary to give it water, it should never come in contact with the foliage or flowers: with these precautions it may be kept very well in a good greenhouse, in which it should remain, even during summer.

It may be raised from seeds, and also from cuttings.

[301]

MELIANTHUS MINOR. SMALL MELIANTHUS, or HONEY- FLOWER.

Class and Order.

DIDYNAMIA ANGIOSPERMIA.

Generic Character.

Cal. 5-phyllus: folio inferiore gibbo. *Petala* 4: nectario infra infima. *Caps.* 4-locularis.

Specific Character and Synonyms.

MELIANTHUS *minor* stipulis geminis distinctis. *Linn. Syst. Vegetab. ed. 14. Murr. p. 581. Ait. Kew. v. 2. p. 368.*

MELIANTHUS *africanus minor foetidus. Comm. rar. 4. t. 4.*



There are few flowers that do not secrete from some kind of a glandular substance, honey, or nectar, to a greater or smaller amount; in those of the present genus, this liquid is particularly abundant, even dropping from the flowers of the *major*, in considerable quantity; in the present species it flows not so copiously, but is retained in the lower part of the blossom, and is of a dark brown colour, an unusual phenomenon.

There are only two species of this genus described, the *major* and the *minor*, both of which are cultivated in our nurseries; the *major* is by far the most common, the most hardy, and the most ornamental plant; its foliage indeed is peculiarly elegant: this species will succeed in the open border, especially if placed at the foot of a wall with a south or south-west aspect, taking care to cover the root to a considerable depth with rotten tan in severe frosts: the *minor* is always kept in the greenhouse, in which, when it has acquired a certain age, it flowers regularly in the spring, and constantly so, as far as we have observed of the plants in Chelsea Garden; Mr. AITON says in August, and COMMELIN the summer through.

The *Melianthus minor* grows to the height of three, four, or five feet; its stem, which is shrubby, during the flowering season is apt to exhibit a naked appearance, having fewer leaves on it at that period, and those not of their full size; but this, perhaps, may in some degree be owing to the plant's being placed at the back of others.

The foliage when bruised has an unpleasant smell.

It is a native of the Cape, and, according to Mr. AITON, was cultivated by the Duchess of BEAUFORT, in 1708; is propagated readily by cuttings.

[302]

MIMOSA MYRTIFOLIA. MYRTLE-LEAVED MIMOSA.

Class and Order.

POLYANDRIA MONOGYNIA.

Generic Character.

Flores mere masculi reliquis intersiti. *Cal.* 5-dentatus. *Cor.* 5-fida aut 0. *Stamina* 4-ocularis. *Legumen.* *Linn. Syst. Nat. ed. 13. Gmel.*

Specific Character.

MIMOSA *myrtifolia* foliis ovato-lanceolatis obliquis undulatis acuminatis margine cartilagineis: primordialibus pinnatis. *Smith Trans. Linn. Soc. v. 1. p. 252.*



The seeds of this species of *Mimosa* having been sent over in plenty, with some of the first vegetable productions of New South-Wales, and growing readily, the plant has been raised by many cultivators in this country; Mr. HOY, gardener to the Duke of NORTHUMBERLAND, produced a specimen of it in flower at a meeting of the Linnean Society in 1790; it is a shrub of quick

growth, and a ready blower: a plant of it in the stove of Chelsea-Garden has this year (May 10, 1795) produced ripe pods, and perfect seeds. In the greenhouse, where it flowers from February to April, the blossoms go off without shewing any tendency to produce fruit.

It is first described by Dr. SMITH, in the Transactions of the Linnean Society; the leaves in the plants that have fallen under our notice have not accorded exactly with those he has described, having neither been of a glaucous green colour, according to the usual acceptation of that term, nor very much undulated; and though those of an individual plant may have presented such an appearance, we are persuaded they do not do so generally when growing and in good health.

The foliage is usually edged with red, and the flowers are fragrant.

[303]

ERICA AMPULLACEA. FLASK HEATH.

Class and Order.

OCTANDRIA MONOGYNIA.

Generic Character.

Cal. 4-phyllus. *Cor.* 4-fida. *Filamenta* receptaculo inserta. *Antheræ* apice bifidæ, pertusæ. *Caps.* 4-locularis, 4-valvis, polysperma.

Specific Character and Synonyms.

ERICA *ampullacea* foliis ciliatis mucronatis, bractæis coloratis, floribus umbellatis subquaternis erecto-patentibus, stylo exserto.



The Erica here figured has some affinity in the form of its flowers to the *E. ventricosa*, as these in their shape resemble a flask or bottle, especially of that kind in which water is usually kept, we have named it *ampullacea*; it is of very modern introduction.

On the 11th of June 1784, we had the pleasure to see a small plant of this species in flower, with Mr. WILLIAMS, Nurseryman, Turnham-Green, an

unwearied and ingenious cultivator of this beautiful tribe of plants in particular, the richness of whose collection will appear in the subsequent list; by him it was raised from Cape seeds, though not more than the height of ten inches, it produced eighteen branches, most of which put forth flowers at their summits; we counted sixty-six blossoms on this small plant.

The leaves are short, linear, somewhat triangular, rigid, edged with fine crooked hairs, very visible when magnified, and terminating in a mucro or point, on the older branches recurved and mostly eight-rowed; each branch is usually terminated by four or five flowers, at first growing closely together, and covered so strongly with a glutinous substance, as to look as if varnished, and which is so adhesive as to catch ants and small flies; as the flowering advances, they separate more widely from each other, and finally a young branch grows out of the centre from betwixt them; the true calyx is composed of four lanceolate leaves, sitting close to and glued as it were to the corolla; besides these, there are several other leaves, which might be mistaken for those of the calyx, but which may with more propriety be called Bracteæ or Floral-leaves; some of these, like the calyx, are wholly red, others red and green mixed together, and broader than the leaves of the plant; the flowers are about an inch and a quarter in length, inflated below, and contracted above into a long narrow neck, dilating again so as to form a kind of knob, in which the antheræ are contained, just below the limb, which divides into four somewhat ovate obtuse segments, the upper side of these segments is of a very pale flesh colour, the under side of them as well as the dilated part just below them bright red, the body of the flower flesh colour, marked with eight longitudinal stripes, of a deeper hue; filaments eight, antheræ within the tube; style projecting about the eighth of an inch beyond the corolla; stigma, a round glutinous head.

The flowers as they decay become of a deeper red colour, and finally pale brown, still retaining their form and appearing to advantage;—hitherto the plant has produced no seeds here, is increased with difficulty either by cuttings or layers, but with most certainty in the latter way.

A CATALOGUE of HEATHS, cultivated and sold by RICHARD WILLIAMS, at his Nursery, *Turnham-Green, Middlesex.*

ERICA.

	* formosa.	planifolia.
	fucata.	* Plukenetii.
* abietina.	* grandiflora.	pubescens.
albens.	* halicacaba.	pyramidalis.
* ampullacea.	herbacea.	quadriflora.
arborea.	incarnata.	ramentacea.
—— var. squarrosa.	incana.	regerminans.
articularis.	lateralis.	scoparia.
australis.	lutea.	* sessiliflora.
baccans.	* mammosa.	* simpliciflora.
* Banksii.	—— var. fl. purp.	* Sparrmanni.
canescens.	margaritacea.	* spicata.
capitata.	—— var. fl. rubro.	stricta.
caffra.	marifolia.	taxifolia.
cernua.	mediterranea.	Tetralix.
* cerinthoides.	* Massoni.	—— var. fl. albo.
cinerea.	minima.	thymifolia.
—— var. fl. albo.	* monadelphica.	triflora.
ciliaris.	* Monsoniana.	—— var. fl. albo.
* coccinea.	mucosa.	* tubiflora.
comosa.	multiflora.	umbellata.
—— var. fl. rubro.	—— var. fl. albo.	urceolaris.
* conspicua.	* muscari.	vagans.
corifolia.	* nudiflora.	* ventricosa.
* cruenta.	parviflora.	* versicolor.
cubica.	* Pattersoni.	* verticillata.
* curviflora.	persoluta.	* vestita.
denticulata.	—— var. fl. rubro.	virgata.
* discolor.	* Petiveri.	vulgaris.
* elata.	physodes.	—— var. fl. albo.
empetrifolia.	* pinifolia.	
* fascicularis.		

N.B. *Those marked with an asterisk have tubular flowers.*

[304]

**HERMANNIA LAVENDULIFOLIA. LAVENDER-LEAVED
HERMANNIA.**

Class and Order.

MONADELPHIA PENTANDRIA.

Generic Character.

Pentagyna. Caps. 5-locularis. *Petala* basi semitubulata, obliqua.

Specific Character and Synonyms.

HERMANNIA *lavendulifolia* foliis lanceolatis obtusis integerrimis. *Linn. Syst. Vegetab. ed. 14. Murr. p. 611. Ait. Kew. v. 2. p. 413.*

HERMANNIA frutescens folio Lavendulæ latiore et obtuso flore parvo aureo BOERH. *Dill. Hort. Elth. t. 147.*



The *Hermannia* here figured is a plant of humble growth, forming a small bushy shrub, a foot or a foot and a half in height, and producing numerous flowers thinly scattered over the branches, the greatest part of the summer; it is this disposition which it has of flowering so freely, that renders it a desirable plant for the greenhouse, in which it is commonly kept, and of which it is an old inhabitant.

DILLENIIUS has figured it in his admirable work the *Hortus Elthamensis*, published in 1732; hence we learn that it was cultivated in Mr. SHERARD'S celebrated garden at Eltham prior to that date.

It is a native of the Cape, and is readily increased by cuttings.

[305]

AMARYLLIS EQUESTRIS. BARBADOES AMARYLLIS, or LILY.

Class and Order.

HEXANDRIA MONOGYNIA.

Generic Character.

Cor. hexapetaloidea, irregularis. *Filamenta* fauci tubi inserta, declinata, inæqualia proportionem vel directionem. *Linn. fil.*

Specific Character and Synonyms.

AMARYLLIS *equestris* spatha subbiflora, pedicellis erectis spatha brevioribus, tubo siliformi horizontali, limbo oblique patulo sursum curvo, fauce, pilosa. *Linn. fil. Ait. Kew. v. 1. p. 417.*

AMARYLLIS dubia *Linn. Am. Ac. 8. p. 254.*

LILIUM americanum puniceo flore Belladonna dictum. *Herm. Par. Bat. p. 194. cum fig.*



Mr. AITON, in his *Hortus Kewensis*, has inserted this species of *Amaryllis*, as named and described by the younger LINNÆUS; he informs us, that it is a native of the West-Indies, and was introduced by Dr. WILLIAM PITCAIRN, in 1778: as its time of flowering is not mentioned, we may presume, that it had not blossomed in the royal garden when the publication before mentioned first made its appearance; it no doubt has since, as we have seen it in that state in the collections of several Nurserymen, particularly those of Mr. GRIMWOOD and Mr. COLVILL.

It flowers towards the end of April.

The flowering stem rises above the foliage, to the height of about a foot or more, produces from one to three flowers, similar to, but not quite so large as those of the Mexican *Amaryllis*, to which it is nearly related; it differs however from that plant essentially in this, that the lower part of the flower projects further than the upper, which gives to its mouth that obliquity which LINNÆUS describes.

The spathe is composed of two leaves, which standing up at a certain period of the plant's flowering like ears, give to the whole flower a fancied resemblance of a horse's head; whether LINNÆUS derived his name of *equestris* from this circumstance or not, he does not condescend to inform us.

Mr. AITON regards it as a greenhouse plant; like those of many of the *Ixias*, however, the bulbs are of the more tender kind.

It is propagated by offsets, but not very readily.

[306]

**OTHONNA PECTINATA. WORMWOOD-LEAVED
OTHONNA.**

Class and Order.

SYNGENESIA POLYGAMIA NECESSARIA.

Generic Character.

Recept. nudum. *Pappus* subnullus, *Cal.* 1-phyllus multifidus subcylindricus.

Specific Character and Synonyms.

OTHONNA *pectinata* foliis pinnatifidis: laciniis linearibus parallelis. *Linn. Syst. Vegetab. ed. 14. Murr. p. 793. Ait. Kew. v. 3. p. 276.*

JACOBÆA africana frutescens, foliis absinthii umbelliferi incanis. *Comm. hort. 2. p. 137. t. 69.*



The *Othonna pectinata* is a native of Africa, a long-established and common plant in greenhouses, having been cultivated by Mr. MILLER, in 1731; it recommends itself chiefly on account of its foliage, which forms a pleasing contrast to the darker greens of other plants.

It flowers in May and June, is moderately hardy, and readily increased by cuttings.

In many collections we meet with old plants of it three or four feet high; formerly, when greenhouse plants were few in numbers and the houses large, it might be proper to keep such; but now there is not that necessity, especially since the vast accession of plants from the Cape and New-Holland, made within these few years.

[307]

HERMANNIA ALTHÆIFOLIA. MARSH-MALLOW-LEAVED HERMANNIA.

Class and Order.

MONADELPHIA PENTANDRIA.

Generic Character.

Pentagyna. Caps. 5-locularis. *Petala* basi semitubulata, obliqua.

Specific Character and Synonyms.

HERMANNIA *althæifolia* foliis ovatis crenatis plicatis tomentosis, calycibus florentibus campanulatis angulatis, stipulis oblongis foliaceis. *Ait. Kew. v. 2. p. 411.*

HERMANNIA *althæifolia* foliis ovatis plicatis crenatis tomentosis. *Linn. Syst. Vegetab. ed. 14. Murr. p. 610.*

HERMANNIA *capensis* *althææ* folio. *Pet. Gaz. 53. t. 34. f. 2.*

KETMIA *africana* frutescens foliis mollibus et incanis. *Comm. hort. 2. p. 151. t. 79.*



The *Hermannia althæifolia*, a native of the Cape, is a plant of much larger growth than the *lavendulifolia*, rising to the height, if suffered to do so, of three, four, or more feet; its blossoms are proportionably large, and of a deep yellow colour, inclined to orange.

It is a plant of free growth, much disposed to produce flowers during most of the summer months; hence it is kept very generally in collections of greenhouse plants: is propagated readily by cuttings.

Was cultivated by Mr. MILLER, in 1728. *Ait. Kew.*

Our readers will see, that the specific description of LINNÆUS has been altered in the *Hortus Kewensis*, and that it now comprizes all the striking features of the plant.

[308]

VERBENA AUBLETIA. ROSE VERVAIN.

Class and Order.

DIANDRIA MONOGYNIA.

Generic Character.

Cor. infundibuliformis subæqualis curva. *Calycis* unico dente truncato. *Semina* 2 s. 4 nuda (*Stam.* 2 s. 4.)

Specific Character and Synonyms.

VERBENA *Aubletia* tetrandra, spicis laxis solitariis, foliis trifidis incis. *Ait. Kew.* v. 1. p. 33.

VERBENA *Aubletia* tetrandra, spicis solitariis, coroliis fasciculatis, foliis cordatis inciso-serratis. *Linn. Syst. Vegetab. ed. 14. Murr. p. 66. Suppl. Pl. p. 86.*

BUCHNERA canadensis *Linn. Syst. Vegetab. ed. 13. p. 478.*

VERBENA *Aubletia.* *Jacq. hort.* v. 2. p. 82. t. 176.

OUBLETIA *Journ. de Rozier introd.* 1. p. 367. t. 2.



It has fallen to the lot of this plant to have an unusual degree of attention bestowed on it by various botanists, and after being regarded as a distinct genus by several, to be finally classed with the *Verbena*; in the *Supplementum Plantarum* of the younger LINNÆUS it is minutely described.

We learn from the *Hortus Kewensis* of Mr. AITON that it is a native of North-America, introduced by Mons. RICHARD in 1774, and that it flowers in June

and July.

The extreme brilliancy of its colours renders it a very ornamental greenhouse plant, it seldom grows above the height of two feet; in favourable seasons ripens its seeds readily, by which it is usually propagated, being a biennial.

[309]

**PELARGONIUM ECHINATUM. PRICKLY-STALKED
GERANIUM.**

Class and Order.

MONADELPHIA HEPTANDRIA.

Generic Character.

Cal. 5-partitus: lacinia suprema definente in tubulum capillarem nectariferum secus pendunculum decurrentem. *Cor.* 5 petala irregularis. *Filamenta* 10, inæqualia, quorum 3 raro 5 castrata. *Fructus* 5 coccus, rostratus: *rostra* spiralia, introrsum barbata.

Specific Character.

PERLARGONIUM *echinatum* caule carnosio, stipulis spinescentibus, foliis cordato-subrotundis 3-5 lobis, floribus umbellatis, umbellis subseptemfloris.



This singular and most beautiful species of *Pelargonium*, recently introduced to this country, this Summer flowered with Mr. COLVILL, Nurseryman, in the King's-Road, Chelsea, from one of whose plants our figure and description have been taken.

Stalk green, surface smooth and somewhat glossy, fleshy, beset with spines which bend back and terminate in brownish somewhat weak points; these appear to have been primarily the stipulæ, which become thus fleshy and rigid, and from this circumstance not altogether peculiar to this species, it takes the name of *echinatum*; the leaves stand on long footstalks, are somewhat heart-shaped, mostly roundish, divided into three or five lobes, veiny, soft, and downy, especially on the under side, which is of a much lighter colour than the upper, the flowering stem proceeds from the summit of the stalk, and is a foot or more in height; as it advances it throws out its branches, or peduncles, ultimately about five in number, each of which has a leaf at its base, similar to the other leaves of the plant, but smaller, and terminates in an umbel of seven or eight flowers; as the umbels blossom in succession, a period of several months usually intervenes betwixt the blowing of the first and the last; when the flower is expanded, the hindmost leaf of the calyx continues upright, the others are reflexed as in other species of this genus, they are all beset with fine long hairs; the three lowermost petals are pure white, with a little gibbosity at the base of each, the two uppermost are marked each with three irregular spots, of a rich purple colour, inclining to carmine, the two lowermost spots narrowest and of the deepest colour; of the stamina there are six filaments which have antheræ, and four of which have none; stigma red, divided into five parts, and a little longer than the fertile filaments.

In its habit this plant resembles somewhat the *Pelargonium cordifolium*, is a native of the Cape, flowers from May to September, in favourable seasons has produced seeds here, but is more usually increased by cuttings.

Varies with petals of a rich purple colour, in which the spots are similar, though not so conspicuous.

[310]

ERINUS ALPINUS. ALPINE ERINUS.

Class and Order.

DIDYNAMIA ANGIOSPERMIA.

Generic Character.

Cal. 5-phyllus. *Cor.* Limbus 5-fidus æqualis. *Caps.* 2-ocularis.

Specific Character and Synonyms.

ERINUS *alpinus* floribus racemosis, foliis spathulatis. *Linn. Syst. Vegetab. ed. 14. Murr. p. 570. Ait. Kew. v. 2. p. 357.*

AGERATUM serratum alpinum. *Bauh. pin. 221.*



The *Erinus alpinus* is a native of Switzerland, Germany, and France; inhabiting the more mountainous parts of those countries.

It is a very desirable little plant for the decoration of rock work, growing in close tufts, and producing numerous flowers of a lively purple colour during most of the summer months.

Is increased without difficulty by parting its roots in Autumn, or from seed; in the winter some plants of it should be kept in pots under a frame or hand-glass, as it is liable to be injured by wet and frost.

Was cultivated here by Mr. MILLER in 1759.

[311]

ROBINIA HISPIDA. ROUGH-STALK'D ROBINIA, or ROSE ACACIA.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Cal. 4-fidus. Legumen gibbum elongatum.

Specific Character and Synonyms.

ROBINIA *hispida* racemis axillaribus, foliis impari pinnatis, caule inermi hispido. *Linn. Mant. p. 668. Ait. Kew. v. 3. p. 53.*

ROBINIA racemis axillaribus, pedicellis unifloris, foliis impari pinnatis, caule inermi. *Jacq. Amer. 211. t. 179. f. 101.*

PSEUDO ACACIA *hispida* floribus roseis. *Catesb. Carol. 3. p. 20. t. 20.*



There are few trees or shrubs which have contributed more to adorn our plantations, and shrubberies, than those of this genus, nine species of which are enumerated in the *Hort. Kew.* of Mr. AITON, most of these are natives either of North-America, or Siberia: the present species, an inhabitant of Carolina, is perhaps the most ornamental of the whole: its large pendant bunches of rose-coloured flowers load the branches in May and June, and sometimes a second crop will be produced late in the season, these with us usually fall off without producing any seed-vessels.

This shrub is not disposed to grow very tall in America, it is most prudent indeed to keep it humble, to the height of four or five feet, and to plant it in a sheltered part of the garden, as its branches are liable to be broken by high winds: MARSHALL (*Arb. Amer.*) describes it as spreading much from its running roots; we have not observed it to do so in any great degree here; it is propagated by layers, by cuttings of the roots, and by grafting; it is of ready growth, disposed to blow even when young, and not nice as to soil, or situation; the flowers afford a good example of the class DIADELPHIA, they are large and beautiful, but without scent.

Was cultivated by Mr. MILLER in 1758. *Ait. Kew.*

[312]

LINUM FLAVUM. YELLOW FLAX.

Class and Order.

PENTANDRIA PENTAGYNIA.

Generic Character.

Cal. 5-phyllus. *Petala* 5. *Caps.* 5-valvis 10 locularis. *Sem.* solitaria.

Specific Character and Synonyms.

LINUM *flavum* calycibus subserrato-scabris lanceolatis subsessilibus, panicula ramis dichotomis. *Linn. Sp. Pl.* v. 1. ed. 3. p. 399. *Mant.* p. 360. *Syst. Vegetab.* ed. 14. *Murr.* p. 303. *Jacq. Fl. Austr.* v. 3. t. 214.

LINUM sylvestre latifolium luteum. *Bauh. pin.* 214.

LINUM sylvestre III. latifolium. *Clus. hist.* 1. p. 317.



There is a considerable similarity betwixt the representation of the present plant and that of the *Linum arboreum* figured on the 234th plate of this work, they are nevertheless two species widely differing, the *flavum* being a hardy herbaceous perennial, a native of Germany, the *arboreum* a greenhouse shrub from the Levant, both possessing considerable beauty, and highly worthy a place in all collections of ornamental plants.

The *Linum flavum* is not mentioned either in the Dictionary of Mr. MILLER, or the Hortus Kewensis of Mr. AITON, and as far as our knowledge extends was a stranger in this country, till we raised it the year before last from seeds sent us by Mr. DAVAL, of Orbe in Switzerland; CLUSIUS gives us a representation of it in flower, and Prof. JACQUIN another much superior; according to the latter, it grows by the sides of hedges and among shrubs in mountainous situations, and rarely exceeds a foot in height.

From the little experience we have had of this plant, it appears to be easy of culture, and to succeed best in a soil moderately stiff and moist; the flowers expand most in a morning when the sun shines, and continue in succession during June, July, and part of August; it appears as if it would ripen its seeds in my garden; these vegetate freely: the plant may also be increased by parting its roots in autumn, or by cuttings of the young shoots.

[313]

DAPHNE CNEORUM. TRAILING DAPHNE.

Class and Order.

OCTANDRIA MONOGYNIA.

Generic Character.

Cal. 0. *Cor.* 4-fida corallacea marcescens stamina includens,
Bacca 1-sperma.

Specific Character and Synonyms.

DAPHNE *Cneorum* floribus congestis terminalibus sessilibus,
foliis lanceolatis nudis mucronatis. *Linn. Syst. Veget. ed. 14.*
Murr. p. 371. Ait. Kew. v. 2. p. 26.

THYMELEÆ affinis facie externa. *Bauh. pin. 463.*

CNEORUM. *Matth. hist. 46. Clus. hist. 89, 90. f. 1.*



This charming little shrub is a native of Switzerland and Austria: CLUSIUS informs us that it grows in great abundance on many of the mountains near Vienna, so much so that women gather it when in flower and sell it in the markets; its beautiful and fragrant blossoms come forth in April and May, the principal season for its flowering, but it frequently blows during most of the Summer, and even in the Autumn; it varies with white blossoms.

It is extremely hardy, thrives remarkably well in road sand in almost any situation; is propagated by seeds, which very rarely ripen with us, by layers, and by grafting it on the stock of the Mezereon, whereby it acquires an elevation superior to what it has naturally.

[314]

GENISTA TRIQUETRA. TRIANGULAR-STALK'D GENISTA.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Cal. 2-labiatus: 2/3. *Vexillum* oblongum a pistillo staminibusque deorsum reflexum.

Specific Character.

GENISTA *triquetra* foliis ternatis, summis simplicibus, ramis triquetris procumbentibus. *L'Herit. Stirp. nov. t. 88. Ait. Kew. v. 3. p. 14.*



Mons. L'HERITIER, author of many modern publications in Botany, distinguished for their accuracy and elegance, was the first who described and figured this species of Genista, a native of Corsica, and cultivated here by JOHN ORD, Esq. as long since as the year 1770.

It is a hardy, evergreen, trailing shrub, producing a vast profusion of bloom; which renders it eminently conspicuous in May and June; its flowers are

rarely succeeded by seed-vessels, so that it is usually propagated by layers.

When tied up properly, and carefully trained to stake, it may vie with most of our ornamental shrubs: for covering a wall, or paling, where the situation is not too shady, it probably would succeed very well, at least it is deserving of trial.

[315]

PELARGONIUM CERATOPHYLLUM. HORN-LEAVED CRANE'S BILL.

Class and Order.

MONADELPHIA HEPTANDRIA.

Generic Character.

Cal. 5-partitus: lacinia suprema definente in tubulum capillarem, nectariferum, secus pendunculum decurrentem. *Cor.* 5-petala, irregularis. *Filam.* 10 inæqualia, quorum 3 raro 5 castrata. *Fructus* 5-coccus, rostratus, rostra spiralia introrsum barbata.

Specific Character and Synonyms.

PELARGONIUM *ceratophyllum* umbellis multifloris, foliis remote pinnatis carnosis teretibus, laciniis canaliculatis obsolete trifidis. *L'Herit. Geran. n. 50. t. 13. Ait. Kew. v. 2 p. 422.*



Mr. AITON informs us that this species of *Pelargonium*, which is one of the more fleshy kinds, is a native of the South-West Coast of Africa, and was introduced to the Royal Garden at Kew by Mr. ANTHONY HOVE in 1786.

It flowers during most of the Summer months, and ripens its seeds, by which it may be increased, as also by cuttings; it is found to be more tender

than many others, and more liable to be injured by damps, and hence it will require a treatment more applicable to a dry stove plant.

[316]

POLYGALA CHAMÆBUXUS. BOX-LEAVED MILK-WORT.

Class and Order.

DIADELPHIA OCTANDRIA.

Generic Character.

Cal. 5-phyllus: foliolis 2 majoribus alæformibus, ante maturitatem seminis coloratis. *Caps.* obcordata, 2-locularis. *Sem.* solitaria.

Specific Character and Synonyms.

POLYGALA *Chamæbuxus* floribus sparsis: carinæ apice subrotundo, foliis lanceolatis. *Linn. Syst. Vegetab. ed. 14. Murr. p. 639. Ait. Kew. v. 3. p. 4. Jacq. Fl. Austr. v. 3. p. 19. t. 233.*

CHAMÆBUXUS flore coluteæ. *Bauh. Pin. 471.*

ANONYMOS flore Coluteæ. *Clus. Hist. p. 105. f.*

POLYGALOIDES procumbens foliis duris ovatis nervo aristato. *Hall. Hist. n. 345.*



CLUSIUS, in his *Hist. Pl. rar.* gives an accurate description and good figure of the present plant, before unnoticed (as he observes) by any author; it has since been particularly described by HALLER and JACQUIN; the former makes a distinct genus of it, by the name of *Polygaloides*.

It is an elegant little evergreen shrub of low growth, rarely exceeding a foot in height, with leaves like those of box, producing flowers from May to October, but most plentifully in May and June; each flower stands on a peduncle proceeding from a kind of triphyllous cup, formed of floral leaves, the true calyx is composed of three leaves, which are nearly white; the two outermost petals, similar to the wings of a papilionaceous flower, are also white, or nearly so; the third petal which forms a kind of tube and contains the eight stamina with the pistillum, is white at the base, but yellow towards the extremity, where it changes by degrees to a bright bay colour: both CLUSIUS and JACQUIN observed a variety of this plant, in which the calyx and wings were of a beautiful purple; this variety, we believe, has not yet been introduced to this country: the common sort was cultivated in the garden at Oxford, in 1658.

MILLER describes it as a plant difficult of cultivation; it is not now regarded as such; both CLUSIUS and JACQUIN describe it as having creeping roots; such plants are generally increased without difficulty, and so is this; planted in bog earth on a shady border, it thrives extremely well, and spawns much, so that there is no necessity for having recourse to its seeds. It grows spontaneously on the Alps of Austria and Switzerland.

[317]

ONONIS FRUTICOSA. SHRUBBY REST-HARROW.

Class and Order.

DIADELPHIA DECANDRIA.

Generic Character.

Cal. 5-partitus: laciniis linearibus. *Vexillum* striatum. *Legumen* turgidum sessile. *Filamenta* connata absque fissura.

Specific Character and Synonyms.

ONONIS *fruticosa* foliis sessilibus ternatis lanceolatis serratis, stipulis vaginalibus, pedunculis subtrifloris. *Linn. Syst. Vegetab. ed. 14. Murr. p. 653. Ait. Kew. 24.*

ONONIS *purpurea verna præcox frutescens, flore rubro amplo. Moris. Hist. 2. p. 170.*



This charming little shrub, highly deserving of being more generally known, is a native of the South of France; and was cultivated here by MILLER in 1748.

In favourable situations, it produces blossoms in great profusion during most of the Summer, and ripens seed in abundance: the situation it affects is dry and sandy, but it is a shrub by no means nice, as to soil or place of

growth, and so hardy as to have borne the severity of last Winter, 1795, without injury.

In the collections about town we frequently find it in pots, kept with greenhouse plants.

It is said to vary with white flowers.

The best mode of raising it is from seed.

[318]

**ANTHERICUM LILIASTRUM. SAVOY ANTHERICUM, or
ST. BRUNO'S LILY.**

Class and Order.

HEXANDRIA MONOGYNIA.

Generic Character.

Cor. 6-petala, patens. *Caps.* ovata.

Specific Character and Synonyms.

ANTHERICUM *Liliastrum* foliis planis, scapo simplicissimo,
corollis campanulatis, staminibus declinatis. *Linn. Syst.*
Vegetab. ed. 14. Murr. p. 330. Ait. Kew. v. 1. p. 449.

HEMEROCALLIS floribus patulis secundis. *Hall. Hist. n. 1230.*

PHALANGIUM magno flore. *Bauh. Pin. 29.*

PHALANGIUM Allobrogicum majus. *Clus. cur. app. alt.*

PHALANGIUM Allobrogicum. The Savoye Spider-wort. *Park.*
Parad. p. 150. tab. 151. f. 1.



Botanists are divided in their opinions respecting the genus of this plant; LINNÆUS considers it as an *Anthericum*, HALLER and MILLER make it an *Hemerocallis*.

It is a native of Switzerland, where, HALLER informs us, it grows abundantly in the Alpine meadows, and even on the summits of the mountains; with us it flowers in May and June.

It is a plant of great elegance, producing on an unbranched stem about a foot and a half high, numerous flowers of a delicate white colour, much smaller but resembling in form those of the common white lily, possessing a considerable degree of fragrance, their beauty is heightened by the rich orange colour of their antheræ; unfortunately they are but of short duration.

MILLER describes two varieties of it differing merely in size.

A loamy soil, a situation moderately moist, with an eastern or western exposure, suits this plant best; so situated, it will increase by its roots, though not very fast, and by parting of these in the autumn, it is usually propagated.

PARKINSON describes and figures it in his *Parad. Terrest.* observing that "divers allured by the beauty of its flowers, had brought it into these parts."

[319]

ANAGALLIS MONELLI. ITALIAN PIMPERNEL.

Class and Order.

PENTANDRIA MONOGYNIA.

Generic Character.

Cor. rotata. *Caps.* circumscissa 1-locularis, polysperma.

Specific Character and Synonyms.

ANAGALLIS *Monelli*, foliis lanceolatis caule erecto. *Linn. Syst. Veget. ed. 14. Murr. p. 196. Ait. Kew. v. 1. p. 201.*

ANAGALLIS *cœrulea* foliis binis ternisve ex adverso nascentibus.
Bauh. Pin. 552.

ANAGALLIS *tenuifolia* Monnelli. *Clus. app. alt.*



In Italy and Spain, where this plant grows spontaneously, it is an annual, producing seed in abundance; with us (as far at least as we have observed) it produces no seed, but like the *Senecio elegans*, and some other annuals, is renewed, and rendered perennial by cuttings, which strike freely, and by which the plant requires to be renovated once or twice in a season; though described as growing with an upright stem, it requires to be tied up to a stick; and if this be neatly and dexterously done, its brilliant azure flowers

springing from every side of the stem, render it a charming ornament for the greenhouse or window: it flowers during most of the year.

CLUSIUS called it *Anagallis Monnelli*, the first knowledge he had of the plant being from his friend JOHANNES MONNELLUS.

On the same plant we find the leaves grow two, three, or four together, with flowers corresponding.

[320]

LOBELIA CARDINALIS. SCARLET LOBELIA, or CARDINAL'S FLOWER.

Class and Order.

SYNGENESIA MONOGAMIA.

Generic Character.

Cal. 5-fidus. *Cor.* 1-petala, irregularis. *Caps.* infera 2, s. 3-ocularis.

Specific Character and Synonyms.

LOBELIA *cardinalis* caule erecto, foliis lato-lanceolatis serratis, racemo terminali secundo. *Linn. Syst. Veg. ed. 14. Murr. p. 801. Ait. Kew. v. 3. p. 284.*

RAPUNTIUM galeatum virginianum, coccineo flore majore. *Moris. Hist. 2. p. 466. s. 5. t. 5. f. 54.*

TRACHELIUM Americarum flore ruberrimo, sive Planta Cardinalis. The rich crimson Cardinal's Flower. *Park. Parad. p. 356. t. 355.*



This species of *Lobelia*, so eminently distinguished for the richness of its scarlet blossoms, is a native of the colder as well as warmer parts of North-America. PARKINSON, who cultivated it in 1629, informs us that he received plants of it from France for his garden, and that "it groweth neere the river of Canada, where the French plantation in America is seated."

It is a hardy herbaceous plant, growing in favourable situations to the height of three or four feet; the main spike of flowers which terminates the stalk, is

often a foot in length; by the time that most of its flowers are blown, side branches shoot out, and flower; so that the plant continues in bloom six weeks or two months: if the Autumn prove favourable, the plant with us produces plenty of seed in the open ground; to insure its ripening, some place pots of it, when blowing, in the greenhouse or stove.

Beautiful and hardy as this plant is, and long as it has been introduced to this country, we do not find it generally in gardens; we attribute this to its having, in a greater degree than many other plants, a partiality for a particular soil; in certain districts, where the soil is stiff and moist, it grows as freely as any weed, in other soils it is perpetually going off: it is also one of those plants whose roots require to be often parted; if this be done every Autumn, and they be planted in a stiff loam, the situation somewhat moist and shady, this very desirable plant may be had to grow and blossom in perfection.

It flowers from the latter end of July to October.

Is increased by parting its roots, by cuttings of the stalk and from seed.

[321]

COTYLEDON ORBICULATA. ROUND-LEAVED NAVEL-WORT.

Class and Order.

DECANDRIA PENTAGYNIA.

Generic Character.

Cal. 5-fidus. *Cor.* 1-petala. *Squamæ* nectariferæ 5 ad basin germinis. *Caps.* 5.

Specific Character and Synonyms.

COTYLEDON *orbiculata* foliis orbiculatis carnosis planis integerrimis, caule fruticoso. *Linn. Syst. Veg. ed. 14. Murr. p. 428. Ait. Kew. var. [delta] v. 2. p. 106.*

COTYLEDON *africanum frutescens incanum orbiculatis foliis. Herm. Lugd. 349. t. 551. Moris. Hist. 3. p. 474. s. 12. t. 7 f. 39.*



The *Cotyledon orbiculata* is one of our oldest succulents, being introduced as long since as 1690, by Mr. BENTICK^[C]: it still retains a place in most collections, deservedly indeed, for it has every claim to our notice; its appearance is magnificent, the glaucous colour of its foliage highly pleasing, its flowers large and of long duration; it blows freely, grows rapidly, is easily increased by cuttings, and will succeed in a house or window, with the common treatment of an African Geranium.

When suffered to grow, it will become a shrub of considerable size; but this is not necessary for its flowering, as young and small plants are disposed to throw out blossoms, which is not the case with a plant extremely similar to, and often confounded with it, viz. the *Crassula Cotyledon*, whose foliage indeed scarcely differs from our plant but in being finely dotted.

It is a native of the Cape, and flowers from June or July to September.

In the *Hort. Kew.* of Mr. AITON, four varieties are enumerated, differing chiefly in the form of their foliage.

[C] AIT. KEW.

[322]

MANULEA TOMENTOSA. WOOLLY MANULEA.

Class and Order.

DIDYNAMIA ANGIOSPERMIA.

Generic Character.

Cal. 5-partitus. *Cor.* limbo 5-partito, subulato: laciniis superioribus 4 magis connexis. *Caps.* 2-locularis, polysperma.

Specific Character and Synonyms.

MANULEA *tomentosa* foliis tomentosis, caulibus foliosis, pedunculis multifloris. *Linn. Mant.* 420. *Syst. Veget. ed.* 14. *Murr. p.* 569. *Ait. Kew. v. 2. p.* 356.

SELAGO *tomentosa* foliis obovatis crenatis, caule prostrato, racemis ramosis. *Linn. Amæn. Acad. v. 6. p.* 90. *Sp. Pl. ed.* 3. *p.* 877.

PLANTA *Pluk. Phyt.* 319. *f.* 2.



LINNÆUS describes this plant in the *Amœnitates Academicæ* under the name of *Selago tomentosa*, by which name he continues to call it in the third edition of the *Spec. Pl.* in his *Mantissa* he describes it more minutely, and changes it to the Genus *Manulea*, first established by him in the said work; he observes, that in this species the corolla is more regular than in the others.

Mr. AITON regards it as a biennial, its stalk is a foot or a foot and a half high, and woolly, its branches are opposite, not alternate as LINNÆUS describes them; in this perhaps they may vary; leaves opposite, sessile, obovate, narrowing to the base, toothed on the edge, edge rolled back a little in the young leaves, flowers grow in a long thyrsus, from two to five proceed from one common short peduncle; they are at first lemon-coloured, or greenish yellow, finally deep orange; LINNÆUS says the whole of the plant except the corolla is woolly, the tube of that even is hairy, the segments are smooth, with their edges rolled back, the upper part of the tube in which the stamina are included is dilated somewhat, as is also the lower part, so that it is narrowest in the middle. The flowers which make their appearance from May to November are usually succeeded by seeds, by which the plant is propagated.

It is a native of the Cape, and, according to Mr. AITON, was introduced by Mr. MASSON, in 1774.

The blossoms have a singular but unpleasant smell, not perceivable at a distance.

The variety of pleasant colours so conspicuous in the flowers, renders this rare plant desirable to such as aim at a general collection.

[323]

RUBUS ODORATUS. FLOWERING RASPBERRY.

Class and Order.

ICOSANDRIA POLYGYNIA.

Generic Character.

Cal. 5-fidus. *Petala* 5. *Bacca* composita acinis monospermis.

Specific Character and Synonyms.

RUBUS *odoratus* foliis simplicibus palmatis, caule inermi
multifolio multifloro. *Linn. Syst. Veg. ed. 14. Murr. p. 475.*
Ait. Kew. v. 2. p. 210.

RUBUS *odoratus*. *Corn. Canad. 149. t. 150.*



Botanists and Gardeners have given to this species of *Rubus* the name of *flowering*, not because it is the only one which produces flowers, but from its being regarded for its flowers merely; they indeed are so shewy, and so plentifully produced, that the plant has long been thought to merit a place in most shrubberies; to the various inhabitants of which, both in the largeness and elegant form of its leaves, and the colour of its blossoms, it forms a pleasing contrast.

It is extremely hardy, and easily propagated by suckers; the only care which it requires, is to keep it within proper bounds: young plants of it produce the largest and finest flowers.

It blossoms from June to September, is a native of different and distant parts of North-America, and was cultivated here by Mr. MILLER, in 1739.

CORNUTUS, who first figured and described this plant, gave it the name of *odoratus*, on account of the fragrance of its foliage; his words are "elegantissimi hujus folia fragrantissima sunt, paremque agrimonio odorato spirant odorem;" the fruit, rarely produced with us, he observes, is like the common Raspberry, but not so pleasant.

[324]

ANTIRRHINUM TRIPHYLLUM. THREE-LEAVED TOAD-FLAX.

Class and Order.

DIDYNAMIA ANGIOSPERMIA.

Generic Character.

Cal. 5-phyllus. *Cor.* basis deorsum prominens nectarifera. *Caps.* 2-ocularis.

Specific Character and Synonyms.

ANTIRRHINUM *triphyllum* foliis ternis ovatis. *Linn. Syst. Vegetab. ed. 14. Murr. p. 555. Ait. Kew.*

LINARIA *triphylla* minor lutea. *Bauh. Pin. 212.*

LINARIA *triphylla* cœrulea. *Bauh. Pin. 212.*

LINARIA *hispanica*. *Clus. Hist. 1. p. 320.*

LINARIA *valentina*. Tode Flaxe of Valentia. *Park. Par. p. 268.*



The *Antirrhinum triphyllum*, so called from the leaves growing by threes on the stalk (a character, by the bye, not very constant) was cultivated by PARKINSON, and described by him in his *Parad. terr.* He appears to have been a stranger to the particoloured variety now so generally cultivated as an ornamental annual in our gardens; in its wild state the flowers of this *Antirrhinum* are of a yellow hue, with little or no purple in them, such indeed are frequently produced from seeds sown in our gardens.

It is a hardy annual, a native of Spain and Sicily, a plant of ready growth, requiring the common treatment of annuals sown in the Spring, and much disposed indeed to come up spontaneously where it has once grown; in sowing its seeds, care should be taken to preserve the produce of such flowers as have the most purple in them.

INDEX.

In which the Latin Names of the Plants contained in the *Ninth Volume* are alphabetically arranged.

Pl.

[295](#) Agrostemma Cœli rosa.
[290](#) Amaryllis lutea.
[294](#) ——— sarniensis.
[305](#) ——— equestris.
[319](#) Anagallis Monelli.
[324](#) Antirrhinum triphyllum.
[318](#) Anthericum Liliastrum.
[293](#) Catananche c[oe]rulea.
[291](#) Capparis spinosa.
[321](#) Cotyledon orbiculata.
[289](#) Convolvulus linearis.
[313](#) Daphne Cneorum.
[297](#) Dianthus superbus.
[303](#) Erica ampullacea.

INDEX.

In which the English Names of the Plants contained in the *Ninth Volume* are alphabetically arranged.

Pl.

[290](#) Amaryllis yellow.
[294](#) ——— Guernsey.
[305](#) ——— Barbadoes.
[318](#) Anthericum Savoy.
[291](#) Caper Shrub.
[293](#) Catananche blue.
[295](#) Cockle smooth-leaved.
[289](#) Convolvulus narrow-leaved.
[315](#) Crane's-bill horn-leaved.
[300](#) Cudweed giant.
[313](#) Daphne trailing.
[298](#) Dittany of Crete.
[310](#) Erinus alpine.
[312](#) Flax yellow.

*** END OF THE PROJECT GUTENBERG EBOOK THE BOTANICAL
MAGAZINE, VOL. 09 ***

Updated editions will replace the previous one—the old editions will be renamed.

Creating the works from print editions not protected by U.S. copyright law means that no one owns a United States copyright in these works, so the Foundation (and you!) can copy and distribute it in the United States without permission and without paying copyright royalties. Special rules, set forth in the General Terms of Use part of this license, apply to copying and distributing Project Gutenberg™ electronic works to protect the PROJECT GUTENBERG™ concept and trademark. Project Gutenberg is a registered trademark, and may not be used if you charge for an eBook, except by following the terms of the trademark license, including paying royalties for use of the Project Gutenberg trademark. If you do not charge anything for copies of this eBook, complying with the trademark license is very easy. You may use this eBook for nearly any purpose such as creation of derivative works, reports, performances and research. Project Gutenberg eBooks may be modified and printed and given away—you may do practically ANYTHING in the United States with eBooks not protected by U.S. copyright law. Redistribution is subject to the trademark license, especially commercial redistribution.

START: FULL LICENSE

THE FULL PROJECT GUTENBERG™ LICENSE

PLEASE READ THIS BEFORE YOU DISTRIBUTE OR USE THIS WORK

To protect the Project Gutenberg™ mission of promoting the free distribution of electronic works, by using or distributing this work (or any other work associated in any way with the phrase “Project Gutenberg”), you agree to comply with all the terms of the Full Project Gutenberg License available with this file or online at www.gutenberg.org/license.

Section 1. General Terms of Use and Redistributing Project Gutenberg electronic works

1.A. By reading or using any part of this Project Gutenberg electronic work, you indicate that you have read, understand, agree to and accept all the terms of this license and intellectual property (trademark/copyright) agreement. If you do not agree to abide by all the terms of this agreement, you must cease using and return or destroy all copies of Project Gutenberg electronic works in your possession. If you paid a fee for obtaining a copy of or access to a Project Gutenberg electronic work and you do not agree to be bound by the terms of this agreement, you may obtain a refund from the person or entity to whom you paid the fee as set forth in paragraph 1.E.8.

1.B. “Project Gutenberg” is a registered trademark. It may only be used on or associated in any way with an electronic work by people who agree to be bound by the terms of this agreement. There are a few things that you can do with most Project Gutenberg electronic works even without complying with the full terms of this agreement. See paragraph 1.C below. There are a lot of things you can do with Project Gutenberg electronic works if you follow the terms of this agreement and help preserve free future access to Project Gutenberg electronic works. See paragraph 1.E below.

1.C. The Project Gutenberg Literary Archive Foundation (“the Foundation” or PGLAF), owns a compilation copyright in the collection of Project Gutenberg electronic works. Nearly all the individual works in the collection are in the public domain in the United States. If an individual work is unprotected by copyright law in the United States and you are

located in the United States, we do not claim a right to prevent you from copying, distributing, performing, displaying or creating derivative works based on the work as long as all references to Project Gutenberg are removed. Of course, we hope that you will support the Project Gutenberg mission of promoting free access to electronic works by freely sharing Project Gutenberg works in compliance with the terms of this agreement for keeping the Project Gutenberg name associated with the work. You can easily comply with the terms of this agreement by keeping this work in the same format with its attached full Project Gutenberg License when you share it without charge with others.

1.D. The copyright laws of the place where you are located also govern what you can do with this work. Copyright laws in most countries are in a constant state of change. If you are outside the United States, check the laws of your country in addition to the terms of this agreement before downloading, copying, displaying, performing, distributing or creating derivative works based on this work or any other Project Gutenberg work. The Foundation makes no representations concerning the copyright status of any work in any country other than the United States.

1.E. Unless you have removed all references to Project Gutenberg:

1.E.1. The following sentence, with active links to, or other immediate access to, the full Project Gutenberg License must appear prominently whenever any copy of a Project Gutenberg work (any work on which the phrase “Project Gutenberg” appears, or with which the phrase “Project Gutenberg” is associated) is accessed, displayed, performed, viewed, copied or distributed:

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg™ License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

1.E.2. If an individual Project Gutenberg electronic work is derived from texts not protected by U.S. copyright law (does not contain a notice indicating that it is posted with permission of the copyright holder), the work can be copied and distributed to anyone in the United States without paying any fees or charges. If you are redistributing or providing access to a work with the phrase “Project Gutenberg” associated with or appearing on the work, you must comply either with the requirements of paragraphs 1.E.1 through 1.E.7 or obtain permission for the use of the work and the Project Gutenberg trademark as set forth in paragraphs 1.E.8 or 1.E.9.

1.E.3. If an individual Project Gutenberg electronic work is posted with the permission of the copyright holder, your use and distribution must comply with both paragraphs 1.E.1 through 1.E.7 and any additional terms imposed by the copyright holder. Additional terms will be linked to the Project Gutenberg License for all works posted with the permission of the copyright holder found at the beginning of this work.

1.E.4. Do not unlink or detach or remove the full Project Gutenberg License terms from this work, or any files containing a part of this work or any other work associated with Project Gutenberg.

1.E.5. Do not copy, display, perform, distribute or redistribute this electronic work, or any part of this electronic work, without prominently displaying the sentence set forth in paragraph 1.E.1 with active links or immediate access to the full terms of the Project Gutenberg License.

1.E.6. You may convert to and distribute this work in any binary, compressed, marked up, nonproprietary or proprietary form, including any word processing or hypertext form. However, if you provide access to or distribute copies of a Project Gutenberg work in a format other than “Plain Vanilla ASCII” or other format used in the official version posted on the official Project Gutenberg website (www.gutenberg.org), you must, at no additional cost, fee or expense to the user, provide a copy, a means of exporting a copy, or a means of obtaining a copy upon request, of the work in its original “Plain Vanilla ASCII” or other form. Any alternate format must include the full Project Gutenberg License as specified in paragraph 1.E.1.

1.E.7. Do not charge a fee for access to, viewing, displaying, performing, copying or distributing any Project Gutenberg works unless you comply with paragraph 1.E.8 or 1.E.9.

1.E.8. You may charge a reasonable fee for copies of or providing access to or distributing Project Gutenberg electronic works provided that:

- You pay a royalty fee of 20% of the gross profits you derive from the use of Project Gutenberg works calculated using the method you already use to calculate your applicable taxes. The fee is owed to the owner of the Project Gutenberg trademark, but he has agreed to donate royalties under this paragraph to the Project Gutenberg Literary Archive Foundation. Royalty payments must be paid within 60 days following each date on which you prepare (or are legally required to prepare) your periodic tax returns. Royalty payments should be clearly marked as such and sent to the Project Gutenberg Literary Archive Foundation at the address specified in Section 4, "Information about donations to the Project Gutenberg Literary Archive Foundation."
- You provide a full refund of any money paid by a user who notifies you in writing (or by e-mail) within 30 days of receipt that s/he does not agree to the terms of the full Project Gutenberg™ License. You must require such a user to return or destroy all copies of the works possessed in a physical medium and discontinue all use of and all access to other copies of Project Gutenberg™ works.
- You provide, in accordance with paragraph 1.F.3, a full refund of any money paid for a work or a replacement copy, if a defect in the electronic work is discovered and reported to you within 90 days of receipt of the work.
- You comply with all other terms of this agreement for free distribution of Project Gutenberg™ works.

1.E.9. If you wish to charge a fee or distribute a Project Gutenberg™ electronic work or group of works on different terms than are set forth in this agreement, you must obtain permission in writing from the Project Gutenberg Literary Archive Foundation, the manager of the Project

Gutenberg™ trademark. Contact the Foundation as set forth in Section 3 below.

1.F.

1.F.1. Project Gutenberg volunteers and employees expend considerable effort to identify, do copyright research on, transcribe and proofread works not protected by U.S. copyright law in creating the Project Gutenberg™ collection. Despite these efforts, Project Gutenberg™ electronic works, and the medium on which they may be stored, may contain “Defects,” such as, but not limited to, incomplete, inaccurate or corrupt data, transcription errors, a copyright or other intellectual property infringement, a defective or damaged disk or other medium, a computer virus, or computer codes that damage or cannot be read by your equipment.

1.F.2. LIMITED WARRANTY, DISCLAIMER OF DAMAGES - Except for the “Right of Replacement or Refund” described in paragraph 1.F.3, the Project Gutenberg Literary Archive Foundation, the owner of the Project Gutenberg™ trademark, and any other party distributing a Project Gutenberg™ electronic work under this agreement, disclaim all liability to you for damages, costs and expenses, including legal fees. YOU AGREE THAT YOU HAVE NO REMEDIES FOR NEGLIGENCE, STRICT LIABILITY, BREACH OF WARRANTY OR BREACH OF CONTRACT EXCEPT THOSE PROVIDED IN PARAGRAPH 1.F.3. YOU AGREE THAT THE FOUNDATION, THE TRADEMARK OWNER, AND ANY DISTRIBUTOR UNDER THIS AGREEMENT WILL NOT BE LIABLE TO YOU FOR ACTUAL, DIRECT, INDIRECT, CONSEQUENTIAL, PUNITIVE OR INCIDENTAL DAMAGES EVEN IF YOU GIVE NOTICE OF THE POSSIBILITY OF SUCH DAMAGE.

1.F.3. LIMITED RIGHT OF REPLACEMENT OR REFUND - If you discover a defect in this electronic work within 90 days of receiving it, you can receive a refund of the money (if any) you paid for it by sending a written explanation to the person you received the work from. If you received the work on a physical medium, you must return the medium with your written explanation. The person or entity that provided you with the defective work may elect to provide a replacement copy in lieu of a refund. If you received the work electronically, the person or entity providing it to

you may choose to give you a second opportunity to receive the work electronically in lieu of a refund. If the second copy is also defective, you may demand a refund in writing without further opportunities to fix the problem.

1.F.4. Except for the limited right of replacement or refund set forth in paragraph 1.F.3, this work is provided to you 'AS-IS', WITH NO OTHER WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR ANY PURPOSE.

1.F.5. Some states do not allow disclaimers of certain implied warranties or the exclusion or limitation of certain types of damages. If any disclaimer or limitation set forth in this agreement violates the law of the state applicable to this agreement, the agreement shall be interpreted to make the maximum disclaimer or limitation permitted by the applicable state law. The invalidity or unenforceability of any provision of this agreement shall not void the remaining provisions.

1.F.6. INDEMNITY - You agree to indemnify and hold the Foundation, the trademark owner, any agent or employee of the Foundation, anyone providing copies of Project Gutenberg™ electronic works in accordance with this agreement, and any volunteers associated with the production, promotion and distribution of Project Gutenberg™ electronic works, harmless from all liability, costs and expenses, including legal fees, that arise directly or indirectly from any of the following which you do or cause to occur: (a) distribution of this or any Project Gutenberg work, (b) alteration, modification, or additions or deletions to any Project Gutenberg work, and (c) any Defect you cause.

Section 2. Information about the Mission of Project Gutenberg

Project Gutenberg is synonymous with the free distribution of electronic works in formats readable by the widest variety of computers including obsolete, old, middle-aged and new computers. It exists because of the

efforts of hundreds of volunteers and donations from people in all walks of life.

Volunteers and financial support to provide volunteers with the assistance they need are critical to reaching Project Gutenberg's goals and ensuring that the Project Gutenberg collection will remain freely available for generations to come. In 2001, the Project Gutenberg Literary Archive Foundation was created to provide a secure and permanent future for Project Gutenberg and future generations. To learn more about the Project Gutenberg Literary Archive Foundation and how your efforts and donations can help, see Sections 3 and 4 and the Foundation information page at www.gutenberg.org.

Section 3. Information about the Project Gutenberg Literary Archive Foundation

The Project Gutenberg Literary Archive Foundation is a non-profit 501(c)(3) educational corporation organized under the laws of the state of Mississippi and granted tax exempt status by the Internal Revenue Service. The Foundation's EIN or federal tax identification number is 64-6221541. Contributions to the Project Gutenberg Literary Archive Foundation are tax deductible to the full extent permitted by U.S. federal laws and your state's laws.

The Foundation's business office is located at 41 Watchung Plaza #516, Montclair NJ 07042, USA, +1 (862) 621-9288. Email contact links and up to date contact information can be found at the Foundation's website and official page at www.gutenberg.org/contact

Section 4. Information about Donations to the Project Gutenberg Literary Archive Foundation

Project Gutenberg™ depends upon and cannot survive without widespread public support and donations to carry out its mission of increasing the number of public domain and licensed works that can be freely distributed in machine-readable form accessible by the widest array of equipment

including outdated equipment. Many small donations (\$1 to \$5,000) are particularly important to maintaining tax exempt status with the IRS.

The Foundation is committed to complying with the laws regulating charities and charitable donations in all 50 states of the United States. Compliance requirements are not uniform and it takes a considerable effort, much paperwork and many fees to meet and keep up with these requirements. We do not solicit donations in locations where we have not received written confirmation of compliance. To SEND DONATIONS or determine the status of compliance for any particular state visit www.gutenberg.org/donate.

While we cannot and do not solicit contributions from states where we have not met the solicitation requirements, we know of no prohibition against accepting unsolicited donations from donors in such states who approach us with offers to donate.

International donations are gratefully accepted, but we cannot make any statements concerning tax treatment of donations received from outside the United States. U.S. laws alone swamp our small staff.

Please check the Project Gutenberg web pages for current donation methods and addresses. Donations are accepted in a number of other ways including checks, online payments and credit card donations. To donate, please visit: www.gutenberg.org/donate.

Section 5. General Information About Project Gutenberg electronic works

Professor Michael S. Hart was the originator of the Project Gutenberg concept of a library of electronic works that could be freely shared with anyone. For forty years, he produced and distributed Project Gutenberg eBooks with only a loose network of volunteer support.

Project Gutenberg eBooks are often created from several printed editions, all of which are confirmed as not protected by copyright in the U.S. unless a

copyright notice is included. Thus, we do not necessarily keep eBooks in compliance with any particular paper edition.

Most people start at our website which has the main PG search facility:
www.gutenberg.org.

This website includes information about Project Gutenberg, including how to make donations to the Project Gutenberg Literary Archive Foundation, how to help produce our new eBooks, and how to subscribe to our email newsletter to hear about new eBooks.

N. 289



Pub. by W. Curtis S^t Geo. Crescent Feb. 1. 1795

[back](#)

N. 290



Pub. by W. Curtis S.^c Geo. Crescent Feb. 1. 1793

[back](#)

N^o 291



Pub. by W. Curtis S^t. Geo. Crescent Feb. 1. 1795

[back](#)

N^o 292



Pub. by W. Curtis S^t Geo. Crescent Mar. 1. 1795

[back](#)

N^o 293



Pub. by W. Curtis S^t Geo. Crescent Mar 1. 1795.

[back](#)

N^o 294



Pub by W Curtis S.^t Geo: Crescent Mar 1. 1795

[back](#)

N. 295



Pub. by W. Curtis N^o 6 Geo. Crescent Apr. 1. 1795.

[back](#)

N^o 296



Pub. by W. Curtis. St Geo. Crescent Apr 1. 1795.

[back](#)



[back](#)

N. 298



Pub. by W. Curtis S.^t Geo. Crescent Apr. 1. 1795

[back](#)

N. 299



Pub. by W. Curtis Sc^{lt} Geo. Crescent May 1. 1795.

[back](#)

N^o 300



Pub. by W. Curtis Sc^{pt} Geo. Crescent May 1. 1795

[back](#)



[back](#)

N: 302



Pub. by W. Curtis Sc^t Geo. Crescent June 1. 1795.

[back](#)

N 303



Pub. by W. Curtis Sc^t Geo. Crescent. June 1. 1795.

[back](#)

N 304



Pub. by W. Curtis S.^t Geo Crescent July 1. 1795.

[back](#)

N. 305



Pub. by W. Curtis Sc. Geo. Crescent July 1. 1795.

[back](#)

N. 306



Pub. by W. Curtis Sc. Geo. Crescent July 1. 1795.

[back](#)

N^o 307



Pub by W. Curtis. S.^c Geo Crescent July 1. 1795.

[back](#)



Pub. by W. Curtis S.^t Geo. Crescent Aug. 1. 1795

[back](#)

N^o 309



Pub. by W. Curtis S^t Geo. Crescent Aug 1 1795

[back](#)

N. 310



Pub. by W. Curtis S.^t Geo. Crescent Sep. 1. 1795.

Edwards del. Sanyer sc.

[back](#)



Edwards del. Senfom sc.

Pub. by W. Curtis S.^t Geo. Crescent Sept. 1795

[back](#)

N 312



Pub by W. Curtis S^t Geo. Crescent Sep. 1. 1795

[back](#)

N^o 313



Pub. by W. Curtis S.^r Geo. Crescent. Oct. 1. 1795.

[back](#)



[back](#)

N. 315



[back](#)

N. 316



Pub. by W. Curtis Sc. Geo. Crescent Nov. 1. 1795

[back](#)

N 317



Pub. by W. Curtis's Geo. Crescent Nov. 1. 1795.

[back](#)



[back](#)

N^o 319



Pub. by W. Curtis Sc^t Geo. Crescent Dec 1 1795.

[back](#)

N^o. 320



Pub. by W. Curtis Sc^{ilicet} Geo: Crescent Decr. 1795.

[back](#)

N. 321



W. Curtis sculp. Geo. Ordant Dec. 1. 1795.

[back](#)

N^o 322



Pub. by W. Curtis Sc Geo. Crescent Jan. 1796

[back](#)

N^o 323



Pub. by W. Curtis, J. Geo. Crescent Jan 1. 1796

[back](#)

N^o 324



Pub. by W. Curtis, S^t Geo. Crescent Jan. 1 1796

[back](#)