

Of the
Botanists Repository
Comprising,



Of the
Botanists Repository
Comprising,



The Project Gutenberg eBook of The botanist's repository for new and rare plants; vol. 10 [of 10]

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

Title: The botanist's repository for new and rare plants; vol. 10 [of 10]

Author: active 1799-1828 Henry Cranke Andrews

Release date: August 13, 2026 [eBook #79358]

Language: English

Original publication: London: The author, 1797

Other information and formats: www.gutenberg.org/ebooks/79358


Credits: Chuck Greif and the Online Distributed Proofreading Team at <https://www.pgdp.net> (This file was produced from images generously made available by Biodiversity Heritage Library.)

*** START OF THE PROJECT GUTENBERG EBOOK THE
BOTANIST'S REPOSITORY FOR NEW AND RARE PLANTS; VOL. 10
[OF 10] ***

INDEX
TO THE PLANTS CONTAINED IN VOL. X
ERRATA VOL. X.

*Alphabetical Index to the 1st. 2d. 3d. 4th. and 5th. Volume of
the Botanist's Repository.*

*Alphabetical Index to the 6th. 7th. 8th. 9th. and 10th. Volume
of the Botanist's Repository.*



Vol. X

of the

Botanist's Repository

Comprising
Colour'd Engravings

of

New and Rare Plants

ONLY

With Botanical Descriptions &c.

—*in*—

Latin and English,

after the

Linnæan System.

by

H. Andrews

Botanical Painter Engraver, &c.

PLATE DCIX.

CITRUS MEDICA, *odoratissima*.

Bergamot Lemon.

CLASS XVIII. ORDER III.

POLYADELPHIA ICOSANDRIA. Many Sets of Chives. Threads from the Calyx or Receptacle.

GENERIC CHARACTER.

CALYX 5-dentatus. Corolla 5-petala. Stamina 20 in cylindrum passim connata. Stylus 1. Bacca 9—12-locularis, pulpa vesiculari.

CUP 5-toothed. Petals five. Stamens about 20, generally united at the base. Shaft 1. Fruit 9-to 12-celled, the pulp bladdery.

SPECIFIC CHARACTER.

CITRUS foliis acuminatis.

CITRUS with pointed leaves.

CITRUS medica β . *Linn. Sp. Pl.* Limon Bergamotta. *Volkamer. Hesperides*, cap. 26. tab. p. 154.

REFERENCE TO THE PLATE.

1. The empalement.
2. Chives spread open.
3. The seed-bud and pointal.
4. A horizontal section of the fruit.

The drawing of this curious and valuable fruit was also taken at Wormleybury last May. Sir Abraham Hume informs us, that the plant was introduced in 1785, by Mrs. Evelyn, of St. Clare in Kent, who brought it with her from Nice. The name Bergamot is said by the Chevalier Lamarck (in the French Encyclopédie) to come from Bergame in Italy, where the

principal cultivation of the plant lies. The valuable perfume called Essence of Bergamot is prepared in Italy from the fruit, but by what method they extract it we have not been able to learn: it is rather remarkable that England, so celebrated for her commerce, her collections of natural history, and spirit in gardening, and annually importing so much of this Essence, should have been so long without the plant that produced it.

In arranging the Bergamot for the present as a variety of the Lemon, we by no means subscribe to the opinion of those who, servilely following Linnæus, include all sorts of apples and pears under his *Pyrus mala* and *communis*; all grape-vines under *Vitis vinifera*; and limes, lemons, citrons, oranges and shaddocks of all kinds, under his *Citrus Medica* and *Aurantium*. Already the last-mentioned genus has been considerably illustrated by the works of Gmelin, Rumphius, Thunberg, and Loureiro; and Professor Willdenow now enumerates six species of Citrus, and the Chevalier Lamarck eleven species; and many more are probably yet latent in the unexplored regions of Asia. Indeed it is only to those who have an opportunity of observing them in their original situations, where the spade of the labourer has never disturbed their repose, that we must look for their complete illustration. The immense tracts on our northern hemisphere, over which apples are naturally scattered, as well as the vast and permanent differences observed in their fruits, give us also much to hope for; even Linnæus himself observes in his *Flora Suecica*, that the apple which he found growing naturally in Smoland was very different from that which he found in other parts of Sweden; the original Paradise-apple (*Malus pumila* of the old authors) has already been recovered by the Russian naturalist Pallas, forming large thickets on the banks of the Wolga and Tanais, (see his *Flora Rossica*, vol. i. p. 22.); and the same author informs us that the apples he found growing about the Terek were of a large size, and excellent even in their wild state; while those that he met with in other parts of Russia were quite worthless. Even in India our indefatigable countryman Dr. Buchanan has discovered five original species of this genus (one of them a quince) growing naturally on the elevated regions towards the snowy mountains. His descriptions of them, however, are yet unpublished, but we have seen his specimens in the collection of A. B. Lambert, esq.

P. 109



Citrus, Medica, Lemnifera

PLATE DCX.

RUELLIA FORMOSA.

Beautiful Ruellia.

CLASS XIV. ORDER II.

DIDYNAMIA ANGIOSPERMIA. Two Chives longer. Seeds covered.

GENERIC CHARACTER.

CALYX 5-partitus. Corolla 1-petala, limbo inæquali 5-lobo. Capsula bilocularis, bivalvis, dentibus elasticis dissiliens.

CUP 5-parted. Blossom of one petal, with the limb unequally 5-lobed. Fruit a two-celled, two-valved capsule bursting with elastic teeth.

SPECIFIC CHARACTER.

RUELLIA caule suffruticoso erecto; foliis pilosis, petiolatis, integerrimis, ovatis, obtusis; pedunculis lateralibus foliis duplo seu triplo longioribus, ramosis; ramulis subtrifloris.

RUELLIA with a suffruticose erect stem; hairy, entire, oval, blunt leaves upon footstalks; the flower-stalks twice or thrice as long as the leaves and branched; the branches three-flowered.

REFERENCE TO THE PLATE.

1. The empalement.
2. A blossom spread open.
3. The seed-bud and pointal.

For this very elegant species we are indebted to Mr. Donn, Curator of the Botanic Garden at Cambridge, who informs us that it is a native of Brazil, and was introduced from Portugal in 1808 by sir Charles Cotton, bart. The plant in the Cambridge garden is now nearly four feet high, with many branches, and has continued flowering profusely since the beginning of May,

and appears as if it would continue to blossom during the summer. It is propagated by cuttings, and has as yet been kept in the hothouse.

We have found no figure or description of the plant in any author, but have seen a very fine dried specimen of it brought from Portugal by sir Thomas Gage, bart. in the herbarium of A. B. Lambert, esq. with this note affixed: "I found this *Ruellia* growing, and ripening seeds abundantly, amongst the hothouse-plants in the Botanic-Garden at Lisbon." So ornamental a plant we trust will soon be equally abundant in the hothouses of our own country.



Psyllia formosa.

by H. A. ...

PLATE DCXI.

DAVIESIA CORYMBOSA.

Corymbes Daviesia.

CLASS X. ORDER I.

DECANDRIA MONOGYNIA. Ten Chives. One Pointal.

GENERIC CHARACTER.

CALYX angulatus, simplex, quinquefidus. Corolla papilionacea. Stylus subulatus. Stigma simplex, acutum. Legumen compressum, monospermum.

CUP angled, simple, 5-cleft. Blossom butterfly-shaped. Shaft awl-shaped. Summit simple, acute. Pod compressed, one-seeded.

SPECIFIC CHARACTER.

D. corymbosa, foliis lineari-oblongis, planis, muticis; pedunculis axillaribus, geminis, corymbosis, multifloris; calyce regulari. *Smith in the Transactions of the Linnean Society, vol. 8. p. 258.*

REFERENCE TO THE PLATE.

1. Empalement.
2. The standard.
3. One of the wings.
4. The keel.
5. The chives and pointal.
6. The seed-bud and pointal.
7. A half-ripe seed-vessel.

This showy species is a native of New Holland, and was raised from seed by Mr. Gibbs, last year, in his nursery at Brompton, and is not as yet, we believe, in any other collection.

It may not be improper to observe, that the calyx of this species appears a little bilabiate, and the germen two-seeded, although one of the seeds in the

fruits we have examined was always abortive.



Daviesia corymbosa

PLATE DCXII.

PÆONIA ALBIFLORA, flore pleno.

Tartarian Pæony, double-flowered variety.

CLASS XIII. ORDER II.—V.

POLYANDRIA DIGYNIA ad PENTAGYNIAM. Many Chives. Two to Five Pointals.

GENERIC CHARACTER.

CALYX 5-phyllus. Petala 5. Styli 0. Capsulæ polyspermæ.

CUP five-leaved. Petals 5. Shafts none. Capsules many-seeded.

SPECIFIC CHARACTER.

PÆONIA foliis biternatis, foliolis ovato-lanceolatis integris nudis, capsulis recurvatis glabris. *Willd. Sp. Pl. vol. 2. p. 1222.*

REFERENCE TO THE PLATE.

1. The empalement and pointals.

The single-flowered *Pæonia albiflora* we have already figured in our first volume (Plate 64.), but never heard of the double variety until we found it in the nursery of Mr. Whitley, who informs us that he raised the plant from seeds received from Mr. Livingstone, surgeon, (in 1808,) who brought them from China under the title of the yellow Pæony. It bloomed last year with only three flowers, but has this year produced five elegant blossoms. Its great beauty certainly entitles it to a place in every flower-garden and curious collection. We shall only add to our former account of this fine species of Pæony above referred to, that Professor Pallas mentions in his *Flora Rossica* that the roots are used as food in Mongolia, where the plant

grows naturally wild, being boiled and eaten by the Tartars, who also powder the seeds to mix with their tea.



PLATE DCXIII.

IPOMŒA pendula.

Pendulous Ipomœa.

CLASS V. ORDER I.

PENTANDRIA MONOGYNIA. Five Stamens. One Style.

GENERIC CHARACTER.

CALYX quinquepartitus. Corolla infundibuliformis. Stigma capitato-globosum. Capsula 2-seu 3-locularis.

CUP five-parted. Blossom funnel-shaped. Stigma with a roundish head. Capsule of 2 or 3 cells.

SPECIFIC CHARACTER.

IPOMŒA pendula, glabra, foliis quinato-digitatis: foliolis lanceolatis, mucronulatis, margine integerrimis; extimis indivisis bifidisve: pedunculis 1-3-floris: calycis foliolis subæqualibus, obtusis, tubo corollæ ter brevioribus. *Brown Prod. Nov. Holl. et Ins. Van-Diem, vol. 1. p. 486.*

PENDULOUS *Ipomœa* with smooth five-fingered leaves: the divisions lanced, sharp-pointed, and entire at the edge; the lowermost being sometimes two-cleft: the peduncles one-to three-flowered: leaflets of the cup nearly equal, blunt, three times shorter than the tube of the blossom.

REFERENCE TO THE PLATE.

1. The empalement.
2. A blossom spread open.
3. Seed-bud and pointal, summit magnified.

This very ornamental species of *Ipomœa* is a native of the north and east coasts of New Holland, according to Mr. Brown's Prodrômus of the plants of

that country above quoted. We have seen wild specimens of it gathered by Dr. White near Port Jackson. The species is so decidedly volubilous as not only to twine round other plants that may come in its way, but twists also its own shoots round each other; and this both in its wild and cultivated state. Towards the root it appears to be woody, and is certainly perennial. The drawing was taken about the middle of July, in the curious collection of the Countess de Vandes at Bayswater, from a plant about five feet high, which was probably the first time the species blossomed in this country. The introducer we have not been able to discover.



Ipomoea pendula

PLATE DCXIV.

FUMARIA NOBILIS.

Noble Fumitory.

CLASS XVI. ORDER II.

DIADELPHIA HEXANDRIA. Two Brotherhoods. Six Stamens.

GENERIC CHARACTER.

CALYX diphyllus. Corolla ringens. Filamenta 2, membranacea, singula antheris tribus.

CUP two-leaved. Blossom gaping. Filaments two, membranaceous, each supporting three anthers.

SPECIFIC CHARACTER.

FUMARIA caule simplici, racemo terminali, bracteis oblongis acutis flore brevioribus, foliis pinnatis, foliolis subtripartito-laciniatis. *Willd. Sp. Pl. vol. 3. p. 858.*

FUMITORY with a simple stem, a terminal bunch of flowers, with oblong acute bracts shorter than the blossoms, and pinnate leaves with the leaflets jagged and three-parted.

REFERENCE TO THE PLATE.

1. A flower.

The *Fumaria nobilis* is a native of the Altaian mountains in Siberia, and was introduced to this country in the year 1783 by Mr. John Græffer, but appears to have been since lost, as we never saw living specimens until the beginning of last May, when that from which our drawing was taken was communicated by Mr. Donn from the Botanic garden at Cambridge. The plant appears well calculated for ornamenting rock-work or curious borders,

the finely cut foliage and flowers being both very graceful. It is said to have been a great favourite with Linnæus.



Fumaria nobilis

PLATE DCXV.

GLOBBA PURPUREA.

Purple Globba.

CLASS I. ORDER I.

MONANDRIA MONOGYNIA. One Stamen. One Style.

GENERIC CHARACTER.

ANTHERA duplex. Filamentum lineare, incurvatum, longissimum, appendiculatum. Stylus laxis filiformis in medio antheræ receptus. Stigma incrassatum. Nectarium utrinque bifidum. *Roscoe in the Transactions of the Linnean Society, vol. 8. p. 355.*

ANTHERS double. Filament linear, incurved, very long and appendaged. Style (or shaft) long thread-shaped, going up between the anthers. Stigma (or summit) swelling. Nectary cleft at both ends.

SPECIFIC CHARACTER.

GLOBBA scapo aphylo paniculato; bracteis calyce multo longioribus subovatis, corollâ trilobâ, filamento medio biappendiculato, nectario cordato-oblongo.

GLOBBA with a leafless paniced stem, the bracts many times longer than the cup and nearly ovate, the blossom of three lobes, the filament with two appendages in the middle, and the nectary between a heart-shape and an oblong.

REFERENCE TO THE PLATE.

1. A leaf.
 2. A flower.
 3. A front view of the same divested of the calyx.
 4. Empalement, seed-bud, and pointal.
-

No genus of plants has more puzzled botanists than the *Globba* of Linnæus, no plant having yet been discovered that agrees with his generic character; but as the possessor of his herbarium has decided that the *Globba marantina*, the only species of this genus Linnæus professes to have seen, was the same plant with that called by Dr. Roxburgh, and also by Mr. Donn in his Catalogue of the Cambridge Botanic Garden, *Colebrookia bulbifera*; (see Smith's *Exotic Botany*, page 85, where he has corrected the generic character;) this plant, also sent from India by Dr. Roxburgh to Sir Abraham Hume, bart. in 1809, now also changes its name to *Globba*. The inflorescence, which is directly from the root, whilst in all the species described by Dr. Smith it is placed above the leaves on a common stem, together with the three-lobed corolla, might doubtless be quite sufficient characters to distinguish it as a separate genus; but another having already been published in honour of Mr. Colebrook, we prefer leaving it with the congeners amongst which it was originally placed by its discoverer Dr. Roxburgh.

The delicacy and elegance of the whole plant, the rich tint of the floral leaves, and the curious construction of the blossoms, (where the long projecting filament drawn back by the style which is held by its summit protruded beyond the anthers, at one period of the inflorescence nearly resembles a half-bent bow,) have been admired by all who have seen it. The leaves only appear as the flowers decay.

The plant blossomed in the middle of last May, for the first time in this country, in the collection of Sir Abraham Hume, bart. where our drawing was taken. From a very useful manuscript catalogue of Indian plants by Dr. Roxburgh in the collection of A. B. Lambert, esq. we learn that it is a native of India, and has been found growing wild about Chittagong.

The ripe fruit we have not seen; but the germ is nearly oval, with three blunt angles and of one cell with many seeds affixed to three lateral receptacles. The plant is perennial, and may be propagated by the roots like other scitamineous plants.



Globba, purpurea

See the description of the species in the text.

PLATE DCXVI.

EUPHORBIA epithymoides.

Broad-leaved Spurge.

CLASS XI. ORDER III.

DODECANDRIA TRIGYNIA. Twelve to Nineteen Stamens. Three Styles.

GENERIC CHARACTER.

CALYX monophyllus, ventricosus. Corolla 4-seu 5-petala calyci insidens. Capsula tricocca.

CUP of one leaf, bellying. Blossom of 4 or 5 petals sitting upon the cup. Fruit a capsule with 3 cells.

SPECIFIC CHARACTER.

EUPHORBIA umbellâ quinquefidâ, radiis sub-bifidis, involucellis ovatis denticulatis, foliis integerrimis lanceolatis obtusis retusis subtus villosis, capsulis papilloso-hispidis. *Willd. Sp. Pl. vol. 2. p. 909.*

EUPHORBIA with a five-cleft umbel, the rays mostly bifid, the involucrets ovate-toothed, the leaves entire, lanced, blunt, retuse and woolly beneath; the capsules with little warts and bristly.

REFERENCE TO THE PLATE.

1. A flower.
2. The same spread open.
3. Seed-bud and pointal.

This ornamental species is a native of Austria, and was introduced, according to Donn's Catalogue, in the year 1805. Professor Jacquin has given an excellent figure and description of it in its wild state in his *Flora Austriaca*, vol. iv. p. 23, tab. 344: but it varies very considerably when

cultivated, as may be seen by comparing the figures. The figure cited to the species by Linnæus, from Columna, and from which it also derives its name, as already remarked by Jacquin, (who was certain of the identity of his species with the Linnean,) has very little resemblance to the plant; but the name of *epithymoides*, however apt to mislead, is now sanctioned by so long use, that were the plant of Columna discovered, and proved to be different, one would hardly venture now to restore to it its original appellation. The specimen was communicated by Mr. Donn last May from the Botanic Garden at Cambridge.

93/66



Euphorbia Epithymoides

PLATE DCXVII.

EUPHORBIA meloformis.

Melon-shaped Euphorbia.

CLASS XI. ORDER III.

DODECANDRIA TRIGYNIA. Twelve to Nineteen Stamens. Three Styles.

GENERIC CHARACTER.

CALYX monophyllus, ventricosus. Corolla 4- seu 5-petala calyci insidens. Capsula tricocca.

CUP of one leaf, bellying. Blossom of 4 or 5 petals sitting upon the cup. Fruit a capsule with 3 cells.

SPECIFIC CHARACTER.

EUPHORBIA subglobosa multangularis.

EUPHORBIA nearly round with many angles.

REFERENCE TO THE PLATE.

1. A flower spread open.
2. A chive magnified.

Those who are unacquainted with the versatility of this extraordinary Genus will be a little surprised at seeing a plant with so little resemblance to that on our last plate under the same generic appellation. Even the variations of Protea itself appear trifling, when compared with those of Euphorbia, which includes species with leaves opposite, alternate, scattered, whorled, and imbricated, and others leafless, like the present; plants round, oval, square, triangular and oblong; annual, herbaceous, succulent, thorny, woody, and even arboreous; and natives of all climates, from Siberia to Dusky Bay. Almost all the species agree in being replete with a milky juice which flows

from them freely when wounded, and is generally of an acrid corrosive quality. An excellent description of the *Euphorbia meloformis* was given in the second volume of the *Hortus Kewensis*, and has been since copied into the *Species Plantarum*. An engraving and description of it by Mons. Desfontaines, one of the professors in the Museum of Natural History at Paris, also ornaments the first volume of their *Annales*. The species is dioicous; and, except in the Royal Gardens at Kew, male flowers only have as yet been produced by the plants cultivated in this country, and their progeny at Paris. The stamens come to maturity at different times, those of the centre being the earliest, the woolly filaments of which persisting after the antheræ are fallen off give to the flowers a singular appearance. The species is a native of Africa towards the Cape of Good Hope, and was introduced by Mr. Masson in 1774, but is still very scarce, and requires to be kept in the dry stove or on a shelf in the hot-house, much moisture being always inimical to it. The propagation is by separating the young bulbs from their parent stock, and planting them in pots of earth nearly dry, as they retain their vegetative powers for months without either earth or water.



PLATE DCXVIII.

ANNESLEA SPINOSA.

Armed Indian Water Lily.

CLASS XIII. ORDER VII.

POLYANDRIA POLYGYNIA. Many Stamens with many Styles or Stigmas.

GENERIC CHARACTER.

CALYX superus, persistens, 4 phyllus. Petala 30-40, oblongo-lanceolata, persistentia. Stamina 60-70, incurva; antheræ subovatæ. Styli nulli. Stigmata: sulculi decem in apice cyathiformi germinis, ad umbonem centralem decurrentes; inconspicui. Germen 10-loculare, supra cyathiforme atque ex columellâ centrali protrusâ umbonatum, margine 10-dentatum. Septa duplicia. Semina loculis in singulis 2, dissepimentis prope parietem alterne affixa, obovata. Fructus: bacca subovata, calyce persistente coronata, spinosissima, 10-20-sperma. Semina matura non vidi.

Anneslea spinosa. Roxburgh.

Calyx above, persistent, 4 leaved. Petals 30 to 40, oblong-lanced, persistent. Stamens 60 to 70, incurved; anthers nearly ovate. Styles none. Stigmas: ten inconspicuous channels in the cup-shaped top of the germen, running down towards the central knob. Germen 10-celled, cup shaped above, with the central column rising like a knob in the middle; the margin 10-toothed. Partitions double. Seeds two in each cell, affixed to the partitions near the outer angle one on each side, inversely ovate. Fruit: a berry nearly ovate, crowned with the persistent calyx, exceeding thorny, 10-to 20-seeded. We have not seen the ripe seeds.

REFERENCE TO THE PLATE.

1. A flower cut open.

Seeds of this wonderful water plant, originally a native of China, were sent preserved in sugar from Calcutta by Dr. Roxburgh, in 1809, to the Most Honourable the Marquis of Blandford, in whose magnificent Aquarium at White Knights it now raises its numerous heads bristling with spines, unrolls its immense leaves, and flourishes in all its grandeur.

In vain we review the plants of its natural order for any analogy to its thorny exterior; the Nymphs and Naiads, *Nymphææ* and *Naiades*, are not more conspicuous for their elegance and beauty than for their mildness; Anneslea, like the panther, seems to unite the extremes of ferocity and beauty. The leaves are nearly orbicular, but sometimes a little extended upon one side with a corresponding notch on the side opposite, the largest being from six to eight feet in circumference; green on their upper surface, and reticulated with purplish branching veins, with a sharp curved thorn at each of their principal ramifications; their under side purple, thinly scattered over with a very fine brown pubescence, and reticulated with high raised, cellular, transparent, brown, branched veins, bearing innoxious thorns of the same form as those of the upper surface at their principal divisions. The compartments between the veins are irregular hexagons, pentagons, and rhomboids, which are again intersected by other minute veins nearly in the same manner. The leaf-stalks are centrally affixed, nearly round, very porous, and bristled with soft spines almost to where they rise from the root. The spines are hollow, of the most simple structure, and appear to be formed by a continuance of the cellular tissue of the plant protruded. See Mirbel's *Anatomie Vegetale*, fig. 24. It is only where they stand out of water, as upon the fruit and upper surface of the leaves, that they become indurated and really formidable. Very different is the nature and structure of the thorns in the genus *Rosa*, where they are proper secreting vessels or abortions of such, indurated; and may be gradually traced in the *R. damascena*, *muscosa*, *ferox*, *rubiginosa*, and many others, from the almost imperceptible hair supporting viscous matter, down to the horny lacerating thorn. In the *Geranium echinatum*, on the contrary, and some *Monsoniæ*, the spines are mere indurated persistent stipules, and in the *Astragali* indurated petioles; while in *Prunus*, *Cratægus*, *Ononis* and *Genista*, the ends of the branches indurate into thorns, and a plant of *Ononis spinosa* deprived of its verdure and dried, would appear to be only one branched thorn. Thorns are also formed from indurated bracts, abortions of roots, leaves, branches, peduncles, flowers, petioles and supernumerary stipules as in the terrible *Gleditschia*, horrid with

huge bunches of three-forked thorns from the very trunk of the tree. All these and other similar parts of plants, their real nature and use often overlooked, are in their aged and indurated state thrown together by Linnæus under the denomination of “Arma, (or Armour,) to prevent animals from injuring the plants.” Camels, however, and asses and goats, are observed to be particularly fond of thorny plants; and even quicks only escape being browsed by the cattle, by becoming woody or growing out of their reach.

The story of the *Anneslea*’s flowering under water may have probably arisen from the very short time the blossoms remain above; as, like those of the *Nymphæa*, they only rise to expand, and again gradually sink to ripen their seeds after the globules of fertilizing pollen have burst from their parent cells, phoenix like, to perish in renovating their race; and by adding circle to circle served to prolong the immoveable and immeasurable chain. That it certainly flowers above water, we can assert from our own observation; but we were informed at White Knights (where our drawing was taken last September) that it had flowered there below; which might have been owing to its artificial treatment, (We have seen the *Nymphæa rubra* flowering at the very bottom of the water in the same Aquarium.) and cannot be its natural state, unless we can bring ourselves to believe that nature has endowed it with the power of propagating itself in both elements.

We hope yet to see its magnificent foliage mantling our ponds. Have we not already taught the *Thea*—the *Camillia*—the *Ta kio*—the *Moutan*—the *Yu lan*, to resist our winters! Our water is not less temperate than our sky. The name *Anneslea* was given by Dr. Roxburgh, in honour of the Right Hon. George Annesley, Viscount of Valentia, who discovered the plant growing in the Gagra River in Oude, and also about Chittagong, when on his travels in India. The fruit abundantly distinguishes the genus from the *Euryale* of Mr. Salisbury.



PLATE DCXIX.

EUGENIA Zeylanica.

Ceylon Eugenia.

CLASS XII. ORDER I.

ICOSANDRIA MONOGYNIA. Stamens from the Cup. One Style.

GENERIC CHARACTER.

CALYX quadripartitus, superus. Petala quatuor. Bacca 1-locularis, 1-sperma.

CUP four-parted, above. Petals four. Berry of one cell, one-seeded.

SPECIFIC CHARACTER.

EUGENIA foliis subovalibus integerrimis coriaceis retusis, pedunculis unifloris solitariis vel geminatis, axillaribus lateralibusque.

EUGENIA with oval entire leathery retuse leaves, and one-flowered peduncles, either solitary or in pairs, axillary or lateral.

Eugenia zeylanica. Willd. *Sp. Pl.* vol. 2. p. 963.

REFERENCE TO THE PLATE.

1. Empalement, seed-bud and pointal.
2. The fruit, with the seed detached.

The Ceylon Eugenia is a low bushy wide-spreading shrub, with both the leaves and branches extending nearly horizontally. The flowers grow from the axils of the leaves and sides of the branches upon simple downy footstalks, which have two small bracts a little above their middle; and open in succession from the lower axils upwards. Professor Willdenow is not quite correct in stating the leaves to be impunctate, a minute dotting being visible

on both surfaces with the naked eye, and very distinctly with the aid of a common eye-glass. The fruit is of the size of a large pea, rough with minute warts, and crowned with the persistent calyx.

We have seldom seen a more striking example of the astonishing resources provided by nature for the continuation of species than this plant, whose little germen when blossoming contains sixteen infant seeds (clustering round a common centre), while the fruit admits of one only coming to maturity! Above eighty stamens are also provided for the fertilization of a solitary stigma. We have often admired the common chesnut, each germen of which when blossoming contains twelve seeds in embryo, but this yet surpasses it.

We were favoured with fresh specimens last July by A. B. Lambert, esq. from his seat at Boyton, and a foreign specimen in his collection has also enabled us to add the fruit, which has not yet ripened in this country. The plant is stated in Donn's Catalogue to have been introduced in 1798, most probably by Sir Abraham Hume, as the only plants of it that we have seen came originally from his collection.



2

Eugenia Seylanica

PLATE DCXX.

SCHINUS DENTATA.

Toothed Schinus.

CLASS XXII. ORDER IX.

DIOECIA s. *POLYGAMIA DECANDRIA*. Shafts and Chives separate on different Plants, or both on the same. Ten Chives.

GENERIC CHARACTER.

MAS. Calyx 5-fidus. Petala 5. Stamina 8-10 sub nectario rotato inserta.

FEM. Calyx et petala maris. Stigmata 3. Drupa sicca, multi-ocularis, submonosperma. Nux subturbinatus, obtusè angulatus. Embryo subreniformis, compressus. Cotyledones 2, æquales, latofalcatae. Radicula supera, cylindrica, incurva.

MALE. Cup 5-cleft. Petals 5. Chives 8-10 affixed under a wheel-shaped nectary.

FEMALE. Cup and blossom as in the male. Summits 3. Berry dry, with many cells and generally one seed. Nut nearly top-shaped with blunt angles. Embryo nearly kidney-shaped, compressed. Seed-leaves 2, equal, broadly falcate. Radicle above, cylindrical, incurved.

SPECIFIC CHARACTER.

SCHINUS foliis simplicibus dentatis.

SCHINUS with simple toothed leaves.

REFERENCE TO THE PLATE.

1. The empalement.
2. A petal magnified.
3. A chive magnified.
4. The nectary magnified.
5. A ripe berry.
6. The nut cleared from the pulp.

Specimens of this curious *Schinus* were communicated in May by the Right Honourable the Marquis of Blandford, from his gardens at White Knights. Its native country, as we are informed, is Owhyhee, an island long to be remembered in English history for the loss of the illustrious Cook;

Who, born to bless mankind, fair Science bore
To Ocean's bounds and isles unknown before;
And, while to guard barbarian lives his care,
Was murder'd by the men he sought to spare.

At what time the *Schinus dentata* was brought to England, or by whom, we have not been able to learn. It thrives very well in sheltered situations here in the open ground, and even ripens fruit in good seasons when trained against a wall. The berries are black, nearly of the size of small currants, with a sweet fleshy pulp, each berry containing a single nut which is bluntly angled, and a flattened two-leaved crooked embryo enclosed in a farinaceous perisperm of unequal thickness, having outwardly the same form as the nut.

Another species agreeing exactly in habit with the present, the toothed leaves excepted, *S. dependens*, (the *Amyris polygama* of Cavanilles and Willdenow,) has before been added to the genus by Ortega: and we cannot help noticing how remarkable it is that three plants, two of which differ so widely in habit, should yet agree so closely in their fructification. The insertion of the stamens, nectary, germen, stigmas and fruit is exactly similar; and the fruit of *Schinus Molle* figured by Gærtner (as far as his figure and description go) coincides exactly with the fruit of *S. dentata*. For the ripe fruit and some observations on the genus we are indebted to A. B. Lambert, esq.



PLATE DCXXI.

JUSSIEUA exaltata.

Tall Jussieua.

CLASS VIII or X. ORDER I.

OCTANDRIA seu DECANDRIA MONOGYNIA. Eight or Ten Stamens.
One Style.

GENERIC CHARACTER.

CALYX 4-seu 5-partitus, superus. Petala 4 seu 5. Capsula 4-seu 5-locularis, angulis dehiscens. Semina numerosa.

CUP 4-or 5-parted, above. Petals four or five. Capsule 4-or 5-celled, splitting at the corners. Seeds numerous.

SPECIFIC CHARACTER.

JUSSIEUA erecta, pubescens, foliis oblongo-lanceolatis pubescentibus punctato-scabris; floribus 4-petalis 8-andris, stigmatibus 4-lobis.

ERECT pubescent Jussieua with oblong-lanced pubescent leaves rough with little dots; flowers with 4 petals and 8 stamens, and the stigma four-lobed.

REFERENCE TO THE PLATE.

1. A petal.
2. Empalement, seed-bud, and pointal.
3. The capsule.
4. A ripe seed.
5. The same magnified.

Linnæus could scarcely have selected a happier genus in honour of the great French botanist; by whose arrangement plants agreeing in general habit

and character are not liable to be thrown to various classes for a slight difference in their number of stamens, as in the Linnæan; where octandrous and decandrous plants can only be brought together by special license.

Of the twelve species of *Jussieua* enumerated by Willdenow, and the two additional species since described by Humboldt and Bonpland, our plant, communicated from Boyton in September, most resembles the *octovalvis*, of which the Professors Swartz and Jacquin appear to have described very different varieties. Ours, however, appears to be specifically distinct from either.

Mr. Lambert informs us that the plant is a native of the East Indies, from whence the seeds were sent to him by Dr. Roxburgh, with a reference to the *Cattu Carambu* of the Hortus Malabaricus, (tom. ii. p. 97, fig. 50,) which is a very fair representation of it.

The name *exaltata*, bestowed upon it by Dr. Roxburgh, may have been suggested by comparing it with the two other Indian species, the *repens* and *suffruticosa*; but the *peruviana* must be a much taller plant, from the account given of it by Father Feuillée.



Zysania, exaltata

PLATE DCXXII.

LEPTOSPERMUM scoparium.

New Zealand Tea.

CLASS XII. ORDER I.

ICOSANDRIA MONOGYNIA. Stamens from the Calyx. One Style.

GENERIC CHARACTER.

CALYX 5-fidus, semisuperus. Petala quinque, unguiculata, staminibus longiora. Stigma capitatum. Capsula 4-seu 5-locularis, polysperma. Semina angulosa.

CUP 5-cleft, free above the middle. Petals five, clawed, and longer than the stamens. Stigma headed. Capsula 4-or 5-celled, many-seeded. Seeds angular.

SPECIFIC CHARACTER.

LEPTOSPERMUM foliis ovatis ovato-lanceolatisque mucronatis obsolete trinerviis; calycibus glabris, dentibus membranaceis coloratis.

LEPTOSPERMUM with ovate and ovate-lanced dagger-pointed faintly 3-nerved leaves, and smooth calyces with membranaceous coloured teeth.

L. scoparium. Willd. sp. pl. 2. p. 948.

Tea Plant. *Cook's Second Voyage, vol. 1. p. 100. tab. 32.*

REFERENCE TO THE PLATE.

1. Empalement, chives, and pointal.
2. The same cut open, one tip magnified.
3. Seed-bud and pointal, summit magnified.
4. A petal.

THE *Leptospermum scoparium* grows naturally in New Zealand, where it was found in Cook's first voyage of discovery in the year 1769, and was first published with an engraving of the fructification in 1776, by the two Forsters, in their *Characteres Generum Plantarum*, or Account of the Plants

they collected in that expedition. The following description of the plant, and accounts of the benefits his people derived from it, are extracted from Captain Cook's Account of his Second Voyage, vol. i. p. 99 to 101.

"The Tea plant is a small tree or shrub, with five white petals or flower-leaves, shaped like those of a rose, having smaller ones of the same figure in the intermediate spaces, and twenty or more filaments or threads. The tree sometimes grows to a moderate height, and is generally bare on the lower part, with a number of small branches growing close together towards the top. The leaves are small and pointed like those of the myrtle; it bears a dry roundish seed-case, and grows commonly in dry places near the shores. The leaves, as I have already observed, were used by many of us as tea, which has a very agreeable bitter and flavour when they are recent, but loses some of both when they are dried. When the infusion was made strong, it proved emetic to some, in the same manner as green tea.

"The beer certainly contributed not a little to the healthiness of our people. As I have already observed, we at first made it of a decoction of the spruce leaves (*Dacrydium cupressinum*); but finding that this alone made the beer too astringent, we afterwards mixed it with an equal quantity of the Tea plant, (a name it obtained in my former voyage from our using it as tea then, as we also did now,) which partly destroyed the astringency of the other, and made the beer exceedingly palatable, and esteemed by every one on board."

In the younger Forster's account of the same expedition, vol. i, p. 128 and 129, a similar account of its utility and beauty is given, with the additional information, that, in a fine soil, in thick forests, it was found from 30 to 40 feet high, and above a foot in diameter; while on a hilly arid situation he found it bearing flowers and seed when only 6 inches high.

The plant was introduced to the Royal Gardens at Kew so early as 1772, and several varieties of it are now in cultivation. It is increased both by seeds and cuttings, and requires to be kept in the Greenhouse or Conservatory.

Specimens were communicated by Mr. Donn from the botanic garden at Cambridge last May, and others in June by Mr. Milne from Fonthill.



Styxpermum, scoparium

PLATE DCXXIII.

ARDISIA ELEGANS.

Elegant Ardisia.

CLASS V. ORDER I.

PENTANDRIA MONOGYNIA. Five Stamens. One Style.

GENERIC CHARACTER.

CALYX 5-phyllus. Corolla hypocrateriformis. Antheræ magnæ, erectæ. Stigma simplex. Drupa sicca, supera, 1-sperma.

CUP five-leaved. Blossom salver-shaped. Anthers large, erect. Stigma simple. Berry dry, above, one-seeded.

SPECIFIC CHARACTER.

ARDISIA corymbis terminalibus, compositis, nutantibus; foliis lato-lanceolatis, crenatis, nitidis, margine reflexis.

ARDISIA with terminal, compound, nodding corymbs, and broad-lanced, crenated, shining leaves, with the margin reflected.

REFERENCE TO THE PLATE.

1. The empalement.
2. A blossom spread open.
3. The seed-bud and pointal.

THIS beautiful species grows naturally in moist situations and by the sides of rivulets in Pulo-Pinang, where it was discovered by the collector sent out by T. Evans, Esq., and brought to England with him on his return in 1809; and our drawing was taken from a plant of it nearly five feet high, which has been flowering beautifully this year in the collection at Stepney, from August to the end of November. The common height of the plant in Pinang, as the collector informs us, is from eight to about twelve feet.

Pl. 623.



Antasia elegans

PLATE DCXXIV

LOTUS australis.

Southern Lotus, or Bird's-foot Trefoil.

CLASS XVII. ORDER IV.

DIADELPHIA DECANDRIA. Stamens in Two Sets, Ten.

GENERIC CHARACTER.

LEGUMEN cylindricum, strictum. Filamenta sub-cuneiformia. Alæ sursum longitudinaliter conniventes.

POD cylindrical, straight. Filaments somewhat wedge-shaped. Wings united lengthways above.

SPECIFIC CHARACTER.

LOTUS herbaceus, foliis stipulisque lanceolatis obovato-lanceolatisque pubescentibus; floribus amplis paucis; capitulis longè pedunculatis: leguminibus tereti-linearibus calyce duplò longioribus.

HERBACEOUS Lotus, with lanceolate and inversely ovate downy leaves and stipules; large flowers few together, in heads upon long footstalks; the pods roundish linear, and double the length of the calyx.

REFERENCE TO THE PLATE.

1. The empalement.
2. The vexillum.
3. One of the wings.
4. The keel.
5. Chives and pointal.
6. The chives spread open and magnified.

NEW HOLLAND, so rich in new families of plants, sometimes (though rarely) also furnishes us with a few species belonging to genera common to our northern regions, as in *Convolvulus*, *Chenopodium*, *Campanula*, and our present subject, of which the specimen was communicated from Fonthill last July by Mr. Milne. By whom the species was introduced we have not been

able to learn; but we have seen dried specimens of it, brought over by the late Governor King, in the herbarium of A. B. Lambert, Esq.

The plant is of humble growth, and rather conspicuous when in blossom from its fine heads of flowers, but has not yet produced seeds in this country; and our account of the fruit is from the foreign specimens above mentioned. Like most other New Holland plants, *Lotus australis* requires to be kept in the Greenhouse, or at least sheltered in winter.

In our account of *Euphorbia meloformis* we omitted to add that the figure was taken from a fine plant in the collection of J. Vere, Esq. Kensington Gore, in August last.



Lotus, australis

PLATE DCXXV.

BARLERIA CRISTATA.

Crested Barleria.

CLASS XIV. ORDER II.

DIDYNAMIA ANGIOSPERMIA. Four Stamens in unequal Pairs. Seeds covered.

GENERIC CHARACTER.

CALYX 4-partitus, inaequalis. Stamina 2, longè minora. Capsula 4-angularis, bilocularis, bivalvis.

CUP 4-parted, unequal. Stamens 2, much smaller. Capsule 4-angled, 2-celled, 2-valved.

SPECIFIC CHARACTER.

BARLERIA foliis oblongis integerrimis; calycis foliolis duobus latioribus ciliatis, duobusque linearibus acutis.

BARLERIA with oblong entire leaves; two of the calyx-leaves broader and ciliated, and two linear acute.

REFERENCE TO THE PLATE.

1. The empalement.
2. A blossom spread open, segments of the border cut off.
3. Seed-bud and pointal.

LINNÆUS's excellent description of this species in the first edition of the *Species Plantarum* is rather at variance with his character of the genus; nor can the plant certainly be a congener of either of the two original species of Plumier, one of which indeed the late Professor Vahl found it necessary to remove to *Ruellia*. To the curious little sheath within the corolla, which encloses the germen about two-thirds of its length, in *Barleria cristata*, and one side of which is shorter than the other, we do not recollect having ever before met with any thing analogous.

We are indebted for fine specimens in blossom to T. Evans, esq., in whose collection it has continued flowering during the autumn, and by whom it was introduced to this country in 1808. The plant seems to grow freely, and is propagated by cuttings. It requires to be kept in the hot-house.



Parleria cristata

Pl. 628

PLATE DCXXVI.

GEODORUM CITRINUM.

Lemon-coloured Geodorum.

CLASS XX. ORDER I.

GYNANDRIA MONANDRIA. Style bearing the Stamens. Stamen One.

ESSENTIAL GENERIC CHARACTER.

PETALA quinque, longitudine subæqualia, patentia. Labellum cymbiforme; carinâ posticè paulò productâ. Anthera terminalis, opercularis, decidua. Massæ pollinis duæ, reniformes, cereaceæ.

PETALS five, about an equal length, spreading. Lip boat-shaped, with the keel a little prolonged behind. Anther terminal, cover-like, deciduous. Masses of pollen two, kidney-shaped, waxy.

REFERENCE TO THE PLATE.

1. A blossom spread open, divested of the lip.
2. The lip detached.

LITTLE more than sixty years ago, Mr. Miller, the best informed gardener of his time, and emphatically styled by foreigners, to whom he was known by his writings, ‘Hortulanorum Princeps;’ after forty years experience in gardening, in the last edition of his Dictionary which he published, when treating of Epidendrums, says: “It would be to little purpose to enumerate them here, as the plants cannot by any art yet known be cultivated in the ground; though, could the plants be brought to thrive by culture, many of them produce very fine flowers of uncommon forms.”—Genius and perseverance, however, have completely vanquished this prejudice, and we now cultivate more species of Epidendrum in England, or what in Mr. Miller’s time would have been considered as such, than he supposed to exist: indeed few plants are at present in greater favour with cultivators than those of the beautiful order of Orchideæ; and our present subject, discovered in Pulo-Pinang, or Prince of Wales’s Island, at the same time with the *Ardisia elegans*, is certainly not one of the least elegant. Both its peculiar habit and

character easily prevent its being confounded with any natural genus yet described, although some species have been enumerated amongst the *Limodorum*s by Dr. Roxburgh, and transposed with no more felicity from thence to *Malaxis* by Professor Willdenow.—The genus, however, is more allied to *Cymbidium* in character than either of the above; and, as far as our knowledge of the order at present extends, ought to be placed near to that in the arrangement. The other species of the genus before alluded to are the *Malaxis nutans* and *cernua* of Willdenow, *Limodorum nutans* of the Plants of Coromandel, and the *L. recurvum* of the same; all of which agree in having the same remarkable flaccidity of foliage, and recurved inflorescence.

The drawing was taken at Stepney last October, and the plant was still in blossom in the end of November.

Some idea of the prodigious extent of this family of plants may be inferred from Dr. Buchanan's having gathered more than fifty new species in his late journey through Napaul, and Mr. Brown above a hundred, (all now described in his *Prodromus*,) during his botanical expedition to New Holland and Van Diemen's Land; and from the statement of the Spanish botanists, Messrs. Ruiz and Pavon, who spent eight years botanizing in South America, that more than a thousand distinct species grow there upon the sides of the Cordilleras!



Geodorum, citrinum

PLATE DCXXVII.

BEGONIA EVANSIANA.

Evans's Begonia.

CLASS XXI. ORDER VII.

MONÆCIA POLYANDRIA. Stamens and Styles separate upon the same Plant. Stamens more than Seven.

GENERIC CHARACTER.

Mas. CALYX 0. Corolla 4-petala, petalis 2 oppositis majoribus. Stamina numerosa.

Fem. CALYX 0. Corolla 6-seu 4-petala, ut mascula. Styli 3, bifidi. Capsula infera, triangularis, alata, 3-locularis, polysperma.

Male. CUP none. Blossom 4-petalled, with two opposite petals larger than the others. Stamens numerous.

Female. CUP none. Blossom 6-or 4-petalled, as the male. Styles 3, bifid. Capsule below, triangular, winged, 3-celled, many-seeded.

SPECIFIC CHARACTER.

BEGONIA foliis inæqualiter cordatis, suprâ hispidulis, nitidis, margine inæqualibus, acutissimè serrulatis: petalis duobus lato-ovatis, duobus obovatis quadruplò majoribus: caulibus flexuoso-erectis, nodosis: axillis bulbiferis.


BEGONIA with unequally heart-shaped leaves a little hispid and shining above, with the margin irregular, most sharply and finely toothed: two of the petals broadly ovate, and four times larger than the two inversely ovate ones: the stem a little zigzag: axils bulbiferous.

REFERENCE TO THE PLATE.

1. One of the large petals.
2. A small petal.
3. The stamens, one of the anthers magnified.

FOR this very ornamental species we are also indebted to T. Evans, esq., whose collector discovered it growing about the sides and clefts of rocks near a waterfall in the interior of the Island of Pulo-Pinang in 1808. As yet it has only produced male flowers at Stepney; but the bulbs in the axils of the leaves always supply abundant means of propagating it, and its beauty and liveliness of colouring well entitle it to a place in every curious collection.

The drawing was taken at Stepney, last October, from a plant about two feet four inches in height.



Begonia, Evansiana.

PLATE DCXXVIII.

CLERODENDRUM PYRAMIDALE.

Pyramidal Clerodendrum.

CLASS XIV. ORDER II.

DIDYNAMIA ANGIOSPERMIA. Two Chives longer. Seeds covered.

GENERIC CHARACTER.

CALYX 5-fidus, campanulatus. Corollæ limbo 5-partito, æquali. Drupa 4-sperma, nuce uniloculari.

CUP 5-cleft, bell-shaped. Limb of the blossom equally 5-parted. Berry 4-seeded, nut one-celled.

SPECIFIC CHARACTER.

CLERODENDRUM foliis basi profundè cordatis, quinquelobis, margine undulatis, edentulis: panícula brachiata; pedunculis pubescentibus, trichotomis.

CLERODENDRUM with the leaves deeply heart-shaped at the base and five-lobed, waved on the margin and toothless; the panicle cross-armed; the peduncles pubescent, three-forked.

REFERENCE TO THE PLATE.

1. The empalement.
2. A blossom spread open.
3. Seed-bud and pointal.

THIS before unnoticed species of *Clerodendrum* was introduced from the Island of Pinang, where it grows naturally wild, by T. Evans, Esq. of Stepney, in 1809. The leaves are very large, a little hispid on their upper surface: the height of the plants, in the specimens which we have seen, is from three to five feet. The plant appears to grow freely, and may be propagated by cuttings, but requires to be kept in the hot-house. The drawing was taken at Stepney last October.

Pl. 622



Clerodendron pyramidale

PLATE DCXXIX.

DESMANTHUS NATANS.

Floating Desmanthus, or Aquatic Sensitive.

CLASS XXIII. ORDER I.

POLYGAMIA MONÆCIA. Various Dispositions upon one Plant.

GENERIC CHARACTER.

Hermaph. CALYX 5-dentatus. Corolla 5-petala vel 5-partita. Stamina 10. Pistillum 1. Legumen bivalve.

Neuter. Calyx 5-dentatus. Corolla 5-petala vel 5-partita vel nulla. Stamina 10, sterilia, lanceolato-dilatata.

Hermaph. CUP 5-toothed. Blossom 5-petalled or 5-parted. Stamens 10. Style 1. Pod 2-valved.

Neuter. Cup 5-toothed. Blossom 5-petalled or 5-parted or none. Stamens 10, sterile, enlarged into a lance-shape.

SPECIFIC CHARACTER.

DESMANTHUS inermis, foliis bipinnatis, partialibus trijugis, propriis tredecimjugis, spicis oblongis, interruptis, pedunculo nudo, caule tereti, radicante. *Willd. Sp. Pl. 4. p. 1044.*

UNARMED Desmanthus, with doubly pinnate leaves, the partial wings in 3 pairs, the proper in 13, the spikes oblong, interrupted, the flower-stalk naked, the stem round and rooting.

REFERENCE TO THE PLATE.

1. Empalement and blossom.
2. Stamens and pointal, one anther magnified.
3. Seed-bud and pointal, summit magnified.

THIS little delicate aquatic, growing naturally in pools and lakes of fresh water in India and Cochinchina, and quivering at every breeze, its roots having no attachment to the soil, and the foliage yielding but little to that of

the *Mimosa pudica* in sensibility, forms a most striking contrast to the *Anneslea spinosa* of our 128th number. Father Loureiro, who calls it *Neptunia oleracea*, informs us that in China and Cochinchina, where they cultivate it as a salad herb in pools and slow-flowing streams, they are accustomed to tie the plants to stakes, to prevent their floating to a distance.

A description of the plant, accompanied by an elegant figure, will be found in the 2d volume of Dr. Roxburgh's *Plants of the Coast of Coromandel*, under the name of *Mimosa natans*; but from the total dissimilarity between the fruit and that of the genus *Mimosa*, we have preferred following the arrangement of Professor Willdenow. The specimens from which the drawing was taken were obligingly communicated by Mr. Milne last September from Fonthill. The plant is annual, and according to Mr. Donn's Catalogue (where see *Mimosa natans*) was introduced to this country in 1802.



Desmanthus, natans

PLATE DCXXX.

ARDISIA LITTORALIS.

Seaside Ardisia.

CLASS V. ORDER I.

PENTANDRIA MONOGYNIA. Five Stamens. One Style.

GENERIC CHARACTER.

CALYX 5-phyllus. Corolla hypocrateriformis. Antherae magnæ, erectæ. Stigma simplex. Drupa supera, 1-sperma.

CUP 5-leaved. Blossom salver-shaped. Anthers large, erect. Stigma simple. Berry above, 1-seeded.

SPECIFIC CHARACTER.

ARDISIA corymbis axillaribus, simplicibus; foliis obovato-ellipticis, integerrimis, coriaceis, planis.

ARDISIA with axillary simple corymbs and inversely ovate-elliptic, entire, leathery, flat leaves.

REFERENCE TO THE PLATE.

1. Empalement.
2. A blossom spread open.
3. Seed-bud and pointal.

THIS species was discovered growing wild on the shores of Pinang, and introduced at the same time with the *Ardisia elegans*; and flowered in the Stepney collection this year in October and November. The plant from which the drawing was taken is about three feet high; and we are informed by the collector, that the species is considerably dwarfer in Pinang than the *A. elegans*, and also much more bushy and compact.



Ardisia, littoralis

PLATE DCXXXI.

STYRAX OFFICINALE.

Storax Tree.

CLASS X. ORDER I.

DECANDRIA MONOGYNIA. Ten Stamens. One Style.

GENERIC CHARACTER.

CALYX inferus. Corolla imo calyci inserta, subcampanulata, 3-8-partita. Antheræ 10-15. Germen triloculare, polyspermum. Drupa coriacea, subglobosa, 1-3-sperma.

CUP below. Blossom adhering to the base of the cup, nearly bell-shaped, 3-to 8-parted. Stamens 10 to 15. Germen of three cells with many seeds. Berry dry, leathery, nearly round. Seeds 1 to 3.

SPECIFIC CHARACTER.

STYRAX foliis ovatis subtus villosis, racemis simplicibus folio brevioribus. *Ait. Kew. 2. p. 75.*

STYRAX with the leaves oval, downy beneath, the racemes (bunches of flowers) simple, shorter than the leaf.

REFERENCE TO THE PLATE.

1. Empalement.
2. A blossom spread open.
3. Seed-bud and pointal.


THE Storax tree is a native of the Levant, of Italy, and of the southern parts of Provence in France. The tree is very ornamental, but requires the shelter of a wall in this climate, and while young either to be kept in the green-house or covered during severe frosts. The time of flowering is June and July, but ripe fruits are rarely produced in this country. The principal importation of Storax is said to be from Turkey, where it is obtained both by gathering that which naturally exudes from the trees, and also by making artificial incisions, to which hollow reeds are affixed to receive it. From the accounts of some

travellers published by Du Hamel, we learn that they generally adulterate it by mixing it with wax. The Storax-tree is by no means a new acquisition in this country. Gerard in his Herball, published in 1597, informs us that he had two small trees of it in his garden. It is however still a very scarce plant, and only to be found in a few of the principal collections, which, we are informed, is occasioned by the great difficulty of propagating it either by cuttings or layers.

The specimen was communicated, in the beginning of last June, from the well known garden of the Company of Apothecaries at Chelsea.



1



2



3

PLATE DCXXXII.

CYTISUS ELONGATUS.

Elongated Cytisus.

CLASS XVII. ORDER IV.

DIADELPHIA DECANDRIA. Two Brotherhoods. Ten Stamens.

GENERIC CHARACTER.

CALYX sub-bilabiatus. Stigma simplex. Legumen oblongum, compressum, polyspermum.

CUP sub-bilabiate. Stigma simple. Pod oblong, compressed, many-seeded.

SPECIFIC CHARACTER.

CYTISUS floribus pedunculatis, lateralibus, subquaternis; caule erecto: ramis elongatis; calycibus tubulosis; foliolis obovatis.—*Waldst. et Kit. Hung. v. 2. p. 200. t. 183.*

CYTISUS with lateral flowers, commonly by fours, on footstalks; the stem erect; branches elongated; cups tubular, and leaflets inversely egg-shaped.

REFERENCE TO THE PLATE.

1. Empalement.
2. The vexillum.
3. One of the alæ.
4. The carina.
5. The chives.
6. The same spread open.
7. Seed-bud and pointal.

THIS ornamental species of *Cytisus* was discovered by Francis Count Waldstein and Dr. Kitaibel, in the county of Bereghi and the Banatian forests in the eastern parts of Hungary, during their botanical excursions in that before nearly uninvestigated region, and published in their *Figures and Descriptions of the rare Plants of Hungary*, above quoted, at Vienna in 1802. The plant was introduced to this country by Mr. Donn of Cambridge in 1808, and from fine flowering specimens which he communicated last May the

present drawing was taken. The common height of the plant is said to be from three to five feet, and the thickness of the stem about an inch. It is perfectly hardy, and has not as yet been enumerated in any catalogue of plants cultivated in this country.



Cytisus elongatus

PLATE DCXXXIII.

LIATRIS ODORATISSIMA.

Sweet-scented Liatris.

CLASS XIX. ORDER I.

SYNGENESIA POLYGAMIA ÆQUALIS. Tips United. Equal Polygamy.

ESSENTIAL GENERIC CHARACTER.

RECEPTACULUM nudum. Calyx oblongus, imbricatus. Pappus plumosus, coloratus.

RECEPTACLE naked. Empalement oblong and tiled. Down feathery and coloured.

SPECIFIC CHARACTER.

LIATRIS foliis oblongis, sub-integerrimis, glabris, caulem amplexantibus. Caulis simplex, glaber. Flores paniculâ corymbosâ, divaricati. Calyx squamosus, lanceolatus, obtusus.

LIATRIS with oblong leaves nearly entire, smooth, and embracing the stem. Stem simple and smooth. Flowers grow in a corymbose panicle, straddling. The squamous empalement lanceolate and obtuse.

REFERENCE TO THE PLATE.

1. The empalement.
2. A flower.
3. The feathery down magnified.
4. A blossom spread open magnified.
5. Seed-bud and pointal.
6. A radical leaf.

WE received the specimen from which our figure is taken from A. B. Lambert, esq., in whose stove it has flowered for the first time in this country. It was brought to England last year by Mr. Fraser from South Carolina. Many attempts have been before made to introduce it, but without success, as the plants either perished on the voyage, or very soon after their

arrival. It is, we think, a great acquisition to our collections, on account of the great fragrance of its leaves when dried, which very much resembles the fine perfume of the Tonquin Bean, (*Dipterix odorata* Linn. Willd.) but more powerful, and which they will retain for many years, as Mr. Lambert has shown us some which have been kept in a box above ten years, and are as highly scented as ever. It seems at present too tender to flower without the assistance of the stove.

Pl. 63



Laurus odoratissima

PLATE DCXXXIV.

PELIOSANTHES HUMILIS.

Humble Peliosanthes.

CLASS VI. ORDER I.

HEXANDRIA MONOGYNIA. Six Stamens. One Style.

GENERIC CHARACTER.

CALYX nullus. Corolla 6-partita, subrotata; laciniis lato-ovatis, obtusis. Nectarium corollâ triplo brevius, incrassatum, integrum. Stamina sub ore nectarii affixa. Filamenta subnulla. Stylus brevissimus, obtusè trigonus. Stigma depressum. Germen inferum, 3-loculare; loculis dispermis. Bacca? subovata, carnosâ.

CUP none. Blossom 6-parted, nearly wheel-shaped; the segments broadly ovate, blunt. Nectary three times shorter than the corolla, fleshy, entire. Stamens affixed under the mouth of the nectary. Filaments scarcely any. Style short, bluntly three-sided. Stigma depressed. Germen below, three-celled; the cells two-seeded. The fruit fleshy, nearly oval.

SPECIFIC CHARACTER.

PELIOSANTHES foliis elliptico-lanceolatis subseptemnerviis scapo longioribus; bracteis flores æquantibus; racemo subovato.

PELIOSANTHES with elliptic-lanced seven-nerved leaves longer than the scape; the bracts equalling the flowers in length, and the bunch ovate.

REFERENCE TO THE PLATE.

1. A segment of the flower magnified.
2. Seed-bud and pointal.
3. Seed-bud cut transversely, magnified.

THIS is one of the species referred to in our description No. 605, and flowered last November in the collection of T. Evans, Esq. at Stepney; where we were informed, that it grows naturally in shady woods in Pinang, or Prince of Wales's Island, and was introduced to this country in 1808. Its

beauty is certainly far inferior to the Bengal species, with which however it agrees very well in essential characters; but the nectary here nearly resembles a little cup, and is not contracted above, as in that species. The height of the plant above ground is only about two inches.



Peliosanthes, humilis

PLATE DCXXXV.

CELOSIA CERNUA.

Nodding Celosia.

CLASS V. ORDER I.

PENTANDRIA MONOGYNIA. Five Stamens. One Pointal.

ESSENTIAL GENERIC CHARACTER.

CALYX triphyllus. Corolla pentapetala. Stamina basi nectarii plicata, conjuncta. Capsula horizontaliter dehiscens.

EMPALEMENT three-leaved. Blossom five-petalled. Stamens at the base of the nectary, folded, and joined together. Capsule splitting horizontally.

SPECIFIC CHARACTER.

CELOSIA floribus in spica aggregata, cernuis; foliis lanceolatis cum petiolis longis. Caulis costatus.

Celosia cernua. Roxb. MSS.

CELOSIA with flowers growing in an aggregate spike, nodding. Leaves lance-shaped, with long footstalks. Stem ribbed.

REFERENCE TO THE PLATE.

1. Empalement and blossom.
2. The stamens.
3. A stamen magnified.
4. Seed-bud and pointal, summit magnified.

THIS elegant little annual was raised from seed last year (1810) in the collection of Sir A. Hume, Bart. who communicated a fine specimen of it, from which our drawing was begun, and finished from another plant of equal beauty sent to us by A. B. Lambert, esq. last month. It is a brilliant addition to the stove. Native of the Raja Mahl hills, and not mentioned by any one but Dr. Roxburgh in his *Plants of the Coast of Coromandel*.



Celosia, cernua

PLATE DCXXXVI.

IPOMŒA INSIGNIS.

Magnificent Ipomœa.

CLASS V. ORDER I.

PENTANDRIA MONOGYNIA. Five Stamens. One Style.

GENERIC CHARACTER.

CALYX 5-partitus. Corolla campanulata v. infundibuliformis, 5-plicata. Stigma capitatum, 2-vel 3-lobum. Capsula 2-3-locularis.

CUP 5-parted. Blossom bell-or funnel-shaped, 5-plicate. Stigma headed, 2-or 3-lobed. Capsule of 2 or 3 cells.

SPECIFIC CHARACTER.

IPOMŒA caule volubili longissimo et ramosissimo, glabro; foliis crassis, glabriusculis, inferioribus palmato-quinquelobis; superioribus cordato-ovatis ovatisque; mediis subtrilobis: umbellis pedunculatis, axillaribus, compositis, multifloris.

IPOMŒA with a turning smooth stem exceedingly long and branching; the leaves fleshy, nearly smooth, the lower palmate-5-lobed; the upper ovate, or a little heart-shaped; the intermediate commonly 3-lobed: the umbels axillary upon footstalks, compound, and bearing many flowers.

REFERENCE TO THE PLATE.

1. A leaf from the base of the plant.
2. A blossom spread open.
3. Seed-bud and pointal.

THIS remarkable species of *Ipomœa* was brought from Mrs. Benyon's hothouse at Englefield, Berkshire, last August, where the plant, after rising to the top of the trellis, extends both right and left to a length of about 30 feet, branching in various directions with hundreds of bunches of flowers. The foliage is singular as to texture, varieties of form, and colour; the lower leaves being of a strong purple on their under side, and dark green above.

The plant is perennial, and continues flowering during the greatest part of the summer, but does not ripen its seed—most probably owing to the extreme luxuriance of its inflorescence. All our attempts to discover its native soil, or time of introduction, have proved abortive; neither have we been able to find that any account of it has before been published, or any specimens of it in herbariums.

The propagation is by cuttings; and we are informed that Mr. Lambert, who favoured us with the specimens, is already in possession of some very fine plants of it.



Ipomoea insignis

PLATE DCXXXVII.

TRICHILIA ODORATA.

Sweet-scented Trichilia.

CLASS X. ORDER I.

DECANDRIA MONOGYNIA. Ten Chives. One Pointal.

ESSENTIAL GENERIC CHARACTER.

CALYX 4-5-dentatus. Petala 4-5. Nectarium cylindricum, in apice dentatum, antheras 8-10 gerens. Capsula 3-locularis, 3-valvis. Semina baccata.

EMPALEMENT 4-5-toothed. Petals 4 or 5. Nectary cylindrical, toothed at the end, and bearing from 8 to 10 stamens. Capsule 3-celled, 3-valved. Seeds berries.

SPECIFIC CHARACTER.

TRICHILIA petalis quatuor et decem antheris: foliis pinnatis: foliolis lanceolatis, undulatis: floribus axillaribus: caule erecto, moschato.

TRICHILIA with four petals and ten stamens. Leaves pinnated: leaflets lance-shaped and undulated. Flowers grow from the axils of the leaves. Stem upright, and musk-scented.

REFERENCE TO THE PLATE.

1. Empalement.
2. Blossom spread open.
3. Seed-bud and pointal.

THE genus *Trichilia* is certainly at present in a confused state, from the latitude annexed to it of 4 to 5 petals, and 8 to 10 stamens. It appears to contain at least two genera. This fine species is said to have been sent over to England from the botanic garden in the Island of St. Vincent's by Doctor A. Anderson about the year 1801, and is not we believe (at present) in any other collection but that of Sir Abraham Hume, who favoured us with the specimen here delineated in fine bloom last summer (1810). It possesses the

same fragrance that is attributed to the *Trichilia moschata*, of which we have never seen specimens.



Nandina odora

PLATE DCXXXVIII.

DAVIESIA LATIFOLIA.

Broad-leaved Daviesia.

CLASS X. ORDER I.

DECANDRIA MONOGYNIA. Ten Chives. One Pointal.

GENERIC CHARACTER.

CALYX angulatus, simplex, quinquefidus. Corolla papilionacea. Stylus subulatus. Stigma simplex, acutum. Legumen compressum, monospermum.

CUP angled, simple, 5-cleft. Blossom butterfly-shaped. Shaft awl-shaped. Summit simple, acute. Pod compressed, one-seeded.

SPECIFIC CHARACTER.

DAVIESIA spicis multifloris, axillaribus: foliis late lanceolatis, undulatis: calyce regulari.

DAVIESIA with spikes of many flowers, axillary: leaves broadly lance-shaped, and undulated. Cup equal.

REFERENCE TO THE PLATE.

1. The empalement.
2. The standard.
3. One of the wings.
4. The keel.
5. The chives spread open.
6. Seed-bud and pointal.

THIS little shrub from New Holland is both new and beautiful, well adapted to grace any collection, being of easy culture, small in size, with abundance of brilliant little flowers growing from the axils of every leaf. The specimen we have delineated was communicated to us by Mr. J. Milne, from the Font Hill gardens, in fine bloom, in the month of April 1811.



Davissia latifolia

PLATE DCXXXIX.

CAREX FRASERI.

Fraser's Carex.

CLASS XXI. ORDER III.

MONÆCIA TRIANDRIA. One House. Three Stamens.

GENERIC CHARACTER.

Masculi flores.

AMENTUM imbricatum. Calyx squamosus, solitarius. Corolla nulla.

Feminei flores.

AMENTUM imbricatum. Calyx squamosus, solitarius. Corolla monopetala, apice bidentata. Stigmata 2 seu 3. Nux triquetra, corolla persistente inclusa.

Male flowers.

CATKIN tiled. Empalement skinny and solitary. Blossom none.

Female flowers.

CATKIN imbricated. Empalement skinny and solitary. Blossom one-petalled, with two teeth at the end. Summits two or three. Nut three-sided, within the persistent blossom.

SPECIFIC CHARACTER.

CAREX floribus androgynis capitibus terminalibus; masculis superioribus, femineis inferioribus. Stigma tripartitum: foliis lanceolatis, undulatis, marginibus crenulatis.

CAREX with hermaphrodite flowers growing in terminal heads; male flowers above, females beneath. Summit three-parted. Leaves lance-shaped, undulated, with finely scalloped margins.

REFERENCE TO THE PLATE.

1. A stamen magnified.
2. A blossom spread open, magnified.

FOR this curious little plant we are indebted to the late Mr. Fraser, nurseryman, Sloane Square, Chelsea; whose ardour in the pursuit of plants carried him thrice over to America in search of botanic novelty. Mr. J. Fraser informs us, that his father and himself discovered this singular species of *Carex*, in the autumn of 1808, near the Table Mountain, and upon the banks of the Catawba River in the neighbourhood of Morgan Town, North Carolina.



Carex, Fraseri

PLATE DCXL.

HELICONIA BIHAI.

Wild Plantain Tree.

CLASS V. ORDER I.

PENTANDRIA MONOGYNIA. Five Chives. One Pointal.

ESSENTIAL GENERIC CHARACTER.

SPATHA universalis partialisque. Calyx 0. Corolla 3-petala. Nectarium 2-phyllum. Capsula 3-locularis, monosperma.

SPATHE common and partial. Empalement none. Blossom 3-petalled. Nectary two-leaved. Capsule 3-celled, one-seeded.

SPECIFIC CHARACTER.

HELICONIA foliis ad apicem acutis, spadice erecto, radicalibus. Spatha disticha, multiflora: nectarii folio superiore trifido.

Heliconia Bihai. Willd. Sp. Pl.

HELICONIA with leaves pointed at the end, with an upright spadix, radical. Spathe two-sided, with many flowers. The upper leaf of the nectary trifid.

REFERENCE TO THE PLATE.

1. The plant in miniature.
2. A flower.
3. A blossom spread open.
4. Seed-bud and pointal, summit magnified.
5. A transverse section of the seed-bud.

THIS fine species of *Heliconia* is known by the common appellation of the Wild Plantain Tree, and grows in mountainous boggy places in South America, and also in most of the West India Islands. It is not generally cultivated with us, on account of the room it occupies, its elegant large foliage being but ill adapted to the confinement of the stove. Our figure was made about the middle of April, from a plant in the collection of the Countess de Vandes.



PLATE DCXLI.

PROSTANTHERA LASIANTHOS.

Downy-flowered Prostanthera.

CLASS XIV. ORDER I.

DIDYNAMIA GYMNOSPERMIA. Two Chives longer. Seeds naked.

GENERIC CHARACTER.

CALYX post florescentiam clausus, utroque labio integerrimo. Corolla monopetala, lanata, ringens; antheris appendiculatis: baccæ quatuor, monospermæ: foliis petiolatis: pedunculis lanceolatis, serratis: floribus paniculis axillaribus terminalibusque: ramulis oppositis, quadrangularibus.

Habitat in Capite Van Diemen.

Labillardiere Nova Hollandia, vol. ii. p. 18. tab. 157.

THE empalement after flowering shuts, both segments entire. Blossom one-petalled, woolly, gaping: chives with an appendage: berries four, one-seeded: leaves with petioles: footstalks lance-shaped, and sawed: flowers terminate the branches in panicles from the axils of the leaves: branches opposite, and four-sided.

Native of Van Diemen's Land.

REFERENCE TO THE PLATE.


1. The empalement, seed-bud and pointal.
2. The same as it appears after flowering.
3. A blossom spread open.
4. A chive magnified.
5. The same shown from the inner side, magnified.

THIS fine plant is perfectly new to the British gardens, and the only species of a genus named and figured by Labillardiere in his *Nova Hollandia*, or Description of New South Wales.

It was raised from seed in the summer of 1807, in the collection of the Right Hon. Lord Grenville, at Dropmore, who favoured us with specimens in full bloom, last June, 1811, being the first time of its flowering in this

country. His Lordship's botanic gardener informs us that the plant is about eight feet high, and the stem one inch and a quarter in diameter; that during the first year it had the appearance of being herbaceous, but afterwards became woody; its growth is very luxuriant, flowering at the termination of every branch, and seems as if it would thrive in any sort of earth, and was nearly if not quite hardy, as the roots remained in the ground one year at Dropmore, and shot up strong again in the summer. It may be increased by cuttings, and will be found not only an acquisition in point of novelty, but ornamental from its numerous light graceful blossoms, which possess an agreeable fragrance.

N. 64.



Prostanthera laevis

PLATE DCXLII.

GOMPHOLOBIUM GRANDIFLORUM.

Large-flowered Air Pod.

CLASS X. ORDER I.

DECANDRIA MONOGYNIA. Ten Chives. One Pointal.

ESSENTIAL GENERIC CHARACTER.

CALYX campanulatus, simplex, alte 5-fidus. Corolla papilionacea. Stigma simplex, acutum. Legumen inflatum, sphæricum, uniloculare, polyspermum.

EMPALEMENT bell-shaped, simple, deeply five-cleft. Blossom butterfly-shaped. Summit simple, acute. Pod inflated, spherical, one-celled, many-seeded.

SPECIFIC CHARACTER.

GOMPHOLBIUM foliis ternis vel quinis, linearibus, apice mucronatis: corollis coccineis: pedunculis basin versus bibracteatis: caule volubili.

GOMPHOLOBIUM with leaves in threes and fives, linear, and mucronated at the end: blossoms of a scarlet colour: peduncles with two floral leaves near the base: stem climbing.



REFERENCE TO THE PLATE.

1. The empalement and floral leaves.
2. The standard.
3. One of the wings.
4. The keel.
5. The chives and pointal.
6. Seed-bud and pointal.
7. A ripe seed-vessel.
8. The same split open.

THIS nondescript species of Gompholobium is, when in flower, the most elegant of the genus, but when out of bloom few plants have less attraction.

It was first raised in the Royal gardens at Kew, about the year 1809, and is as yet in very few collections. Our figure represents an entire plant from

the conservatory of the Countess de Vandes, which flowered in the month of June 1811, and perfected its seed about the middle of September, at which period its slender branches had grown nearly six inches longer (than they were when in bloom), and, twining round a small stick placed for their support, had every appearance of becoming a graceful climbing plant.



Mimulus, grandiflorum

PLATE DCXLIII.

JUSTICIA BICOLOR.

Two-coloured-flowered Justicia.

CLASS II. ORDER I.

DIANDRIA MONOGYNIA. Two Chives. One Pointal.

ESSENTIAL GENERIC CHARACTER.

CALYX simplex seu duplex. Corolla 1-petala, irregularis. Capsula ungue elastico dissiliens: dissepimentum contrarium adnatum.

EMPALEMENT simple or double. Blossom one-petalled, irregular. Capsule splitting with an elastic claw: partition contrary to the valve affixed.

SPECIFIC CHARACTER.

JUSTICIA fruticosa: foliis ellipticis: floribus plerumque ternis: pedicellis calycibusque hirsutis: caule terragono.

Habitat in Insula Jamaicæ.

JUSTICIA with a shrubby stem: leaves elliptic: flowers mostly by threes: footstalks and calyx hairy: stem four-sided.

Native of Jamaica.

REFERENCE TO THE PLATE.

1. The empalement.
2. A blossom spread open.
3. Seed-bud and pointal, summit magnified.

THIS ornamental stove plant was raised from West India seed, by Mr. J. Milne, nurseryman at Fonthill, about the year 1807; and from a fine specimen communicated by him in June 1811 our figure was delineated.

It is a perfectly new species, and has not hitherto, we believe, been either figured or described.



Justicia, bicolor

PLATE DCXLIV.

LOBELIA SPECULUM.

Lobel's Venus's Looking-glass.

CLASS XIX. ORDER VI.

SYNGENESIA MONOGAMIA. Tips united. Flowers simple.

ESSENTIAL GENERIC CHARACTER.

CALYX 5-fidus. Corolla 1-petala, irregularis. Capsula infera, 2-seu 3-locularis.

CUP 5-cleft. Blossom 1-petalled, irregular. Capsule beneath, 2-or 3-celled.

SPECIFIC CHARACTER.

LOBELIA foliis linearibus, integerrimis, superne acutis, ad basin dentatis, et ad apicem latioribus: corollis atro-purpureis, laciniis subequalibus: frutex pygmeus, ramulis numerosis tenuissimis patentibus.

LOBELIA with linear leaves, entire, and pointed on the upper part of the plant, but toothed on the lower part, and broader at the end: Blossoms of a dark purple colour, with nearly equal segments. A dwarf shrub, with numerous slender spreading branches.

REFERENCE TO THE PLATE.

1. A blossom spread open.
2. The empalement, chives, and pointal, tips and summit magnified.

OF this perfectly new species of Lobelia we do not find either figure or description in any author ancient or modern. The resemblance it bears to the *Campanula Speculum*, or Bell-flowered Venus's Looking-glass, induced us to give it the same specific title. It is a delicate slender little annual, with an abundant succession of dark purple flowers, which appear to great advantage when the sun shines. In the regularity of its corolla it differs essentially from the genus Lobelia, and in a manuscript of the late Dr. Solander we find an intention to separate it by the generic title of *Speculare*.



Lobelia speculum

PLATE DCXLV.

EPIDENDRUM FRAGRANS.

Sweet-scented Epidendrum.

CLASS XX. ORDER I.

GYNANDRIA DIANDRIA. Chives on the Pointal. Two Chives.

ESSENTIAL GENERIC CHARACTER.

NECTARIUM turbinatum, obliquum, reflexum.

HONEY-CUP top-shaped, oblique, and reflexed. See Pl. XIII. Vol. I.
Epidendrum cochleatum.

SPECIFIC CHARACTER.

EPIDENDRUM folio lanceolate bulbo innato, scapo abbreviato multifloro,
lamina labelli cordata acuta.

Willdenow's Sp. Pl.

EPIDENDRUM with a lance-shaped leaf upon the bulb, with a short stalk,
many-flowered.

REFERENCE TO THE PLATE.

1. The empalement, chives, and pointal.
2. The honey-cup.

THIS species of *Epidendrum* is an old inhabitant of the hot-house, and much esteemed for the fragrance of its flowers, and the only one at present known to us (the *angustifolium* excepted) with a single leaf. It is enumerated in the *Species Plantarum* of Willdenow, as described by Swartz; but we do not find that any figure of it has before been published.



Epidendrum fragrans

PLATE DCXLVI.

PROTEA RADIATA.

Radiated Protea.

CLASS IV. ORDER I.

TETRANDRIA MONOGYNIA. Four Chives. One Pointal.

ESSENTIAL GENERIC CHARACTER.

COROLLA 4-fida seu 4-petala. Antheræ lineares, petalis infra apices insertæ. Calyx proprius nullus. Semina solitaria.

BLOSSOM four-cleft, or of four petals. Tips linear, inserted into the petals below the points. Cup proper none. Seeds solitary.

SPECIFIC CHARACTER.

PROTEA foliis obtuse lanceolatis, obliquis, horizontaliter sitis: squamis calycinis spathulatis, rubris, tomentosis, marginibus pilosis: caule erecto, bipedali.

PROTEA with leaves obtusely lance-shaped, standing sideways in an horizontal direction: scales of the empalement spathula-shaped, red, and downy, with hairy margins. Stem upright, and about two feet high.

REFERENCE TO THE PLATE.

1. A flower complete.
2. The empalement and flowers of a green variety.

THIS Protea is nearest allied to the *P. coronata* published in the Seventh Volume of this Work, Pl. 469, but differs essentially from it (and also from every other species of Protea at present known to us) in having so few flowers, that the centre looks like an empty cup, the flowers being spread out towards the imbricated scales of the empalement in a circular manner; whence its specific title of radiata. In the bud state it does not promise that splendid appearance which it exhibits when expanded. Our figure represents a plant from the nursery of Mr. Knight, of the King's Road, Chelsea. At the

base we have added the head of flowers of a green variety, communicated to us from the Hammersmith Nursery, raised from Cape seed, at the same time as the red, about the year 1809, and which flowered for the first time in this country last autumn (1811).

In our last Number we forgot to mention that the figure of the *Lobelia Speculum* was taken from a plant in the nursery of Messrs. Colville, and the *Epidendrum fragrans* from the collection of J. Vere, esq.



Protea radiata

PLATE DCXLVII.

ANDROSACE CORONOPIFOLIA.

Buck's-horn Plantain-leaved Androsace.

CLASS V. ORDER I.

PENTANDRIA MONOGYNIA. Five Stamens. One Style.

GENERIC CHARACTER.

INVOLUCRUM umbellatum. Corolla hypocrateriformis, 5-loba, ore glanduloso. Capsula quinquevalvis.

UMBEL with an involucre. Corolla salver-shaped, five-lobed, with glands at the mouth. Capsule with five valves.

SPECIFIC CHARACTER.

ANDROSACE foliis lineari-lanceolatis dentatis, umbellæ radiis laxis subcapillaribus numerosis, corollis calyce duplo longioribus.

ANDROSACE with linear-lanced toothed leaves: the rays of the umbel hair-like, loose, very numerous, and the blossoms double the length of the calyx.

REFERENCE TO THE PLATE.

1. The empalement.
2. A blossom spread open.
3. Seed-bud and pointal, summit magnified.
4. A ripe seed-vessel.
5. The same opened to expose the seeds.

OUR drawing of this elegant little plant was taken at the nursery of Messrs. Malcolm and Co., at Kensington, July 1811. No figure or description of the species has before been published; but we have seen wild specimens of it in the herbarium of A. B. Lambert, esq., gathered by the Swedish naturalist Laxman, near the Lake Baical in Siberia, and marked *Androsace angustifolia*. The height of the plants in cultivation with us varies from three to ten inches, the number of rays in the central umbel are from twelve to twenty, and in the lateral ones from six to fourteen; the plant is annual, and

generally sows itself upon the borders where it grows. We are informed it was introduced about 1806, by Mr. Bell, of Sion Gate near Brentford, who received seeds of it from Siberia, under the name of *Androsace lactea*.



Androsace coronopifolia

PLATE DCXLVIII.

CROTOLARIA SALTIANA.

Salt's Crotonaria.

CLASS XVII. ORDER IV.

DIADELPHIA DECANDRIA. Two Brotherhoods. Ten Chives.

ESSENTIAL GENERIC CHARACTER.

LEGUMEN turgidum, inflatum, pedicellatum: filamenta connata, cum fissura dorsali.

POD turgid, inflated, pedicelled. Chives conjoined, with a dorsal fissure.

SPECIFIC CHARACTER.

CROTOLARIA frutescens, floribus spicatis: ramis teretibus, tomentosis: foliis ternatis: foliolis ovatis, tomentosis: petiolis uncialibus.

SHRUBBY Crotonaria. Flowers grow in spikes: branches are round and downy. Leaves ternate. Leaflets ovate and downy. Petioles an inch long.

REFERENCE TO THE PLATE.

1. The empalement.
2. The standard.
3. The under-side of the same.
4. One of the wings.
5. The keel.
6. The chives.
7. Seed-bud and pointal.

THIS nondescript species of Crotonaria was introduced by Mr. Salt last April (1811) from the distant region of Abyssinia. We have therefore named it after him, in compliment to that ardour, which in the pursuit of natural history has twice impelled him to undertake a journey to so remote a country. Our figure represents a fine specimen communicated to us by A. B. Lambert, esq., who raised it from seed in his hot-stove at Boyton, and where it flowered last summer (1811) for the first time in this country.

Pl. 640



brodiaea, Salicaria

PLATE DCXLIX.

ALSTROMERIA EDULIS.

Eatable Alstromeria.

CLASS VI. ORDER I.

HEXANDRIA MONOGYNIA. Six Chives. One Pointal.

GENERIC CHARACTER.

COROLLA 6-petala, sub-bilabiata: petalis 2 inferioribus basi-tubulosis. Stamina declinata.

BLOSSOM 6-petalled, nearly bilabiate: two of the lower petals are tubular at the base. Stamens standing sideways.

SPECIFIC CHARACTER.

ALSTROMERIA foliis alternis, ellipticis, lanceolatis, acuminatis, glabris: floribus terminalibus, cernuis, pedunculis longis. Caulis volubilis. Radices tuberosæ, globosæ.

ALSTROMERIA with alternate elliptic leaves, lance-shaped, pointed, and smooth: flowers terminate the branches upon long footstalks, nodding. Stem twining. Roots tuberous and round.

REFERENCE TO THE PLATE.

1. A flower spread open.
2. The same shown from the outer side.
3. A half ripe seed-vessel.

THIS elegant plant was introduced to this country by T. Evans, Esq., who received it in 1801 from the late Dr. Anderson, of the Botanic Garden in the Island of St. Vincent, and which flowered for the first time in the hot-house of the Countess de Vandes at Bayswater last autumn (1811).

The only figure of it extant is a good coloured one recently published in the *Flore des Antilles*, by F. R. Tussac, a colonist of Saint Domingo; who describes it as being eagerly sought after by the Negroes, not for the beauty

of its flowers, but for the sake of the roots, which not only serve them for food, but for an article of commerce, which they take with them when they travel to the Cape of Good Hope, and sell under the title of white Jerusalem artichokes.

This species very much resembles the *A. Salsilla*. The principal and almost only distinction is, that the roots of the *Salsilla* are long and ligneous, while those of the *edulis* are round and succulent, about the size of young potatoes, and when boiled are said to be a light and delicate food. A farinaceous or mealy substance is also made of them, from which cream is made, wholesome and very agreeable to the taste.—All the species which compose this fine genus are to be found in the superb gardens of the Incas of Peru, where it is indigenous.



Alstromeria, edulis

PLATE DCL.

XERANTHEMUM FASCICULATUM, *var. flore rubro.*

Bundled-leaved Everlasting Flower; red-flowered variety.

CLASS XIX. ORDER II.

SYNGENESIA POLYGAMIA SUPERFLUA. Tips united. Superfluous Pointals.

ESSENTIAL GENERIC CHARACTER.

RECEPTACULUM paleaceum. Pappus setaceus. Calyx imbricatus, radiatus; radio colorato.

RECEPTACLE chaffy. Feather bristly. Cup tiled, rayed; the ray coloured.

See *Xeranthemum speciosissimum*, Pl. LI. Vol. I.

SPECIFIC CHARACTER.


XERANTHEMUM foliis caulinis linearibus, fasciculatis, longissimis: floribus solitariis; squamis calycinis lanceolatis, rubris.

EVERLASTING FLOWER with the stem-leaves linear, roundish, bundled, and very long. Flowers solitary; the scales of the cup lance-shaped and red.

REFERENCE TO THE PLATE.

1. An hermaphrodite floret with its seed magnified.
2. Seed-bud, chives, and pointal, magnified.

THIS *Xeranthemum* was raised from seed brought over from the Cape of Good Hope by Mr. Niven in 1809, and resembles in its habit and foliage the *X. fasciculata* figured in the Fourth Volume of this Work, Plates 242 and 279, but is much more esteemed than either of them for the brilliance of its fine red flowers. It requires no other treatment than what is common to most of the genus, and continues in flower from May till October.



Ceranthemum, fasciculatum. Vår flore rubro

PLATE DCLI.

CYMBIDIUM ANDERSONII.

Anderson's Cymbidium.

CLASS XX. ORDER I.

GYNANDRIA DIANDRIA. Chives on the Pointals. Two Chives.

ESSENTIAL GENERIC CHARACTER.

COROLLA 4-5-petala, erecta vel patens, labello basi concavo. Anthera opercularis, decidua.

BLOSSOM 4-5-petalled, upright or spreading, with a nectary concave at the base. Tips covered, deciduous.

SPECIFIC CHARACTER.

CYMBIDIUM caule elato: foliis vaginatis, lanceolatis, costatis: floribus in spicis longis ramosis: corollis flavis.

CYMBIDIUM with a tall stem: leaves sheathed, lance-shaped, and ribbed. Flowers grow in long branched spikes: blossoms yellow.

REFERENCE TO THE PLATE.

1. A flower without the nectarium.
2. The nectarium.

OUR figure represents a nondescript species of *Cymbidium* communicated to us by T. Evans, Esq., in whose collection at Stepney it flowered last summer (1811) for the first time in this country. We have given it the specific title of *Andersonii*, as a tribute of respect to the late Dr. Anderson, by whom it was introduced in 1804, with many other fine new plants, from the botanic garden in the island of St. Vincent, and may be considered as a valuable addition to the hot stove, from its great height and numerous large yellow blossoms.

✱ In the description of the *Alstromeria edulis* of our last Number, Cape of Good Hope is inserted instead of Cape François, a mistake occasioned by our writing it off in a hurry along with some pages of the *Erica*, a tribe of plants (with few exceptions) all natives of the Cape of Good Hope. We detected the error as soon as printed, but did not think it of sufficient consequence to cancel the page and prolong its publication, as it had already been too long delayed; and should have only noticed it in the Errata at the end of the Volume, (which is nearly terminated,) were it not for the partial criticism in the Monthly Magazine given by an interested friend to the Botanical Magazine, who probably imagines the sale of our work is detrimental to that publication, and reminds us of the story of the man with a very short nose, who was always railing at the long nose of another person; which led to the conclusion that he thought it was made at the expense of his short one. But as our readers may think these observations long enough, they will doubtless think those of the botanical report still more so, as in the comments on the *Protea radiata*, fifteen lines are nearly filled with technical phrases, in an ineffectual attempt to explain the meaning of two botanical terms which he allows to be differently accepted by different authors, and at last leaves off just where he began. But meeting with a better opportunity in *Alstromeria edulis*, he has not only made the most of that, but rather more than enough of it, verifying the old adage, that those who can make something of nothing will make a little go a great way.



PLATE DCLII.

XERANTHEMUM HUMILE.

Low-growing Xeranthemum.

CLASS XIX. ORDER II.

SYNGENESIA POLYGAMIA SUPERFLUA. Tips united. Superfluous Pointals.

ESSENTIAL GENERIC CHARACTER.

RECEPTACULUM paleaceum, Pappus setaceus. Calyx imbricatus, radiatus, radio colorato.

RECEPTACLE chaffy. Feather bristly. Cup tiled, rayed, the ray coloured.

See *Xeranthemum speciosissimum*, Pl. LI. Vol. I.

SPECIFIC CHARACTER.

XERANTHEMUM foliis subulatis sub-vestitum. Caulis humilis, robustus, ramulis procumbentibus: floribus magnis rubris.

XERANTHEMUM with awl-shaped leaves nearly clothing the stem. Stem low and stout, with branches hanging down. Flowers large, and of a red colour.

REFERENCE TO THE PLATE.

1. An hermaphrodite floret with its seed magnified.

THIS fine dwarf species of *Xeranthemum* was introduced from the Cape of Good Hope at the same time with the *X. fasciculatum rubrum* by Mr. Niven. It is a hardy robust green-house shrub, of humble growth, and appears as if it would always retain that character. It requires an airy situation, and when watered (which should not be often) to be thoroughly wetted, as partial watering is a general destroyer of plants, by suffering the atmosphere to exhale the humidity before it has half penetrated to the bottom; which by repeated wetting rots the upper part, and leaves the roots below to starve for want of moisture.



Ceranthemum, humilis

PLATE DCLIII.

CORRÆA SPECIOSA.

Showy Corræa.

CLASS VIII. ORDER I.

OCTANDRIA MONOGYNIA. Eight Chives. One Pointal.

ESSENTIAL GENERIC CHARACTER.

CALYX monophyllus, quadri-dentatus. Corolla 4-petala. Stamina octo. Germen superum. Capsula quadrilocularis. Semina quatuor.

EMPALEMENT one leafed, four-toothed. Blossom four-petalled. Eight chives. Seed-bud above. Capsule four cells. Seeds four.

SPECIFIC CHARACTER.

CORRÆA foliis oblongis, oppositis, undulatis, hirsutis, supra viridibus, subtus tomentosis, ferrugineis: ramulis oppositis, alternis, ferrugineis, scabre pilosis: corollis bicoloribus, scabris.

CORRÆA with oblong opposite leaves, waved, and harshly haired, green on the upper surface and downy and rusty beneath. Branches opposite, alternate, rusty, and roughly haired. Blossoms two-coloured, and rough.

REFERENCE TO THE PLATE.

1. A blossom spread open.
2. One of the chives.
3. Empalement, seed-bud, and pointal.
4. Seed-bud magnified.

THIS elegant species of Corræa, a native of New Holland, is enumerated in Donn's catalogue by the specific title of speciosa, but has been overlooked in the new edition of the Hortus Kewensis. It is a hardy green-house plant with brilliant flowers in great succession, and may be propagated by cuttings. It requires an airy situation, and must not be too often watered. Our figure represents one half of a fine plant in the collection of Messrs. Colvilles.



Correa, speciosa

PLATE DCLIV.

GNAPHALIUM EXIMIUM.

Showy Gnaphalium.

CLASS XIX. ORDER II.

SYNGENESIS POLYGAMIA SUPERFLUA. Tips united. Superfluous Pointals.

ESSENTIAL GENERIC CHARACTER.

RECEPTACULUM nudum. Pappus pilosus vel plumosus. Calyx, imbricatus, radiatus, radio colorato.

RECEPTACLE naked. Down hairy or feathery. Empalement imbricated, rayed, with the ray coloured.

SPECIFIC CHARACTER.

GNAPHALIUM foliis sessilibus, ovatis, confertis, erecto-patentibus, lanatis: floribus terminalibus, paniculatis.

GNAPHALIUM with sessile ovate leaves crowded together, erect, spreading, and woolly: flowers paniculated and terminal.

REFERENCE TO THE PLATE.

1. One of the scales of the empalement.
2. A floret.
3. The chives and pointal.
4. Seed-bud and pointal.

THIS magnificent species of *Gnaphalium* was introduced to the British gardens from the Cape of Good Hope by Capt. W. Paterson in 1704, and is said to be found in a wild state on the borders of the Caffre country 500 miles from the Cape. It flowers in perfection during the months of July and August, and when in the bud state much resembles a rich ripe fruit, with leaves like flannel. The only figure we have seen of it is a small one given in the Botanical Magazine many years ago, before it was well naturalized, and on that account better adapted to the scale of their work. Our figure was

taken from the Clapham collection when in its greatest splendour, and where several fine plants of it flowered with a luxuriance fully equal to what we have represented. It should be kept in an airy part of the greenhouse; and when watered, care should be taken not to wet the leaves, as that frequently destroys the plant.

N. 652



Gnaphalium Eximium

PLATE DCLV.

CERBERA MANGHAS.

Poisonous-fruited Cerbera.

CLASS V. ORDER I.

PENTANDRIA MONOGYNIA. Five Stamens. One Pointal.

ESSENTIAL GENERIC CHARACTER.

FLORES contorti. Corolla infundibuliformis, pulposa. Drupa monosperma.

FLOWERS contorted. Blossom funnel-shaped, pulpy. Seed-vessel one-seeded.

SPECIFIC CHARACTER.

CERBERA foliis alternis, lato-lanceolatis, glabris: ramis floriferis terminalibus: corolla alba, infundibuliformi: perianthio 5-phyllo: foliolis lanceolatis, deciduis: fructu persistente.

CERBERA with leaves alternate, broadly lance-shaped, and smooth: flower-branches terminal: blossoms white and funnel-shaped: empalement 5-leaved: leaflets lance-shaped and deciduous: fruit persistent.

REFERENCE TO THE PLATE.

1. A blossom spread open.
2. Empalement, summit magnified.

OUR drawing represents a new species of *Cerbera* introduced from Pulo-Pinang in 1809, and which flowered late in the autumn of 1812, for the first time in this country, in the hot stove of T. Evans, Esq. whose collector informs us that it grows in abundance by the sea-side, and is a handsome shrub about three feet high. By the natives it is called Devil's flower, a title they apply indiscriminately to all noxious plants. But although it is considered as poisonous (as are all the *Cerberas* at present known) the natives of Amboyna are said by Rumphius to use it medicinally as a purgative, but that it is too powerful to be taken with safety by strangers. In

Macassar an oil is expressed from the fruit, which is burnt in lamps, and emits a disagreeable smell. It bears a variety of names. By the Dutch it is called Mùkhout; in Portugal and Malay, Manga Brava; at Amboyna, Wabba; in Macassar, Lambuto Baleyice, Bintaro and Sassura Utan, and by some Caju Mattu, Buta or Blinhout. There are three uncoloured figures of it; one in Burmann's *Zeylanica*, page 150, tab. 70. figure 1; another in Rumphius's *Amboynensis*, vol. ii. page 243, figure 81; and in Rheede's *Hortus Malabancus*, page 71, tab. 3. there is a representation of it by the title of Odallam; and, although a rough draft, it is certainly not the worst figure of the plant.

Pl. Ind.



Cerbera manghas

PLATE DCLVI.

CRASSULA PERFOLIATA.

Perfoliate-leaved Crassula.

CLASS V. ORDER V.

PENTANDRIA PENTAGYNIA. Five Chives. Five Pointals.

ESSENTIAL GENERIC CHARACTER.

CALYX pentaphyllus. Petala quinque. Squamæ quinque nectariferæ ad basin germinis.

EMPALEMENT five-leaved. Petals five. Five honey-bearing pores at the base of the germ.

SPECIFIC CHARACTER.

CRASSULA altissima: foliis subulatis, crassis, amplexicaulibus, subtus convexis, glaucis.

CRASSULA with a tall stem: leaves awl-shaped, thick, and surrounding the stem, convex beneath, and of a sea-green colour.

REFERENCE TO THE PLATE.

1. A flower.
2. The pointals.

OF this species of *Crassula* we know of no other figure extant, but a very small uncoloured one in the *Hortus Elthamensis* of Dillenius. Our specimen was taken from the hot stove of the Countess de Vandes, where it flowered for the first time in this country in 1809—in size and grandeur like the *C. obliqua*, with an equal profusion of blossoms, and bending down as if the weight of the terminal mass of flowers was too much for the stem to support; which we do not consider as the positive character of it, but only arising from the situation in which it stood; being too near the glass, which most likely drew too much of its strength out in length of stalk, and in all

probability occasioned the loss of the plant, as it damped off very soon after flowering.



PLATE DCLVII.

PASSIFLORA LUNATA.

Crescent-leaved Passion-flower.

CLASS XX. ORDER V.

GYNANDRIA PENTANDRIA. Chives on the Pointals. Five Chives.

ESSENTIAL GENERIC CHARACTER.

TRIGYNIA. Calyx 5-phyllus. Petala 5. Nectarium coroniforme. Bacca pedicellata.

THREE styles. Cup 5-leaved. Petals 5. Honey-cup forming a crown. Berry standing on a footstalk.

SPECIFIC CHARACTER.

PASSIFLORA floribus axillaribus, binis: foliis lunatis: flore parvo, albo: fructu succulento, ovato.

PASSIFLORA. Flowers axillary, by twos: leaves crescent-formed; flower small and white: fruit oval and succulent.

REFERENCE TO THE PLATE.

1. A flower spread open.

THIS Passiflora is (like every one of the genus) a graceful meandering plant, and calculated to make an ugly wall look beautiful. It bears an abundant succession of delicate small flowers, which are fragrant, and open early in the day, and during the summer and autumnal months is a great ornament to the hot stove. Our drawing was made from the collection of the Countess de Vandes at Bayswater.

Pl. 057



Passiflora lunata

From the Bot. Garden, Vienna
by H. Androsch & H. Knappe

PLATE DCLVIII.

STYLEDIUM GRAMINIFOLIUM.

Grass-leaved Styledium.

CLASS XX. ORDER II.

GYNANDRIA DIANDRIA. Chives on the Pointals. Two Chives.

ESSENTIAL GENERIC CHARACTER.

CALYX bilabiatus. Cor. tubulosa, irregulariter 5-partita, laciniis interioribus tripartitis. Capsula bilocularis, bivalvis, polysperma.

EMPALEMENT two-lipped. Blossom tabular, irregularly 5-parted: segments of the interior 3-parted. Capsule two-locular, two-valved, many-seeded.

SPECIFIC CHARACTER.

STYLEDIUM foliis radicalibus lineari-lanceolatis: floribus spicatis: corollis 4-partitis. *Swartz. Nov. Act. Soc. Natur. Scrut. Berol. vol. 5. fig. 1. Billardiére, Nova Hollandia, tab. 215.*

STYLEDIUM with radical leaves linear and lance-shaped: flowers grow in spikes: blossom 4-parted.

REFERENCE TO THE PLATE.

1. The empalement.
2. A blossom spread open.
3. Chives and pointals, summit magnified, both from the upper and under surface.
4. A half-ripe seed-vessel.

THIS curious genus, in its style of action, displays an appearance of locomotive power superior to the Mimosas; for, whilst they shrink from the touch, this boldly comes forward with singular elasticity; and although it may not exceed in beauty, it certainly possesses a more moving power than the utmost grace and beauty of other plants can boast. Like most of the productions of New Holland, it bears a strict appearance, and is seen to most advantage in a plant about three years old, such as our figure represents, as it flowered for the first time in this country at the Hammersmith nursery.



Stylidium, graminifolium

PLATE DCLIX.

LOBELIA FULGENS.

Refulgent-flowered Lobelia.

CLASS XIX. ORDER VI.

SYNGENESIA MONOGAMIA. Tips united. Flowers simple.

ESSENTIAL GENERIC CHARACTER.

CALYX 5-fidus. Corolla 1-petala, irregularis. Capsula infera, 2-seu 3-ocularis.

CUP 5-cleft. Blossom 1-petalled, irregular. Capsule beneath, 2-or 3-celled.

SPECIFIC CHARACTER.

LOBELIA foliis lanceolatis obsolete denticulatis, supra tomentosis, glabris, subtus lucidis: racemis compositis, terminalibus.

LOBELIA with lance-shaped leaves obscurely toothed, downy on the upper surface, smooth, and shining beneath: the racemes compound and terminal.

REFERENCE TO THE PLATE.

1. The chives.
2. Empalement and pointal, summit magnified.

AMONGST the Lobelias, this elegant species claims altogether precedence; superior in colour even to the *L. cardinalis*, large in its flowers, tall in growth, and so easily cultivated that their abundance and beauty will ensure them admission into every collection. Our figure was made from a plant in the collection of J. Vere, Esq. in the summer of 1810. It was raised from Botany Bay seed at the Hammersmith Nursery in 1809.



Pl. 634

Lobelia fulgens.

1

2

PLATE DCLX.

CAMELLIA JAPONICA. *Var. flore pleno roseo.*

Rose-coloured Camellia.

CLASS XVI. ORDER VI.

MONADELPHIA POLYANDRIA. Threads united. Many Chives.

ESSENTIAL GENERIC CHARACTER.

CALYX imbricatus, polyphyllus: foliolis interioribus majoribus.

EMPALEMENT tiled, many-leaved: the inner leaves the largest.

SPECIFIC CHARACTER.

CAMELLIA floribus roseis: petalis exterioribus magnis, patentibus: interioribus parvis, erectis: foliis latis, nitidis.

CAMELLIA with rose-coloured flowers: the outer petals are large and spreading: the inner ones small and upright: the leaves broad and shining.

CAMELLIA JAPONICA. *Var. flore pleno incarnato.*

Flesh-coloured Camellia.

SPECIFIC CHARACTER.

CAMELLIA foliis conspicuè nervosis: floribus petalis numerosis, pallide carneis.

CAMELLIA with leaves strongly nerved: flowers with numerous petals, of a pale flesh colour.

So high in estimation is every variation of the Camellia Japonica, that even a new and rare plant is not considered more interesting, nor of so much value as one of these beautiful varieties; the number of which is now extended to sixteen, at least so enumerated by cultivators, and all considered as perfectly distinct. The rose-coloured variety was introduced to the royal gardens at Kew about 1808, and the pale variety was first cultivated in the conservatory of the late Lady Amelia Hume, we believe, in 1806. It is nearly

allied to the white, and in severe winters so little colour is visible, that Mrs. Beaumont (with whom this flowered last December at Bretton Hall) informed me it was only by comparing it with the white that it could positively be said to possess any colour; but, upon comparison, a blush was evident. So delicate a test is only requisite in rigid seasons from the want of sun. Both these varieties flower from December till March.



PLATE DCLXI.

MIMULUS LUTEUS.

Yellow-flowered Mimulus.

CLASS XIV. ORDER II.

DIDYNAMIA ANGIOSPERMIA. Two Chives longer. Seeds covered.

ESSENTIAL GENERIC CHARACTER.

CALYX prismaticus, 5-dentatus. Corolla ringens, labio superiore lateribus replicato. Stigma crassum. Capsula bilocularis, polysperma.

EMPALEMENT prismatical, 5-toothed. Blossom ringent, the upper lip folded back at the sides. Shaft fleshy. Seed-vessel two-celled, many-seeded.

SPECIFIC CHARACTER.

MIMULUS foliis subcordatis, multi-nervosis, dentatis.

Habitat in Peru.

MONKEY-FLOWER with nearly heart-shaped leaves, many-nerved and toothed.

Native of Peru.

REFERENCE TO THE PLATE.

1. A radical leaf.
2. The empalement.
3. A blossom spread open.
4. Seed-bud and pointal.

THE *Mimulus luteus* flowers the same year it is sown, like an annual; but from its habit and manner of throwing out round the base of the flowering stem prostrate branches, which take root, and from the extremities of some of them other flowering stems shooting up, we are inclined to think it biennial. It was long ago described and figured by Father Feuillée in his *Flora Peruviana*, and quoted from him by Linnæus, who inserted it in his *Species Plantarum* under the specific title of *luteus*. It flowers in the

autumnal months, and enlivens the borders with its successively numerous bright yellow blossoms; is easily propagated by seeds; and, according to Donn's Catalogue, was introduced in the year 1812.



Mimulus, luteus.

PLATE DCLXII.

CAMELLIA JAPONICA. *Var. petalis plicatis.*

Plicate-petalled or Warratta Camellia.

CLASS XVI. ORDER VI.

MONADELPHIA MONOGYNIA. Threads united. Many Chives.

ESSENTIAL GENERIC CHARACTER.

CALYX imbricatus, polyphyllus: interioribus majoribus.

EMPALEMENT tiled, many-leaved; the inner ones the largest.

SPECIFIC CHARACTER.

CAMELLIA foliis lato-lanceolatis apice reflexis, nitidissimis: corollis rubris, fulgentibus: petalis quinque exterioribus magnis, patentibus: interioribus numerosis, parvis, plicatis, erectis.

CAMELLIA with broadly lance-shaped leaves reflexed at the end, very shining: blossoms red and brilliant: the five outer petals are large and spreading; the inner ones are numerous, small, folded together, and upright.

The flowers of this Camellia very much resemble those of the genus Anemone, and it is certainly the most distinct variety at present known. By cultivators it is denominated the Warratta Camellia, from the brilliance of its flowers resembling the *Embothrium speciosum*, or Warratta. Our drawing was made from a plant in the conservatory of Mr. Alnutt of Clapham Common, where it flowered in February 1813 in great perfection. It also blossomed again last winter, but, from the severity of the season, inferior in colour, and much reduced in size.



Camellia Japonica,
V. petalis plicatis.

PLATE DCLXIII.

LILIUM CONCOLOR.

Self-coloured Lily.

CLASS VI. ORDER I.

HEXANDRIA MONOGYNIA. Six Chives. One Pointal.

ESSENTIAL GENERIC CHARACTER.

COROLLA 6-partita, campanulata. Nectarium linea longitudinalis. Capsulæ valvæ pilis cancellatis connexæ.

BLOSSOM 6-parted, bell-shaped. Nectary a longitudinal line. Valves of the seed-pod connected with a lattice-work of hair.

SPECIFIC CHARACTER.

LILIUM foliis sparsis, subverticillatis, lanceolatis: floribus erectis, fasciculis terminalibus 3-6-floris: petalis recurvis, patentibus: stylo brevissimo.

LILY with scattered leaves, nearly in whorls, and lance-shaped: flowers upright, in terminal bunches of 3-6 flowers: petals recurved and spreading. Shaft very short.

REFERENCE TO THE PLATE.

1. Chives and pointal.

THE *Lilium concolor* was first raised in the conservatory of the late Sir Charles Greville at Paddington in 1806, and still continues to be one of the rarest of this showy genus. Our figure was made from a plant in the nursery of Messrs. Lee and Kennedy, late in the autumn of 1813. Of what country it is a native is not positively known, but it is supposed to be of Chinese origin.



Lilium, concolor

PLATE DCLXIV.

ARBUTUS LONGIFOLIA.

Long-leaved Arbutus.

CLASS X. ORDER I.

DECANDRIA MONOGYNIA. Ten Chives. One Pointal.

ESSENTIAL GENERIC CHARACTER.

CALYX 5-partitus. Corolla ovata, basi pellucida. Bacca 5-locularis.

EMPALEMENT 5-parted. Blossom egg-shaped, transparent at the base. Berry 5-celled.

SPECIFIC CHARACTER.

ARBUTUS foliis longis, lanceolatis, serratis: floribus paniculis verticillatis: pedunculis hispidis, glutinosis: caule arboreo.

ARBUTUS with long lance-shaped leaves, sawed: flowers grow in panicles, whorled: footstalks hispid and glutinous: stem woody.

REFERENCE TO THE PLATE.

1. A blossom.
2. The same spread open.
3. A chive magnified.
4. A side view of the same magnified.
5. Seed-bud and pointal, summit magnified.

OUR drawing of the *Arbutus longifolia* was made from a fine specimen, obligingly communicated by Mrs. Beaumont, from the superb collection of Colonel Beaumont at Bretton Hall, in March 1814, whose gardener, Mr. Benning, informs us that it is kept in the conservatory, is eleven feet high, bears eight heads of flowers like the one figured, and sheds the old bark annually. It was first raised at the Hammersmith nursery from seeds received from the Island of Madeira. It is also said to be found in the Canary Isles, where the berries are made into a sweetmeat. Although sixteen years in

cultivation with us, it has very rarely flowered. May be increased by cuttings or layers.



INDEX

TO THE PLANTS CONTAINED IN VOL. X.

Plate 609	Citrus medica, <i>odoratissima</i>	Bergamot Lemon	H. H.	Shrub.	May.
610	Ruellia formosa	Beautiful Ruellia	H. H.	Shrub.	June.
611	Daviesia corymbosa	Corymbed Daviesia	G. H.	Shrub.	August.
612	Pæonia albiflora, <i>flore pleno</i>	Tartarian Pæony, <i>double-flowered</i> <i>variety</i>	G. H.	Shrub.	August.
613	Ipomæa pendula	Pendulous Ipomæa	G. H.	Shrub.	July.
614	Fumaria nobilis	Noble Fumitory	Har.	Herb.	May.
615	Globba purpurea	Purple Globba	H. H.	Herb.	May.
616	Euphorbia epithymoides	Broad-leaved Spurge	H. H.	Shrub.	May.
617	Euphorbia meloformis	Melon-shaped Euphorbia	H. H.	Bulb.	July.
618	Anneslea spinosa	Armed Indian Water Lily	H. H.	Aquatic.	September.
619	Eugenia Zeylanica	Ceylon Eugenia	H. H.	Shrub.	August.
620	Schinus dentata	Toothed Schinus	Har.	Shrub.	September.
621	Jussieua exaltata	Tall Jussieua	H. H.	Shrub.	August.
622	Leptospermum scoparium	New Zealand Tea	G. H.	Tree.	June.
623	Ardisia elegans	Elegant Ardisia	G. H.	Shrub.	August.

624	<i>Lotus australis</i>	Southern Lotus, or Bird's Foot Trefoil	G. H.	Shrub.	August.
625	<i>Barleria cristata</i>	Crested Barleria	H. H.	Shrub.	September.
626	<i>Geodorum citrinum</i>	Lemon-coloured Geodorum	H. H.	Shrub.	October.
627	<i>Begonia Evansiana</i>	Evans's Begonia	H. H.	Shrub.	October.
628	<i>Clerodendrum pyramidale</i>	Pyramidal Clerodendrum	H. H.	Shrub.	October.
629	<i>Desmanthus natans</i>	Floating Desmanthus, or Aquatic Sensitive	H. H.	Aquatic.	September.
630	<i>Ardisia littoralis</i>	Seaside Ardisia	G. H.	Shrub.	November.
631	<i>Styrax officinale</i>	Storax Tree	Har.	Tree.	June.
632	<i>Cytisus elongatus</i>	Elongated Cytisus	G. H.	Shrub.	May.
633	<i>Liatris odoratissima</i>	Sweet-scented Liatris	H. H.	Shrub.	July.
634	<i>Peliosanthes humilis</i>	Humble Peliosanthes	G. H.	Shrub.	August.
635	<i>Celosia cernua</i>	Nodding Celosia	G. H.	Annual.	August.
636	<i>Ipomæa insignis</i>	Magnificent Ipomæa	G. H.	Shrub.	August.
637	<i>Trichilia odorata</i>	Sweet-scented Trichilia	H. H.	Shrub.	July.
638	<i>Daviesia latifolia</i>	Broad-leaved Daviesia	G. H.	Shrub.	April.
639	<i>Carex Fraseri</i>	Fraser's Carex	Har.	Shrub.	September.
640	<i>Heliconia Bihai</i>	Wild Plantain Tree	H. H.	Herb.	April.

641	Prostanthera lasianthos	Downy-flowered Prostanthera	Har.	Shrub.	June.
642	Gompholobium grandiflorum	Large-flowered Air Pod	G. H.	Shrub.	June.
643	Justicia bicolor	Two-coloured flowered Justicia	H. H.	Shrub.	June.
644	Lobelia Speculum	Lobel's Venus's Looking-glass	G. H.	Annual.	August.
645	Epidendrum fragrans	Sweet-scented Epidendrum	H. H.	Herb.	September.
646	Protea radiata	Radiated Protea	G. H.	Shrub.	August.
647	Androsace coronopifolia	Buck's Horn Plaintain-leaved Androsace	Har.	Annual.	July.
648	Crotalaria Saltiana	Salt's Crotalaria	H. H.	Shrub.	July.
649	Alstromeria edulis	Eatable Alstromeria	H. H.	Shrub.	September.
650	Xeranthemum fasciculatum, var. <i>flore rubro</i>	Bundled-leaved Everlasting Flower, <i>red-fl. variety</i>	G. H.	Shrub.	July.
651	Cymbidium Andersonii	Anderson's Cymbidium	H. H.	Herb.	August.
652	Xeranthemum humile	Low-growing Xeranthemum	G. H.	Shrub.	September.
653	Corræa speciosa	Showy Corræa	G. H.	Shrub.	September.
654	Gnaphalium eximium	Showy Gnaphalium	G. H.	Shrub.	August.
655	Cerbera Manghas	Poisonous-fruited Cerbera	H. H.	Shrub.	September.
656	Crassula perfoliata	Perfoliate-leaved Crassula	H. H.	Shrub.	September.

657	Passiflora lunata	Crescent-leaved Passion Flower	H. H.	Shrub.	August.
658	Styledium graminifolium	Grass-leaved Styledium	G. H.	Shrub.	September.
659	Lobelia fulgens	Refulgent-flowered Lobelia	G. H.	Shrub.	August.
660	Camellia Japonica, <i>var. rosea et incarnata</i>	Camellia Japonica, <i>rose and flesh coloured varieties</i>	G. H.	Shrub.	February.
661	Mimulus luteus	Yellow-flowered Mimulus	Har.	Herb.	September.
662	Camellia Japonica, <i>var. petiolis plicatis</i>	Camellia Japonica, <i>plicate-petalled variety</i>	G. H.	Shrub.	February.
663	Lilium concolor	Self-coloured Lily	Har.	Bulb.	September.
664	Aibutus longiflora	Long-leaved Arbutus	G. H.	Shrub.	April.

ERRATA.

Justicia bicolor, for 663, read 643.

Plate 649, description, line 7., for Cape of Good Hope, read Cape Francois; and in the last line, for it is indigenous, read they are indigenous.

*Alphabetical Index to the 1st. 2d. 3d. 4th. and 5th. Volume of the Botanist's
Repository.*

	Pl. vol
Aizoon canariense	201 3
Aletris fragrans	306 5
farmentosa	54 1
Allium gracile	107 2
Amaryllis Brasilienis	358 5
Fothergillia	163 3
radiata	95 2
reticulata	179 3
Anchusa capensis	336 5
Anemone palmata	172 3
Anthericum costatum	298 5
Anthillis erinacea	15 1
Antholyza Æthiopica	210 3
fulgens	192 3
ringens	32 1
spicata	56 1
tubulosa	174 3
Aponogeton distachyon	290 5
Arctotis paradoxa	357 5
Arethusa plicata	321 5
Aristea cyanea	10 1
major	160 3
Arum orixensis	356 5
Asclepias gigantea	271 4
Aspalathus crassifolius	353 5
Aster dentatus	61 1
reflexus	93 2

Atrogene Austriaca	80	3
capensis	9	1
Azalea Pontica	16	1
Banksia ericafolia	156	3
præmorsa	258	4
serrata	82	2
Baurea rubioides	198	3
Bignonia Leucoxylon	43	1
pandorana	86	2
Blandfordia cordata	343	5
Borbonia cordata	31	1
Boronia pinnata	58	1
Brunsfelsia undulata	167	3
Buchnera fœtida	80	2
pedunculata	84	2
Camellia japonica, <i>alb. pl.</i>	25	1
japonica, <i>pl. vg. to.</i>	91	2
japonica, <i>rub. pl.</i>	199	3
Cameraria latifolia	201	4
Casuarina stricta	346	5
Cerbera Ahouai	231	4
undulata	130	2
Chelone campanuloides	40	1
Ruelloides	34	1
Chiococca racemosa	284	4
Chrysanthemum tricolor	109	2
Cineraria aurita	24	1
Clematis viorna	71	1
Clusia flava	223	4
Cobbea scandens	342	5

Codon Royeni	325	5
Cordia sebestena	157	3
Cortusa Mathioli	1	1
Corræa alba	17	1
Crassula odoratissima	26	1
Crateva Capparoides	176	3
Crinum giganteum	169	3
spirale	92	2
Crowea saligna	79	2
Crotolaria verrucosa	308	5
Cyanella capensis	141	2
Cyrtanthus obliquus	265	4
Cytisus tomentosus	237	4
Daphne Pontica	73	2
Daviesia ulicifolia	304	5
Diosma latifolia	33	1
Dodonæa triquetra	230	4
Dracena borealis	206	3
Echites suberecta	187	3
Echium argenteum	154	3
ferocissimum	39	1
glaucophyllum	165	3
grandiflorum	20	1
Embothrium buxifolium	218	4
lineare	272	4
salignum	215	3
sericeum	100	2
Epidendrum cochleatum	13	1
ensifolium	344	5
sinense	216	3

Epigæa repens	102	2
Eranthemum pulchellum	88	2
Falkia repens	257	4
Ferraria pavonia	178	3
viridiflora	285	4
Fuchsia lycioides	120	2
Galaxia grandiflora	164	3
ovata	94	2
Gardenia tubiflora	183	3
Gaultheria procumbens	116	2
Gentiana purpurea	117	2
Geranium astragalifolium	190	3
barbatum	303	5
barbatum minor	323	5
bubonifolium	328	5
ciliatum	247	4
coronillæfolium	305	5
coronopifolium	338	5
cortusæfolium	121	2
crassifolium	136	2
echinatum	158	3
elegans	28	1
fragile	37	1
grandiflorum	12	1
incisum	67	1
incrassatum	246	4
laciniatum	131	2
flo. purp.	204	3
bicolor	269	4
lineare	193	3

	melananthum	209	3
	oxalidifolium	300	5
	pictum	168	3
	pilosum	259	4
	pinnatum	311	5
	præmorsum	150	3
	procumbens	254	4
	punctatum	60	1
	quinquevulnerum	114	2
	radiatum	222	4
	reflexum	224	4
	reniforme	108	2
	revolutum	354	5
	roseum	173	3
	selinum	239	4
	spathulatum	152	3
	curviflo.	282	4
	tomentosum	115	2
	undulatum	292	5
	minor	317	5
Gladiolus	alatus	8	1
	abreviatus	166	3
	blandas	99	2
	campanulatus	188	3
	carneus	240	4
	crispus	142	2
	cuspidatus	147	3
	pet. crisp.	219	4
	galeatus	122	2
	gramineus	62	1
	grandiflorus	118	2
	longiflorus	5	1

nanus	137	2
orchidiflorus	241	4
plicatus	268	4
polystachius	66	4
præcox flo. rub.	38	1
pyramidatus	335	5
ringens cin. odor.	27	1
multiflo.	227	4
undul.	275	4
roseus	11	1
striatus	111	2
versicolor	19	1
Gloriosa superba	129	2
Gnidia lævigata	89	2
opposætifolia	225	4
pinifolia	52	1
simplex	70	1
Goodenia calendulacea	22	1
ovata	68	1
 Hæmanthus multiflorus	 318	 5
Hebenstreitia aurea	252	4
Heliconia psittacorum	124	2
Helonias bullata	352	5
Hemerocallis alba	194	3
cærulea	6	1
graminea	244	4
Hermannia pulverata	161	3
Hibbertia volubilis	126	2
Hibiscus mutabilis fl. pl.	228	4
Patersonius	286	4
Hillia longiflora	145	3

Houstonia coccinea	106	2
Hyacinthus corymbosus	345	5
Hypoxis linearis	171	3
obliqua	195	3
stellata	101	2
fl. alb.	236	4
Iris longifolia	45	1
Itea spinosa	314	5
Ixia aristata	87	2
bulbifera	48	1
bulbocodium	170	3
capitata ovat.	68	1
fl. aur.	50	1
fl. al. fun. nig.	159	3
stellata	232	4
cinnamonea	44	1
columnaris	203	3
versic.	211	3
latifol.	213	3
grandiflo.	250	4
crispifolia fl. cærul.	35	1
crocata nig. mac.	134	2
fistulosa	59	1
maculata	196	3
min. fl. pur.	256	4
polystachia	155	3
incar.	128	2
punctata	177	3
pusilla	245	4
reflexa	14	1
speciosa	186	3

spicata	29	1
Ixora Pavetta virid. nig.	76	2
Jasminum gracile	127	2
Jatropha panduræfolia	267	4
Justicea lucida	313	5
Lachenalia fragrans	302	5
pendula	41	1
purpureo cærul.	257	4
pustulata	350	5
quadricolor	148	3
reflexa	299	5
rosea	296	5
Lachnæa eriocephala	104	2
purpurea	293	5
Lambertia formosa longi flor.	69	1
Laseopetalum ferrugineum	208	3
Leea pinnata	355	5
Limonia trifoliata	143	2
Lobelia coronopifolia	339	5
gracilis	340	5
pinifolia	273	4
Magnolia fuscata	229	4
pumila	226	4
purpurea	324	5
Mahernia odorata	85	2
Malpighia crassifolia	49	1
Malva divaricata	182	3
reflexa	135	2
Massonia scabra	220	4

violacea	46	1
Melaleuca coronata	278	4
ericæfolia	175	3
hypericifolia	200	3
Melanthium viride	233	4
Mesembryanthemum glabrum	57	1
Metrosidera hirsuta	281	4
Mimosa discolor	235	4
longifolia	207	3
stricta	53	1
Monsonia filia	276	4
Moræa Northiana	255	4
tricolor	83	2
Musa coccinea	47	1
Neotia speciosa	3	1
Nymphæa cærulea	197	3
ordorata	297	5
stellata	330	5
Olea apetala	316	5
Ophrys lilifolia	65	1
Orchis bicornis	315	5
ciliaris	42	1
Ornithogalum lacteum	274	4
odoratum	260	4
Pæonia albiflora	64	1
Passiflora aurantia	295	5
maliformis	217	4
Pergularia minor	184	3
odoratissima	185	3

Persoonia lanceolata	74	2
Persoonia latifolia	280	4
linearis	77	2
Pharnaceum lineare	326	5
fl. alb.	329	5
Phaseolus Caracalla	341	5
Physalis prostrata	75	2
Pitcairnia latifolia	322	5
sulphurea	249	4
Pittosporum coriaceum	151	3
Platylobium lanceolatum	205	3
ovatum	266	4
scolopendrum	191	3
Pogonia debilis	212	3
glabra	283	4
Primula Cortusoides	7	1
Protea anemonifolia	332	5
candicans	294	5
cordata	289	5
cynaroides	288	4
decumbens	349	5
formosa	17	1
globosa	307	5
glomerata	264	4
grandiflora fol. und.	301	5
lagopus	243	4
longifolia nigra	132	2
ferr. purp.	133	2
cono turb.	144	2
pinifolia	76	2
pulchella	270	4
speciosa latifolia	110	2

nigra	103	2
fol. glab.	277	4
spicata	234	4
triternata	337	5
umbellata	248	4
Psoralea aculeata	146	3
Pultenæa dapnoides	98	2
ilcifolia	320	5
rubiaefolia	331	5
villosa	309	5
Punica granatum fl. alb.	96	2
Renealmia nutans	360	5
Rhododendron Dauricum	4	1
punctatum	36	1
Roëlla decurrens	238	4
Sagittaria lancifolia	333	5
Samyda serrulata	202	3
Senecio pseudo China	291	5
Septas capensis	90	2
Schotia speciosa	348	5
Sophora capensis	347	5
Sowerbia juncea	81	2
Sprengelia incarnata	2	1
Stachys coccinea	310	5
Struthiola ciliata	149	3
fl. rub.	139	2
imbricata	133	2
ovata	119	2
tomentosa	334	5
Styphelia parviflora	287	4

triflora	72	1
viridis	312	5
Tagetas lucida	359	5
Talinum patens	253	4
Thunbergia fragrans	123	2
Usteria scandens	63	1
Vaccinium amænum	138	2
arctostaphyllus	30	1
crassifolium	105	2
dumosum	112	2
formosum	97	2
frondosum	140	2
parviflorum	125	2
staminium	263	4
virgatum	181	3
Verbascum ferrugineum	162	3
Vereia crenata	21	1
Vicia galegifolia	319	5
Viola pedata	153	3
Westeringia rosmarinacea	214	3
Wurmbea capensis β	221	4
Xeranthemum argenteum	327	5
fasciculatum	242	4
fl. alb.	279	4
speciocissimum	51	1
spirale	262	4
Xylophylla falcata	331	5

Zinnia verticillata	189	3
violacea	55	1

*Alphabetical Index to the 6th. 7th. 8th. 9th. and 10th. Volume of the
Botanist's Repository.*

	Pl. vol
<i>Achania mollis</i>	452 7
<i>Achyranthes porrigens</i>	380 6
<i>Ægiphila diffusa</i>	578 9
<i>Agave Americana</i>	433 7
<i>Albuca fastigiata</i>	450 7
<i>Allium Chamæ-Moly</i>	377 6
<i>Aloe arborescens</i>	468 7
<i>Alstromeria edulis</i>	649 10
<i>Amaryllis spectabilis</i>	390 6
<i>Anagallis grandiflora</i>	367 6
<i>Androsace coronopifolia</i>	647 10
<i>Anneslea spinosa</i>	618 10
<i>Anthericum paniculatum</i>	396 6
<i>pugioniforme</i>	386 6
<i>Aponogeton monostachyon</i>	406 6
<i>Arbutus longifolia</i>	665 10
<i>Ardisia elegans</i>	623 10
<i>littoralis</i>	630 10
<i>Aspalathus globosus</i>	510 8
<i>Astragalus villosus</i>	516 8
<i>Bæckia virgata</i>	598 9
<i>Banksia spinulosa</i>	457 7
<i>Barleria cristata</i>	625 10
<i>Begonia Evansiana</i>	627 10
<i>Bignonia grandiflora</i>	493 8
<i>uncata</i>	530 8

Broussonetia papyrifera	488	7
Cactus coccinellifer	533	8
grandiflorus	508	8
hexagonus	513	8
Cæsulea axillaris	429	6
Calendula viscosa	412	6
Calycanthus fertilis	539	8
Camellia Japonica, <i>semi-duplex</i>	559	9
Japonica, <i>pleno</i> Vars.	660	10
Japonica, <i>Var., pe. pli.-s.</i>	663	10
Campanula laciniata	385	6
versicolor	396	6
Cantua Coronopifolia	415	6
Calendula dentata	407	6
Callicoma serratifolia	566	9
Carex Fraseri	639	10
Ceanothus laniger	569	9
Celosia cernua	635	10
Ceratonia siliqua	567	9
Cerbera manghas	655	10
Chamærops humilis	599	9
Cinchona caribæa	481	7
Citrus medica, odoratissima	609	10
nobilis	608	9
Clematis florida	402	6
Clerodendrum pyramidale	628	10
tomentosum	607	9
Commelina tuberosa	399	6
Commersonia echinata	519	8
dasyphylla	603	9
Corchorus Japonicus, <i>flore pleno</i>	587	9

Corræa speciosa	653	10
viridiflora	436	7
Crassula obliqua	414	6
perfoliata	656	10
Cratægus Azarolus	579	9
Crinum latifolium	478	7
Crocus biflorus	362	6
Crotolaria juncea	422	6
pulchella	417	6
pulchra	601	9
Saltiana	648	10
tetragona	593	9
Cucumis Dudaim	548	8
Cymbidium Andersonii	651	10
Cynanchum bicolor	562	9
mucronatum	515	8
undatum	410	6
Cytisus elongatus	632	10
Dahlia pinnata	408	6
nana	483	7
Daviesia corymbosa	611	10
Daviesia latifolia	638	10
Desmanthus natans	629	10
Dianthus alpinus	482	7
Diosma ericæfolia	451	7
ovata	464	7
Dolichos hirsutus	446	7
Dombeya Erythroxyton	389	6
Elychrisum Stæhelina	428	6
Epidendrum fuscatum	441	7

fragrans	645	10
lineare	445	7
Eriospermum folioliferum	521	8
Erythina speciosa	443	7
Eucalyptus resinifera	400	6
Eucomis purpureocaulis	369	6
Eugenia malaccensis	458	7
Zeylanica	619	10
Euosma albiflora	520	8
Euphorbio epithymoides	616	10
meloformis	617	10
 Fragaria indica	 479	 7
Ferula Persica	558	9
Fumaria formosa	393	6
nobilis	614	10
 Gærtnera racemosa	 600	 9
Gardenia radicans	491	7
Genista lusitanica	419	6
Gentiana Catesbæi	418	6
fimbriata	509	8
Geodorum citrinum	626	10
Geranium barbatum. <i>Var. undulatum</i>	366	6
fissifolium	378	6
floribundum	420	6
hymenodes	413	6
luteum	423	6
Gladiolus angustus, <i>minor</i>	589	9
Globba purpurea	615	10
Glycine Comptoniane	602	9
Gnaphalium eximium	654	10

grandiflorum	489	7
Gompholobium grandiflorum	642	10
maculatum	427	6
Goodenia tenella	466	7
Gorteria pavonia	528	8
Heliconia bihai	640	10
Hellenia Allugas	501	8
Hermannia flammea	550	8
Hibbertia crenata	472	7
Hibiscus pruriens	498	8
Hypericum Virginicum	552	8
Ipomæa coccinea	499	8
grandiflora	403	6
insignis	636	10
pendula	613	10
Iris pavonia	364	6
Ixia columnaris. <i>Var. angustifolia</i>	392	6
curta	564	9
Jasminum multiflorum	496	8
sambac, <i>Var. flore pleno</i>	497	8
Juniperus Daurica	534	8
Jussieua exaltata	621	10
Justicia bicolor	643	10
nitida	570	9
Lachenalia flava	456	7
sessiliflora	460	7
Lachnæa Buxifolia	524	8
Lasiopetalum ferrugineum	459	7

Laurus Cinnamomum	596	9
Leptospermum scoparium	622	10
Liatris odoratissima	633	10
Lilium concolor	662	10
speciosum	586	9
Limodorum Tankervilliae	426	6
Linum trigynum	449	7
venustum	477	7
Liparia sphærica	568	9
villosa	382	6
Lithospermum Tinctorium	576	9
Lobelia assurgens	553	9
fulgens	659	10
hirsuta	444	7
speculum	644	10
surinamensis, <i>Var. flore rubro</i>	502	8
Lonicera Japonica	583	9
Lopezia coronata	551	8
Lotus australis	624	10
Lythrum fruticosum	467	7
Magnolia auriculata	573	9
grandiflora	518	8
Malpighia polystachia	604	9
Malus Japonica	462	7
Magnifera indica	425	6
Martynia diandra	575	9
Melaleuca diosmæfolia	476	7
salicifolia	485	7
Melanthium Massoniæfolium	368	6
Mesembryanthemum acinaciforme	580	9
includens	388	6

heterophyllum	540	8
Mespilus odoratissima	590	9
tanacetifolia	591	9
Mimosa elegans	563	9
grandiflora	592	9
Linifolia	394	6
pudica	544	8
purpurea	372	6
Mimulus luteus	661	10
Monarda punctata	546	8
Morea miniata	404	6
Neottia minor	376	6
Nicotiana glutinosa	484	7
Nitraria Schoberi	529	8
Nymphæa Lotus	391	6
rubra	503	8
Ophrys arachnoides	470	7
myodes	471	7
Ornithogalum elatum	528	8
flavissimum	505	8
Origanum Tournefortii	537	8
Oxylobium cordifolium	492	7
Pæonia albiflora, <i>flore pleno</i>	612	10
anomala	514	8
Daurica	486	7
papavaracea	463	7
suffruticosa	373	6
Var. <i>flore purpureo</i>	448	7
Panax fruticosum	595	9

Pancratium amœnum	556	9
Pascalìa glauca	549	8
Passiflora lunata	657	10
perfoliata	547	8
Peliosanthes humilis	634	10
Teta	605	9
Periploca Africana	557	9
Phlomis Samia	584	9
Pittosporum undulatum	583	6
Plectranthus barbatus	594	9
Podalyria hirsuta	525	8
Polygala alopecuroides	371	6
micrantha	424	6
mixta	455	7
stipulacea	363	6
teretifolia	370	6
Pontederia dilatata	490	7
Primula dentiflora	405	6
Prostranthera lasianthos	641	10
Protea abrotanifolia	507	8
hirta	522	8
<i>minor</i>	536	8
odorata	545	8
Protea argentifolia	447	7
canaliculata	437	7
cespitosa	526	8
conifera	541	8
coronata	469	7
corymbosa	495	8
divaricata	465	7
humiflora	532	8
imbricata	517	8

incurva	431	6
mellifera albiflora	582	9
mucronifolia	500	8
pinnata	512	8
pulchella. <i>Var. speciosa</i>	442	7
radiata	646	10
repens	453	7
Scolymus	409	6
saligna	572	9
speciosa	438	7
speciosa patens	543	8
teretifolia	461	7
virgata	577	9
Psoralea pinnata	474	7
spicata	411	6
Pultenæa nana	434	7
obcordata	574	9
 Renealmia calcarata	 421	 6
Ricinus armatus	430	6
Rhododendron ponticum. <i>Var. deciduum</i>	379	6
Ruellia cristata	506	8
formosa	610	10
fulgida	527	8
infundibuliformis	542	8
Ruta linifolia	565	9
 Sansevieria carnea	 361	 6
Salix violacea	581	9
Salvia Chamædrifolia	416	6
Sarracenia flava	381	6
Schinus dentata	620	10

Scilla siberica	365	6
Scutellaria serrata	494	8
Serapias cordigera	475	7
Serratula spicata	401	6
Sida patens	571	9
hastata	588	9
Solanum betaceum	511	8
Seaforthianum	504	8
Sophora Japonica	585	9
sericea	440	7
Stapelia orbicularis	439	7
Stewartia marilandica	397	6
Strelitzia Reginae	432	6
Styledium graminifolium	658	10
Styrax officinale	631	10
Trichilia odorata	637	10
Tropæolum peregrinum	597	9
pinnatum	535	8
Urtica baccifera	454	7
Vaccinium nitidum	480	7
Vanilla planifolia	538	8
Verbena mutabilis	535	7
Veronica Derwentia	531	8
Verea acutiflora	560	9
Volkameria angustifolia	554	9
Wachendorfia villosa	398	6
Xeranthemum bracteatum	375	6

fasciculatum, <i>Var. flore rubro</i>	650	10
fragrans	561	10
herbaceam	487	7
humile	652	10
proliferum	374	6
variegatum	384	6
rigidum	387	6
 Yucca gloriosa	 473	 7
 Zieria Smithii	 606	 9
Zingiber Cliffordiæ	555	9

*** END OF THE PROJECT GUTENBERG EBOOK THE BOTANIST'S
REPOSITORY FOR NEW AND RARE PLANTS; VOL. 10 [OF 10] ***

Updated editions will replace the previous one—the old editions will be renamed.

Creating the works from print editions not protected by U.S. copyright law means that no one owns a United States copyright in these works, so the Foundation (and you!) can copy and distribute it in the United States without permission and without paying copyright royalties. Special rules, set forth in the General Terms of Use part of this license, apply to copying and distributing Project Gutenberg™ electronic works to protect the PROJECT GUTENBERG™ concept and trademark. Project Gutenberg is a registered trademark, and may not be used if you charge for an eBook, except by following the terms of the trademark license, including paying royalties for use of the Project Gutenberg trademark. If you do not charge anything for copies of this eBook, complying with the trademark license is very easy. You may use this eBook for nearly any purpose such as creation of derivative works, reports, performances and research. Project Gutenberg eBooks may be modified and printed and given away—you may do practically ANYTHING in the United States with eBooks not protected by U.S. copyright law. Redistribution is subject to the trademark license, especially commercial redistribution.

START: FULL LICENSE

THE FULL PROJECT GUTENBERG™ LICENSE

PLEASE READ THIS BEFORE YOU DISTRIBUTE OR USE THIS WORK

To protect the Project Gutenberg™ mission of promoting the free distribution of electronic works, by using or distributing this work (or any other work associated in any way with the phrase “Project Gutenberg”), you agree to comply with all the terms of the Full Project Gutenberg License available with this file or online at www.gutenberg.org/license.

Section 1. General Terms of Use and Redistributing Project Gutenberg electronic works

1.A. By reading or using any part of this Project Gutenberg electronic work, you indicate that you have read, understand, agree to and accept all the terms of this license and intellectual property (trademark/copyright) agreement. If you do not agree to abide by all the terms of this agreement, you must cease using and return or destroy all copies of Project Gutenberg electronic works in your possession. If you paid a fee for obtaining a copy of or access to a Project Gutenberg electronic work and you do not agree to be bound by the terms of this agreement, you may obtain a refund from the person or entity to whom you paid the fee as set forth in paragraph 1.E.8.

1.B. “Project Gutenberg” is a registered trademark. It may only be used on or associated in any way with an electronic work by people who agree to be bound by the terms of this agreement. There are a few things that you can do with most Project Gutenberg electronic works even without complying with the full terms of this agreement. See paragraph 1.C below. There are a lot of things you can do with Project Gutenberg electronic works if you follow the terms of this agreement and help preserve free future access to Project Gutenberg electronic works. See paragraph 1.E below.

1.C. The Project Gutenberg Literary Archive Foundation (“the Foundation” or PGLAF), owns a compilation copyright in the collection of Project Gutenberg electronic works. Nearly all the individual works in the collection are in the public domain in the United States. If an individual work is unprotected by copyright law in the United States and you are

located in the United States, we do not claim a right to prevent you from copying, distributing, performing, displaying or creating derivative works based on the work as long as all references to Project Gutenberg are removed. Of course, we hope that you will support the Project Gutenberg mission of promoting free access to electronic works by freely sharing Project Gutenberg works in compliance with the terms of this agreement for keeping the Project Gutenberg name associated with the work. You can easily comply with the terms of this agreement by keeping this work in the same format with its attached full Project Gutenberg License when you share it without charge with others.

1.D. The copyright laws of the place where you are located also govern what you can do with this work. Copyright laws in most countries are in a constant state of change. If you are outside the United States, check the laws of your country in addition to the terms of this agreement before downloading, copying, displaying, performing, distributing or creating derivative works based on this work or any other Project Gutenberg work. The Foundation makes no representations concerning the copyright status of any work in any country other than the United States.

1.E. Unless you have removed all references to Project Gutenberg:

1.E.1. The following sentence, with active links to, or other immediate access to, the full Project Gutenberg License must appear prominently whenever any copy of a Project Gutenberg work (any work on which the phrase “Project Gutenberg” appears, or with which the phrase “Project Gutenberg” is associated) is accessed, displayed, performed, viewed, copied or distributed:

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg™ License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

1.E.2. If an individual Project Gutenberg electronic work is derived from texts not protected by U.S. copyright law (does not contain a notice indicating that it is posted with permission of the copyright holder), the work can be copied and distributed to anyone in the United States without paying any fees or charges. If you are redistributing or providing access to a work with the phrase “Project Gutenberg” associated with or appearing on the work, you must comply either with the requirements of paragraphs 1.E.1 through 1.E.7 or obtain permission for the use of the work and the Project Gutenberg trademark as set forth in paragraphs 1.E.8 or 1.E.9.

1.E.3. If an individual Project Gutenberg electronic work is posted with the permission of the copyright holder, your use and distribution must comply with both paragraphs 1.E.1 through 1.E.7 and any additional terms imposed by the copyright holder. Additional terms will be linked to the Project Gutenberg License for all works posted with the permission of the copyright holder found at the beginning of this work.

1.E.4. Do not unlink or detach or remove the full Project Gutenberg License terms from this work, or any files containing a part of this work or any other work associated with Project Gutenberg.

1.E.5. Do not copy, display, perform, distribute or redistribute this electronic work, or any part of this electronic work, without prominently displaying the sentence set forth in paragraph 1.E.1 with active links or immediate access to the full terms of the Project Gutenberg License.

1.E.6. You may convert to and distribute this work in any binary, compressed, marked up, nonproprietary or proprietary form, including any word processing or hypertext form. However, if you provide access to or distribute copies of a Project Gutenberg work in a format other than “Plain Vanilla ASCII” or other format used in the official version posted on the official Project Gutenberg website (www.gutenberg.org), you must, at no additional cost, fee or expense to the user, provide a copy, a means of exporting a copy, or a means of obtaining a copy upon request, of the work in its original “Plain Vanilla ASCII” or other form. Any alternate format must include the full Project Gutenberg License as specified in paragraph 1.E.1.

1.E.7. Do not charge a fee for access to, viewing, displaying, performing, copying or distributing any Project Gutenberg works unless you comply with paragraph 1.E.8 or 1.E.9.

1.E.8. You may charge a reasonable fee for copies of or providing access to or distributing Project Gutenberg electronic works provided that:

- You pay a royalty fee of 20% of the gross profits you derive from the use of Project Gutenberg works calculated using the method you already use to calculate your applicable taxes. The fee is owed to the owner of the Project Gutenberg trademark, but he has agreed to donate royalties under this paragraph to the Project Gutenberg Literary Archive Foundation. Royalty payments must be paid within 60 days following each date on which you prepare (or are legally required to prepare) your periodic tax returns. Royalty payments should be clearly marked as such and sent to the Project Gutenberg Literary Archive Foundation at the address specified in Section 4, "Information about donations to the Project Gutenberg Literary Archive Foundation."
- You provide a full refund of any money paid by a user who notifies you in writing (or by e-mail) within 30 days of receipt that s/he does not agree to the terms of the full Project Gutenberg™ License. You must require such a user to return or destroy all copies of the works possessed in a physical medium and discontinue all use of and all access to other copies of Project Gutenberg™ works.
- You provide, in accordance with paragraph 1.F.3, a full refund of any money paid for a work or a replacement copy, if a defect in the electronic work is discovered and reported to you within 90 days of receipt of the work.
- You comply with all other terms of this agreement for free distribution of Project Gutenberg™ works.

1.E.9. If you wish to charge a fee or distribute a Project Gutenberg™ electronic work or group of works on different terms than are set forth in this agreement, you must obtain permission in writing from the Project Gutenberg Literary Archive Foundation, the manager of the Project

Gutenberg™ trademark. Contact the Foundation as set forth in Section 3 below.

1.F.

1.F.1. Project Gutenberg volunteers and employees expend considerable effort to identify, do copyright research on, transcribe and proofread works not protected by U.S. copyright law in creating the Project Gutenberg™ collection. Despite these efforts, Project Gutenberg™ electronic works, and the medium on which they may be stored, may contain “Defects,” such as, but not limited to, incomplete, inaccurate or corrupt data, transcription errors, a copyright or other intellectual property infringement, a defective or damaged disk or other medium, a computer virus, or computer codes that damage or cannot be read by your equipment.

1.F.2. LIMITED WARRANTY, DISCLAIMER OF DAMAGES - Except for the “Right of Replacement or Refund” described in paragraph 1.F.3, the Project Gutenberg Literary Archive Foundation, the owner of the Project Gutenberg™ trademark, and any other party distributing a Project Gutenberg™ electronic work under this agreement, disclaim all liability to you for damages, costs and expenses, including legal fees. YOU AGREE THAT YOU HAVE NO REMEDIES FOR NEGLIGENCE, STRICT LIABILITY, BREACH OF WARRANTY OR BREACH OF CONTRACT EXCEPT THOSE PROVIDED IN PARAGRAPH 1.F.3. YOU AGREE THAT THE FOUNDATION, THE TRADEMARK OWNER, AND ANY DISTRIBUTOR UNDER THIS AGREEMENT WILL NOT BE LIABLE TO YOU FOR ACTUAL, DIRECT, INDIRECT, CONSEQUENTIAL, PUNITIVE OR INCIDENTAL DAMAGES EVEN IF YOU GIVE NOTICE OF THE POSSIBILITY OF SUCH DAMAGE.

1.F.3. LIMITED RIGHT OF REPLACEMENT OR REFUND - If you discover a defect in this electronic work within 90 days of receiving it, you can receive a refund of the money (if any) you paid for it by sending a written explanation to the person you received the work from. If you received the work on a physical medium, you must return the medium with your written explanation. The person or entity that provided you with the defective work may elect to provide a replacement copy in lieu of a refund. If you received the work electronically, the person or entity providing it to

you may choose to give you a second opportunity to receive the work electronically in lieu of a refund. If the second copy is also defective, you may demand a refund in writing without further opportunities to fix the problem.

1.F.4. Except for the limited right of replacement or refund set forth in paragraph 1.F.3, this work is provided to you 'AS-IS', WITH NO OTHER WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR ANY PURPOSE.

1.F.5. Some states do not allow disclaimers of certain implied warranties or the exclusion or limitation of certain types of damages. If any disclaimer or limitation set forth in this agreement violates the law of the state applicable to this agreement, the agreement shall be interpreted to make the maximum disclaimer or limitation permitted by the applicable state law. The invalidity or unenforceability of any provision of this agreement shall not void the remaining provisions.

1.F.6. INDEMNITY - You agree to indemnify and hold the Foundation, the trademark owner, any agent or employee of the Foundation, anyone providing copies of Project Gutenberg™ electronic works in accordance with this agreement, and any volunteers associated with the production, promotion and distribution of Project Gutenberg™ electronic works, harmless from all liability, costs and expenses, including legal fees, that arise directly or indirectly from any of the following which you do or cause to occur: (a) distribution of this or any Project Gutenberg work, (b) alteration, modification, or additions or deletions to any Project Gutenberg work, and (c) any Defect you cause.

Section 2. Information about the Mission of Project Gutenberg

Project Gutenberg is synonymous with the free distribution of electronic works in formats readable by the widest variety of computers including obsolete, old, middle-aged and new computers. It exists because of the

efforts of hundreds of volunteers and donations from people in all walks of life.

Volunteers and financial support to provide volunteers with the assistance they need are critical to reaching Project Gutenberg's goals and ensuring that the Project Gutenberg collection will remain freely available for generations to come. In 2001, the Project Gutenberg Literary Archive Foundation was created to provide a secure and permanent future for Project Gutenberg and future generations. To learn more about the Project Gutenberg Literary Archive Foundation and how your efforts and donations can help, see Sections 3 and 4 and the Foundation information page at www.gutenberg.org.

Section 3. Information about the Project Gutenberg Literary Archive Foundation

The Project Gutenberg Literary Archive Foundation is a non-profit 501(c)(3) educational corporation organized under the laws of the state of Mississippi and granted tax exempt status by the Internal Revenue Service. The Foundation's EIN or federal tax identification number is 64-6221541. Contributions to the Project Gutenberg Literary Archive Foundation are tax deductible to the full extent permitted by U.S. federal laws and your state's laws.

The Foundation's business office is located at 41 Watchung Plaza #516, Montclair NJ 07042, USA, +1 (862) 621-9288. Email contact links and up to date contact information can be found at the Foundation's website and official page at www.gutenberg.org/contact

Section 4. Information about Donations to the Project Gutenberg Literary Archive Foundation

Project Gutenberg™ depends upon and cannot survive without widespread public support and donations to carry out its mission of increasing the number of public domain and licensed works that can be freely distributed in machine-readable form accessible by the widest array of equipment

including outdated equipment. Many small donations (\$1 to \$5,000) are particularly important to maintaining tax exempt status with the IRS.

The Foundation is committed to complying with the laws regulating charities and charitable donations in all 50 states of the United States. Compliance requirements are not uniform and it takes a considerable effort, much paperwork and many fees to meet and keep up with these requirements. We do not solicit donations in locations where we have not received written confirmation of compliance. To SEND DONATIONS or determine the status of compliance for any particular state visit www.gutenberg.org/donate.

While we cannot and do not solicit contributions from states where we have not met the solicitation requirements, we know of no prohibition against accepting unsolicited donations from donors in such states who approach us with offers to donate.

International donations are gratefully accepted, but we cannot make any statements concerning tax treatment of donations received from outside the United States. U.S. laws alone swamp our small staff.

Please check the Project Gutenberg web pages for current donation methods and addresses. Donations are accepted in a number of other ways including checks, online payments and credit card donations. To donate, please visit: www.gutenberg.org/donate.

Section 5. General Information About Project Gutenberg electronic works


Professor Michael S. Hart was the originator of the Project Gutenberg concept of a library of electronic works that could be freely shared with anyone. For forty years, he produced and distributed Project Gutenberg eBooks with only a loose network of volunteer support.

Project Gutenberg eBooks are often created from several printed editions, all of which are confirmed as not protected by copyright in the U.S. unless a

copyright notice is included. Thus, we do not necessarily keep eBooks in compliance with any particular paper edition.

Most people start at our website which has the main PG search facility:
www.gutenberg.org.

This website includes information about Project Gutenberg, including how to make donations to the Project Gutenberg Literary Archive Foundation, how to help produce our new eBooks, and how to subscribe to our email newsletter to hear about new eBooks.



[back](#)

P. 109



[back](#)



[back](#)



Daviesia corymbosa

[back](#)



[back](#)

Fig. 503



Ipomoea pendula

[back](#)



[back](#)



Globba, purpurea

See the description of the species in the text.

[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



[back](#)



Styxpermum, scoparium

[back](#)

Pl. 6252



Antasia elegans

[back](#)




Lotus, australis

[back](#)





[back](#)



[back](#)

Pl. 020



Clerodendron pyramidale

[back](#)

Pl. 699

Desmanthus, natans

[back](#)



[back](#)



[back](#)




[back](#)

Pl. 63



Lantana odoratissima

[back](#)



Peliosanthes, humilis



[back](#)



Ipomoea insignis

[back](#)

1469



Trichilia catappa

[back](#)



[back](#)



Carex, Fraseri





[back](#)

N. 64



Prostanthera laevis

[back](#)



Mimulus, grandiflorum

Pl 643



Justicia, bicolor

[back](#)

N^o 44



Lobelia speculum

[back](#)

Pl. 64



Epidendrum fragrans

[back](#)



Protea nerioides

[back](#)



Androsace coronopifolia

[back](#)



[back](#)



[back](#)



Ceranthemum, fasciculatum. Vår flore rubro

[back](#)



[back](#)

Pl. 672



Ceranthemum, humilis

[back](#)



Correa, speciosa


[back](#)

N. 655



Gnaphalium Eximium

[back](#)



Nerium Manghas

[back](#)



[back](#)



Pl. 037

Passiflora lunata

From the Bot. Garden, Vienna
by H. Androsch & H. H. H. H.

[back](#)



Stylidium, graminifolium

[back](#)



[back](#)



[back](#)



Mimulus, luteus.

[back](#)



Camellia Japonica,
V. petal. plicatis.



Lilium, concolor

[back](#)



[back](#)