

Curiosa Mathematica

PART II

PILLOW-PROBLEMS

*THOUGHT OUT DURING
WAKEFUL HOURS*

BY

CHARLES L. DODGSON, M.A.

*Student and late Mathematical Lecturer
of Christ Church, Oxford*

FOURTH EDITION

PRICE TWO SHILLINGS

London

MACMILLAN AND CO.

1895

[All rights reserved]

Curiosa Mathematica

PART II

PILLOW-PROBLEMS

*THOUGHT OUT DURING
WAKEFUL HOURS*

BY

CHARLES L. DODGSON, M.A.

*Student and late Mathematical Lecturer
of Christ Church, Oxford*

FOURTH EDITION

PRICE TWO SHILLINGS

London

MACMILLAN AND CO.

1895

[All rights reserved]

The Project Gutenberg eBook of Curiosa mathematica, part II

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

Title: Curiosa mathematica, part II

Pillow-problems thought out during wakeful hours

Author: Lewis Carroll

Release date: July 12, 2026 [eBook #79080]

Most recently updated: July 28, 2026

Language: English


Original publication: London: Macmillan and Co., 1895

Other information and formats: www.gutenberg.org/ebooks/79080

Credits: Aaron Adrignola, Laura Natal and the Online Distributed Proofreading Team at <https://www.pgdp.net> (This book was produced from images made available by the HathiTrust Digital Library.)

*** START OF THE PROJECT GUTENBERG EBOOK CURIOSA
MATHEMATICA, PART II ***

PILLOW-PROBLEMS



[See [p. 100](#).

Curiosa Mathematica

PART II

PILLOW-PROBLEMS

*THOUGHT OUT DURING
WAKEFUL HOURS*

BY
CHARLES L. DODGSON, M.A.

Student and late Mathematical Lecturer of Christ Church Oxford

FOURTH EDITION

PRICE TWO SHILLINGS

London
MACMILLAN AND CO.
1895

[All rights reserved]

Oxford
HORACE HART, PRINTER TO THE UNIVERSITY


PREFACE TO FOURTH EDITION.

I TAKE this opportunity of explaining *why* it is that (as stated in the Note to [p. xix](#)) I have used the symbols $\cap$ and $\frown$ to represent the words 'sine' and 'cosine'.

The use of *some* symbols needs, I suppose, no more justification than the use of + and - to represent 'plus' and 'minus'.

These particular symbols are derived from the old theory of Trigonometry, in which sines, cosines, &c. were actual *lines*.

In this diagram, OP being taken as the unit of length, PN is the *sine* of the angle NOP , and ON its *cosine*.



In each of my two symbols I have retained the semicircle in the symbol sin, I have merely moved PN to the middle; and, in the symbol cos, I have lengthened ON , taking it a little *beyond* the curve, in order to avoid confusion with the existing symbol for 'semicircle'.

I also take this opportunity of adding a sort of Corollary (lately thought out) to the solution of Problem 59 (see [p. 84](#)).

If a, b, c be given lengths, they must, in order that the Tetrahedron may be *possible*, fulfil certain conditions, as follows:—

(1) they have to form the sides of a Triangle: hence any two of them must be greater than the third;

(2) the three angles of this Triangle have to form a *solid* angle: hence any two of these angles must be together greater than the third: hence any two of them must be together greater than 90° : hence any *one* of them must be *less* than 90° : hence the *cosine* of any one of them must be greater than 0: i.e. $b^2 + c^2 - a^2$ must be greater than 0, &c.: hence a, b, c must be such

that the *squares* of any two of them are together greater than the *square* of the third.

For example, the lengths 2, 3, 4 would *not* do as the given lengths, since, although fulfilling the *first* condition, by having $2 + 3 > 4$, they fail to fulfil the *second*, as $2^2 + 3^2$ is *not* $> 4^2$.

C. L. D.

CH. CH., OXFORD.

March, 1895.

PREFACE TO SECOND EDITION.

THE principal changes, made in this Second Edition of "Pillow-Problems", are as follows:—

(1) After the numeral, which precedes each Question, Answer, or Solution, references are given to the pages at which the corresponding matter may be found.

(2) Some of the Solutions have been re-arranged, and duplicate-diagrams have been inserted, in order that every portion of text may have its illustrative diagram visible along with it, and that the reader may thus be saved the trouble, and the strain on his temper, involved in turning a leaf backwards and forwards while referring from the one to the other.

(3) In the title of the book, the words "sleepless nights" have been replaced by "wakeful hours".

This last change has been made in order to allay the anxiety of kind friends, who have written to me to express their sympathy in my broken-down state of health, believing that I am a sufferer from chronic "insomnia", and that it is as a remedy for that exhausting malady that I have recommended mathematical calculation.

The title was not, I fear, wisely chosen; and it certainly *was* liable to suggest a meaning I did not intend to convey, viz. that my "nights" are very often *wholly* "sleepless". This is by no means the case: I have never suffered from "insomnia": and the over-wakeful hours, that I have had to spend at night, have often been simply the result of the over-sleepy hours I have spent during the preceding evening! Nor is it as a remedy for *wakefulness* that I have suggested mathematical calculation; but as a remedy for the *harassing thoughts* that are apt to invade a wholly-unoccupied mind. I hope the new title will express my meaning more lucidly.

To state the matter logically, the dilemma which my friends *suppose* me to be in has, for its two horns, the endurance of a sleepless night, and the

adoption of some recipe for inducing sleep. Now, so far as *my* experience goes, no such recipe has any effect, unless when you are sleepy: and mathematical calculation would be more likely to delay, than to hasten, the advent of sleep.

The *real* dilemma, which I have had to face, is this: given that the brain is in so wakeful a condition that, do what I will, I am *certain* to remain awake for the next hour or so, I must choose between two courses, viz. either to submit to the fruitless self-torture of going through some worrying topic, over and over again, or else to dictate to myself some topic sufficiently absorbing to keep the worry at bay. A mathematical problem *is*, for me, such a topic; and is a benefit, even if it lengthens the wakeful period a little. I believe that an hour of calculation is much better for me than half-an-hour of worry.

The reader will, I think, be interested to see a curiously illogical solution which has been proposed, by a correspondent of the *Educational Times*, for Problem 61, viz. "Prove that, if any 3 Numbers be taken, which cannot be arranged in *A.P.*, and whose sum is a multiple of 3, the sum of their squares is also the sum of another set of 3 squares, the 2 sets having no common term."

The proposed solution is as follows:—

"Let $3m$, $21m$, $30m$ be the three Numbers; then

$$3m + 21m + 30m = 3 \times 18m.$$

Also
$$\begin{aligned}(3m)^2 + (21m)^2 + (30m)^2 &= (6m)^2 + (15m)^2 + (33m)^2 \\ &= (5m)^2 + (13m)^2 + (34m)^2 = (10m)^2 + (17m)^2 + (31m)^2 \\ &= (14m)^2 + (23m)^2 + (25m)^2.\end{aligned}$$

Now, if we denote, by ' α ', the property "which cannot be arranged in *A.P.*, and whose sum is a multiple of 3," and, by ' β ', the property "the sum of whose squares is also the sum of another set of 3 squares, the 2 sets having no common term," we see that all, that this writer has succeeded in proving, is that *certain selected Numbers*, which have property ' α ', have also property ' β ': but this does not prove my Theorem, viz. that *any Numbers whatever*, which have property ' α ', have also property ' β '. If his

argument were arranged in a syllogistic form, it would be found to assume a quite untenable Major Premiss, viz. "that, which is true of *certain selected Numbers* which have property ' α ', is true of *any Numbers whatever* which have property ' α '."

C. L. D.

CH. CH., OXFORD.
September, 1893.

INTRODUCTION.

NEARLY all of the following seventy-two Problems are veritable "Pillow-Problems", having been solved, in the head, while lying awake at night. (I have put on record the exact dates of some.) No. 37 and one or two others belong to the daylight, having been solved while taking a solitary walk; but every one of them was worked out, to the very end, before drawing any diagram or writing down a single word of the solution. I generally wrote down the *answer*, first of all: and *afterwards* the question and its solution. For example, in No. 70, the very first words I wrote down were as follows:—"(1) down back-edge; up again; down again; and so on; (2) about $\cdot 7$ of the way down the back-edge; (3) about $18^{\circ} 18'$; (4) about 14° ." These answers are not quite correct; but at least they are *genuine*, as the results of *mental work only*. "A poor thing, Sir, but mine own!"

My motive, for publishing these Problems, with their mentally-worked solutions, is most certainly *not* any desire to display powers of mental calculation. Mine, I feel sure, are nothing out-of-the-way; and I have no doubt there are many mathematicians who could produce, mentally, much shorter and better solutions. It is not for such persons that I intend my little book; but rather for the much larger class of *ordinary* mathematicians, who perhaps have never tried this resource, when mental occupation was needed, and who will, I hope, feel encouraged—by seeing what can be done, after a little practice, by one of *average* mathematical powers—to try the experiment for themselves, and find in it as much advantage and comfort as I have done.

The word "comfort" may perhaps sound out of place, in connection with so entirely *intellectual* an occupation; but it will, I think, come home to many who have known what it is to be haunted by some worrying subject of thought, which no effort of will is able to banish. Again and again I have said to myself, on lying down at night, after a day embittered by some vexatious matter, "I will *not* think of it any more! I have gone through it all, thoroughly. It can do no good whatever to go through it again. I *will* think

of something else!" And in another ten minutes I have found myself, once more, in the very thick of the miserable business, and torturing myself, to no purpose, with all the old troubles.

Now it is not possible—this, I think, all psychologists will admit—by any effort of volition, to carry out the resolution "I will *not* think of so-and-so." (Witness the common trick, played on a child, of saying "I'll give you a penny, if you'll stand in that corner for five minutes, and *not once* think of strawberry-jam." No human child ever yet won the tempting wager!) But it *is* possible—as I am most thankful to know—to carry out the resolution "I *will* think of so-and-so." Once fasten the attention upon a subject so chosen, and you will find that the worrying subject, which you desire to banish, is *practically* annulled. It may recur, from time to time—just looking in at the door, so to speak; but it will find itself so coldly received, and will get so little attention paid to it, that it will, after a while, cease to be any worry at all.

Perhaps I may venture, for a moment, to use a more serious tone, and to point out that there are mental troubles, much worse than mere worry, for which an absorbing subject of thought may serve as a remedy. There are sceptical thoughts, which seem for the moment to uproot the firmest faith; there are blasphemous thoughts, which dart unbidden into the most reverent souls; there are unholy thoughts, which torture, with their hateful presence, the fancy that would fain be pure. Against all these some real mental *work* is a most helpful ally. That "unclean spirit" of the parable, who brought back with him seven others more wicked than himself, only did so because he found the chamber "swept and garnished", and its owner sitting with folded hands: had he found it all alive with the "busy hum" of active *work*, there would have been scant welcome for him and his seven!

My purpose—of giving this encouragement to others—would not be so well fulfilled had I allowed myself, in writing out my solutions, to *improve* on the work done in my head. I felt it to be much more important to set down *what had actually been done in the head*, than to supply shorter or neater solutions, which perhaps would be much harder to do without paper. For example, a Long-Multiplication sum (say the multiplying together of two numbers of 7 digits) is no doubt best done, on *paper*, by beginning at the unit-end, and writing out 7 rows of figures, and adding up the columns in the usual way. But it would be very difficult indeed—to *me* quite

impossible—to do such a thing in the *head*. The only chance seems to be to begin with the *millions*, and get *them* properly grouped; then the hundred-thousands, adding the results to the previous one; and so on. Very often it seems to happen, that the easiest *mental* process looks decidedly lengthy and round-about when committed to paper.

When I first tried this plan, easy geometrical problems were all I could manage; and, even in these, I had to pause from time to time, in order to re-draw the diagram, which *would* persist in getting 'rubbed-out'. Algebraical problems I avoided at first, owing to the provoking fact that, if one single co-efficient escaped the memory, there was no resource but to begin the calculation all over again. But I soon got over both these difficulties, and found myself able to remember fairly large numerical co-efficients, and also to retain, in the mind's eye, fairly complex diagrams, even to the extent of *finding my way* from one part of the diagram to another. The *lettering* of the diagrams proved such a troublesome thing to keep in the memory, that I almost gave up using it, and learned to recognise Points by their *situation* only. In my MS. of No. 53, I find the following memorandum:—

"I had never set myself this Problem before the week ending Ap. 6, 1889. I tried it, two or three nights, lying awake; and finally worked it out on the night of Ap. $\frac{6}{7}$. All the conclusions were worked out mentally before any use was made of pen and paper. While working it, I did not give *names* to any Points, except *A*, *B*, *C*, and *P*: I merely thought of them by their positions (e. g. 'the foot of the perpendicular from *P* on *BC*')."

If any of my readers should feel inclined to reproach me with having worked too uniformly in the region of Common-place, and with never having ventured to wander out of the beaten tracks, I can proudly point to my one Problem in 'Transcendental Probabilities'—a subject in which, I believe, *very* little has yet been done by even the most enterprising of mathematical explorers. To the casual reader it may seem abnormal, and even paradoxical; but I would have such a reader ask himself, candidly, the question "Is not Life itself a Paradox?"

To give the Reader some idea of the process of construction of these Problems, I will give the biography of No. 63. The history of one is, to a great extent, the history of all.

It was begun during the night of Sept. $\frac{3}{4}$, 1890, and completed during the following night. The idea had occurred to me, a short time previously, that something interesting might be found in the subject of what I may call 'partially-regular' Solids. The 'regular' Solids are provokingly few in number; and it would be hopeless to find any question, connected with them, that has not already been exhaustively analysed: some also of the 'partially-regular' Solids (e. g. rhomboidal crystals) have probably been similarly treated; but there seemed to be room for the invention of other such Solids.

Accordingly, I devised a Solid enclosed, above and below, by 2 equal and parallel Squares, having their centres in the same vertical line, and the upper one twisted round so that its sides should be parallel to the diagonals of the lower Square. Then I imagined the upper one raised until its corners formed the vertices of 4 equilateral Triangles, whose bases were the sides of the lower one. The Solid, thus obtained, was evidently enclosed by 2 Squares and 8 equilateral Triangles: and the Problem I set myself was to obtain its *Volume*.

There was no great difficulty in proving that the distance between the 2 Squares (taking each side as equal to '2') was $2\frac{3}{4}$. But, when I looked about for some Trigonometrical method for calculating the Volume, despair soon seized upon me! A calculable Prism could be cut out of the *middle* of the Solid, I saw: but the outlying projections completely baffled me. After a while, the happy idea occurred to me of trying Algebraical Geometry, and regarding each facet as the base of a Pyramid, having its vertex at the centre of the Solid, which I decided to take as the Origin. I saw at once that I could calculate the co-ordinates of all the vertical Points, thence obtain equations to the Planes containing the facets, and thence calculate their distances from the Origin, which would be the altitudes of the Pyramids. Also it was evident that a sample Pyramid would suffice. I worked out a value for the Volume, that first night; but the thing got into a tangle, and I felt pretty sure I had got it wrong.

The next night I began again, and worked it all through from the beginning. In the morning the *answer* was clear in my memory, and I wrote it down at once; and did not write out the Problem, and its solution, until

later in the day, when I was well pleased to find the written proof confirm the result I had arrived at in the hours of darkness.

It is not, perhaps, much to be wondered at that, when these Problems came to be re-written and arranged for publication, a good many mistakes were discovered. Some were so bad as quite to spoil the solutions in which they occurred: these Problems I have omitted altogether. The others I have corrected, in the solutions as given in Chapter III: but, that I may not be credited with an amount of accuracy, as a computator, which I am well aware I do not possess, I here append a list of them.

In No. 7, in the denominator ' $2 \sin A$ ', I forgot the '2'.^[1]

In No. 10, I failed to notice that the 3 coins might *also* be a half-crown and 2 shillings.

In No. 13, in the last line but one, I put ' $2bc.ca$ ', instead of ' $4bc.ca$ '.

In No. 32, I brought out the arithmetical value as '358520', instead of '358550'.

In No. 38, I got the decimal wrong, making it $\cdot 476$ instead of $\cdot 478$, and thus brought out the answer as $\cdot 042$ instead of $\cdot 044$.

In No. 44, I said that the denominator would be of the form $(10^n - 1) \cdot 10^m$. This last factor is superfluous: i. e. $m = 0$.

In No. 50, I made a mistake near the end, bringing out $\frac{41}{108}$, instead of $\frac{50}{108}$.

In No. 55, I put 'tan' for 'sin'.

In No. 57, in the last paragraph, I replaced the denominator ' $a \sin B \sin C$ ' by (what I imagined to be its equivalent) ' $2m$ '. Apparently I was under the delusion that ' $a \sin B \sin C$ ' was the same thing as ' $\sin A \cdot bc$ '!

In No. 70, section (3), I forgot to add in the $\cdot 45$, thus making the answer half a degree wrong. And, in section (4), I forgot to add in the $\cdot 53$, thus again making the answer half a degree wrong.

Let me, in conclusion, gratefully acknowledge the valuable assistance I have received from Mr. F. G. Brabant, M.A., of Corpus Christi College, Oxford, who has most patiently and carefully gone through my proofs, first

working out each result independently, and has thus detected many mistakes which had escaped my notice. He has also supplied, for No. 59, a much neater answer than mine, viz. $\frac{abc}{3} \cdot \sqrt{\cos A \cos B \cos C}$.

Other mistakes may perchance, having eluded us both, await the penetrating glance of some critical reader, to whom the joy of discovery, and the intellectual superiority which he will thus discern, in himself, to the author of this little book, will, I hope, repay to some extent the time and trouble its perusal may have cost him!

C. L. D.

CH. CH., OXFORD.
May, 1893.

FOOTNOTES:

[1] In the trigonometrical Problems, I have used the symbols $\cap$ and $\cup$, to represent the words 'sine' and 'cosine'.

CONTENTS.



	PAGE
CHAPTER I. QUESTIONS	1
CHAPTER II. ANSWERS	19
CHAPTER III. SOLUTIONS	28

SUBJECTS CLASSIFIED.

ARITHMETIC. No. [31](#).

ALGEBRA:—

Equational Problems. Nos. [8](#), [25](#), [39](#), [52](#), [68](#).

Series. Nos. [21](#), [32](#).

Indeterminate Equations. No. [47](#).

Properties of Numbers. Nos. [1](#), [14](#), [29](#), [44](#), [61](#).

Chances. Nos. [5](#), [10](#), [16](#), [19](#), [23](#), [27](#), [38](#), [41](#), [45](#), [50](#), [58](#), [66](#).

PURE GEOMETRY, PLANE. Nos. [2](#), [3](#), [9](#), [15](#), [17](#), [18](#), [20](#), [24](#), [26](#), [30](#), [34](#), [35](#),
[36](#), [40](#), [46](#), [51](#), [57](#), [62](#), [64](#), [71](#).

TRIGONOMETRY:—

Plane. Nos. [4](#), [6](#), [7](#), [11](#), [12](#), [13](#), [18](#), [22](#), [28](#), [37](#), [42](#), [43](#), [48](#), [54](#), [55](#), [56](#),
[57](#), [60](#), [65](#), [69](#).

Solid. Nos. [49](#), [59](#), [63](#), [70](#).

ALGEBRAICAL GEOMETRY:—

Plane. No. [53](#).

Solid. No. [67](#).

DIFFERENTIAL CALCULUS:—

Maxima and Minima. No. [33](#).

TRANSCENDENTAL PROBABILITIES. No. [72](#).

PILLOW-PROBLEMS.

CHAPTER I.

Questions.

—

1. (28)^[2]

Find a general formula for two squares whose sum = 2.

[24/3/84

2. (29)

In a given Triangle to place a line parallel to the base, such that the portions of sides, intercepted between it and the base, shall be together equal to the base.

3. (30)

If the sides of a Tetragon pass through the vertices of a Parallelogram, and if three of them are bisected at those vertices: prove that the fourth is so also.

4. (30)

In a given acute-angled Triangle inscribe a Triangle, whose sides make, at each of the vertices, equal angles with the sides of the given Triangle.

[19/4/76

5. (19, 31)

A bag contains one counter, known to be either white or black. A white counter is put in, the bag shaken, and a counter drawn out, which proves to be white. What is now the chance of drawing a white counter?

[8/9/87]

6. (19, 32)

Given lengths of lines drawn, from vertices of Triangle, to middle points of opposite sides, to find its sides and angles.

7. (19, 33)

Given 2 adjacent sides, and the included angle, of a Tetragon; and that the angles, at the other ends of these 2 sides, are right: find (1) remaining sides, (2) area.

[4 or 5/89]

8. (20, 34)

Some men sat in a circle, so that each had 2 neighbours; and each had a certain number of shillings. The first had 1/ more than the second, who had 1/ more than the third, and so on. The first gave 1/ to the second, who gave 2/ to the third, and so on, each giving 1/ more than he received, as long as possible. There were then 2 neighbours, one of whom had 4 times as much as the other. How many men were there? And how much had the poorest man at first?

[3/89]

9. (35)

Given two Lines meeting at a Point, and given a Point lying within the angle contained by them: draw, from the given Point, two lines, at right angles to each other, and forming with the given Lines and the line joining their intersection to the given Point, two equal Triangles.

[11/76]


10. (20, 36)

A triangular billiard-table has 3 pockets, one in each corner, one of which will hold only one ball, while each of the others will hold two. There are 3 balls on the table, each containing a single coin. The table is tilted up, so that the balls run into one corner, it is not known which. The 'expectation', as to the contents of the pocket, is $\frac{2}{6}$. What are the coins?

[8/90]

11. (20, 36)

A Triangle ABC has another $A'B'C'$ inscribed in it, so that $\angle BA'C' = \angle CB'A' = \angle AC'B' = \theta$; thus making it similar to the first Triangle. Find ratio between homologous sides. And solve for " $\theta = 90^\circ$ ".



The Triangles can be proved similar thus:—

$$\begin{aligned}\angle C'A'B' + \angle B'A'C &= \text{supp. of } \theta, \\ \angle B'A'C + \angle A'CB' &= \text{supp. of } \theta; \\ \therefore \text{ these pairs are equal; } \therefore \angle C'A'B' &= C.\end{aligned}$$

Hence $\angle A'B'C' = A$, and $\angle B'C'A' = B$.

Let $C'A' = ka$; $\therefore A'B' = kb$, and $B'C' = kc$. We have to find k .

[31/3/82]

12. (20, 37)

Given the semi-perimeter and the area of a Triangle, and also the volume of the cuboid whose edges are equal to the sides of the Triangle: find the sum of the squares of its sides.

[23/1/91]

13. (20, 38)

Given the lengths of the radii of two intersecting Circles, and the distance between their centres: find the area of the Tetragon formed by the tangents at the points of intersection.

[3/89]

14. (39)

Prove that 3 times the sum of 3 squares is also the sum of 4 squares.

[2/12/81]

15. (39)

If a Figure be such that the opposite angles of every inscribed Tetragon are supplementary: the Figure is a Circle.

[3/91]

16. (20, 40)

There are two bags, one containing a counter, known to be either white or black; the other containing 1 white and 2 black. A white is put into the first, the bag shaken, and a counter drawn out, which proves to be white. Which course will now give the best chance of drawing a white—to draw from one of the two bags without knowing which it is, or to empty one bag into the other and then draw?

[10/87]

17. (40)

In a given Triangle place a line parallel to the base, such that if, from its ends, lines be drawn, parallel to the sides and terminated by the base, they shall be together equal to the first line.

[3/89]

18. (21, 41)

Find a Point, in the base of a given Triangle, such that, if from it perpendiculars be dropped upon the sides, the line joining their extremities shall be parallel to the base. (1) Trigonometrically. (2) Geometrically.

[11/89]

19. (21, 42)

There are 3 bags; one containing a white counter and a black one, another two white and a black, and the third 3 white and a black. It is not known in what order the bags are placed. A white counter is drawn from one of them, and a black from another. What is the chance of drawing a white counter from the remaining bag?

20. (43)

In the base of a given Triangle find a Point such that if from it two lines be drawn, terminated by the sides, one being perpendicular to the base and one to the left-hand side, they shall be equal.

[5/88]

21. (21, 44)

Sum, (1) to n terms, (2) to 100 terms, the series

$$1 \cdot 3 \cdot 5 + 2 \cdot 4 \cdot 6 + \&c.$$

[7/4/89]

22. (21, 45)

Given the 3 'altitudes' of a Triangle: find its (1) sides, (2) angles, (3) area.

[4/6/89]


23. (21, 46)

A bag contains 2 counters, each of which is known to be black or white. 2 white and a black are put in, and 2 white and a black drawn out. Then a white is put in, and a white drawn out. What is the chance that it now contains 2 white?

[25/9/87]

24. (21, 47)

If, from the vertices of a triangle ABC , the lines AD , BE , CF be drawn, intersecting at O : find the ratio $\frac{DO}{DA}$ in terms of the two ratios $\frac{EO}{EB}$, $\frac{FO}{FC}$.



[5/86]

25. (22, 48)

If ' ϵ ', ' α ', ' λ ' represent proper fractions; and if, in a certain hospital, ' ϵ ' of the patients have lost an eye, ' α ' an arm, and ' λ ' a leg: what is the least possible number who have lost all three?

[7/2/76]

26. (48)

Within a given Triangle place a similar Triangle, whose area shall have to its area a given ratio less than unity, whose sides shall be parallel to its sides, and whose vertices shall be equidistant from its vertices.

[4/89]

27. (22, 50)

There are 3 bags, each containing 6 counters; one contains 5 white and one black; another, 4 white and 2 black; the third, 3 white and 3 black. From two of the bags (it is not known which) 2 counters are drawn, and prove to be black and white. What is the chance of drawing a white counter from the remaining bag?

[4/3/80]

28. (22, 50)

If the sides of a given Triangle, taken cyclically, be divided in extreme and mean ratio; and if the Points be joined: find the ratio which the area of the Triangle, so formed, has to the area of the given Triangle.

[12/78]

29. (51)

Prove that the sum of 2 different squares, multiplied by the sum of 2 different squares, gives the sum of 2 squares in 2 different ways.

[3/12/81]

30. (52)

In a given Triangle, to place a line parallel to the base, such that if from its extremities lines be drawn, to the base, parallel to the sides, they shall be together double of the inscribed Line.

[15/3/89]

31. (22, 53)

On July 1, at 8 a.m. by my watch, it was 8*h.* 4*m.* by my clock. I took the watch to Greenwich, and, when it said 'noon', the true time was 12*h.* 5*m.* That evening, when the watch said '6*h.*', the clock said '5*h.* 59*m.*'

On July 30, at 9 a.m. by my watch, it was 8*h.* 57*m.* by my clock. At Greenwich, when the watch said '12*h.* 10*m.*', the true time was 12*h.* 5*m.* That evening, when the watch said '7*h.*' the clock said '6*h.* 58*m.*'

My watch is only wound up for each journey, and goes uniformly during any one day: the clock is always going, and goes uniformly.

How am I to know when it is *true* noon on July 31?

[14/3/89]

32. (22, 53)

Sum the Series $1 \cdot 5 + 2 \cdot 6 + \&c.$ (1) to n terms; (2) to 100 terms.

[7/4/89]

33. (54)

Inscribe in a given Circle the maximum Tetragon having 2 parallel sides, one double the other.

34. (55)

From a given Point draw 2 Lines, one to the centre of a given Circle, and the other cutting off from it a Segment containing an angle equal to that

between the Lines.

[21/12/74]

35. (56)

With a given Triangle, to describe a Circle, cutting each side in two points, such that, if radii be drawn perpendicular to the sides, they are divided by the sides in given ratios.

[11/76]

36. (57)

In a given Triangle, to draw a line, from a Point on one side of it, to a Point on the other side, perpendicular to one of these sides, and equal to the sum of the portions, of these sides, intercepted between it and the base.

[3/89]

37. (22, 58)

Two given Circles intersect, so that their common chord subtends angles of 30° and 60° at their centres. What fraction of the smaller Circle is within the larger?

[12/91]

38. (22, 60)

There are 3 bags, '*A*', '*B*', and '*C*'. '*A*' contains 3 red counters, '*B*' 2 red and one white, '*C*' one red and 2 white. Two bags are taken at random, and a counter drawn from each: both prove to be red. The counters are replaced, and the experiment is repeated with the same two bags: one proves to be red. What is the chance of the other being red?

[3/76]

39. (22, 60)

A and *B* begin, at 6 a. m. on the same day, to walk along a road in the same direction, *B* having a start of 14 miles, and each walking from 6 a. m. to 6 p. m. daily. *A* walks 10 miles, at a uniform pace, the first day, 9 the second, 8 the third, and so on: *B* walks 2 miles, at a uniform pace, the first day, 4 the second, 6 the third, and so on. When and where are they together?

[16/3/78]

40. (61)

In a given Triangle, whose base-angles are acute, draw two lines, at right angles to the base, and together equal to the line drawn, from the vertex, at right angles to the base, and such that

- (1) they are equidistant from the line drawn from the vertex;
- (2) they are equidistant from the ends of the base.

[5/76]

41. (23, 62)

My friend brings me a bag containing four counters, each of which is either black or white. He bids me draw two, both of which prove to be white. He then says "I meant to tell you, before you began, that there was at least *one* white counter in the bag. However, you know it now, without my telling you. Draw again."

- (1) What is now my chance of drawing white?
- (2) What would it have been, if he had not spoken?

[9/87]

42. (23, 63)

If the angles of a given Triangle be bisected, and if lines be drawn, through its vertices, at right angles to the bisectors, so as to form a fresh

Triangle: find the ratio of the area of this Triangle to the area of the given Triangle.

[17/5/78]

43. (65)

From the ends of the base of a given Triangle draw two lines, intersecting, terminated by the sides, and forming an isosceles Triangle at the base, and a Tetragon, equal to it, at the vertex.

[2/82]

44. (66)

If a, b be two numbers prime to each other, a value may be found for n which will make $(a^n - 1)$ a multiple of b .

[18/3/81]

45. (23, 67)

If an infinite number of rods be broken: find the chance that one at least is broken in the middle.

[5/84]

46. (68)

In a given Triangle, whose base is divided at a given Point, inscribe a Triangle, having its angles equal to given angles, and having an assigned vertex at the given Point.

[19/11/87]

47. (23, 69)

Solve the 2 Indeterminate Equations

$$\left. \begin{aligned} \frac{x}{y} &= x - z; \\ \frac{x}{z} &= x - y; \end{aligned} \right\} \quad \begin{aligned} (1) \\ (2) \end{aligned}$$

and find the limits, if any, between which the *real* values lie.

[12/90]

48. (70)

If semicircles be described, externally, on the sides of a given Triangle; and if their common tangents be drawn; and if their lengths be α , β , γ : prove that

$$\left(\frac{\beta\gamma}{\alpha} + \frac{\gamma\alpha}{\beta} + \frac{\alpha\beta}{\gamma} \right)$$

is equal to the semiperimeter of the Triangle.

[9/2/81]

49. (23, 72)

If four equilateral Triangles be made the sides of a square Pyramid: find the ratio which its volume has to that of a Tetrahedron made of the Triangles.

[16/11/86]

50. (23, 72)

There are 2 bags, H and K , each containing 2 counters: and it is known that each counter is either black or white. A white counter is added to bag H , the bag is shaken up, and one counter transferred (without looking at it) to bag K , where the process is repeated, a counter being transferred to bag H . What is now the chance of drawing a white counter from bag H ?

51. (74)

From a given Point, in one side of a given Triangle, to draw a line, terminated by the other side, so that, if from its ends lines be drawn at right angles to the base, their sum shall be equal to the first line.

[12/81

52. (23, 75)

Five beggars sat down in a circle, and each piled up, in a heap before him, the pennies he had received that day: and the five heaps were equal.

Then spake the eldest and wisest of them, unfolding, as he spake, an empty sack.

"My friends, let me teach you a pretty little game! First, I name myself 'Number One,' my left-hand neighbour 'Number Two,' and so on to 'Number Five.' I then pour into this sack the whole of my earnings for the day, and hand it on to him who sits next but one on my left, that is, 'Number Three.' *His* part in the game is to take out of it, and give to his two neighbours, so many pennies as represent their names (that is, he must give four to 'Number Four' and two to 'Number Two'); he must then put *into* the sack half as much as it contained when he received it; and he must then hand it on just as I did, that is, he must hand it to him who sits next but one on his left—who will of course be 'Number Five.' *He* must proceed in the same way, and hand it on to 'Number Two,' from whom the sack will find its way to 'Number Four,' and so to me again. If any player cannot furnish, from his own heap, the whole of what he has to put into the sack, he is at liberty to draw upon any of the other heaps, *except mine!*"

The other beggars entered into the game with much enthusiasm: and in due time the sack returned to 'Number One,' who put into it the two pennies he had received during the game, and carefully tied up the mouth of it with a string. Then, remarking "it is a *very* pretty little game," he rose to his feet, and hastily quitted the spot. The other four beggars gazed at each other with rueful countenances. Not one of them had a penny left!

How much had each at first?

[16/2/89

53. (24, 76)

In a triangular billiard-table, a Point is given by its trilinear co-ordinates. A ball, starting from the given Point, strikes the three sides, and returns to the starting-point. Find, in terms of the trilinear co-ordinates and of the angles of the Triangle, the Point where the ball strikes the second side.

[6/4/89]

54. (24, 78)

Cut off, from a given Triangle, by lines parallel to the sides, 3 Triangles, so that the remaining Hexagon may be equilateral. Also find the lengths of its sides in terms of the sides of the given Triangle: and the ratios in which the sides of the given Triangle are divided.

[18/4/86]

55. (79)

Given three cylindrical towers on a Plane: find a Point, on the Plane, from which they shall look the same width.

[20/12/74]

56. (24, 80)

Given the 3 altitudes of a Triangle: construct it.

[27/6/84]

57. (25, 80)

In a given Triangle describe three Squares, whose bases shall lie along the sides of the Triangle, and whose upper edges shall form a Triangle;

(1) geometrically; (2) trigonometrically.

[27/1/91]

58. (25, 83)

Three Points are taken at random on an infinite Plane. Find the chance of their being the vertices of an obtuse-angled Triangle.

[20/1/84]

59. (25, 84)

Given a Tetrahedron, having every edge equal to the opposite edge, so that its facets are all (when looked at from the outside) identically equal: find its volume in terms of its edges.

[8/90]

60. (25, 87)

Given a Triangle ABC , and that its base BC is divided at D in the ratio m to n : find the angles BAD , CAD .

[21/3/90]

61. (89)

Prove that, if any 3 Numbers be taken, which cannot be arranged in A. P., and whose sum is a multiple of 3, the sum of their squares is also the sum of another set of 3 squares, the 2 sets having no common term.

[1/12/81]

62. (91)

Given two Lines meeting at a Point, and given a Point lying within the angle contained by them: draw a line, through the given Point, and forming, with the given Lines, the least possible Triangle.

[12/76]

63. (26, 92)

Given 2 equal Squares, in different horizontal planes, having their centres in the same vertical line, and so placed that the sides of each are parallel to the diagonals of the other, and at such a distance apart that, by joining neighbouring vertices, 8 equilateral Triangles are formed: find the volume of the solid thus enclosed.

[3, 4/9/90]

64. (94)

Given a Triangle, and a Point within it such that its distance from one of the sides is less than its distance from either of the others: describe a Circle, with given Point as centre, such that its intercepts on the sides may be equal to the sides of a right-angled Triangle.

[18/12/74]

65. (95)

How many shapes are there for Triangles which have all their angles aliquot parts of 360° ?

[5/89]

66. (26, 97)

Given that there are 2 counters in a bag, as to which all that was originally known was that each was either white or black. Also given that the experiment has been tried, a certain number of times, of drawing a counter, looking at it, and replacing it; that it has been white every time; and that, as a result, the chance of drawing white, next time, is $\frac{\alpha}{\alpha + \beta}$. Also given that the same experiment is repeated m times more, and that it still continues to be white every time. What would then be the chance of drawing white?

[9/89]

67. (26, 100)

If a regular Tetrahedron be placed, with one vertex downwards, in a socket which exactly fits it, and be turned round its vertical axis, through an angle of 120° , raising it only so much as is necessary, until it again fits the socket: find the Locus of one of the revolving vertices.

[27/1/72]

68. (26, 101)

Five friends agreed to form themselves into a Wine-Company (Limited). They contributed equal amounts of wine, which had been bought at the same price. They then elected one of themselves to act as Treasurer; and another of them undertook to act as Salesman, and to sell the wine at 10% over cost-price.

The first day the Salesman drank one bottle, sold some, and handed over the receipts to the Treasurer.

The second day he drank none, but pocketed the profits on one bottle sold, and handed over the rest of the receipts to the Treasurer.

That night the Treasurer visited the Cellars, and counted the remaining wine. "It will fetch just £11," he muttered to himself as he left the Cellars.

The third day the Salesman drank one bottle, pocketed the profits on another, and handed over the rest of the receipts to the Treasurer.

The wine was now all gone: the Company held a Meeting, and found to their chagrin that their profits (i. e. the Treasurer's receipts, less the original value of the wine) only cleared 6d. a bottle on the whole stock. These profits had accrued in 3 equal sums on the 3 days (i. e. the Treasurer's receipts for the day, less the original value of the wine taken out during the day, had come to the same amount every time); but of course only the Salesman knew this.

(1) How much wine had they bought? (2) At what price?

[28/2/89]

69. (26, 102)

If, from each of the angles of a given Triangle ABC , taken cyclically, a certain proper fraction of it be cut off, the arithmetical values of the 3 fractions being represented by ' k, l, m '; and if it be given that the Triangle, formed by the lines so drawn, is similar to the given one, the angle, formed by the lines drawn from B and C , being equal to A , and so on: find k, l, m , as similar functions of a single variable. Also find the ratio which each side of the second Triangle bears to the corresponding side of the first.

[8/89]

70. (27, 105)

Let an equilateral and equiangular Tetrahedron be placed with one facet in front: and suppose a series of triangles, equal to that facet, constructed in the Plane containing that facet, and having a base common with it; and that they are all wrapped round the Tetrahedron as far as they will go. Find (1) the locus of their vertices; (2) the situation of the vertex of the one whose left-hand base-angle is 15° ; (3) the left-hand base-angle of the one which (wrapped round towards the right) covers portions of all four facets of the Tetrahedron, and whose vertex coincides with *its* vertex; (4) the left-hand base-angle of the one which (similarly treated) occupies all four facets, and then the front and right-hand facet for the second time, and whose vertex coincides with the distal vertex of the base of the Tetrahedron.

71. (108)

In a given Triangle place a Hexagon having its opposite sides equal and parallel, and three of them lying along the sides of the Triangle, and such that its diagonals intersect in a given Point.

[14/12/74]

72. (27, 109)

A bag contains 2 counters, as to which nothing is known except that each is either black or white. Ascertain their colours without taking them

out of the bag.

[8/9/87

FOOTNOTES:

[\[2\]](#) The numerals, placed in parentheses, indicate the pages where the corresponding matter may be found.

CHAPTER II.

Answers.

5. (2, 31)

Two-thirds.


6. (2,32)

Calling the sides ' $2a$ ', ' $2b$ ', ' $2c$ ' and the lines ' α ', ' β ', ' γ ', we have

$$a^2 = \frac{-\alpha^2 + 2\beta^2 + 2\gamma^2}{9},$$

$$\cos A = \frac{5\alpha^2 - \beta^2 - \gamma^2}{2\sqrt{2\alpha^2 - \beta^2 + 2\gamma^2} \cdot \sqrt{2\alpha^2 + 2\beta^2 - \gamma^2}}.$$

7. (2, 33)



Let AB , AD be given sides, and B , D the right $\angle$ s; and let $AB = b$, $AD = d$.

$$(1) \quad BC = \frac{d - b \cos A}{\sin A}; \quad CD = \frac{b - d \cos A}{\sin A};$$

$$(2) \quad \text{area} = \frac{2bd - (b^2 + d^2) \cos A}{2 \sin A}.$$

8. (2, 34)

7 men; 2 shillings.

10. (3, 36)

Either 2 florins and a sixpence; or else a half-crown and 2 shillings.

11. (3, 36)

The required ratio is equal to

$$\frac{\sin A \sin B \sin C}{\sin \theta (1 + \cos A \cos B \cos C) + \cos \theta \sin A \sin B \sin C}.$$

$$\text{If } \theta = 90^\circ, \text{ this} = \frac{\sin A \sin B \sin C}{1 + \cos A \cos B \cos C}.$$

12. (4, 37)

If s = semi-perimeter, m = area, v = volume; then

$$a^2 + b^2 + c^2 = 2.\left(s^2 - \frac{v}{s} - \frac{m^2}{s^2}\right).$$

13. (4, 38)

If ' $2M$ ' = area of Tetragon whose vertices are the Centres and the Points of intersection; and if its sides be ' a ', ' b ', and its diagonal, joining the Centres, ' c ': required area

$$= \frac{32M^3}{(b^2 + c^2 - a^2).(c^2 + a^2 - b^2)}.$$

16. (4, 40)

The first course gives chance = $\frac{1}{2}$; the second, $\frac{5}{12}$. Hence the first is best.

18. (5, 41)

(1) Divide base BC , at E , so that $\frac{BE}{EC} = \frac{\sin 2C}{\sin 2B}$.

(2) At B, C , make right angles ABD, ACD ; and join AD cutting BC at E , which is the required Point.

19. (5, 42)

Eleven-seventeenths.

21. (5, 44)

(1) $\frac{\overline{n.n + 1.n + 4.n + 5}}{4}$; (2) 27, 573, 000.

22. (5, 45)

Calling the given altitudes ' α, β, γ '; and the fraction

$$\frac{2\alpha^2\beta^2\gamma^2.(\alpha^2 + \beta^2 + \gamma^2) - (\beta^4\gamma^4 + \gamma^4\alpha^4 + \alpha^4\beta^4)}{4\alpha^4\beta^4\gamma^4}, k^2,$$

(1) $a = \frac{1}{k\alpha}$, &c.

(2) $\sin A = k\beta\gamma$, &c.

(3) $\text{area} = \frac{1}{2k}$.

23. (5, 46)

Two-fifths.

24. (6, 47)

$\frac{DO}{DA} + \frac{EO}{EB} + \frac{FO}{FC} = 1$; whence any one can be found in terms of the other two.

25. (6, 48)

$$\epsilon + \alpha + \lambda - 2.$$

27. (6, 50)

Seventeen-twentyfifths.

28. (7, 50)

$$7 - 3\sqrt{5}.$$

31. (7, 53)

When the clock says '12h. 2m. $29\frac{277}{288}$ sec.'

32. (8, 53)

$$(1) \quad \frac{n.(n+1).(2n+13)}{6};$$

$$(2) \quad 358550.$$

37. (8, 58)

$$\frac{4 + \sqrt{3}}{12} - \frac{1 + \sqrt{3}}{2\pi}; \quad \text{i.e. about } .044$$

38. (9, 60)

Fortynine-seventytwoths.

39. (9, 60)

They meet at end of $2d$. $6h.$, and at end of $4d.$: and the distances are 23 miles, and 34 miles.

41. (9, 62)

(1) Seven-twelfths. (2) One-half.

42. (10, 63)

$$\frac{abc}{2(s-a).(s-b).(s-c)}.$$

45. (10, 67)

.6321207 &c.

47. (11, 69)

One set of values is 0, 0, 0.

A 2nd set is $x = y = 0$; z has any value.

A 3rd is $x = z = 0$; y has any value.

And the 4th set is $x = \frac{k^2}{k-1}$, $y = z = k$; where k has any value.

If x has any positive value less than 4, y and z are unreal.

49. (11, 72)

Two.

50. (11, 72)

Seventeen-twentysevenths.

52. (12, 75)

2l. 18s. 0d.

53. (13, 76)

The portion, cut off from the second side, is equal to

$$\frac{(\alpha \sin C + \gamma \sin A)(2\gamma \cos A + \beta)}{\alpha \cos C + \gamma \cos A + \beta} + \frac{\beta \cos A + \gamma \cos 2A}{\sin A}.$$

54. (13,78)


Side AB must be divided at D, G , so that

$$AD : DG : GB :: \frac{1}{a} : \frac{1}{c} : \frac{1}{b};$$

and similarly for the other sides. Also each side of the Hexagon =

$$\frac{1}{\frac{1}{a} + \frac{1}{b} + \frac{1}{c}}.$$

56. (13, 80)



Draw BC , CE , BD equal to the given altitudes, so as to form right $\angle$ s at B and C : and produce DB , EC . Join DC , and draw $CF \perp$ to it. Join EB , and draw $BG \perp$ to it. With centre B , and distance BF , describe a circle: with centre C , and distance CG describe another: let them meet at A : and join AB , AC . Triangle ABC may be proved to be similar to required Triangle. The rest of the construction is obvious.

57. (14, 80)

(1) *Geometrically.*

If Squares be described externally on the sides of the given Triangle; and if their outer edges be produced to form a new Triangle; and if the sides of the given Triangle be divided similarly to those of the new Triangle: their central portions will be the bases of the required Squares.

(2) *Trigonometrically.*

If a, b, c be the sides of the given Triangle, and m its area; and if x, y, z be the sides of the required Squares: then

$$\frac{a}{x} = \frac{b}{y} = \frac{c}{z} = \frac{a^2 + b^2 + c^2}{2m} + 1.$$

58. (14, 83)

$$\frac{3}{8 - \frac{6\sqrt{3}}{\pi}}.$$

59. (14, 84)

Calling lengths of the 3 pairs of edges ' a, b, c ', and the corresponding $\angle$ s, in each facet, ' A, B, C '; volume =

$$\frac{abc}{6} \cdot \sqrt{1 - (\cos^2 A + \cos^2 B + \cos^2 C) + 2 \cos A \cos B \cos C}.$$

60. (14, 87)

$$\cot BAD = \frac{(m+n) \cot A + n \cot B}{m};$$

$$\text{similarly, } \cot CAD = \frac{(m+n) \cot A + m \cot C}{n}.$$

63. (15, 92)

If each side of each Square = 2, the volume =

$$\frac{8.2^{\frac{1}{4}} \cdot (\sqrt{2} + 1)}{3}.$$

66. (15, 97)

$$\frac{2^m \cdot (\alpha - \beta) + \beta}{2^m \cdot (\alpha - \beta) + 2\beta}.$$

67. (16, 100)

If the centre of the horizontal facet be taken as the Origin, and if the X -axis pass through one of the vertices of that facet, and the Y -axis be parallel to the opposite edge of that facet, and the Z -axis be perpendicular to that facet: and if the altitude (measured downwards) of the Tetrahedron be called ' h ', and the intercept on the X -axis be called ' a ': the Equations to the Locus are

$$(x + \sqrt{3}.y).(h - z) = ah;$$
$$x^2 + y^2 = a^2.$$

68. (16, 101)

(1) 5 dozen; (2) 8/4 a bottle.

69. (17, 102)

$$(1) k = \frac{\theta - B}{A}; l = \frac{\theta - C}{B}; m = \frac{\theta - A}{C}.$$

(2) Calling new Triangle ' $A'B'C'$ ',

$$\frac{a'}{a} = \frac{b'}{b} = \frac{c'}{c} = 2 \cos \theta.$$

70. (17, 105)

(1) Down the back-edge; up again; and so on. (2) about $\cdot 7$ of the way down the back-edge. (3) About $18 \cdot 65^\circ$. (4) About $14 \cdot 53^\circ$.

72. (18, 109)

One is black, and the other white.

CHAPTER III.

Solutions.

1. (1)

Let u, v be the Nos.

Then $u^2 + v^2 = 2$.

Evidently ' $(1 + k), (1 - k)$ ' is a form for the squares.

Also, if we write ' $2m^2$ ' for ' 2 ' (which will not interfere with the problem, as we can divide by m^2 , and get $\frac{u^2}{m^2} + \frac{v^2}{m^2} = 2$), the above form becomes ' $(m^2 + k), (m^2 - k)$ '.

Now, as these are *squares*, their resemblance to

$$'(a^2 + b^2 + 2ab), (a^2 + b^2 - 2ab)'$$

at once suggests itself; so that the problem depends on the known one of finding a, b , such that $(a^2 + b^2)$ is a square; and we can then take $2ab$ as k .

A general form for this is

$$a = x^2 - y^2,$$

$$b = 2xy;$$

$$\therefore a^2 + b^2 = (x^2 + y^2)^2;$$


$\therefore$ the formula ' $u^2 + v^2 = 2m^2$ ' becomes

$$(x^2 - y^2 + 2xy)^2 + (x^2 - y^2 - 2xy)^2 = 2(x^2 + y^2)^2;$$

$$\text{i. e. } \left(\frac{x^2 - y^2 + 2xy}{x^2 + y^2}\right)^2 + \left(\frac{x^2 - y^2 - 2xy}{x^2 + y^2}\right)^2 = 2.$$

Q. E. F.

2. (1)



(Analysis.)

Let ABC be the Triangle, and DE the required line, so that $BD + CE = BC$.

From BC cut off BF equal to BD ; then $CF = CE$.

Join DF, EF .

Now $\angle BDF = \angle BFD = [\text{by I. 29}] \angle FDE$;

Similarly $\angle CEF = \angle FED$;

$\therefore \angle$ s BDE, CED , are bisected by DF, EF , and F is centre of $\odot$ escribed to $\triangle ADE$.

Drop, from F , $\perp$ s on BD, DE, EC ; then these $\perp$ s are equal.

Hence, if AF be joined, it bisects $\angle A$.

Hence construction.

(Synthesis.)

Bisect $\angle A$ by AF : from F draw $FB', FC', \perp AC, AB$: also draw $FA' \perp BC$ and equal to FB' : and through A' draw $DE \perp FA'$, i. e. $\parallel BC$. Then DE shall be line required.

$\therefore \angle$ s at A', B', C' , are right, and $FA' = FB' = FC'$,


$\therefore \angle$ s BDE, CED , are bisected by DF, EF .

Now $\angle BFD = \angle FDA'$; $\therefore$ it $= \angle BDF$; $\therefore BF = BD$;

Similarly $CF = CE$; $\therefore BC = BD + CE$.

Q. E. F.

3. (1)



Let $ABCD$ be the Tetragon; and let the 3 sides, AB , BC , CD , be bisected by vertices of the Parallelogram $EFGH$.

Join BD .

$\therefore$, in Triangle BCD , sides BC , CD are bisected at F and G ,

$\therefore FG$ is parallel to BD ;

but EH is parallel to FG ;

$\therefore EH$ is parallel to BD ;


$\therefore$ Triangles AEH , ABD are similar;

now AE is half of AB ;

$\therefore AH$ is half of AD .

Q. E. D.

4. (1)



Let ABC be the given Triangle, and $A'B'C'$ the required Triangle, so that $\angle BA'C' = \angle CA'B'$, &c.

Evidently $A'C'$, $A'B'$ are equally inclined to a line drawn, from A' , $\perp BC$; and so of the others: i. e. these $\perp$ s bisect the $\angle$ s at A' , B' , C' ;

$\therefore$ they meet in the same Point. Draw them; let them meet at O ; and call the $\angle C'A'B'$ ' 2α ', and so on.

Now $(\beta + \gamma) = \pi - \angle B'OC' = A$;

$\therefore 2A = 2(\beta + \gamma) = \pi - 2\alpha$;

$\therefore \alpha = 90^\circ - A$;

$\therefore \angle BA'C' = A$.

Similarly, $\angle BC'A' = C$.

$\therefore$ Triangle $BC'A'$ is similar to Triangle BCA ; and so of the others;

$$\begin{aligned}
\therefore BA' &= \frac{c}{a} \cdot BC' = \frac{c}{a} \cdot (c - AC'), \\
&= \frac{c}{a} \cdot \left(c - \frac{b}{c} \cdot AB'\right), \\
&= \frac{c^2}{a} - \frac{b}{a} \cdot (b - CB'), \\
&= \frac{c^2}{a} - \frac{b^2}{a} + \frac{b}{a} \cdot \frac{a}{b} \cdot CA', \\
&= \frac{c^2}{a} - \frac{b^2}{a} + a - BA'; \\
\therefore 2BA' &= \frac{c^2 + a^2 - b^2}{a} = \frac{2ca \cos B}{a}; \\
\therefore BA' &= c \cos B;
\end{aligned}$$

$\therefore A'$ is foot of $\perp$ drawn, from A , to BC . Hence the construction is obvious.

Q. E. F.

5. (2, 19)

At first sight, it would appear that, as the state of the bag, *after* the operation, is necessarily identical with its state *before* it, the chance is just what it then was, viz. $\frac{1}{2}$. This, however, is an error.


The chances, *before* the addition, that the bag contains (a) 1 white (b) 1 black, are (a) $\frac{1}{2}$ (b) $\frac{1}{2}$. Hence the chances, *after* the addition, that it contains (a) 2 white (b) 1 white, 1 black, are the same, viz. (a) $\frac{1}{2}$ (b) $\frac{1}{2}$. Now the probabilities, which these 2 states give to the observed event, of drawing a white counter, are (a) certainty (b) $\frac{1}{2}$. Hence the chances, after drawing the white counter, that the bag, before drawing, contained (a) 2 white, (b) 1 white, 1 black, are proportional to (a) $\frac{1}{2} \cdot 1$ (b) $\frac{1}{2} \cdot \frac{1}{2}$; i. e. (a) $\frac{1}{2}$ (b) $\frac{1}{4}$; i. e. (a) 2 (b) 1. Hence the chances are (a) $\frac{2}{3}$ (b) $\frac{1}{3}$. Hence, after the removal of a white counter, the chances, that the bag now contains (a) 1 white (b) 1 black, are for (a) $\frac{2}{3}$ and for (b) $\frac{1}{3}$.

Thus the chance, of now drawing a white counter, is $\frac{2}{3}$.

Q. E. F.

6. (2, 19)

Call sides ' $2a, 2b, 2c$ ', and lines in question ' α, β, γ '.



Now $\cos ADB + \cos ADC = 0$;

$$\therefore \frac{\alpha^2 + a^2 - 4c^2}{2\alpha a} + \frac{\alpha^2 + a^2 - 4b^2}{2\alpha a} = 0;$$

$$\therefore 2\alpha^2 + 2a^2 - 4b^2 - 4c^2 = 0;$$

$$\therefore \alpha^2 = -a^2 + 2b^2 + 2c^2.$$

Similarly, $\beta^2 = 2a^2 - b^2 + 2c^2$;

$$\gamma^2 = 2a^2 + 2b^2 - c^2.$$

To eliminate b, c , let us multiply by k, l, m , so taken that

$$2k - l + 2m = 0,$$

and $2k + 2l - m = 0$;

$$\therefore 3(l - m) = 0; \text{ i. e. } l = m;$$

$$\therefore 2k = -l = -m;$$

hence we may make $k = -1, \quad l = 2, \quad m = 2$;

$$\therefore -\alpha^2 + 2\beta^2 + 2\gamma^2 = 9a^2;$$

$$\text{i. e. } a^2 = \frac{-\alpha^2 + 2\beta^2 + 2\gamma^2}{9};$$

$$\therefore BC \text{ (which} = 2a) = \frac{2}{3} \sqrt{-\alpha^2 + 2\beta^2 + 2\gamma^2}, \text{ \&c.,}$$

which gives lengths of sides.

$$\begin{aligned}
\text{Also } \cos A &= \frac{b^2 + c^2 - a^2}{2bc} \\
&= \frac{2\alpha^2 - \beta^2 + 2\gamma^2 + 2\alpha^2 + 2\beta^2 - \gamma^2 + \alpha^2 - 2\beta^2 - 2\gamma^2}{2 \cdot \sqrt{2\alpha^2 - \beta^2 + 2\gamma^2} \cdot \sqrt{2\alpha^2 + 2\beta^2 - \gamma^2}} \\
&= \frac{5\alpha^2 - \beta^2 - \gamma^2}{\text{den.}}; \quad \text{and so for other angles.}
\end{aligned}$$

Q. E. F.

7. (2, 19)

Let AB , AD be given sides, and B , D the right $\angle$ s; and let $AB = b$, $AD = d$.



Produce DC to meet AB -produced at E .

Now $AE = AD \cdot \sec A = d \sec A$;

$\therefore BE = d \sec A - b$.

Also $BC = BE \cdot \tan E = (d \sec A - b) \cdot \cot A$,
 $= \frac{d - b \cos A}{\sin A}$;

similarly, $CD = \frac{b - d \cos A}{\sin A}$; which answers (1).



$$\begin{aligned}
\text{Also area} &= \frac{1}{2} \cdot (AB \cdot BC + AD \cdot DC), \\
&= \frac{1}{2} \cdot \frac{b \cdot (d - b \cos A) + d \cdot (b - d \cos A)}{\sin A}, \\
&= \frac{2bd - (b^2 + d^2) \cos A}{2 \sin A}; \quad \text{which answers (2).}
\end{aligned}$$

Q. E. F.

8. (2, 20)

Let m = No. of men, k = No. of shillings possessed by the last (i. e. the poorest) man. After one circuit, each is a shilling poorer, and the moving heap contains m shillings. Hence, after k circuits, each is k shillings poorer, the last man now having nothing, and the moving heap contains mk shillings. Hence the thing ends when the last man is again called on to hand on the heap, which then contains $(mk + m - 1)$ shillings, the penultimate man now having nothing, and the first man having $(m - 2)$ shillings.

It is evident that the first and last man are the only 2 neighbours whose possessions can be in the ratio '4 to 1'. Hence either

$$\begin{aligned}
mk + m - 1 &= 4(m - 2), \\
\text{or else } 4(mk + m - 1) &= m - 2.
\end{aligned}$$


The first equation gives $mk = 3m - 7$, i. e. $k = 3 - \frac{7}{m}$, which evidently gives no integral values other than $m = 7, k = 2$.

The second gives $4mk = 2 - 3m$, which evidently gives no positive integral values.

Hence the answer is '7 men; 2 shillings'.

9. (2)

Let AB, AC , be the given Lines, and P the given Point; and join AP .



Through A draw EAF , $\perp AP$, and bisected at A ; from E, F , draw EG, FH , parallel to AP , and meeting AB, AC , at G, H ; join GH , and on it describe a semicircle cutting AP at K ; and join KG, KH . Then $\angle GKH$ is a right angle. From P draw PL, PM , parallel to KG, KH .

Now Triangle APL has, to Triangle AKG , the duplicate ratio of AP to AK ;

but so also has triangle APM to Triangle AKH ;

also Triangles AKG, AKH , are equal, being on the same base AK , and having equal altitudes AE, AF ;

$\therefore$ Triangles APL, APM are equal: and $\angle LPM$ is evidently equal to $\angle GKH$; $\therefore$ it is a right angle.

Q. E. F.

10. (3, 20)

Call them x, y, z ; and let $x + y + z = s$.

The chance, that the pocket contains 2 balls, is $\frac{2}{3}$; and, if it does, the 'expectation' is the average value of

$$(y + z), (z + x), (x + y); \text{ i.e. it is } \frac{2s}{3}.$$

Also the chance, that it contains only one, is $\frac{1}{3}$; and, if it does, the 'expectation' is $\frac{s}{3}$.


$$\text{Hence total 'expectation'} = \frac{4s}{9} + \frac{s}{9} = \frac{5s}{9}.$$

$$\therefore \frac{5s}{9} = 30d.; \quad \therefore s = 54d. = 4/6.$$

Hence the coins must be 2 florins and a sixpence; or else a half-crown and 2 shillings.

Q. E. F.

11. (3, 20)



$$\text{Now } \frac{BA'}{A'C'} = \frac{\sin(B + \theta)}{\sin B}; \quad \text{and } \frac{A'C}{A'B'} = \frac{\sin \theta}{\sin C}.$$

$$\therefore BA = \frac{\sin(B + \theta)}{\sin B} \cdot ka; \quad \text{and } A'C = \frac{\sin \theta}{\sin C} \cdot kb$$

$$\text{but } BA' + A'C = a; \quad \therefore k$$

$$= \frac{a}{\frac{a \cdot \sin(B + \theta)}{\sin B} + \frac{b \sin \theta}{\sin C}} = \frac{\sin A}{\frac{\sin A \sin(B + \theta)}{\sin B} + \frac{\sin B \sin \theta}{\sin C}}$$

$$\begin{aligned}
&= \frac{\sin A \sin B \sin C}{\sin A \sin(B + \theta) \sin C + \sin^2 B \sin \theta} \\
&= \frac{\sin A \sin B \sin C}{\sin A \sin C (\sin B \cos \theta + \cos B \sin \theta) + (1 - \cos^2 B) \sin \theta} \\
&= \frac{\sin A \sin B \sin C}{\left. \begin{array}{l} \sin \theta + \sin \theta (\sin A \sin C \cos B - \cos^2 B) \\ + \cos \theta \sin A \sin B \sin C \end{array} \right\}} \\
&= \frac{\sin A \sin B \sin C}{\left. \begin{array}{l} \sin \theta + \sin \theta \cos B (\sin A \sin C + \cos(A+C)) \\ + \cos \theta \sin A \sin B \sin C \end{array} \right\}} \\
&= \frac{\sin A \sin B \sin C}{\sin \theta (1 + \cos A \cos B \cos C) + \cos \theta \sin A \sin B \sin C}.
\end{aligned}$$

Q. E. F.

COR. Let $\theta = 90^\circ$; then $k = \frac{\sin A \sin B \sin C}{1 + \cos A \cos B \cos C}$.

12. (4, 20)


Let s = semi-perimeter, m = area, v = volume.

We know that $m = \sqrt{s.(s-a).(s-b).(s-c)}$;

$$\begin{aligned}
\therefore m^2 &= s.(s-a).(s-b).(s-c); \\
\therefore \frac{m^2}{s} &= s^3 - s^2.(a+b+c) + s.(bc+ca+ab) - abc; \\
&= s^3 - 2s^3 + s.(bc+ca+ab) - v; \\
\therefore \frac{m^2}{s^2} + \frac{v}{s} + s^2 &= bc+ca+ab; \\
\therefore 2. \left(\frac{m^2}{s^2} + \frac{v}{s} + s^2 \right) &= (a+b+c)^2 - (a^2+b^2+c^2); \\
&= 4s^2 - (a^2+b^2+c^2); \\
\therefore a^2+b^2+c^2 &= 2.(s^2 - \frac{v}{s} - \frac{m^2}{s^2}).
\end{aligned}$$

Q. E. F.

13. (4, 20)



Let A, B , be the centres of the Circles; C, D , their points of intersection; and $CFDE$ the Tetragon whose area is required.

Let the sides of the Triangle ABC be a, b, c ; and its $\angle$ s α, β, γ .

Then $CE = b \cdot \tan \alpha$, and $CF = a \cdot \tan \beta$.

Also $\angle FCE = \angle ACE + \angle FCB - \gamma = \pi - \gamma$;

$\therefore \sin FCE = \sin \gamma$.

Hence area of Triangle $FCE = \frac{1}{2} \cdot ab \cdot \tan \alpha \cdot \tan \beta \cdot \sin \gamma$;

$\therefore$ area of Tetragon $= \frac{ab \sin \alpha \sin \beta \sin \gamma}{\cos \alpha \cos \beta}$.

Now, writing ' M ' for area of Triangle ABC , we have

$$\sin \alpha = \frac{2M}{bc}, \quad \sin \beta = \frac{2M}{ca}, \quad \sin \gamma = \frac{2M}{ab};$$

$$\begin{aligned} \therefore \text{area of Tetragon} &= ab \cdot \frac{8M^3}{a^2 b^2 c^2} \cdot \frac{4bc \cdot ca}{(b^2 + c^2 - a^2)(c^2 + a^2 - b^2)}; \\ &= \frac{32M^3}{(b^2 + c^2 - a^2) \cdot (c^2 + a^2 - b^2)}. \end{aligned}$$

Q. E. F.

14. (4)

This simply expresses the identity


$$\begin{aligned}
& 3(a^2 + b^2 + c^2) \\
&= (a + b + c)^2 + (b^2 - 2bc + c^2) + (c^2 - 2ca + a^2) + (a^2 - 2ab + b^2); \\
&= (a + b + c)^2 + (b - c)^2 + (c - a)^2 + (a - b)^2.
\end{aligned}$$

Q. E. D.

Numerical Examples (not thought out).

$$\begin{aligned}
3(1^2 + 2^2 + 3^2) &= 6^2 + 1^2 + 2^2 + 1^2. \\
3(1^2 + 3^2 + 7^2) &= 11^2 + 4^2 + 6^2 + 2^2.
\end{aligned}$$

15. (4)



Let $ABCD$ be an inscribed Tetragon. Join AC : and about Triangle ACD describe a Circle.

Now, if this Circle does not pass through B , let it cut CB , or CB produced, in B' or B'' . Join AB' , AB'' .

Then $\angle AB'C$, or $\angle AB''C$, is supplementary to $\angle ADC$;

$\therefore$ it $= \angle ABC$; which is absurd;

$\therefore$ this Circle does pass through B .

The same thing may be proved for any other Point on that portion, of the perimeter of the given Figure, which lies on the same side of AC as the Point D .

Similarly for the other portion.

Hence the Figure is a Circle.

Q. E. D.

16. (4, 20)

The 'a priori' chances of possible states of first bag are ' $W, \frac{1}{2}; B, \frac{1}{2}$ '. Hence chances, after putting W in, are ' $WW, \frac{1}{2}; WB, \frac{1}{2}$ '. The chances, which these give to the 'observed event', are $1, \frac{1}{2}$. Hence chances of possible states ' W, B ', after the event, are proportional to $1, \frac{1}{2}$; i. e. to $2, 1$; i. e. their actual values are $\frac{2}{3}, \frac{1}{3}$.

Now, in first course, chance of drawing W is $\frac{1}{2} \cdot \frac{2}{3} + \frac{1}{2} \cdot \frac{1}{3}$; i. e. $\frac{1}{2}$.

And, in second course, chances of possible states ' $WWBB, WBBB$ ' are $\frac{2}{3}, \frac{1}{3}$; hence chance of drawing W is $\frac{2}{3} \cdot \frac{1}{2} + \frac{1}{3} \cdot \frac{1}{4}$; i. e. $\frac{5}{12}$.


Hence *first* course gives best chance.

Q. E. F.

17. (4)

(Analysis.)

Let ABC be the given Triangle, and DE the line required.



From D, E , draw DF, EG , parallel to the sides. Then $DF + EG = DE$.

Because BE is a Parallelogram, $\therefore DB = EG$;

similarly $EC = DF$;

$\therefore DB + EC = DE$.

Hence construction.

(Synthesis.)

Bisect $\angle s B, C$, by BH, CH , meeting at H ; through H draw DE parallel to BC ; and from D, E , draw DF, EG , parallel to AC, AB .

Because DE is parallel to BC ,

$$\therefore \angle DHB = \text{alternate } \angle HBF = \angle DBH;$$

$$\therefore DB = DH.$$

Similarly $EC = EH$.

$$\therefore DB + EC = DE.$$


Because BE, DC are Parallelograms,

$$\therefore EG = DB, \text{ and } DF = EC;$$

$$\therefore DF + EG = DE.$$

Q. E. F.

18. (5, 21)



(1) Call required Point E . From E draw $EF, EG \perp$ sides. Join FG . From F, G , draw $FH, GK \perp BC$. Call BE ' x ', and EC ' y '.

Now FH must $= GK$.

$$\begin{aligned} \text{Also } EF &= x \sin B; \text{ and } FH &= EF \sin FEH, \\ & &= EF \cos B, \end{aligned}$$


$$= x \sin B \cos B;$$

$$\text{similarly, } GK = y \sin C \cos C.$$

But $FH = GK$; $\therefore x \sin B \cos B = y \sin C \cos C$;

$$\therefore \frac{x}{y} = \frac{\sin 2C}{\sin 2B}.$$

Q. E. F.



(2) At B, C , make right angles ABD, ACD ; and join AD , cutting BC at E . From E draw $EF, EG \perp$ sides; and join FG .

$\therefore BD, FE$ are $\perp AB$, $\therefore$ they are $\parallel$; $\therefore AF : FB :: AE : ED$;

$\therefore CD, GE$ are $\perp AC$, $\therefore$ they are $\parallel$; $\therefore AG : GC :: AE : ED$;

$\therefore AF : FB :: AG : GC$;

$\therefore FG$ is parallel to BC .

Q. E. F.

19. (5, 21)

Call the bags A, B, C ; so that A contains a white counter and a black one; &c.

The chances of the orders $ABC, ACB, BAC, BCA, CAB, CBA$, are, a priori, $\frac{1}{6}$ each. Since they are equal, we may, instead of multiplying each by the probability it gives to the observed event, simply assume those probabilities as being proportional to the chances *after* the observed event.

These probabilities are:—

$$\begin{array}{llll}
 \text{for } ABC, & \frac{1}{2} \times \frac{1}{3}; & \text{i. e.} & \frac{1}{6}. \\
 ACB, & \frac{1}{2} \times \frac{1}{4}; & \text{i. e.} & \frac{1}{8}. \\
 BAC, & \frac{2}{3} \times \frac{1}{2}; & \text{i. e.} & \frac{1}{3}. \\
 BCA, & \frac{2}{3} \times \frac{1}{4}; & \text{i. e.} & \frac{1}{6}. \\
 CAB, & \frac{3}{4} \times \frac{1}{2}; & \text{i. e.} & \frac{3}{8}. \\
 CBA, & \frac{3}{4} \times \frac{1}{3}; & \text{i. e.} & \frac{1}{4}.
 \end{array}$$

Hence the chances are proportional to 4, 3, 8, 4, 9, 6; i. e. they are these Nos. divided by 34.

Hence the chance, of drawing a white counter from the remaining bag; is

$$\begin{aligned}
 & \frac{1}{34} \cdot \left\{ 4 \times \frac{3}{4} + 3 \times \frac{2}{3} + 8 \times \frac{3}{4} + 4 \times \frac{1}{2} + 9 \times \frac{2}{3} + 6 \times \frac{1}{2} \right\}; \\
 \text{i. e. } & \frac{1}{34} \times \{ 3 + 2 + 6 + 2 + 6 + 3 \}; \quad \text{i. e. } \frac{22}{34}; \quad \text{i. e. } \frac{11}{17}.
 \end{aligned}$$

20. (5)

(Analysis.)

Let ABC be the given Triangle, and P the required Point. Draw $PQ \perp BC$, and $PR \perp AB$. Then $PQ = PR$.

Hence $PC \tan C = PB \sin B$;


$\therefore PC : PB :: \sin B : \tan C$, (draw $AD \perp BC$.)

$$\begin{aligned}
 & \therefore \frac{AD}{AB} : \frac{AD}{DC}, \\
 & \therefore DC : AB.
 \end{aligned}$$

Hence construction.

(Synthesis.)

From A draw $AD \perp BC$. Produce BA to E , making AE equal to DC . Join EC . From A draw AP parallel to EC ; and from P draw $PQ \perp BC$, and $PR \perp AB$.



$$\begin{aligned}
\text{Then } \frac{PQ}{PC} &= \frac{AD}{DC} = \frac{AD}{AB} \cdot \frac{AB}{DC}, \\
&= \frac{PR}{PB} \cdot \frac{AB}{AE}, \\
&= \frac{PR}{PB} \cdot \frac{PB}{PC} = \frac{PR}{PC}; \\
\therefore PQ &= PR.
\end{aligned}$$

Q. E. F.

21. (5, 21)

$$\begin{aligned}
(1) \text{ The } n\text{th term is } \overline{n.n+2.n+4}; \\
\therefore \text{ the } (n+1)\text{th term is } \overline{n+1.n+3.n+5}; \\
= (n+1).(\overline{n+2+1}).(n+5); \\
= \overline{n+1.n+2.n+5} + \overline{n+1.n+5} \\
= \overline{n+1.n+2.(\overline{n+3+2})} + \overline{n+1.(\overline{n+2+3})}; \\
= \overline{n+1.n+2.n+3} + 2.\overline{n+1.n+2} + \overline{n+1.n+2} + 3.\overline{n+1}; \\
= \overline{n+1.n+2.n+3} + 3.\overline{n+1.n+2} + 3.\overline{n+1}. \\
\therefore S = \frac{\overline{n.n+1.n+2.n+3}}{4} + n.\overline{n+1.n+2} + \frac{3}{2}.n.\overline{n+1} + C; \\
\text{and } C = 0. \\
\therefore S = n.\overline{n+1} \cdot \left(\frac{n^2 + 5n + 6}{4} + n + 2 + \frac{3}{2} \right); \\
= n.\overline{n+1} \cdot \frac{n^2 + 9n + 20}{4} = \frac{\overline{n.n+1.n+4.n+5}}{4}.
\end{aligned}$$

Q. E. F.

$$\begin{aligned}
(2) S, \text{ to 100 terms,} \\
= \frac{100.101.104.105}{4} = 100.101.26.105; \\
\text{now } 101.105 = 10,605;
\end{aligned}$$


$$\therefore 101.105.13 = 130,000 + 7800 + 65 = 137,865;$$

$$\text{and twice this} = 274,000 + 1730 = 275,730;$$

$$\therefore S = 27,573,000.$$

Q. E. F.

22. (5, 21)



Call given altitudes ' α, β, γ '.

$$\text{Now } a\alpha = b\beta = c\gamma;$$

$$\therefore \alpha \sin A = \beta \sin B = \gamma \sin C;$$

$$\therefore \frac{\sin A}{\beta\gamma} = \frac{\sin B}{\gamma\alpha} = \frac{\sin C}{\alpha\beta} = k \text{ (say);}$$

$$\therefore \sin A = k\beta\gamma, \quad \sin B = k\gamma\alpha, \quad \sin C = k\alpha\beta.$$

$$\text{Now } \sin(A + B) = \sin C;$$

$$\therefore \sin A \cos B + \cos A \sin B = \sin C;$$

$$\therefore \sin A \cos B = \sin C - \cos A \sin B;$$

$$\therefore \sin^2 A(1 - \sin^2 B) = \sin^2 C + \sin^2 B(1 - \sin^2 A) - 2 \sin C \cos A \sin B$$


$$\therefore \sin^2 A - \sin^2 A \sin^2 B = \sin^2 C + \sin^2 B - \sin^2 A \sin^2 B - 2 \sin B \sin C \cos A;$$

$$\therefore \sin^2 A - \sin^2 B - \sin^2 C = -2 \sin B \sin C \cos A;$$

$$\therefore, \text{squaring, } (\sin^2 A + \sin^2 B + \sin^2 C) - 2 \sin^2 A \sin^2 B - 2 \sin^2 A \sin^2 C + 2 \sin^2 B \sin^2 C = 4 \sin^2 B \sin^2 C (1 - \sin^2 A)$$

$$\therefore (\sin^4 A + \&c.) - 2(\sin^2 B \sin^2 C + \&c.) + 4 \sin^2 A \sin^2 B \sin^2 C = 0;$$

$$\therefore, \text{ substituting for } \sin A, \&c., \text{ and dividing by } k^4, \\ (\beta^4 \gamma^4 + \&c.) - 2\alpha^2 \beta^2 \gamma^2 . (\alpha^2 + \&c.) + 4k^2 \alpha^4 \beta^4 \gamma^4 = 0;$$



$$\therefore k^2 = \frac{2\alpha^2 \beta^2 \gamma^2 (\alpha^2 + \&c.) - (\beta^4 \gamma^4 + \&c.)}{4\alpha^4 \beta^4 \gamma^4}.$$

Now $\sin A = k\beta\gamma$, &c.; which answers (2).

Also $\alpha = b \sin C$; and similarly $\gamma = a \sin B$;

$$\therefore a = \frac{\gamma}{\sin B} = \frac{\gamma}{k\gamma\alpha} = \frac{1}{k\alpha}, \text{ \texttt{\&c.}}; \text{ which answers (1).}$$

$$\text{Also area} = \frac{bc \sin A}{2} = \frac{1}{2} \cdot \frac{1}{k\beta} \cdot \frac{1}{k\gamma} \cdot k\beta\gamma = \frac{1}{2k};$$

which answers (3).

Q.E.F.

23. (5, 21)

The original chances, as to states of bag, are

$$\begin{array}{ll} \text{for } 2W & \frac{1}{4}; \\ & 1W, 1B & \frac{1}{2}; \\ & 2B & \frac{1}{4}. \end{array}$$

∴ the chances, after adding 2 W and 1 B, are

$$\begin{array}{ll} \text{for } 4W, 1B & \frac{1}{4}; \\ 3W, 2B & \frac{1}{2}; \\ 2W, 3B & \frac{1}{4}. \end{array}$$

Now the chances, which these give to the observed event, drawing 2 W and 1 B, are $\frac{3}{5}, \frac{3}{5}, \frac{3}{10}$.

∴ the chances, after this event, are proportional to $\frac{3}{20}, \frac{3}{10}, \frac{3}{40}$; i. e. to 2, 4, 1. Hence they are $\frac{2}{7}, \frac{4}{7}, \frac{1}{7}$.

Hence the chances, as to states, now are

$$\begin{array}{ll} \text{for } 2W & \frac{2}{7}; \\ 1W, 1B & \frac{4}{7}; \\ 2B & \frac{1}{7}. \end{array}$$

∴ the chances, after adding 1W, are

$$\begin{array}{ll} \text{for } 3W & \frac{2}{7}; \\ 2W, 1B & \frac{4}{7}; \\ 1W, 2B & \frac{1}{7}. \end{array}$$


Now the chances, which these give to the observed event, of drawing 1 W, are 1, $\frac{2}{3}, \frac{1}{3}$.

∴ the chances, after this event, are proportional to $\frac{2}{7}, \frac{8}{21}, \frac{1}{21}$;

i. e. to 6, 8, 1. Hence they are $\frac{6}{15}, \frac{8}{15}, \frac{1}{15}$.

Hence the chance, that the bag now contains 2 white, is $\frac{6}{15}$; i. e. $\frac{2}{5}$.

Q. E. F.



Because $\frac{DO}{OA} = \frac{\triangle DOC}{\triangle OAC} = \frac{\triangle DOB}{\triangle OAB} = \frac{\triangle OBC}{\triangle OCA + \triangle OAB};$

$$\therefore \frac{DO}{DA} = \frac{\triangle OBC}{\triangle ABC}.$$

Similarly, $\frac{EO}{EB} = \frac{\triangle OCA}{\triangle ABC},$ and $\frac{FO}{FC} = \frac{\triangle OAB}{\triangle ABC}.$

Hence $\frac{DO}{DA} + \frac{EO}{EB} + \frac{FO}{FC} = 1.$

Q. E. F.

25. (6, 22)

Let '*E*' mean 'having lost an eye', '*A*' 'having lost an arm', and '*L*' 'having lost a leg'.



Then the state of things which gives the least possible number of those who, being *E* and *A*, are also *L*, may evidently be found by arranging the patients in a row, so that the *EA*-class may begin from one end of the row, and the *L*-class from the other end, and counting the portion where they overlap; and, the smaller the *EA*-class, the smaller will be this common portion: hence we must make the *EA*-class a minimum.

This may be done by re-arranging the patients, so that the *E*-class may begin from one end of the row, and the *A*-class from the other: and the least

$$\frac{(\epsilon + \alpha - 1)}{(1 - \lambda) \quad \lambda}$$

possible number for the EA -class is the common portion, i. e. $(\epsilon - 1 - \alpha)$, i. e. $(\epsilon + \alpha - 1)$.

Then, as already shown, the least possible number for the EAL -class is the common portion, i. e. $(\epsilon + \alpha - 1 - 1 - \lambda)$, i. e. $(\epsilon + \alpha + \lambda - 2)$.

Q. E. F.


26. (6)

(Analysis.)

Let ABC be the given Triangle, and $A'B'C'$ the required one; and let the ratio, which $B'C'$ has to BC , be ' k '; so that k is less than 1.

Since $BB' = CC'$, and that $BC, B'C'$, are parallel, it may easily be proved, by dropping perpendiculars from B', C' , upon BC , which must necessarily be equal, that $\angle s B'BC, C'CB$, are equal.

Similarly, $\angle s A'AC, C'CA$, are equal; and so are $\angle s A'AB, B'BA$.



Call $\angle B'BC$ ' θ '; then $\angle C'CB = \theta$;

$\therefore \angle C'CA = C - \theta = \angle A'AC$;

$\therefore \angle A'AB = A - (C - \theta) = \angle B'BA$.

Now $\angle s B'BC, B'BA$, together = B ;

$\therefore \theta + A - (C - \theta) = B$;

$\therefore 2\theta = B + C - A = 180^\circ - 2A$;

$$\therefore \theta = 90^\circ - A.$$

Hence, if BB' , CC' , be produced to meet at D , Triangle DBC will be isosceles, with a vertical $\angle$ equal to $2A$.

Now, if a Circle be drawn about ABC , and its centre joined to B and C , the Triangle, so formed, will fulfil the same conditions;

hence the centre of this Circle will be D ;

hence the construction.

(*Synthesis.*)

Bisect the sides, and draw perpendiculars, meeting at D . Join D to the vertices B , C . From DB cut off $DB' = k.DB$. From B' draw $B'C'$ parallel to BC .

Then $B'C'$ is easily proved equal to $k.BC$.

And if, from B' , C' , parallels to AB , AC , be drawn, it may easily be proved that they meet on DA , and that they are respectively equal to $k.AB$, $k.AC$.

Q. E. F.

27. (6, 22)

Call the bags A , B , C .

If remaining bag be A , chance of observed event = $\frac{1}{2}$ chance of drawing white from B and black from C + $\frac{1}{2}$ chance of drawing black from B and white from C :

$$\text{i. e. it} = \frac{1}{2} \cdot \left\{ \frac{2}{3} \times \frac{1}{2} + \frac{1}{3} \times \frac{1}{2} \right\} = \frac{1}{4}.$$

Similarly, if remaining bag be B , it is $\frac{1}{2} \cdot \left\{ \frac{5}{6} \cdot \frac{1}{2} + \frac{1}{6} \cdot \frac{1}{2} \right\} = \frac{1}{4}$; and, if it be C , it is $\frac{1}{2} \cdot \left\{ \frac{5}{6} \cdot \frac{1}{3} + \frac{1}{6} \cdot \frac{2}{3} \right\} = \frac{7}{36}$.


$\therefore$ chances of remaining bag being A , B , or C , are as $\frac{1}{4}$ to $\frac{1}{4}$ to $\frac{7}{36}$; i. e. as 9 to 9 to 7. $\therefore$ they are, in value, $\frac{9, 9, 7}{25}$.

Now, if remaining bag be A , chance of drawing white from it is $\frac{5}{6}$; $\therefore$ chance, on this issue, is $\frac{5}{6} \cdot \frac{9}{25} = \frac{3}{10}$; similarly, for B , it is $\frac{2}{3} \cdot \frac{9}{25} = \frac{6}{25}$; and, for

C , $\frac{1}{2} \cdot \frac{7}{25} = \frac{7}{50}$. And entire chance of drawing white from the remaining bag is the sum of these; i. e. $\frac{15+12+7}{50} = \frac{34}{50} = \frac{17}{25}$.

28. (7, 22)

Let ABC be the given Triangle; and let its sides be divided internally at A' , B' , C' , in extreme and mean ratio.



And let M be the area of ABC .

Let $BA' = x$; then $x^2 = a.(a - x)$;

i. e. $x^2 + ax - a^2 = 0$;

$\therefore x = \frac{-a \pm a\sqrt{5}}{2} = \frac{a}{2}(\sqrt{5} - 1)$, the other sign being excluded by the terms of the question.

Then area of Triangle $AB'C'$

$$\begin{aligned} &= \frac{1}{2} \cdot \frac{c}{2} \cdot (\sqrt{5} - 1) \cdot \left\{ b - \frac{b}{2} \cdot (\sqrt{5} - 1) \right\} \cdot \sin A, \\ &= \frac{1}{8} \cdot (\sqrt{5} - 1)(3 - \sqrt{5})bc \cdot \sin A, \\ &= \frac{1}{4} \cdot (4\sqrt{5} - 8) \cdot M = (\sqrt{5} - 2) \cdot M. \end{aligned}$$

Similarly for $BC'A'$ and $CA'B'$.

Hence the sum of these 3 Triangles $= 3 \cdot (\sqrt{5} - 2) \cdot M$, and area of Triangle $A'B'C' = (7 - 3\sqrt{5}) \cdot M$.

Q. E. F.

29. (7)

This may be deduced from the identity

$$(a^2 + b^2).(c^2 + d^2) = a^2c^2 + b^2d^2 + a^2d^2 + b^2c^2.$$

$$\begin{aligned} (a^2 + b^2).(c^2 + d^2) &= a^2c^2 + b^2d^2 + a^2d^2 + b^2c^2; \\ \text{or else} &= a^2c^2 + b^2d^2 + 2acbd + a^2d^2 + b^2c^2 - 2adbc, \\ &= a^2c^2 + b^2d^2 - 2acbd + a^2d^2 + b^2c^2 + 2adbc; \\ \text{i. e.} &= (ac + bd)^2 + (ad - bc)^2, \\ \text{or else} &= (ac - bd)^2 + (ad + bc)^2. \end{aligned} \quad \left. \vphantom{\begin{aligned} (a^2 + b^2).(c^2 + d^2) &= a^2c^2 + b^2d^2 + a^2d^2 + b^2c^2; \\ \text{or else} &= a^2c^2 + b^2d^2 + 2acbd + a^2d^2 + b^2c^2 - 2adbc, \\ &= a^2c^2 + b^2d^2 - 2acbd + a^2d^2 + b^2c^2 + 2adbc; \\ \text{i. e.} &= (ac + bd)^2 + (ad - bc)^2, \\ \text{or else} &= (ac - bd)^2 + (ad + bc)^2. \end{aligned}} \right\}$$

Now, if these last 2 sets are *identical*, $(ac + bd)$ must = $(ad + bc)$; for it cannot = $(ac - bd)$;

i. e., $a(c - d) - b(c - d)$ must = 0;


i. e., $(a - b).(c - d)$ must = 0;

i. e., one or other of the first 2 sets is the sum of 2 *identical* squares.

Hence, contranominally, if *each* of the original sets consists of 2 *different* squares, their product gives the sum of 2 squares in 2 *different* ways.

Q. E. D.

30. (7)



(Analysis.)

Let ABC be the Triangle: and suppose $B'C'$ so placed that $B'D$, $C'E$, drawn parallel to the sides, shall together $= 2B'C'$.

By Euc. I. 34, $B'D = C'B$, and $C'E = B'C$:

$$\therefore B'C + C'B = 2B'C'.$$

Hence, if $B'L$ be cut off equal to half $B'C$, $C'L =$ half $C'B$.

Hence construction.

(Synthesis.)

In BC' take any point F : draw FG , $\parallel BC$, and equal to half BF : and join BG .

Similarly, in CB' take any point H : draw HK , $\parallel BC$, and equal to half HC : and join CK .

Produce BG , CK , to meet at L : and through L draw $C'B' \parallel BC$: and from B' , C' , draw $B'D$, $C'E$, $\parallel$ the sides.

$\therefore FG = \text{half } FB$; $\therefore$, by similar Triangles, $C'L = \text{half } C'B$;

Similarly $B'L = \text{half } B'C$;

$\therefore C'B' = \text{half sum of } C'B, B'C$; i. e. $C'B + B'C = 2B'C'$.

But, by Euc. I. 34, $C'B = B'D$, and $B'C = C'E$;

$\therefore B'D + C'E = 2B'C'$.

Q. E. F.

31. (7, 22)

On July 1, watch gained on clock 5*m.* in 10*h.*; i. e. $\frac{1}{2}$ *m.* per hour; i. e. 2*m.* in 4*h.* Hence, when watch said 'noon', clock said '12*h.* 2*m.*'; i. e. clock was 3*m.* slow of true time, when *true* time was 12*h.* 5*m.*

On July 30, watch lost on clock 1*m.* in 10*h.*; i. e. 6*sec.* per hour; i. e. 19*sec.* in 3*h.* 10*m.* Hence, when watch said '12*h.* 10*m.*', clock said '12*h.* 7*m.* 19*sec.*'; i. e. clock was 2*m.* 19*sec.* fast of *true* time, when true time was 12*h.* 5*m.*

Hence clock gains, on *true* time, 5*m.* 19*sec.* in 29 days; i. e. 319*sec.* in 29 days; i. e. 11*sec.* per day; i. e. $\frac{11}{24 \times 12}$ *sec.* in 5*m.*

Hence, while *true* time goes 5*m.*, watch goes 5*m.* $\frac{11}{288}$ *sec.*

Now, when *true* time is 12*h.* 5*m.* on July 31, clock is (2*m.* 19*sec.* + 11*sec.*) fast of it; i. e. says '12*h.* 7 $\frac{1}{2}$ *m.*' Hence, if *true* time be put 5*m.* back, clock must be put 5*m.* $\frac{11}{288}$ *sec.* back; i. e. must be put back to 12*h.* 2*m.* 29 $\frac{277}{288}$ *sec.*

Hence, on July 31, when clock indicates this time, it is *true* noon.

Q. E. F.

32. (8, 22)

The *n*th term is $n.(n + 4)$;

$$\begin{aligned}\therefore \text{the } (n+1)\text{th term is } (n+1).(n+5) &= (n+1).\{(n+2)+3\}, \\ &= (n+1).(n+2)+3(n+1);\end{aligned}$$

$$\therefore S_n = \frac{n.(n+1).(n+2)}{3} + 3 \cdot \frac{n.(n+1)}{2} + C; \text{ and } C = 0;$$


$$\therefore S_n = n.(n+1). \left(\frac{n+2}{3} + \frac{3}{2} \right) = \frac{n.(n+1).(2n+13)}{6}.$$

Q. E. F.

$$\begin{aligned}\text{Also} \\ S_{100} &= \frac{100.101.213}{6} = \frac{100.101.71}{2} = \frac{100.7171}{2} = \frac{717100}{2} = 358550.\end{aligned}$$

Q. E. F.

33. (8)



$$\text{Let } DE = x; \therefore BC = 2x.$$

$$\text{Area} = 3x.(\sqrt{r^2 - x^2} + \sqrt{r^2 - 4x^2}) = \text{max.}$$

$$\text{let } v = x.(\sqrt{r^2 - x^2} + \sqrt{r^2 - 4x^2}) = \text{max.}$$

$$\begin{aligned}
\therefore \frac{dv}{dx} &= \sqrt{r^2 - x^2} + \sqrt{r^2 - 4x^2} - x^2 \cdot \left(\frac{1}{\sqrt{r^2 - x^2}} + \frac{4}{\sqrt{r^2 - 4x^2}} \right) = 0; \\
\therefore (r^2 - x^2) \cdot \sqrt{r^2 - 4x^2} + (r^2 - 4x^2) \cdot \sqrt{r^2 - x^2} &= x^2 \cdot (4\sqrt{r^2 - x^2} + \sqrt{r^2 - 4x^2}); \\
\therefore (r^2 - 2x^2) \cdot \sqrt{r^2 - 4x^2} &= -(r^2 - 8x^2) \cdot \sqrt{r^2 - x^2}; \\
\therefore r^4 - 4(r^2x^2 + 4x^4) \cdot (r^2 - 4x^2) &= (r^4 - 16r^2x^2 + 64x^4) \cdot (r^2 - x^2); \\
\therefore r^6 - 8r^4x^2 + 20r^2x^4 - 16x^6 &= r^6 - 17r^4x^2 + 80r^2x^4 - 64x^6;
\end{aligned}$$


$\therefore$, omitting r^6 , and dividing by x^2 ,

$$\begin{aligned}
48x^4 - 60r^2x^2 + 9r^4 &= 0; \\
\text{i. e. } 16x^4 - 20r^2x^2 + 3r^4 &= 0;
\end{aligned}$$

$$\therefore \frac{x^2}{r^2} = \frac{20 \pm \sqrt{208}}{32} = \frac{5 \pm \sqrt{13}}{8} \quad (\text{upper sign being inadmissible, though this was not thought out.})$$

Q. E. F.

34. (8)




(Analysis.)

Let A be the given Point, and C the centre of the given Circle. Join AC , and let ABD be the required Line. From B draw the Chord BE parallel to AC . Then $\angle DBE = \angle A$. Hence Arc $DE =$ Arc BD ; i. e. Arc BE is bisected by D ; i. e. D is on perpendicular from C .


(Synthesis.)

Join AC . From C draw CD perpendicular to AC . Join AD cutting Circle at B . From B draw BE parallel to AC .

It is easily proved that $\text{Arc } BD = \text{Arc } DE$. Hence $\text{Arc } BD$ subtends, in the Circle, an angle $= \angle DBE = \angle A$.



35. (8)



Let ABC be the given Triangle; and let the portions of the radii, outside the Triangle, have to the radius the given ratios $k : 1, l : 1, m : 1$. (N.B. k, l, m , are supposed to be proper fractions.)

From B draw $BD \perp BA$, and $BE \perp BC$; and make BD have, to BE , the ratio $1 - m : 1 - k$. Through D draw DF parallel to BA , and EF parallel to BC ; and join BF . From F draw $FG \perp BA$, and $FH \perp BC$.

Then $FG : FH :: 1 - m : 1 - k$.

Similarly, draw CO so that the $\perp$ s, drawn from any Point of it to CA and CB , are in the ratio $1 - l : 1 - k$; and produce BF to meet it at O .

From O draw OA', OB', OC' , $\perp$ the sides.

Then $OA' : OB' : OC' :: 1 - k : 1 - l : 1 - m$.

Produce OA' to K , so that $OK : OA' :: 1 : 1 - k$.

With centre O , and distance OK , describe a Circle; and produce OB' , OC' , to meet it at L , M .

$$\begin{aligned}\text{Now } OK : OA' &:: 1 : 1 - k; \\ OA' : OB' &:: 1 - k : 1 - l; \\ \therefore OK : OB' &:: 1 : 1 - l;\end{aligned}$$


Similarly, $OK : OC' :: 1 : 1 - m$.

But $A'K : OK :: OK - OA' : OK :: k : 1$.

Similarly $B'L : \text{radius} :: l : 1$, and $C'M : \text{radius} :: m : 1$.

Q. E. F.

36. (8)



(Analysis.)

Let $B'C'$ be required Line: and let $\angle$ at C' be right.

Cut off $C'D$ equal to $C'B$: then $DB' = B'C$.

Join DB , DC : then $\angle DBC' = 45^\circ$, and $\angle B'DC = \angle B'CD$.

From C draw $CE \perp AB$.

Then $\angle B'DC = \angle DCE$; $\therefore \angle B'CD = \angle DCE$.

(Synthesis.)

Hence construction. Draw $CE \perp AB$: bisect $\angle ACE$: at B make $\angle ABD = 45^\circ$. Let these lines meet at D . Through D draw $B'DC' \perp AB$.


Then $\angle C'DB = \pi - (\angle DC'B + \angle C'BD) = 45^\circ = \angle C'BD$;

$\therefore C'D = C'B$.

Also $\angle B'DC = \angle DCE = \angle DCB'$;

$\therefore DB' = B'C$;

$\therefore C'B' = \text{sum of } BC', CB'$.



Limits of possibility:—

$\angle A$ must not be $> 90^\circ$;


$\angle B$ must not be $< 45^\circ$;

$\angle C$ must not be $< \text{half complement of } A$,

i. e. not $< (45^\circ - \frac{A}{2})$.

37. (8, 22)

Let BC be the common chord, and A, D , the centres.



Let $\angle A = 30^\circ$, and $\angle D = 60^\circ$.

And let BC (which $= DB = DC$) $= 1$.

And let $AB = x$.

$$\text{Now } \cos A = \frac{\sqrt{3}}{2} = \frac{2x^2 - 1}{2x^2};$$

$$\therefore \frac{\sqrt{3}}{2} = 1 - \frac{1}{2x^2}; \therefore \frac{1}{2x^2} = \frac{2 - \sqrt{3}}{2};$$

$$\therefore x^2 = \frac{1}{2 - \sqrt{3}} = 2 + \sqrt{3};$$

$\therefore$ areas of Circles are $\pi \cdot (2 + \sqrt{3})$ and π ;

$\therefore$ areas of Sectors are $\pi \cdot \frac{2 + \sqrt{3}}{12}$ and $\frac{\pi}{6}$;

$\therefore$ their sum $= \pi \cdot \frac{4 + \sqrt{3}}{12}$.

Again, area of Triangle $ABC = \frac{1}{2} \cdot (2 + \sqrt{3}) \cdot \frac{1}{2}$,

$$= \frac{2 + \sqrt{3}}{4};$$

also area of Triangle $DBC = \frac{\sqrt{3}}{4}$;

$$\therefore \text{their sum} = \frac{2 + 2\sqrt{3}}{4} = \frac{1 + \sqrt{3}}{2}.$$

Now the portion, of the smaller Circle, that is within the larger one, is the difference between these two sums;

$$\therefore \text{it} = \pi \cdot \frac{4 + \sqrt{3}}{12} - \frac{1 + \sqrt{3}}{2}.$$

Hence its ratio, to the area of the smaller Circle, is this sum divided by π ;

$$\begin{aligned} \therefore \text{it} &= \frac{4 + \sqrt{3}}{12} - \frac{1 + \sqrt{3}}{2\pi}, \\ &= \frac{5 \cdot 732}{12} - \frac{2 \cdot 732}{\left(\frac{44}{7}\right)} = .478 - \frac{.248}{\left(\frac{4}{7}\right)}, \\ &= .478 - \frac{1 \cdot 736}{4} = .478 - .434 = .044. \end{aligned}$$

Q. E. F.

38. (9, 22)

Taking, in order, the bag from which this unknown counter is drawn, the bag from which a red one was twice drawn, and the remaining bag, we see that there are six possible arrangements of 'A', 'B', and 'C': viz.—

- | | |
|-------------|-------------|
| (1) ABC , | (4) BCA , |
| (2) ACB , | (5) CAB , |
| (3) BAC , | (6) CBA . |

Now the chance of the observed event is, in case (1), $1 \times \frac{4}{9} = \frac{4}{9}$; in case (2), $1 \times \frac{1}{9} = \frac{1}{9}$; in case (3), $\frac{2}{3} \times 1 = \frac{2}{3}$; in case (4), $\frac{2}{3} \times \frac{1}{9} = \frac{2}{27}$; in case (5), $\frac{1}{3} \times 1 = \frac{1}{3}$; and in case (6), $\frac{1}{3} \times \frac{4}{9} = \frac{4}{27}$.

Hence the chances of existence, for these 6 states, are proportional to '12, 3, 18, 2, 9, 4'. Hence their actual values are ' $\frac{1}{4}, \frac{1}{16}, \frac{3}{8}, \frac{1}{24}, \frac{3}{16}, \frac{1}{12}$ '.

Hence the chance of the unknown counter being red is the sum of $\frac{1}{4} \times 1, \frac{1}{16} \times 1, \frac{3}{8} \times \frac{2}{3}, \frac{1}{24} \times \frac{2}{3}, \frac{3}{16} \times \frac{1}{3}, \frac{1}{12} \times \frac{1}{3}$;

$$\text{i. e. it is } \frac{36 + 9 + 36 + 4 + 9 + 4}{9 \times 16}; \text{ which } = \frac{98}{9 \times 16} = \frac{49}{72}.$$

Q. E. F.

39. (9, 22)

Let x = no. of days.

$$\text{Then } (2 \times 10 - \overline{x - 1}) \cdot \frac{x}{2} = 14 + \{2 \times 2 + \overline{x - 1} \cdot 2\} \cdot \frac{x}{2};$$

$$\text{i. e. } \frac{21x}{2} - \frac{x^2}{2} = 14 + x + x^2;$$

$$\therefore 3x^2 - 19x + 28 = 0; \therefore x = \frac{19 \pm 5}{6} = 4 \text{ or } \frac{7}{3}.$$

Now the above solution has taken no account of the *discontinuity* of increase, or decrease of pace, and is the true solution only on the supposition that the increase or decrease is *continuous*, and such as to coincide with the above data at the end of each day. Hence '4' is a correct answer; but ' $\frac{7}{3}$ ' only indicates that a meeting occurs *during the third day*. To find the hour of this, let y = no. of hours.

Now in 2 days A has got to the end of 19 miles, B to the end of $(14 + 6)$, i. e. 20.

$$\therefore 19 + y \cdot \frac{8}{12} = 20 + y \cdot \frac{6}{12}.$$


$$\text{i. e. } y \cdot \frac{2}{3} = 1 + y \cdot \frac{1}{2}; \therefore y = 6.$$

Hence they meet at end of 2d. 6h., and at end of 4d.: and the distances are 23 miles, and 34 miles.

Q. E. F.

40. (9)

(1) Let ABC be the given Triangle, and AD the line from the vertex.



From D draw DE , DF , parallel to the sides; and from E and F draw EG , FH , $\perp BC$.

Then Triangles FBD , EDC , are similar to ABC ;

$\therefore FH : AD :: BD : BC$,

and $EG : AD :: DC : BC$;

$\therefore (FH + EG) : AD :: BC : BC$;


$\therefore FH + EG = AD$.

Also, $\because$ Triangles AED , AFD , are equal and on the same base AD ,

$\therefore$ their altitudes are equal; i. e. $DH = DG$.

Q. E. F.

(2) Let ABC be the given Triangle, and AD the line from the vertex.



Make $CE = BD$; from E draw EF , EG , parallel to the sides; and from F , G , draw FH , GK , $\perp BC$.

Then Triangles GBE , FEC , are similar to ABC ;

$$\therefore GK : AD :: BE : BC,$$

and $FH : AD :: EC : BC$;

$$\therefore (GK + FH) : AD :: BC : BC;$$

$$\therefore GK + FH = AD.$$

Also $BK : BE :: BD : BC$;

$$\therefore BK : DC :: EC : BC;$$

$$:: HC : DC;$$

$$\therefore BK = HC.$$

Q. E. F.

41. (9, 23)

(1) As there was certainly at least one W in the bag at first, the 'a priori' chances for the various states of the bag, ' WWW , WWB , WBW , WBB , BWW , BWB , BBW , BBB ,' were ' $\frac{1}{8}$, $\frac{3}{8}$, $\frac{3}{8}$, $\frac{1}{8}$ '.

These would have given, to the observed event, the chances ' 1 , $\frac{1}{2}$, $\frac{1}{6}$, 0 '.

Hence the chances, after the event, for the various states, are proportional to ' $\frac{1}{8} \cdot 1$, $\frac{3}{8} \cdot \frac{1}{2}$, $\frac{3}{8} \cdot \frac{1}{6}$ '; i. e. to ' $\frac{1}{8}$, $\frac{3}{16}$, $\frac{1}{16}$ '; i.e. to ' 2 , 3 , 1 '. Hence their actual values are ' $\frac{1}{3}$, $\frac{1}{2}$, $\frac{1}{6}$ '.

Hence the chance, of now drawing W , is ' $\frac{1}{3} \cdot 1 + \frac{1}{2} \cdot \frac{1}{2}$ '; i. e. it is $\frac{7}{12}$.

Q. E. F.

(2) If he had not spoken, the 'a priori' chances for the states ' $WWWW$, $WWWB$, $WWBB$, $WBBB$, $BBBB$ ', would have been ' $\frac{1, 4, 6, 4, 1}{16}$ '.

These would have given, to the observed event, the chances ' $1, \frac{1}{2}, \frac{1}{6}, 0, 0$ '.


Hence the chances, after the event, for the various states, are proportional to ' $\frac{1}{16} \cdot 1, \frac{1}{4} \cdot \frac{1}{2}, \frac{1}{6} \cdot \frac{3}{8}$ '; i. e. to ' $1, 2, 1$ '. Hence their actual values are ' $\frac{1}{4}, \frac{1}{2}, \frac{1}{4}$ '.

Hence the chance, of now drawing W , is ' $\frac{1}{4} \cdot 1 + \frac{1}{2} \cdot \frac{1}{2}$ '; i. e. it is $\frac{1}{2}$.

Q. E. F.

42. (10, 23)

Let ABC be the given Triangle. Bisect its angles, and draw $\perp$ s to them, forming the Triangle $A'B'C'$.



Now $\angle CBA' = 90^\circ - \frac{B}{2}$; and so of the others.

$$\therefore A' = 180^\circ - (CBA' + BCA') = \frac{B + C}{2} = 90^\circ - \frac{A}{2};$$

$$\therefore BA' = a \cdot \frac{\cos \frac{C}{2}}{\cos \frac{A}{2}}.$$


$$\text{Similarly, } BC' = c \cdot \frac{\cos \frac{A}{2}}{\cos \frac{C}{2}};$$

$$\begin{aligned} \therefore A'C' &= \frac{a \cos^2 \frac{C}{2} + c \cos^2 \frac{A}{2}}{\cos \frac{A}{2} \cos \frac{C}{2}} = \frac{a \cdot \frac{s(s-c)}{ab} + c \cdot \frac{s(s-a)}{bc}}{\frac{s}{b} \cdot \sqrt{\frac{(s-a)(s-c)}{ac}}}, \\ &= \frac{s-c + s-a}{\sin \frac{B}{2}} = \frac{b}{\sin \frac{B}{2}}. \end{aligned}$$

$$\text{Similarly, } A'B' = \frac{c}{\sin \frac{C}{2}};$$


$$\therefore \text{area of } A'B'C' = \frac{bc \cos \frac{A}{2}}{2 \sin \frac{B}{2} \sin \frac{C}{2}};$$

$$\begin{aligned} \therefore \frac{\text{area of } A'B'C'}{\text{area of } ABC} &= \frac{bc \cos \frac{A}{2}}{2 \sin \frac{B}{2} \sin \frac{C}{2}} \cdot \frac{2}{bc \sin A}, \\ &= \frac{\cos \frac{A}{2}}{\sin \frac{B}{2} \sin \frac{C}{2} \cdot 2 \sin \frac{A}{2} \cos \frac{A}{2}}, \\ &= \frac{1}{2 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}}, \\ &= \frac{abc}{2(s-a)(s-b)(s-c)}. \end{aligned}$$



Q. E. F.

43. (10)



Let ABC be the given Triangle; and let BFD, CFE , be the required lines, so that $FB = FC$, and Tetragon $AEFD = \text{Triangle } FBC$. And call the angle FBC ' θ '. It will suffice to calculate this angle.

Because Triangle $FBC = \text{Tetragon } AEFB$,

$$\begin{aligned} \therefore \text{Triangle } DBC &= \text{Triangle } AEC, \\ &= \text{Triangle } ABC - \text{Triangle } EBC; \end{aligned}$$

$\therefore$ Triangles DBC, EBC , together = Triangle ABC ;

$$\therefore \frac{1}{2} \cdot \frac{a^2}{\cot \theta + \cot C} + \frac{1}{2} \cdot \frac{a^2}{\cot \theta + \cot B} = \frac{1}{2} \cdot \frac{a^2}{\cot B + \cot C};$$

$$\therefore \frac{1}{\cot \theta + \cot C} + \frac{1}{\cot \theta + \cot B} = \frac{1}{\cot B + \cot C};$$

$$\therefore \frac{2 \cot \theta + (\cot B + \cot C)}{\cot^2 \theta + \cot \theta \cdot (\cot B + \cot C) + \cot B \cot C} = \text{do.};$$

$$\therefore \cot^2 \theta + \cot \theta \cdot (\cot B + \cot C) + \cot B \cot C = 2 \cot \theta \cdot (\cot B + \cot C) + (\cot B + \cot C);$$

$$\therefore \cot^2 \theta - \cot \theta \cdot (\cot B + \cot C) - (\cot^2 B + \cot B \cot C + \cot^2 C) =$$

$$\therefore \cot \theta = \frac{1}{2} \cdot \{\cot B + \cot C \pm \sqrt{(5 \cot^2 B + 6 \cot B \cot C + 5 \cot^2 C)}\}$$

Q. E. F.

44. (10)

Let k be a No. not containing 2 or 5 as a factor, i. e. let it be prime to 10. Then, if $\frac{1}{k}$ be reduced to a circulating decimal, and that to a vulgar fraction, the digits of the denominator will be a certain number of 9's; i. e. it will be of the form $(10^n - 1)$.

And since this fraction $= \frac{1}{k}$, and that k is prime to 10, and so prime to 10^m , the factor $(10^n - 1)$ must be a multiple of k .

This evidently holds good in any other scale of notation. Hence, if a be the radix of the scale of notation, and b a No. prime to a , a value may be found for n , which will make $(a^n - 1)$ a multiple of b .

Q. E. D.

EXAMPLES (not thought out).

(1) With radix 10, find a value, for n , which will make $(10^n - 1)$ a multiple of 7.

$$\frac{1}{7} = .\dot{1}4285\dot{7} = \frac{142857}{10^6 - 1}.$$

Ans. $n = 6$.

(2) Let the two given Nos. be 8, 9.

Taking 8 as radix, we get $\frac{1}{9} = .\dot{0}\dot{7} = \frac{7}{8^2 - 1}.$

Ans. $n = 2$.

(3) Let the two given Nos. be 7, 13.

Taking 7 as radix, we get

$$\frac{1}{13} = .\dot{0}3524563142\dot{1} = \frac{35245631421}{7^{12} - 1}.$$

Ans. $n = 12$.

45. (10, 23)

Divide each rod into $(n + 1)$ parts, where n is assumed to be odd, and the n points of division are assumed to be the only points where the rod will break, and to be equally frangible.

$$\begin{aligned} \text{The chance of one failure} & \text{ is } \frac{n-1}{n}; \\ \therefore \text{ " " } n \text{ failures} & \text{ is } \left(\frac{n-1}{n}\right)^n \\ & = \left(1 - \frac{1}{n}\right)^n. \end{aligned}$$

Now, if $m = \frac{1}{n}$; then, when $n = \frac{1}{0}$, $m = 0$;

$\therefore$ the chance that no rod is broke in the middle $= (1 - m)^{\frac{1}{m}}$, when $m = 0$;

i. e. it approaches the limit $(1 - 0)^{\frac{1}{0}}$.

And Ans. $= 1 - (1 - 0)^{\frac{1}{0}}$.

Now $(1 + 0)^{\frac{1}{0}} = e$. Hence if, in the series for e , we call the sum of the odd terms 'a', and of the even terms 'b'; then $e = a + b$; and $(1 - 0)^{\frac{1}{0}} = a - b = 2a - e$.

Q. E. F.

[N. B. What follows here was *not* thought out.]

$$\text{Now } a = 1 + \frac{1}{\underline{2}} + \frac{1}{\underline{4}} + \&c.$$

$$1 = 1$$

$$\frac{1}{\underline{2}} = .5$$

$$\frac{1}{\underline{4}} = .04166666 \&c.$$

$$\frac{1}{\underline{6}} = .00138888 \&c.$$

$$\frac{1}{\underline{8}} = .00002480 \&c.$$

$$\frac{1}{\underline{10}} = .00000027$$

$c.$

$$\therefore a = 1.5430806 \&c.$$

$$\therefore 2a = 3.0861612 \&c.$$



$$e = 2.7182818 \&c.$$

$$\therefore (1 - 0)^{\frac{1}{0}} = .3678793 \&c.$$

$$\therefore \text{Ans.} = 1 - (1 - 0)^{\frac{1}{0}} = .6321207 \&c.$$

46. (10)


Let ABC be the given Triangle, and D the given Point.



If we make a Triangle $D'E'F'$, having its angles equal to the given angles, and having D' as its assigned vertex, the Problem may be solved, if we can circumscribe, about the Triangle $D'E'F'$, a Triangle similar to ABC .

Now we can construct, on $E'F'$, $F'D'$, $D'E'$, segments of Circles containing angles equal to A , B , C . Hence the Problem may be solved, if we can place, in these Circles, a line $B'D'C'$, divided in the same proportion as BDC .

This Lemma may be solved as follows. Let G , H , be the centres of the Circles. Join GH , and divide it, at K , proportionally to BDC .



Join KD' ; through D' draw $B'D'C' \perp KD'$; and from G , H , draw GL , HM , $\perp B'C'$.

Now it may be easily proved that

$$LD' : D'M :: GK : KH :: BD : DC.$$

But $B'D'$, $D'C'$, are doubles of LD' , $D'M$;

$$\therefore B'D' : D'C' :: BD : DC.$$

Q. E. F.

[The construction is now obvious, viz. to join $B'F'$, $C'E'$, and produce them to meet, on the third Circle (as they may be easily proved to do), at A' ; then to divide AB , AC , at F and E , proportionally to $A'F'B'$, $A'E'C'$; and then to join DE , DF .]

By inspection, '0, 0, 0' are one set of values.

Subtracting, we get $x \cdot \left(\frac{1}{y} - \frac{1}{z} \right) = y - z$;

$$\therefore x = yz \cdot \frac{y - z}{z - y} = -yz, \text{ unless } y = z, \text{ in which case } x = \frac{0}{0}.$$

Now, by (1), $x = xy - yz$;

$\therefore$, when $y \neq z$, $x = xy + x$;

$\therefore xy = 0$, unless x be infinite.

Similarly, by (2), $xz = 0$, unless x be infinite.

Hence, if x be finite, and if $y \neq z$, either x or $y = 0$, and also either x or $z = 0$; i. e. either $x = 0$, or else $y = z = 0$. But the latter is excluded by our hypothesis. Hence $x = 0$. Hence $yz = 0$; i. e. either y or $z = 0$, and the other may take any value.

This gives us 2 more sets of values, viz.

$$x = y = 0; z \text{ has any value};$$

$$x = z = 0; y \text{ has any value}.$$

We have now to ascertain what happens when $y = z$.

By (1), $\frac{x}{y} = x - y$;

$$\therefore y^2 = x \cdot (y - 1); \text{ i.e. } x = \frac{y^2}{y - 1}.$$

Similarly, by (2), $x = \frac{z^2}{z - 1}.$

This gives us a 4th set of values, viz. $x = \frac{k^2}{k - 1}$, $y = z = k$; where k has any value.

Now y and z may evidently have *any* real values, but x is restricted by the equation


$$y^2 - xy + x = 0,$$

in which y cannot be real, unless $(x^2 - 4x) > 0$. Hence x may have any negative value, and any positive value that is not less than 4; but it cannot have any positive value, less than 4, without making y unreal.

Q. E. F.

48. (11)

Let ABC be the given Triangle, A', B', C' , the centres of the semicircles, and DE, FG, HJ , the common tangents; so that $DE = \alpha$, $FG = \beta$, and $HJ = \gamma$.



Join $B'D, C'E$; and from C' draw $C'K \perp B'D$. Hence $CK = \alpha$.

Call sides of given Triangle ' $2a, 2b, 2c$ '.

Then $B'C' = a$, and $B'K = b - c$;

$$\therefore C'K = \sqrt{a^2 - (b - c)^2};$$

$$\text{i.e. } \alpha = \sqrt{(a - b + c) \cdot (a + b - c)};$$

$$\text{similarly, } \beta = \sqrt{(a + b - c) \cdot (-a + b + c)},$$

$$\text{and } \gamma = \sqrt{(-a + b + c) \cdot (a - b + c)};$$

$$\therefore \frac{\beta\gamma}{\alpha} = -a + b + c;$$

$$\text{similarly, } \frac{\gamma\alpha}{\beta} = a - b + c,$$

$$\text{and } \frac{\alpha\beta}{\gamma} = a + b - c;$$

$$\therefore \text{ their sum} = a + b + c, \\ = \text{semi-perimeter of } ABC.$$

Q.E.D.

49. (11, 23)

Take, as unit, a side of one of the Triangles.

If the Tetrahedron be cut by a vertical Plane containing one of the slant edges, the section is a Triangle whose base is $\frac{\sqrt{3}}{2}$, and whose sides are $\frac{\sqrt{3}}{2}, 1$;

hence cosine of smaller base-angle

$$= \left(\frac{3}{4} + 1 - \frac{3}{4} \right) \cdot \frac{1}{\sqrt{3}} = \frac{1}{\sqrt{3}};$$

$$\therefore \text{ its sine} = \frac{\sqrt{2}}{\sqrt{3}} = \text{its altitude};$$

and this is the altitude of the Tetrahedron;

$$\therefore \text{ volume of Tetrahedron} = \frac{1}{3} \cdot \frac{\sqrt{2}}{\sqrt{3}} \cdot \frac{\sqrt{3}}{4} = \frac{\sqrt{2}}{12}.$$

Also altitude of Pyramid = altitude of Triangle whose base is $\sqrt{2}$, and whose sides are 1, 1;

$$\text{i.e. it} = \frac{\sqrt{2}}{2};$$

$$\therefore \text{ volume of Pyramid} = \frac{1}{3} \cdot \frac{\sqrt{2}}{2} = \frac{\sqrt{2}}{6}.$$

$$\text{Hence required ratio} = \frac{\sqrt{2}}{6} \cdot \frac{12}{\sqrt{2}} = 2.$$

Q. E. F.

50. (11,23)

At first, the chance that bag H shall contain

2 W counters, is $\frac{1}{4}$.

1 W and 1 B , is $\frac{1}{2}$.

2 B , is $\frac{1}{4}$.

$\therefore$, after adding a W , the chance that it shall contain

3 W , is $\frac{1}{4}$.

2 W , 1 B , is $\frac{1}{2}$.

1 W , 2 B , is $\frac{1}{4}$.

hence the chance of drawing a W from it is

$$\frac{1}{4} \times 1 + \frac{1}{2} \times \frac{2}{3} + \frac{1}{4} \times \frac{1}{3} : \text{i. e. } \frac{2}{3}.$$

$\therefore$ the chance of drawing a B is $\frac{1}{3}$.

After transferring this (unseen) counter to bag K , the chance that it shall contain

3 W , is $\frac{2}{3} \times \frac{1}{4}$; i. e. $\frac{1}{6}$.

2 W , and 1 B , is $\frac{2}{3} \times \frac{1}{2} + \frac{1}{3} \times \frac{1}{4}$; i. e. $\frac{5}{12}$.

1 W , 2 B , is $\frac{2}{3} \times \frac{1}{4} + \frac{1}{3} \times \frac{1}{2}$; i. e. $\frac{1}{3}$.

3 B , is $\frac{1}{3} \times \frac{1}{4}$; i. e. $\frac{1}{12}$;

$\therefore$ the chance of drawing a W from it is

$$\frac{1}{6} \times 1 + \frac{5}{12} \times \frac{2}{3} + \frac{1}{3} \times \frac{1}{3}; \text{ i. e. } \frac{5}{9}.$$

$\therefore$ the chance of drawing a B is $\frac{4}{9}$.

Before transferring this to bag H , the chance that bag H shall contain

$$\begin{array}{lll}
2W, & \text{is} & \frac{1}{4} \times 1 + \frac{1}{2} \times \frac{1}{3}; \quad \text{i. e. } \frac{5}{12}. \\
1W, 1B, & & \frac{1}{2} \times \frac{2}{3} + \frac{1}{4} \times \frac{2}{3}; \quad \text{i. e. } \frac{1}{2}. \\
2B, & & \frac{1}{4} \times \frac{1}{3}; \quad \text{i. e. } \frac{1}{12}.
\end{array}$$

∴, after transferring it, the chance that bag H shall contain

$$\begin{array}{lll}
3W, & \text{is} & \frac{5}{12} \times \frac{5}{9}; \quad \text{i. e. } \frac{25}{108}. \\
2W, 1B, & & \frac{5}{12} \times \frac{4}{9} + \frac{1}{2} \times \frac{5}{9}; \quad \text{i. e. } \frac{50}{108}. \\
1W, 2B, & & \frac{1}{2} \times \frac{4}{9} + \frac{1}{12} \times \frac{5}{9}; \quad \text{i. e. } \frac{29}{108}. \\
3B, & & \frac{1}{12} \times \frac{4}{9}; \quad \text{i. e. } \frac{4}{108}.
\end{array}$$

Hence the chance of drawing a W is


$$\frac{1}{108} \times \{25 \times 1 + 50 \times \frac{2}{3} + 29 \times \frac{1}{3}\}; \text{ i. e. } \frac{17}{27}.$$

i. e. the odds are 17 to 10 on its happening.

Q. E. F.

51. (12)

Let ABC be the given Triangle, and D the given Point.



(Analysis.)

Let DE be the line required. Draw DF , EG , $\perp$ the base. Then their sum is equal to DE .

Bisect DE at H , and draw $HK \perp$ the base: then it is evident that HK is the $A.M.$ of DF , EG , and is equal to half their sum; i. e. it is equal to half of DE . Hence a Circle, drawn with centre H and at distance HD , will pass through E and K , and will touch the base at K .

Through H draw LHM parallel to AC . Then DA is evidently bisected at L . Also LM passes through the centre of the Circle. Hence, if DN be drawn $\perp$ LM (or CA), it is a chord of the Circle, and is bisected at R . Produce ND to meet the base produced at S . Hence SDN cuts the Circle, and SK touches it at K . But S can be found, and SK can then be taken, so that sq. of SK may be equal to rect. of SD , SN .

(*Synthesis.*)

From D draw $DN \perp AC$, and produce it to meet the base produced at S . Take SK , so that its square may be equal to rect. of SD , SN .

Bisect DA at L , and from L draw LM parallel to AC ; and from K draw $KH \perp$ the base, to meet LM at H . Join DH , and produce it to meet AC at E , and draw DF , EG , $\perp$ the base.

Because $DL = LA$, and that LM is parallel to AC ,

$\therefore DH = HE = HK$; $\therefore DE = 2HK$.

But $DF + EG = 2HK$; $\therefore DF + EG = DE$.

Q. E. F.

[N. B. This proof is incomplete. I have assumed, without proving it, that $DH = HK$. It may be proved thus. Because sq. of $SK =$ rect. of SD , SN , $\therefore DN$ is a chord of a Circle which touches the base at K ; $\therefore LM$, which bisects it at right angles, passes through the centre. But KH also passes through the centre; $\therefore H$ is the centre; $\therefore HD = HK$.]

52. (12, 23)

Let x be the number of pennies each had at first.

No. (3) received x , took out $(2 + 4)$, and put in $\frac{x}{2}$; so that the sack then contained $(x \cdot \frac{3}{2} - 6)$. Let us write ' a ' for ' $\frac{3}{2}$ '.

No. (5) received $(xa - 6)$, took out $(4 + 1)$, and put in enough to multiply, by a , its contents when he received it. The sack now contained

$$(xa^2 - 6a - 5).$$

No. (2) took out $(1 + 3)$, and handed on $(xa^3 - 6a^2 - 5a - 4)$.

No. (4) took out $(3 + 5)$, and handed on

$$(xa^4 - 6a^3 - 5a^2 - 4a - 8).$$

No. (1) put in 2. The sack now contained $5x$.


Hence $xa^4 - 6a^3 - 5a^2 - 4a - 6 = 5x$;

$$\begin{aligned} \therefore x &= \frac{6a^3 + 5a^2 + 4a + 6}{a^4 - 5}; \\ &= \frac{(6.3^3 + 5.3^2.2 + 4.3.2^2 + 6.2^3).2}{3^4 - 5.2^4}; \\ &= \frac{(162 + 90 + 48 + 48).2}{81 - 80} = 696 = 2l. 18s. 0d. \end{aligned}$$

Q. E. F.

53. (13,24)

Let ABC be the given Triangle, and P the given Point; and call its trilinear co-ordinates ' α, β, γ '.



From P draw PA' , PB' , PC' , $\perp$ the sides, and therefore equal to α , β , γ . Produce PA' and PC' to A'' and C'' , making $A'A'' = PA'$, and $C'C'' = PC'$. From C'' draw $C''D \perp AC$, and produce it to E , making $DE = C''D$. Join EA'' , cutting AC in R , and BC in S . Join $C''R$, cutting AB in Q . Join PQ , PS .

The path of the ball is evidently $PQRSP$; and we have to calculate the length of AR .

$$\text{Now } AR = DR + AD = DR + AB' - DB'.$$

First, to calculate DR .

From P draw PU , PV , parallel to AB , AC ; from C' draw $C'W \perp PV$; and from A'' draw $A''F \perp AC$.

By similar Triangles, $DR : RF :: DE : A''F :: C''D : A''F$;

$$\therefore DR : DF :: C''D : (C''D + A''F);$$

$$\therefore DR = \frac{DF \cdot C''D}{C''D + A''F}.$$

$$\text{Now } \angle C'VP = A; \therefore \angle C'PV = 90^\circ - A;$$

$$\therefore WP = \gamma \sin A;$$

$$\therefore DB', \text{ which } = 2WP, = 2\gamma \sin A.$$

$$\text{Similarly, } B'F = 2\alpha \sin C;$$

$$\therefore DF = 2(\alpha \sin C + \gamma \sin A).$$

$$\text{Again, } C'W = \gamma \cos A;$$

$$\therefore C''D, \text{ which } = 2C'W + PB', = 2\gamma \cos A + \beta.$$

$$\text{Similarly, } A''F = 2\alpha \cos C + \beta;$$

$$\therefore C''D + A''F = 2(\alpha \cos C + \gamma \cos A + \beta);$$

$$\therefore DR = \frac{(\alpha \sin C + \gamma \sin A) \cdot (2\gamma \cos A + \beta)}{\alpha \cos C + \gamma \cos A + \beta}.$$

$$\text{Now } AB' = B'U + UA = B'U + PV,$$

$$= \beta \cot A + \gamma \operatorname{cosec} A = \frac{\beta \cos A + \gamma}{\sin A};$$

$$\begin{aligned}
\therefore AB' - DB' &= \frac{\beta \cos A + \gamma}{\sin A} - 2\gamma \sin A, \\
&= \frac{\beta \cos A + \gamma(1 - 2 \sin^2 A)}{\sin A} \\
&= \frac{\beta \cos A + \gamma \cos 2A}{\sin A}.
\end{aligned}$$

Now $AR = DR + AB' - DB'$;

$$\therefore AR = \frac{(\alpha \sin C + \gamma \sin A) \cdot (2\gamma \cos A + \beta)}{\alpha \cos C + \gamma \cos A + \beta} + \frac{\beta \cos A + \gamma \cos 2A}{\sin A}.$$

Q. E. F.

54. (13, 24)


It is evident that Triangle ADE is similar to ABC .

Let ' k ' = ratio $\frac{DE}{a} = \frac{AE}{b} = \frac{AD}{c}$.

Now $DG = DE$; $\therefore DG = ka$;

$\therefore GB = c - ka - kc$;

$\therefore \frac{GB}{c} = 1 - k - k \cdot \frac{a}{c}$;



$$\therefore GF \text{ (which } = GB \cdot \frac{b}{c}) = b - kb - k \cdot \frac{ab}{c};$$

$$\text{but } GF = DE = ka;$$

$$\therefore b - kb - k \cdot \frac{ab}{c} = ka;$$

$$\therefore bc = k.(bc + ca + ab);$$

$$\therefore k = \frac{bc}{bc + ca + ab} = \frac{\frac{1}{a}}{\frac{1}{a} + \frac{1}{b} + \frac{1}{c}} = \frac{\frac{1}{a}}{m} \text{ (say).}$$


$$\text{Hence } AD = \frac{c \cdot \frac{1}{a}}{m}; DG = \frac{1}{m} = \frac{c \cdot \frac{1}{c}}{m}.$$

$$\therefore GB \text{ (which } = c - AD - DG) = \frac{c.(m - \frac{1}{a} - \frac{1}{c})}{m} = \frac{c \cdot \frac{1}{b}}{m};$$

$$\therefore AD : DG : GB :: \frac{1}{a} : \frac{1}{c} : \frac{1}{b}.$$

$$\text{Also } DE = ka = \frac{1}{m} = \frac{1}{\frac{1}{a} + \frac{1}{b} + \frac{1}{c}}.$$

Q. E. F.



Let A, B, C be the centres of the bases of the towers; and a, b, c their radii. Suppose P the required Point; and from P draw a pair of tangents to each circle, and lines to the centres, which will evidently bisect the angles contained by the pairs of tangents.

Hence angles APD, BPE, CPF are equal;

$$\therefore \sin APD = \sin BPE = \sin CPF;$$

$$\text{i.e. } \frac{a}{AP} = \frac{b}{BP} = \frac{c}{CP};$$

$$\therefore AP : BP : CP :: a : b : c.$$

Draw a Line through A, B , and on it take Points G, H , such that $AG : GB :: AH : HB :: a : b$.

Then the Semicircle, described on GH , is the locus of all Points whose distances, from A and B , are proportional to a, b .

Hence, if a Line be drawn through B, C , and a Semicircle described which shall be the locus of all Points whose distances, from B and C , are proportional to b, c ; the intersection of these two Semicircles will be the Point required.

Q. E. F.

[Note. "The locus of all Points whose distances &c.," if represented algebraically, is evidently a Circle, whose centre is on the Line through A, B , and which passes through G and H .]

Draw BC, CE, BD , equal to the given altitudes, so as to form right $\angle$ s at B and C : and produce DB, EC . Join DC , and draw $CF \perp$ to it. Join EB , and draw $BG \perp$ to it. With centre B , and distance BF , describe a circle: with centre C , and distance CG , describe another: let them meet at A : and join AB, AC .

Call the altitudes of ABC , ' α, β, γ '.

$$\text{Now } \alpha.BC = \beta.CA = \gamma.AB$$

= twice area of ABC ;

also, taking BC as unit-line,

$$BC = \frac{1}{BC}, \quad CA = CG = \frac{1}{CE},$$

$$AB = BF = \frac{1}{BD};$$

$$\therefore \frac{\alpha}{BC} = \frac{\beta}{CE} = \frac{\gamma}{BD};$$

i.e. α, β, γ are proportional to given altitudes;

$\therefore$ Triangle ABC is similar to required Triangle.

The rest of the construction is obvious.

Q. E. F.

57. (14, 25)

(1) *Geometrically.*

Let ABC be given Triangle.


(*Analysis.*)

Suppose the 3 Squares described, and that their upper edges form the Triangle $A'B'C'$. Join AA', BB', CC' .

Now it is evident that, if BB' be produced, the perpendiculars dropped, from any Point of it, upon AB, BC , will be proportional to $B'F, B'D$.

Similarly for AA' and CC' .

Hence these 3 Lines will meet at the Point from which the perpendiculars, dropped upon the sides of ABC , are proportional to $B'C'$, $C'A'$, $A'B'$.



Hence, if Squares be described externally on the sides of ABC , and if their outer edges be produced to form a new Triangle $A''B''C''$: this Triangle, with these 3 Squares, will form a Diagram wholly similar to that formed by the Triangle ABC , with the 3 Squares inside it.

(*Synthesis.*)

Hence, if Squares be described externally on the sides of the given Triangle; and if their outer edges be produced to form a new Triangle; and if the sides of the given Triangle be divided similarly to those of the new Triangle: their central portions will be the bases of the required Squares.

Q. E. F.

(2) *Trigonometrically.*

Let a, b, c be the sides of the given Triangle, and m its area; and let x, y, z be the sides of the required Squares.

It is evident that a Circle can be described about the Tetragon $BDB'F$.

Hence $\angle B'BD = \angle B'FD$.

Now, in Triangle $B'FD$, we know that

$$B'D \sin D = B'F \sin F;$$


$$\text{i. e. } x \sin(B' + F) = z \sin F;$$

$$\therefore x \sin B' \cos F + x \cos B' \sin F = z \sin F.$$

Now $\angle B$ is supplementary to $\angle B'$;

$$\therefore x \sin B \cos F = (z + x \cos B) \sin F;$$

$$\therefore \cot F = \frac{z + x \cos B}{x \sin B} = \cot B' BD.$$



Now $BD = x \cot B' BD$;

$$\therefore BD = \frac{z + x \cos B}{\sin B}.$$

$$\text{Similarly, } EC = \frac{y + x \cos C}{\sin C}.$$

But $BD + EC = a - x$;

$$\therefore \frac{z + x \cos B}{\sin B} + \frac{y + x \cos C}{\sin C} = a - x;$$

$$\therefore \frac{x \sin(B + C) + y \sin B + z \sin C}{\sin B \sin C} = a - x;$$

$$\text{i. e. } \frac{x \sin A + y \sin B + z \sin C}{\sin B \sin C} = a - x.$$

Now it is evident that these Triangles are similar; so that

$$\frac{a}{x} = \frac{b}{y} = \frac{c}{z}.$$

Hence, multiplying the last equation, throughout, by one or other of these equal fractions, we get

$$\begin{aligned}\frac{a \sin A + b \sin B + c \sin C}{\sin B \sin C} &= \frac{a^2}{x} - a; \\ \therefore \frac{a \sin A + b \sin B + c \sin C}{a \sin B \sin C} &= \frac{a}{x} - 1; \\ \therefore \frac{a}{x} &= \frac{a \sin A + b \sin B + c \sin C}{a \sin B \sin C} + 1.\end{aligned}$$


Hence, multiplying above and below by one or other of the equal fractions $\frac{a}{\sin A}$, $\frac{b}{\sin B}$, $\frac{c}{\sin C}$,

$$\begin{aligned}\frac{a}{x} &= \frac{a^2 + b^2 + c^2}{ab \sin C} + 1; \\ &= \frac{a^2 + b^2 + c^2}{2m} + 1 = \frac{b}{y} = \frac{c}{z}.\end{aligned}$$

Q. E. F.

58. (14, 25)

It may be assumed that the 3 Points form a Triangle, the chance of their lying in a straight Line being (practically) *nil*.




Take the longest side of the Triangle, and call it ' AB ': and, on that side of it, on which the Triangle lies, draw the semicircle AFB . Also, with centres A , B , and distances AB , BA , draw the arcs BDC , AEC , intersecting at C .

Then it is evident that the vertex of the Triangle cannot fall outside the Figure $ABDCE$.

Also, if it fall inside the semicircle, the Triangle is obtuse-angled: if outside it, acute-angled. (The chance, of its falling *on* the semicircle, is practically *nil*.)

$$\text{Hence required chance} = \frac{\text{area of semicircle}}{\text{area of fig. } ABDCE}.$$



Now let $AB = 2a$: then area of semicircle $= \frac{\pi a^2}{2}$; and area of Fig. $ABDCE = 2 \times \text{sector } ABDC - \text{Triangle } ABC$;

$$= 2 \cdot \frac{4\pi a^2}{6} - \sqrt{3} \cdot a^2 = a^2 \cdot \left(\frac{4\pi}{3} - \sqrt{3} \right);$$

$$\therefore \text{chance} = \frac{\frac{\pi}{2}}{\frac{4\pi}{3} - \sqrt{3}} = \frac{3}{8 - \frac{6\sqrt{3}}{\pi}}.$$

Q. E. F.

59. (14, 25)

Let $KL = MN = a$,

$KN = LM = b$,


$KM = LN = c$;

and let $\angle$ s LMK , MKL , KLM be equal to 'A, B, C'; and similarly for the $\angle$ s of the other facets.

From K draw $KT \perp$ base-facet LMN . Also draw $KR, KS, \perp LM, MN$. And join TR, TM, TS .

It is easily proved that $\angle s TRM, TSM$ are right.

The required volume is $\frac{1}{3} \cdot KT \cdot LMN$. The area of LMN is of course known. All we need is the length of KT . Now $KT^2 = KS^2 - TS^2$; and KS evidently $= c \cdot \sin B$. Hence all we need is the length of TS .




Now this requires a preliminary Lemma, in itself a very pretty problem, viz.—

LEMMA (1).

Given, in Tetragon $RMST$, sides RM, MS , and $\angle RMS$, and that $\angle s TRM, TSM$ are right: find TS .

$$\text{Now } \frac{TS}{\sin TRS} = \frac{TR}{\sin TSR};$$

$$\text{also } TS \cos TSR + TR \cos TRS = RS;$$



$$\begin{aligned}
\therefore \frac{TS}{\sin TRS} &= \frac{TR}{\sin TSR}, \\
&= \frac{TS \cos TSR + TR \cos TRS}{\sin TRS \cos TSR + \sin TSR \cos TRS}, \\
&= \frac{RS}{\sin RMS} = \frac{MS}{\sin MRS}; \\
\therefore \frac{TS}{\cos MRS} &= \frac{MS}{\sin MRS}; \text{ i.e. } TS = MS \cot MRS.
\end{aligned}$$


Q. E. F.

Hence this requires another Lemma, in order to find the value of $\cot MRS$ (or $\tan MRS$, which will do as well, and makes a prettier problem).

LEMMA (2).

Given, in Triangle RMS , sides RM , MS , and $\angle RMS$: find $\tan MRS$.

$$\begin{aligned}
\tan MRS &= \frac{\sin MRS}{\cos MRS} = \frac{RS \sin MRS}{RS \cos MRS}, \\
&= \frac{MS \sin RMS}{RM - MS \cos RMS}.
\end{aligned}$$



Q. E. F.

Hence, in Tetragon $RMST$, we have by Lemma (1),

$$TS = MS \cot MRS;$$

$$\text{and, by Lemma (2), } \cot MRS = \frac{RM - MS \cos RMS}{MS \sin RMS},$$

$$= \frac{c \cos A - c \cos B \cos C}{c \cos B \sin C} = \frac{\cos A - \cos B \cos C}{\cos B \sin C};$$

$$\therefore TS = \frac{c}{\sin C} \cdot (\cos A - \cos B \cos C).$$

$$\text{Now } KT^2 = KS^2 - TS^2;$$

$$\therefore \text{it} = (c \sin B)^2 - \frac{c^2}{\sin^2 C} \cdot (\cos A - \cos B \cos C)^2,$$

$$= \frac{c^2}{\sin^2 C} \cdot \{(\sin B \sin C)^2 - (\cos A - \cos B \cos C)^2\};$$


$$\text{therefore} \quad KT = \frac{c}{\sin C} \quad \text{multiplied} \quad \text{by} \quad \sqrt{\sin^2 B \sin^2 C - \cos^2 B \cos^2 C - \cos^2 A + 2 \cos A \cos B \cos C},$$

$$= \frac{c}{\sin C} \text{ multiplied by}$$

$$\sqrt{(1 - \cos^2 B) \cdot (1 - \cos^2 C) - \cos^2 B \cos^2 C - \cos^2 A + 2 \cos A \cos B \cos C},$$

$$= \frac{c}{\sin C} \cdot \sqrt{1 - (\cos^2 A + \cos^2 B + \cos^2 C) + 2 \cos A \cos B \cos C},$$

which is symmetrical, as it ought to be.



$$\text{Now area of } LMN = \frac{ab \sin C}{2};$$

hence volume of Tetrahedron

$$= \frac{abc}{6} \cdot \sqrt{1 - (\cos^2 A + \cos^2 B + \cos^2 C) + 2 \cos A \cos B \cos C}.$$

Q. E. F.

60. (14, 25)

Let $\angle BAD = \theta$, $\angle CAD = \phi$.



$$\begin{aligned} \text{Now } \frac{\sin(B + \theta)}{\sin \theta} &= \frac{c}{\left(\frac{ma}{m+n}\right)} \\ &= \frac{c(m+n)}{ma}; \end{aligned}$$

$$\therefore \sin B \cot \theta + \cos B = \frac{c(m+n)}{ma};$$

$$\therefore \cot \theta = \frac{c(m+n)}{ma \cdot \sin B} - \cot B,$$

$$= \frac{(m+n) \cdot (a \cos B + b \cos A) - ma \cos B}{ma \sin B},$$

$$= \frac{(m+n)b \cos A + na \cos B}{ma \sin B};$$



$$\begin{aligned} \text{i. e. } \cot \theta &= \frac{(m+n) \cdot \frac{b}{\sin B} \cdot \cos A + na \cot B}{ma}, \\ &= \frac{(m+n)a \cot A + na \cot B}{ma}, \end{aligned}$$

$$= \frac{(m+n) \cot A + n \cot B}{m}.$$

Similarly, $\cot \phi = \frac{(m+n) \cot A + m \cot C}{n}.$

Q. E. F.

COROLLARIES.

(1) $m \cot \theta - n \cot \phi = n \cot B - m \cot C.$

(2) $\frac{\cot B + \cot \phi}{\cot C + \cot \theta} = \frac{m}{n}.$

(3) If Triangle be equilateral,

$$\cot \theta = \frac{m+2n}{m} \cdot \frac{1}{\sqrt{3}},$$

$$\cot \phi = \frac{n+2m}{n} \cdot \frac{1}{\sqrt{3}};$$

$$\therefore \frac{\cot \theta}{\cot \phi} = \frac{mn+2n^2}{mn+2m^2};$$

$$\therefore \frac{\tan \theta}{\tan \phi} = \frac{mn+2m^2}{mn+2n^2};$$

i. e., if $CD'B'$ be drawn $\perp$ to AD , $\frac{B'D'}{D'C} = \frac{mn+2m^2}{mn+2n^2};$

e. g., if $\frac{m}{n} = \frac{1}{2}$, $\frac{B'D'}{D'C} = \frac{2}{5}.$

(4) Let $\tan A = 1, \tan B = 2, \tan C = 3;$

then $\cot \theta = \frac{m+n+n \cdot \frac{1}{2}}{m} = \frac{2m+3n}{2m},$

$$\cot \phi = \frac{m+n+m \cdot \frac{1}{3}}{n} = \frac{3n+4m}{3n};$$

$$\therefore \frac{\tan \theta}{\tan \phi} = \frac{6mn + 8m^2}{6mn + 9n^2};$$

from which, if $\frac{\tan \theta}{\tan \phi}$ were given, we could find $\frac{m}{n}$ from a Quadratic Equation.

I tried various values, to find one which would give rational values for m and n , and found that $\frac{2}{3}$ would do, as it leads to the Quadratic

$$2(6mn + 9n^2) - 3(6mn + 8m^2) = 0,$$

in which $(B^2 - 4AC)$ becomes, after dividing all through by 6, $(1^2 + 4.4.3)$, i. e. 49.

The Quadratic is $4m^2 + mn - 3n^2 = 0$;

whence $\frac{m}{n} = \frac{-1 \pm 7}{8} = \frac{3}{4}$; which solves the Problem 'Given a Triangle ABC , having the tangents of its angles equal to 1, 2, 3: divide BC at D , so that, if AD be joined, and $CD'B'$ drawn $\perp$ to it, the ratio $\frac{B'D'}{D'C}$ may be $\frac{2}{3}$ '. The answer is 'Divide it so that $\frac{BD}{DC} = \frac{3}{4}$ '.

61. (14)

We know that the equation

$$\begin{aligned} &'(a^2 + 4b^2 + 4c^2) + (4a^2 + b^2 + 4c^2) \\ &\quad + (4a^2 + 4b^2 + c^2) = 9(a^2 + b^2 + c^2)' \end{aligned}$$

is identically true.

Hence $a^2 + b^2 + c^2$

$$\begin{aligned}
&= \frac{1}{9} \cdot \{(a^2 + 4b^2 + 4c^2) + (4a^2 + b^2 + 4c^2) + (4a^2 + 4b^2 + c^2)\}; \\
&= \frac{1}{9} \cdot \{(a^2 + 4b^2 + 4c^2 + 8bc - 4ca - 4ab) \\
&\quad + (4a^2 + b^2 + 4c^2 - 4bc + 8ca - 4ab) \\
&\quad + (4a^2 + 4b^2 + c^2 - 4bc - 4ca + 8ab)\}; \\
&= \frac{1}{9} \cdot \{(-a + 2b + 2c)^2 + (2a - b + 2c)^2 + (2a + 2b - c)^2\}; \\
&= \left(\frac{-a + 2b + 2c}{3}\right)^2 + \left(\frac{2a - b + 2c}{3}\right)^2 + \left(\frac{2a + 2b - c}{3}\right)^2.
\end{aligned}$$

Now $(-a + 2b + 2c) = 3(b + c) - (a + b + c)$;

$\therefore$, if $(a + b + c)$ be a multiple of 3, so also is $(-a + 2b + 2c)$;

$\therefore \frac{-a + 2b + 2c}{3}$ is an integer;

and similarly for the other 2 fractions.

Also it may be proved that, if $\frac{-a + 2b + 2c}{3}$ be equal to a , or b , or c , then a, b, c can be arranged in $A.P.$

First, let $\frac{-a + 2b + 2c}{3} = a$;

then $-a + 2b + 2c = 3a$; i. e. $b + c = 2a$;

secondly, let $\frac{-a + 2b + 2c}{3} = b$;

then $-a + 2b + 2c = 3b$; i. e. $2c = a + b$;

thirdly, let $\frac{-a + 2b + 2c}{3} = c$;

then $-a + 2b + 2c = 3c$; i. e. $2b = c + a$.

And similarly for the other 2 fractions.

Hence, contranominally, if a, b, c can *not* be arranged in $A.P.$, the 2 sets of squares have no common term.


Q. E. D.

Numerical Examples (not thought out).

a^2	b^2	c^2	$\left(\frac{-a + 2b + 2c}{3}\right)^2$	$\left(\frac{2a - b + 2c}{3}\right)^2$	$\left(\frac{2a + 2b - c}{3}\right)^2$
1^2	4^2	4^2	5^2	2^2	2^2
3^2	4^2	8^2	7^2	6^2	2^2
4^2	5^2	9^2	8^2	7^2	3^2

62. (14)

Let AB , AC , be the given Lines, and P the given Point.



Through P draw PD parallel to AB ; from DC cut off DE equal to AD ; join EP , and produce it to meet AB at F .

Because $AD = DE$, and that DP is parallel to AB ,
 $\therefore FP = PE$.

Now let GPH be any other line through P ;
 then $\angle PFH > \angle PEG$.

Because, in Triangles PFH , PEG , $PF = PE$, and

$$\angle FPH = \angle GPE, \quad \text{and} \quad \angle PFH > \angle PEG,$$

$\therefore PH > PG$, and Triangle $PFH >$ Triangle PGE .

To each add Tetragon $AFPG$;


$\therefore$ Triangle $AGH >$ Triangle AEF .

And so of any other line through P .

Hence AEF is the least possible Triangle.

Q. E. F.

63. (15, 26)



Let each side of each Square = 2.

Then $LG = \sqrt{3}$, $MG = (\sqrt{2} - 1)$;

$$\therefore LM(= JK) = \sqrt{3 - (2 + 1 - 2\sqrt{2})}$$

$$= 2^{\frac{3}{4}};$$

$$\therefore OJ = OK = \frac{1}{2^{\frac{1}{4}}}.$$

Take O as origin, the X -axis $\parallel$ to AD , and the Y -axis to AB ; and let JK be part of the Z -axis.

Let equation to plane containing Triangle CDG be

$$x \cos \alpha + y \cos \beta + z \cos \gamma - p = 0,$$

where p is length of perpendicular dropped, from O , upon this plane, and meeting it somewhere in LG .

Hence we can find p from equation to LG , in the XZ -plane, which will be

$$x \cos \alpha + z \cos \gamma - p = 0;$$

now this line contains L , whose co-ordinates are $\left(1, \frac{1}{2^{\frac{1}{4}}}\right)$, and G , whose

co-ordinates are $\left(\sqrt{2}, -\frac{1}{2^{\frac{1}{4}}}\right)$;

$$\therefore \cos \alpha + \frac{1}{2^{\frac{1}{4}}} \cdot \cos \gamma - p = 0,$$

$$\text{and } \sqrt{2} \cdot \cos \alpha - \frac{1}{2^{\frac{1}{4}}} \cdot \cos \gamma - p = 0;$$

$$\therefore (\sqrt{2} - 1) \cdot \cos \alpha = \frac{2}{2^{\frac{1}{4}}} \cdot \cos \gamma = 2^{\frac{3}{4}} \cdot \cos \gamma;$$

$$\therefore \frac{\cos \alpha}{2^{\frac{3}{4}}} = \frac{\cos \gamma}{\sqrt{2} - 1} = \frac{1}{\sqrt{2^{\frac{3}{2}} + 3 - 2^{\frac{3}{2}}}} = \frac{1}{\sqrt{3}};$$

$$\therefore \cos \alpha = \frac{2^{\frac{3}{4}}}{\sqrt{3}}, \cos \gamma = \frac{\sqrt{2} - 1}{\sqrt{3}};$$

$$\therefore p = \frac{2^{\frac{3}{4}}}{\sqrt{3}} + \frac{\sqrt{2} - 1}{2^{\frac{1}{4}} \cdot \sqrt{3}} = \frac{\sqrt{2} + 1}{2^{\frac{1}{4}} \cdot \sqrt{3}}.$$

Now area of $CDG = \sqrt{3}$;

$\therefore$ volume of pyramid, whose base is CDG and whose vertex is O , $= \frac{\sqrt{2} + 1}{3 \cdot 2^{\frac{1}{4}}}$;

and there are eight such pyramids in the solid;

$\therefore$ their sum $= \frac{8(\sqrt{2} + 1)}{3 \cdot 2^{\frac{1}{4}}}$.

Also volume of pyramid, whose base is $ABCD$, and whose vertex is O , $= \frac{4}{3 \cdot 2^{\frac{1}{4}}}$;


and there are 2 such pyramids in the solid;

$\therefore$ their sum $= \frac{8}{3 \cdot 2^{\frac{1}{4}}}$;

$\therefore$ volume of solid $= \frac{8(2 + \sqrt{2})}{3 \cdot 2^{\frac{1}{4}}} = \frac{8 \cdot 2^{\frac{1}{4}} \cdot (\sqrt{2} + 1)}{3}$

Q. E. F.

64. (15)



Let ABC be the given Triangle, and O the given Point; and let OD , its distance from BC , be less than either OE or OF , its distances from CA , AB .

Draw a line GH equal to OE , and $GK \perp$ it and equal to OF ; and join HK ; and about the Triangle GHK describe a Circle; and place in it a line KL equal to OD ; and join LH .

Because sqs of $KL, LH =$ sqs of KG, GH , and that KL is less than either KG or GH , $\therefore LH$ is greater than either;

$\therefore$ a Circle, with centre O , and radius equal to LH , will cut all three Lines, in two Points each. Describe this Circle.

Then sqs of $MD, DO =$ sqs of PE, EO ;

also sq. of $LH =$ sqs of RF, FO ;

$\therefore$ sqs of $MD, DO, LH =$ sqs of PE, RF, EO, FO ;

but sqs of $DO, LH =$ sqs of KL, LH ,

$=$ sqs of $GH, GK =$ sqs of EO, FO ;

$\therefore$ sq. of $MD =$ sqs of PE, RF ;

$\therefore$ 4 times sq. of $MD =$ 4 times sqs of PE, RF ;

i. e. sq. of $MN =$ sqs of PQ, RS .

Hence MN, PQ, RS , can be sides of a right-angled Triangle.

Q. E. F.

65. (15)

Calling the angles $\frac{1}{x}, \frac{1}{y}, \frac{1}{z}$, of 360° , we must have

$$\frac{1}{x} + \frac{1}{y} + \frac{1}{z} = \frac{1}{2};$$

an Indeterminate Equation with 3 unknowns.

Evidently none of them can be so small as 2.

(1) Let $x = 3$; then $\frac{1}{y} + \frac{1}{z} = \frac{1}{6}$.

Now, if $\frac{1}{y} = \frac{k}{k+l} \times \frac{1}{6}$, $\frac{1}{z}$ will = $\frac{l}{k+l} \times \frac{1}{6}$:

hence k can only be 1, or 2, or 3, or 6; and the same is true of l .

(N.B. It is assumed that the fractions $\frac{k}{k+l}$, $\frac{l}{k+l}$, are in their lowest terms.)

Let $\frac{1}{y}$ be $\neq \frac{1}{z}$. Then $\frac{k}{k+l} \neq \frac{1}{2}$.

Then its possible values are $\frac{1}{2}$, so that $\frac{l}{k+l} = \frac{1}{2}$

$$\begin{array}{cc} \frac{2}{3}, & \frac{1}{3} \\ \frac{3}{4}, & \frac{1}{4} \\ \frac{3}{5} & \frac{2}{5} \\ \frac{6}{7}, & \frac{1}{7}. \end{array}$$

This gives 5 sets of values for $\frac{1}{x}$, $\frac{1}{y}$, $\frac{1}{z}$, viz.:

$$\frac{1}{3}, \frac{1}{12}, \frac{1}{12}; \quad \frac{1}{3}, \frac{1}{9}, \frac{1}{18}; \quad \frac{1}{3}, \frac{1}{8}, \frac{1}{24}; \quad \frac{1}{3}, \frac{1}{10}, \frac{1}{15}; \quad \frac{1}{3}, \frac{1}{7}, \frac{1}{42}.$$

(2) Let $x = 4$. Then $\frac{1}{y} + \frac{1}{z} = \frac{1}{4}$, and, as before, k can only be 1, or 2,

or 4, and the same is true of l . Hence the possible values for $\frac{k}{k+l}$ are $\frac{1}{2}$,

so that $\frac{l}{k+l} = \frac{1}{2}$

$$\begin{array}{cc} \frac{2}{3}, & \frac{1}{3} \\ \frac{4}{5}, & \frac{1}{5}. \end{array}$$

This gives 3 more sets of values for $\frac{1}{x}$, $\frac{1}{y}$, $\frac{1}{z}$, viz.:

$$\frac{1}{4}, \frac{1}{8}, \frac{1}{8}; \quad \frac{1}{4}, \frac{1}{6}, \frac{1}{12}; \quad \frac{1}{2}, \frac{1}{5}, \frac{1}{20}.$$

(3) Let $x = 5$; then $\frac{1}{y} + \frac{1}{z} = \frac{3}{10}$.

Hence denominator must contain factor "3", and k can be only 1, or 2, or 5, or 10; and the same is true of l .

Hence possible values of $\frac{k}{k+l}$ are $\frac{1}{2}$, so that $\frac{l}{k+l} = \frac{1}{2}$

$$\frac{2}{3}, \quad \frac{1}{3}$$

$$\frac{5}{6}, \quad \frac{1}{6}.$$

This gives 2 sets of values for $\frac{1}{x}, \frac{1}{y}, \frac{1}{z}$, viz.:—

$$\frac{1}{5}, \frac{1}{5}, \frac{1}{10}; \quad \frac{1}{5}, \frac{1}{4}, \frac{1}{20};$$

but the latter (a fact overlooked in thinking out) we have had already.

(4) Let $x = 6$; then $\frac{1}{y} + \frac{1}{z} = \frac{1}{3}$.

Hence k can be only 1, or 3, and the same is true of l .

Hence possible values for $\frac{k}{k+l}$ are $\frac{1}{2}$, so that $\frac{l}{k+l} = \frac{1}{2}$

$$\frac{3}{4}, \quad \frac{1}{4}.$$

This gives 2 sets of values, viz.:—

$$\frac{1}{6}, \frac{1}{6}, \frac{1}{6}; \quad \frac{1}{6}, \frac{1}{4}, \frac{1}{12};$$

but the latter (a fact overlooked in thinking out) we have had already.

There is no use in giving, to x , any values greater than 6; for these would make $\frac{1}{y} + \frac{1}{z} > \frac{1}{3}$; so that one or other must be $> \frac{1}{6}$; i. e. either y or z must < 6 , and we should get old values over again.

Hence there are 10 different shapes.

Q. E. F.

The 10 sets of angles (I am not certain that they were all thought out) are

(1)	120°,	30°,	30°;
(2)	120°,	40°,	20°;
(3)	120°,	45°,	15°;
(4)	120°,	36°,	24°;
(5)	120°	$51\frac{3}{7}^\circ$,	$8\frac{4}{7}^\circ$;
(6)	90°,	45°,	45°;
(7)	90°,	60°,	30°;
(8)	90°,	72°,	18°;
(9)	72°,	72°,	36°;
(10)	60°,	60°,	60°.

66. (15, 26)

Write k for $\frac{\alpha}{\alpha + \beta}$. Now the counters must be either both white, or one white and one black. Let chance of first condition be x ; hence chance of second is $(1 - x)$. Hence chance of drawing white is $x \times 1 + (1 - x) \times \frac{1}{2}$.

$$\begin{aligned}\therefore x + \frac{1 - x}{2} &= k; & \therefore x &= 2k - 1; \\ & & \therefore (1 - x) &= 2 - 2k.\end{aligned}$$

Let a counter now be drawn and prove white; then chance of 'observed event,' in 1st condition, is 1, and, in 2nd condition, $\frac{1}{2}$;

Hence the chances, of the existence of these two conditions, are proportional to $(2k - 1) \times 1$, $(2 - 2k) \times \frac{1}{2}$; i. e. are proportional to $2k - 1$, $1 - k$;

hence these chances actually are $\frac{2k-1}{k}, \frac{1-k}{k}$;

hence the chance of now drawing white,

$$\begin{aligned} & \text{is } \frac{2k-1}{k} \times 1 + \frac{1-k}{k} \times \frac{1}{2}; \\ & \text{i. e. } \frac{3k-1}{2k}. \end{aligned}$$

Hence the effect of *one* repetition of the experiment has been to change k into $\frac{3k-1}{2k}$.

Hence a second repetition of it will change

$$\frac{3k-1}{2k} \text{ into } \frac{3 \times \frac{3k-1}{2k} - 1}{2 \times \frac{3k-1}{2k}}; \text{ i. e. into } \frac{7k-3}{6k-2}.$$

We have now to discover the law (if there is one) for the series

$$k, \quad \frac{3k-1}{2k}, \quad \frac{7k-3}{6k-2},$$

regarding these as identical functions of 1, 2, 3.

We can write the 1st and 2nd term in the form of the 3rd, thus:—

$$\frac{k-0}{0 \times k - (-1)}, \quad \frac{3k-1}{2k-0}, \quad \frac{7k-3}{6k-2}$$

and, by inspection, we see that each is of the form

$$\frac{(2^n - 1) \times k - (2^{n-1} - 1)}{(2^n - 2) \times k - (2^{n-1} - 2)},$$

where n denotes the place of the term.

Suppose this law to hold for n terms, what will be the effect of repeating the experiment once more?

We know that it changes k into $\frac{3k-1}{2k}$. Hence the new chance will be

$$\frac{3 \times \frac{(2^n-1) \times k - (2^{n-1}-1)}{(2^n-2) \times k - (2^{n-1}-2)} - 1}{2 \times \frac{(2^n-1) \times k - (2^{n-1}-1)}{(2^n-2) \times k - (2^{n-1}-2)}};$$

i.e. $\frac{k \times (3 \cdot 2^n - 3 - 2^n + 2) - 3 \cdot 2^{n-1} + 3 + 2^{n-1} - 2}{(2^{n+1} - 2) \times k - (2^n - 2)};$

i.e. $\frac{(2^{n+1} - 1) \times k - (2^n - 1)}{(2^{n+1} - 2) \times k - (2^n - 2)};$

i. e. the $\overline{n+1}^{\text{th}}$ term of the series will follow the same law. But we know that the law holds for the 1st, 2nd, and 3rd terms. Hence it holds universally.

Hence, after $\overline{m}$ repetitions of the experiment, the chance of drawing white will be the $\overline{m+1}^{\text{th}}$ term of the above series; i. e. it will be

$$\frac{(2^{m+1} - 1) \times k - (2^m - 1)}{(2^{m+1} - 2) \times k - (2^m - 2)}.$$

Now, for k , write $\frac{\alpha}{\alpha + \beta}$.

Then chance is $\frac{(2^{m+1} - 1) \times \alpha - (2^m - 1) \cdot (\alpha + \beta)}{(2^{m+1} - 2) \times \alpha - (2^m - 2) \cdot (\alpha + \beta)};$

i.e. $\frac{(2^{m+1} - 2^m)\alpha - (2^m - 1) \cdot \beta}{(2^{m+1} - 2^m)\alpha - (2^m - 2) \cdot \beta};$

i.e. $\frac{2^m \cdot (\alpha - \beta) + \beta}{2^m \cdot (\alpha - \beta) + 2\beta}.$


Q. E. F.

EXAMPLE—Let chance be $\frac{9}{10}$; and then let experiment be repeated 5 times more.

Here $\alpha = 9, \beta = 1$;

$\therefore$ chance becomes $\frac{32 \times 8 + 1}{32 \times 8 + 2}$, i.e. $\frac{257}{258}$.

67. (16, 26)



Let $ABCD$ be the socket. Revolve the Tetrahedron until the plane, in it, DOA has taken the new position $D'QA'$; and let edge DA , in its new position, meet the socket-rim AC at R . From A' draw $A'L \perp XY$ -plane. Join OR , and produce it to L . And draw RM, LN , the y -ordinates of R and L .

Then co-ordinates of A' are ON, NL, LA' .

Call $OM, MR, 'x', y'$; and $OA, OR, OD, 'a, a', h'$; and $\angle XOR ' \theta'$.

It is evident that the vertical axis of the Tetrahedron always coincides with the Z -axis.

Hence A moves on the surface of a cylinder,

$$\text{i. e. } x^2 + y^2 = a^2 \quad (1)$$

Now $\angle XAC = 150^\circ$;

$$\therefore \text{Equation to } AC \text{ is } y = -\frac{1}{\sqrt{3}} \cdot (x - a);$$

$$\text{i. e. } x + \sqrt{3} \cdot y = a \quad (2)$$

Also Equation to OR is $\frac{x}{\cos \theta} = \frac{y}{\sin \theta} = a$;

$$\therefore, \text{ at } R, \frac{x'}{\cos \theta} = \frac{y'}{\sin \theta} = a'; \quad (3)$$

$$\therefore, \text{ by (2), } a' \cdot \cos \theta + \sqrt{3} \cdot a' \cdot \sin \theta = a;$$

$$\therefore a' = \frac{a}{\cos \theta + \sqrt{3} \cdot \sin \theta} \quad (4)$$

Also, by similar $\triangle$ s $D'QA'$, $D'OR$, $QA':QD':OR:OD'$

$$\text{i. e. } a:h:: \frac{a}{\cos \theta + \sqrt{3} \cdot \sin \theta} : h - z;$$

$$\therefore h - z = \frac{h}{\cos \theta + \sqrt{3} \cdot \sin \theta};$$

$$\text{but } \cos \theta = \frac{x}{a}, \text{ and } \sin \theta = \frac{y}{a};$$

$$\therefore h - z = \frac{ah}{x + \sqrt{3} \cdot y};$$

$$\text{i. e. } (x + \sqrt{3} \cdot y) \cdot (h - z) = ah \quad (5)$$

Equations (1) and (5) give the required Locus.

Q. E. F.

68. (16,26)

Let the Nos of bottles, taken out of the 3 days, be ' x, y, z '. Let each bottle have cost $10v$ pence, and therefore be sold for $11v$ pence.

Then the Treasurer's receipts, on the 3 days, were $(x - 1) \cdot 11v$, $y \cdot 11v - v$, $(z - 1) \cdot 11v - v$; yielding, as profits (i. e. as remainders after deducting cost-price of bottles taken out), $xv - 11v$, $yv - v$, $zv - 11v$. Then these 3 quantities are equal.

Hence $y = x - 10$, and $z = x + 1$;

$\therefore$ total No. of bottles, being $(x + y + z)$, $= 3x - 9$.

Now total profits are $(x + y + z) \cdot v - 24v$; i. e. $(3x - 33)v$;

$\therefore$ profit, per bottle $= \frac{(3x - 33) \cdot v}{3x - 9}$; and this must $= 6$;

$\therefore (x - 11) \cdot v = (x - 3) \cdot 6$.

Also $z \cdot 11v = 11 \times 240$; i. e. $(x + 1) \cdot 11v = 11 \times 240$;

$\therefore \frac{x - 11}{x + 1} = \frac{6 \cdot (x - 3)}{240}$;

$\therefore (x + 1) \cdot (x - 3) = 40 \cdot (x - 11)$;

$\therefore x^2 - 2x - 3 = 40x - 440$;

$\therefore x^2 - 42x + 437 = 0$.

Now $42^2 - 4 \times 437 = 1764 - 1748 = 16$;

$\therefore x = \frac{42 \pm 4}{2} = 23 \text{ or } 19$;

$\therefore$ No. of bottles $= 60 \text{ or } 48$; but it is a multiple of 5;

$\therefore$ it $= 60$.

Also $(x + 1) \cdot 11v = 11 \times 240$; i. e. $24v = 240$;

$\therefore v = 10$;

i. e. the wine was bought @ $8/4$ a bottle, and sold @ $9/2$ a bottle.

Q. E. F.

69. (17, 26)

§ 1. Let $\angle BAD = k.A$, $\angle CBE = l.B$, $\angle ACF = m.C$.

Then $\angle ABE = (1 - l).B$.

Now $\angle BC'D = \angle C'AB + \angle C'BA$.


$$\text{i.e. } k.A + (1 - l).B = C. \quad (1)$$

$$\text{Similarly, } l.B + (1 - m).C = A; \quad (2)$$

$$\text{and } m.C + (1 - k).A = B. \quad (3)$$

From equations (1) and (3), l and m may be found in terms of k : but these, taken along with k , will not be *similar* functions of the single variable k . We must have k a certain function of A, B, C , and θ (say); l a similar function of B, C, A , and θ ; and m a similar function of C, A, B , and θ ; i. e. we must have

$$\begin{aligned} k &= f(A, B, C, \theta), \\ l &= f(B, C, A, \theta), \\ m &= f(C, A, B, \theta). \end{aligned}$$



Now we know, by (1), that $kA - lB = C - B$;

$$\text{i. e. } A.f(A, B, C, \theta) - B.f(B, C, A, \theta) = C - B.$$

Now, as an experiment, let

$$\begin{aligned} k.A &= xA + yB + zC + \theta, \\ l.B &= xB + yC + zA + \theta; \end{aligned}$$

$$\text{then } kA - lB = (x - z).A + (y - x).B + (z - y).C;$$

$$\begin{aligned}\therefore x - z &= 0; \text{ i. e. } x = z; \\ z - y &= 1; \text{ i. e. } z = y + 1.\end{aligned}$$


These conditions will be fulfilled, if we make $y = 1$, and $x = z = 2$; so that

$$\begin{aligned}kA &= 2A + B + 2C + \theta, \\ lB &= 2B + C + 2A + \theta;\end{aligned}$$

which would make

$$f(A, B, C, \theta) \text{ mean } \frac{2A + B + 2C + \theta}{A}.$$

Now this may evidently be simplified by omitting $(A + B + C)$, which is constant; and we then have $k = \frac{A + C + \theta}{A}$; or, in a yet simpler form, by again subtracting 180° , $k = \frac{\theta - B}{A}$.



$$\begin{aligned}\text{Similarly } l &= \frac{\theta - C}{B}, \\ m &= \frac{\theta - A}{C}.\end{aligned}$$

Q. E. F.

§ 2. We see that $kA = \theta - B$, so that $\angle ADC$ is evidently equal to θ ; and so are $\angle s BEA, CFB$.

This gives us a geometrical construction, viz. to draw lines from A, B, C , so that each makes the same angle θ with the opposite side.

§ 3. Let us now ascertain the limits within which the value of θ must lie.

We know that $kA = \theta - B$.

Now $kA \not\geq A$; $\therefore \theta - B \not\geq A$; i. e. $\theta \not\geq A + B$;

i. e. $\theta \not\geq$ the supplement of C ;

and of course this is true for *each* of the three angles A, B, C ; i. e. if A, B, C , be the order of the angles in a descending order of magnitude, $\theta \not\geq$ supplement of A .

Again $kA \not\leq 0$.

Hence $\theta - B \not\leq 0$; i. e. $\theta \not\leq B$;

and of course this is true for *each* angle.

Hence if A, B, C , be the order in a descending order of magnitude, $\theta \not\leq A$, and $\not\leq 180^\circ - A$.

Q. E. F.

§ 4. We have now to ascertain the ratio which $B'C'$ bears to BC .

In Triangle ABC' , whose $\angle s$ are $(\theta - B)$, $(180^\circ - \theta - A)$, $(180^\circ - C)$, we have

$$AC' = \frac{AB}{\sin AC'B} \cdot \sin ABC' = \frac{c}{\sin C} \cdot \sin(\theta + A) = \frac{a}{\sin A} \cdot \sin(\theta + A);$$

$$BC' = \frac{AB}{\sin AC'B} \cdot \sin BAC' = \frac{c}{\sin C} \cdot \sin(\theta - B) = \frac{a}{\sin A} \cdot \sin(\theta - B)$$

$$\therefore, \text{ by symmetry, } AB' = \frac{a}{\sin A} \cdot \sin(\theta - A).$$

$$\text{Now } B'C' = AC' - AB';$$

$$\begin{aligned}\therefore \text{it} &= \frac{a}{\sin A} \{\sin(\theta + A) - \sin(\theta - A)\}, \\ &= \frac{a}{\sin A} \cdot 2 \cos \theta \sin A = a 2 \cos \theta.\end{aligned}$$

$$\text{Hence } \frac{a'}{a} = \frac{b'}{b} = \frac{c'}{c} = 2 \cos \theta.$$

Q. E. F.

70. (17, 27)

Before folding the Plane containing the Triangles, the locus of their vertices is evidently a Line parallel to their common base. Hence, if the base of the Tetrahedron = 1, we may imagine a slip of paper, whose width is $\frac{\sqrt{3}}{2}$, attached to the front facet of the Tetrahedron, and wrapped round towards the right; and the upper edge of this slip will evidently be the locus of the vertices. This slip may be conveniently regarded as divided into equilateral Triangles, placed base-downwards and base-upwards alternately, and it is evident that these Triangles will successively cover the facets of the Tetrahedron, in the order 'front, right side, base, left side, front, &c.'; and its *upper* edge, made up of the bases of the inverted constituent Triangles, will evidently run as follows. Calling the successive Triangles, after the first (which occupies the front facet of the Tetrahedron), ' α ' (base-up), ' β ' (base-down), ' γ ' (base-up), ' δ ' (base-down), ' ϵ ' (base-up), and so on, the locus consists of the bases of α , γ , &c. Now ' α ' will occupy the right facet, its base coinciding with the back-edge of the Tetrahedron; ' β ' will occupy the base of the Tetrahedron, its base coinciding with the front-edge; ' γ ' will occupy the left facet, its base coinciding with the back-edge; and so on. Hence the locus runs down the back-edge; up again; and so on. Which answers Question (1).

Q. E. F.

We may therefore, in answering the other three questions, consider the slip *before* it is folded, and calculate the positions of the vertices along its *upper* edge: and the problems thus become '*plane*' ones.

(2) Gives us a right-angled Triangle, whose left-hand base-angle is 15° , and whose altitude is $\frac{\sqrt{3}}{2}$. We must calculate its base, and then, deducting half the base of the initial Triangle (i. e. deducting $\frac{1}{2}$), we shall get the distance, measured along the upper edge of the slip, from the vertex of the initial Triangle to the vertex of the given Triangle; and from that we can calculate how many times we must go down and up the back-edge to reach it. Call the base of this right-angled Triangle ' x .' Then $\frac{\sqrt{3}}{2} \div x = \tan 15^\circ$.

$$\text{Now call } \tan 15^\circ \text{ 't'; then } \frac{2t}{(1-t^2)} = \tan 30^\circ = \frac{1}{\sqrt{3}};$$

$$\therefore 1 - t^2 = 2\sqrt{3}.t; \quad t^2 + 2\sqrt{3}.t - 1 = 0;$$

$$\therefore t = \frac{-2\sqrt{3} \pm 4}{2} = (\text{rejecting negative value}) 2 - \sqrt{3}.$$

$$\therefore x = \frac{\sqrt{3}}{2(2 - \sqrt{3})} = \frac{\sqrt{3}}{2}(2 + \sqrt{3}) = \sqrt{3} + \frac{3}{2}.$$

Deducting $\frac{1}{2}$, we get $(\sqrt{3} + 1)$ as the required distance.

Now $\sqrt{3} = 1.7$ &c.; $\therefore$ distance $= 2.7$ &c.

Hence we must go *down* back-edge, up again, and then about $.7$ down again. This answers question (2).

(3) We need to go down the back-edge, and up again; i. e. we must use up the upward bases of ' α ' and ' γ '. Hence the base of the required right-angled Triangle is $2\frac{1}{2}$. Hence the required left-hand base-angle is

$$\tan^{-1}\left(\frac{\sqrt{3}}{2} \div \frac{5}{2}\right); \text{ i. e. } \tan^{-1} \frac{\sqrt{3}}{5}.$$

Hence, for the required base-angle, we have $\frac{\sin}{\cos} = \frac{\sqrt{3}}{5}$;

$$\begin{aligned} \therefore \frac{\sin}{\sqrt{3}} &= \frac{\cos}{5} = \frac{1}{\sqrt{28}}; & \therefore \sin &= \sqrt{\frac{3}{28}} = \frac{\sqrt{84}}{28}, \\ &= \frac{\text{rather over } 9}{28}; & & \frac{7 \mid 9.}{\underline{4 \mid 1 \cdot 28 \text{ \&c.} } \\ & & & \cdot 32 \text{ \&c.} \end{aligned}$$

$$\begin{aligned} \text{Now (by mem. tech.) } \sin^{-1} \cdot 3 &= 17 \cdot 45 \text{ \&c.}^\circ. \\ \sin^{-1} \cdot 4 &= 23 \cdot 57 \text{ \&c.}^\circ. \end{aligned}$$

and the required angle is about $\frac{1}{5}$ of the way from one to the other. But the difference is almost exactly 6° . Hence we must add, to the lesser, about $1 \frac{1}{5}$ degrees, or $1 \cdot 20^\circ$. And the total will be about $18 \cdot 65^\circ$.

(4) Here the right-angled Triangle has, for its base, $3\frac{1}{2}$.

$\therefore$ the required base-angle has, for its tangent,


$$\left(\frac{\sqrt{3}}{2} \div \frac{7}{2} \right); \text{ i. e. } \frac{\sqrt{3}}{7};$$

$$\begin{aligned} \therefore \frac{\sin}{\sqrt{3}} &= \frac{\cos}{7} = \frac{1}{\sqrt{52}}; & \therefore \sin &= \sqrt{\frac{3}{52}} = \text{nearly } \sqrt{\frac{1}{17}}, \\ &= \text{nearly } \frac{\sqrt{17}}{17}. & \text{Now } \sqrt{17} &= 4 \cdot 12 \text{ \&c. } \therefore \sin = \cdot 24 \text{ \&c.} \end{aligned}$$

Now $\sin^{-1} \cdot 2 = 11 \cdot 53 \text{ \&c.}^\circ$; and we must go about half-way to the next angle, viz. $17 \cdot 45 \text{ \&c.}^\circ$. The difference is about 6° ; $\therefore$ we must add about 3° . Hence the answer is about $14 \cdot 53^\circ$.

71. (18)

Let ABC be the given Triangle, and P the given Point.



Bisect the sides of ABC at D, E, F ; and join these Points.

First, let P be within the Triangle DEF .

Draw HG parallel to BC , so that its distance from BC may be double the distance of P from BC ; join GP, HP , and produce them to meet BC in L, M . From L draw LK parallel to AC ; join KP , and produce it to meet AC at N : join MN .

Because HG is parallel to LM ,

$\therefore GP = PL$, and $HP = PM$;

$\therefore KL$ is parallel to GN , and that $LP = PG$,

$\therefore KP = PN$; $\therefore MN$ is parallel to HK .

Now the Triangles PGH, PLM , are equal in all respects;

$\therefore GH = LM$. Similarly $KL = GN$, and $MN = HK$.

If P lies on FE , HG and LM vanish, and the Hexagon becomes a Parallelogram.

If P lies at D , the Hexagon becomes the line BC .

If P lies outside the Triangle DEF , the Problem is insoluble.

Q. E. F.

We know that, if a bag contained 3 counters, 2 being black and one white, the chance of drawing a black one would be $\frac{2}{3}$; and that any *other* state of things would *not* give this chance.

Now the chances, that the given bag contains (α) BB , (β) BW , (γ) WW , are respectively $\frac{1}{4}$, $\frac{1}{2}$, $\frac{1}{4}$.

Add a black counter.

Then the chances, that it contains (α) BBB , (β) BWB , (γ) WWB , are, as before, $\frac{1}{4}$, $\frac{1}{2}$, $\frac{1}{4}$.

Hence the chance, of now drawing a black one,

$$= \frac{1}{4} \cdot 1 + \frac{1}{2} \cdot \frac{2}{3} + \frac{1}{4} \cdot \frac{1}{3} = \frac{2}{3}.$$

Hence the bag now contains BBW (since any *other* state of things would *not* give this chance).

Hence, before the black counter was added, it contained BW , i. e. one black counter and one white.

Q. E. F.

THE END.

WORKS BY C. L. DODGSON.

PUBLISHED BY
MACMILLAN & CO., LONDON.

THE FORMULÆ OF PLANE TRIGONOMETRY, printed with symbols, instead of words, to express the goniometrical ratios. (First published in 1861.) Crown 4to, sewed. Price 1s.

EUCLID AND HIS MODERN RIVALS. (First published in 1879.)
Second Edition, 1885. Crown 8vo, cloth. Price 6s.

SUPPLEMENT TO FIRST EDITION OF "EUCLID AND HIS MODERN RIVALS," containing a Notice of Henrici's Geometry, together with Selections from the Reviews. (First published in 1885.)
Crown 8vo, sewed. Price 1s.

EUCLID, BOOKS I, II, with Notes. (First published in 1882.) Sixth Edition, 1888. Crown 8vo, cloth. Price 2s.

CURIOSA MATHEMATICA.

Part I. A New Theory of Parallels. (First published in 1888.) Fourth Edition, 1895. Crown 8vo, cloth. Price 2s.

Part II. Pillow-Problems, thought out during Wakeful Hours. (First published in 1893.) Fourth Edition, 1895. Crown 8vo, cloth. Price 2s.

TRANSCRIBER'S NOTES

For the sake of clarity, the notation used by the author for $\cap$ and $\supset$ has been replaced with modern terminology.

Due to a lack of distinction between 'a' and ' α ' in the printed version, certain expressions have been corrected to avoid potential mathematical errors.

*** END OF THE PROJECT GUTENBERG EBOOK CURIOSA
MATHEMATICA, PART II ***

Updated editions will replace the previous one—the old editions will be renamed.

Creating the works from print editions not protected by U.S. copyright law means that no one owns a United States copyright in these works, so the Foundation (and you!) can copy and distribute it in the United States without permission and without paying copyright royalties. Special rules, set forth in the General Terms of Use part of this license, apply to copying and distributing Project Gutenberg™ electronic works to protect the PROJECT GUTENBERG™ concept and trademark. Project Gutenberg is a registered trademark, and may not be used if you charge for an eBook, except by following the terms of the trademark license, including paying royalties for use of the Project Gutenberg trademark. If you do not charge anything for copies of this eBook, complying with the trademark license is very easy. You may use this eBook for nearly any purpose such as creation of derivative works, reports, performances and research. Project Gutenberg eBooks may be modified and printed and given away—you may do practically ANYTHING in the United States with eBooks not protected by U.S. copyright law. Redistribution is subject to the trademark license, especially commercial redistribution.

START: FULL LICENSE

THE FULL PROJECT GUTENBERG™ LICENSE

PLEASE READ THIS BEFORE YOU DISTRIBUTE OR USE THIS WORK

To protect the Project Gutenberg™ mission of promoting the free distribution of electronic works, by using or distributing this work (or any other work associated in any way with the phrase “Project Gutenberg”), you agree to comply with all the terms of the Full Project Gutenberg License available with this file or online at www.gutenberg.org/license.

Section 1. General Terms of Use and Redistributing Project Gutenberg electronic works

1.A. By reading or using any part of this Project Gutenberg electronic work, you indicate that you have read, understand, agree to and accept all the terms of this license and intellectual property (trademark/copyright) agreement. If you do not agree to abide by all the terms of this agreement, you must cease using and return or destroy all copies of Project Gutenberg electronic works in your possession. If you paid a fee for obtaining a copy of or access to a Project Gutenberg electronic work and you do not agree to be bound by the terms of this agreement, you may obtain a refund from the person or entity to whom you paid the fee as set forth in paragraph 1.E.8.

1.B. “Project Gutenberg” is a registered trademark. It may only be used on or associated in any way with an electronic work by people who agree to be bound by the terms of this agreement. There are a few things that you can do with most Project Gutenberg electronic works even without complying with the full terms of this agreement. See paragraph 1.C below. There are a lot of things you can do with Project Gutenberg electronic works if you follow the terms of this agreement and help preserve free future access to Project Gutenberg electronic works. See paragraph 1.E below.

1.C. The Project Gutenberg Literary Archive Foundation (“the Foundation” or PGLAF), owns a compilation copyright in the collection of Project Gutenberg electronic works. Nearly all the individual works in the collection are in the public domain in the United States. If an individual work is unprotected by copyright law in the United States and you are

located in the United States, we do not claim a right to prevent you from copying, distributing, performing, displaying or creating derivative works based on the work as long as all references to Project Gutenberg are removed. Of course, we hope that you will support the Project Gutenberg mission of promoting free access to electronic works by freely sharing Project Gutenberg works in compliance with the terms of this agreement for keeping the Project Gutenberg name associated with the work. You can easily comply with the terms of this agreement by keeping this work in the same format with its attached full Project Gutenberg License when you share it without charge with others.

1.D. The copyright laws of the place where you are located also govern what you can do with this work. Copyright laws in most countries are in a constant state of change. If you are outside the United States, check the laws of your country in addition to the terms of this agreement before downloading, copying, displaying, performing, distributing or creating derivative works based on this work or any other Project Gutenberg work. The Foundation makes no representations concerning the copyright status of any work in any country other than the United States.

1.E. Unless you have removed all references to Project Gutenberg:

1.E.1. The following sentence, with active links to, or other immediate access to, the full Project Gutenberg License must appear prominently whenever any copy of a Project Gutenberg work (any work on which the phrase “Project Gutenberg” appears, or with which the phrase “Project Gutenberg” is associated) is accessed, displayed, performed, viewed, copied or distributed:

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg™ License included with this eBook or online at www.gutenberg.org. If you are not located in the United States, you will have to check the laws of the country where you are located before using this eBook.

1.E.2. If an individual Project Gutenberg electronic work is derived from texts not protected by U.S. copyright law (does not contain a notice indicating that it is posted with permission of the copyright holder), the work can be copied and distributed to anyone in the United States without paying any fees or charges. If you are redistributing or providing access to a work with the phrase “Project Gutenberg” associated with or appearing on the work, you must comply either with the requirements of paragraphs 1.E.1 through 1.E.7 or obtain permission for the use of the work and the Project Gutenberg trademark as set forth in paragraphs 1.E.8 or 1.E.9.

1.E.3. If an individual Project Gutenberg electronic work is posted with the permission of the copyright holder, your use and distribution must comply with both paragraphs 1.E.1 through 1.E.7 and any additional terms imposed by the copyright holder. Additional terms will be linked to the Project Gutenberg License for all works posted with the permission of the copyright holder found at the beginning of this work.

1.E.4. Do not unlink or detach or remove the full Project Gutenberg License terms from this work, or any files containing a part of this work or any other work associated with Project Gutenberg.

1.E.5. Do not copy, display, perform, distribute or redistribute this electronic work, or any part of this electronic work, without prominently displaying the sentence set forth in paragraph 1.E.1 with active links or immediate access to the full terms of the Project Gutenberg License.

1.E.6. You may convert to and distribute this work in any binary, compressed, marked up, nonproprietary or proprietary form, including any word processing or hypertext form. However, if you provide access to or distribute copies of a Project Gutenberg work in a format other than “Plain Vanilla ASCII” or other format used in the official version posted on the official Project Gutenberg website (www.gutenberg.org), you must, at no additional cost, fee or expense to the user, provide a copy, a means of exporting a copy, or a means of obtaining a copy upon request, of the work in its original “Plain Vanilla ASCII” or other form. Any alternate format must include the full Project Gutenberg License as specified in paragraph 1.E.1.

1.E.7. Do not charge a fee for access to, viewing, displaying, performing, copying or distributing any Project Gutenberg works unless you comply with paragraph 1.E.8 or 1.E.9.

1.E.8. You may charge a reasonable fee for copies of or providing access to or distributing Project Gutenberg electronic works provided that:

- You pay a royalty fee of 20% of the gross profits you derive from the use of Project Gutenberg works calculated using the method you already use to calculate your applicable taxes. The fee is owed to the owner of the Project Gutenberg trademark, but he has agreed to donate royalties under this paragraph to the Project Gutenberg Literary Archive Foundation. Royalty payments must be paid within 60 days following each date on which you prepare (or are legally required to prepare) your periodic tax returns. Royalty payments should be clearly marked as such and sent to the Project Gutenberg Literary Archive Foundation at the address specified in Section 4, "Information about donations to the Project Gutenberg Literary Archive Foundation."
- You provide a full refund of any money paid by a user who notifies you in writing (or by e-mail) within 30 days of receipt that s/he does not agree to the terms of the full Project Gutenberg™ License. You must require such a user to return or destroy all copies of the works possessed in a physical medium and discontinue all use of and all access to other copies of Project Gutenberg™ works.
- You provide, in accordance with paragraph 1.F.3, a full refund of any money paid for a work or a replacement copy, if a defect in the electronic work is discovered and reported to you within 90 days of receipt of the work.
- You comply with all other terms of this agreement for free distribution of Project Gutenberg™ works.

1.E.9. If you wish to charge a fee or distribute a Project Gutenberg™ electronic work or group of works on different terms than are set forth in this agreement, you must obtain permission in writing from the Project Gutenberg Literary Archive Foundation, the manager of the Project

Gutenberg™ trademark. Contact the Foundation as set forth in Section 3 below.

1.F.

1.F.1. Project Gutenberg volunteers and employees expend considerable effort to identify, do copyright research on, transcribe and proofread works not protected by U.S. copyright law in creating the Project Gutenberg™ collection. Despite these efforts, Project Gutenberg™ electronic works, and the medium on which they may be stored, may contain “Defects,” such as, but not limited to, incomplete, inaccurate or corrupt data, transcription errors, a copyright or other intellectual property infringement, a defective or damaged disk or other medium, a computer virus, or computer codes that damage or cannot be read by your equipment.

1.F.2. LIMITED WARRANTY, DISCLAIMER OF DAMAGES - Except for the “Right of Replacement or Refund” described in paragraph 1.F.3, the Project Gutenberg Literary Archive Foundation, the owner of the Project Gutenberg™ trademark, and any other party distributing a Project Gutenberg™ electronic work under this agreement, disclaim all liability to you for damages, costs and expenses, including legal fees. YOU AGREE THAT YOU HAVE NO REMEDIES FOR NEGLIGENCE, STRICT LIABILITY, BREACH OF WARRANTY OR BREACH OF CONTRACT EXCEPT THOSE PROVIDED IN PARAGRAPH 1.F.3. YOU AGREE THAT THE FOUNDATION, THE TRADEMARK OWNER, AND ANY DISTRIBUTOR UNDER THIS AGREEMENT WILL NOT BE LIABLE TO YOU FOR ACTUAL, DIRECT, INDIRECT, CONSEQUENTIAL, PUNITIVE OR INCIDENTAL DAMAGES EVEN IF YOU GIVE NOTICE OF THE POSSIBILITY OF SUCH DAMAGE.

1.F.3. LIMITED RIGHT OF REPLACEMENT OR REFUND - If you discover a defect in this electronic work within 90 days of receiving it, you can receive a refund of the money (if any) you paid for it by sending a written explanation to the person you received the work from. If you received the work on a physical medium, you must return the medium with your written explanation. The person or entity that provided you with the defective work may elect to provide a replacement copy in lieu of a refund. If you received the work electronically, the person or entity providing it to

you may choose to give you a second opportunity to receive the work electronically in lieu of a refund. If the second copy is also defective, you may demand a refund in writing without further opportunities to fix the problem.

1.F.4. Except for the limited right of replacement or refund set forth in paragraph 1.F.3, this work is provided to you 'AS-IS', WITH NO OTHER WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR ANY PURPOSE.

1.F.5. Some states do not allow disclaimers of certain implied warranties or the exclusion or limitation of certain types of damages. If any disclaimer or limitation set forth in this agreement violates the law of the state applicable to this agreement, the agreement shall be interpreted to make the maximum disclaimer or limitation permitted by the applicable state law. The invalidity or unenforceability of any provision of this agreement shall not void the remaining provisions.

1.F.6. INDEMNITY - You agree to indemnify and hold the Foundation, the trademark owner, any agent or employee of the Foundation, anyone providing copies of Project Gutenberg™ electronic works in accordance with this agreement, and any volunteers associated with the production, promotion and distribution of Project Gutenberg™ electronic works, harmless from all liability, costs and expenses, including legal fees, that arise directly or indirectly from any of the following which you do or cause to occur: (a) distribution of this or any Project Gutenberg work, (b) alteration, modification, or additions or deletions to any Project Gutenberg work, and (c) any Defect you cause.

Section 2. Information about the Mission of Project Gutenberg

Project Gutenberg is synonymous with the free distribution of electronic works in formats readable by the widest variety of computers including obsolete, old, middle-aged and new computers. It exists because of the

efforts of hundreds of volunteers and donations from people in all walks of life.

Volunteers and financial support to provide volunteers with the assistance they need are critical to reaching Project Gutenberg's goals and ensuring that the Project Gutenberg collection will remain freely available for generations to come. In 2001, the Project Gutenberg Literary Archive Foundation was created to provide a secure and permanent future for Project Gutenberg and future generations. To learn more about the Project Gutenberg Literary Archive Foundation and how your efforts and donations can help, see Sections 3 and 4 and the Foundation information page at www.gutenberg.org.

Section 3. Information about the Project Gutenberg Literary Archive Foundation

The Project Gutenberg Literary Archive Foundation is a non-profit 501(c)(3) educational corporation organized under the laws of the state of Mississippi and granted tax exempt status by the Internal Revenue Service. The Foundation's EIN or federal tax identification number is 64-6221541. Contributions to the Project Gutenberg Literary Archive Foundation are tax deductible to the full extent permitted by U.S. federal laws and your state's laws.

The Foundation's business office is located at 41 Watchung Plaza #516, Montclair NJ 07042, USA, +1 (862) 621-9288. Email contact links and up to date contact information can be found at the Foundation's website and official page at www.gutenberg.org/contact

Section 4. Information about Donations to the Project Gutenberg Literary Archive Foundation

Project Gutenberg™ depends upon and cannot survive without widespread public support and donations to carry out its mission of increasing the number of public domain and licensed works that can be freely distributed in machine-readable form accessible by the widest array of equipment

including outdated equipment. Many small donations (\$1 to \$5,000) are particularly important to maintaining tax exempt status with the IRS.

The Foundation is committed to complying with the laws regulating charities and charitable donations in all 50 states of the United States. Compliance requirements are not uniform and it takes a considerable effort, much paperwork and many fees to meet and keep up with these requirements. We do not solicit donations in locations where we have not received written confirmation of compliance. To SEND DONATIONS or determine the status of compliance for any particular state visit www.gutenberg.org/donate.

While we cannot and do not solicit contributions from states where we have not met the solicitation requirements, we know of no prohibition against accepting unsolicited donations from donors in such states who approach us with offers to donate.

International donations are gratefully accepted, but we cannot make any statements concerning tax treatment of donations received from outside the United States. U.S. laws alone swamp our small staff.

Please check the Project Gutenberg web pages for current donation methods and addresses. Donations are accepted in a number of other ways including checks, online payments and credit card donations. To donate, please visit: www.gutenberg.org/donate.

Section 5. General Information About Project Gutenberg electronic works

Professor Michael S. Hart was the originator of the Project Gutenberg concept of a library of electronic works that could be freely shared with anyone. For forty years, he produced and distributed Project Gutenberg eBooks with only a loose network of volunteer support.

Project Gutenberg eBooks are often created from several printed editions, all of which are confirmed as not protected by copyright in the U.S. unless a

copyright notice is included. Thus, we do not necessarily keep eBooks in compliance with any particular paper edition.

Most people start at our website which has the main PG search facility:
www.gutenberg.org.

This website includes information about Project Gutenberg, including how to make donations to the Project Gutenberg Literary Archive Foundation, how to help produce our new eBooks, and how to subscribe to our email newsletter to hear about new eBooks.