Drug A / Drug B Dosing in a 96-well Plate

Fictional demo protocol for Glassistant testing. Not a validated lab method. Uses no real samples or data.

Purpose

Compare the response of adherent cells to Drug A and Drug B against a vehicle control after a short incubation.

Experimental design

Group	Treatment	Replicates
Control	Vehicle (PBS)	5
Drug A	Drug A, 10 µM working solution	5
Drug B	Drug B, 10 µM working solution	5

Use one row of the plate per group. Leave the outer wells empty to avoid edge effects. Keep two spare wells per group in case a well has to be redone.

Materials

- 96-well flat-bottom plate, cells seeded the day before
- Culture medium (pre-warmed to 37 °C)
- Vehicle (PBS), Drug A and Drug B working solutions
- Single-channel P200 and P20 pipettes, sterile tips

Procedure

1. Add medium. Add 100 µL of fresh medium to every well that will be used. *Tip: dispense slowly against the well wall so the cell layer is not disturbed.*

2. Add vehicle. Add 10 µL of vehicle (PBS) to each Control well.

3. Add Drug A. Add 10 µL of Drug A working solution to each Drug A well. *Tip: do not pipette up and down after adding the drug.*

4. Add Drug B. Add 10 µL of Drug B working solution to each Drug B well.

5. Incubate. Incubate the plate for 30 min at 37 °C before reading.

Notes

Record any well that is mis-dosed, has bubbles, or looks damaged, and exclude it from analysis. Each group needs at least 5 usable wells.