TABLE S1

Outcome classification by interview year and prediction horizon. Counts are mutually exclusive at the main category level. Indented rows are subsets, not additional participants. Death before the final wave excludes a participant regardless of prior dementia. Unknown dementia or vital status is excluded. Positive and negative correspond to outcome Y equal to 1 and 0, respectively.

Classification / outcome rule	**Outcome status**	**2000: 6-year (2000–2006)**	**2000: 8-year (2000–2008)**	**2010: 6-year (2010–2016)**	**2010: 8-year (2010–2018)**
Eligible sample size at baseline	Baseline	17,336	17,336	19,243	19,243
Incident dementia: dementia at any follow-up assessment and alive	Positive	1,190	1,313	1,267	1,304
Missing intermediate assessment, dementia identified later	Positive	55	113	67	90
Adequately observed non-dementia: confirmed dementia-free at final wave	Negative	12,065	10,927	12,960	10,564
Missing assessments, subsequently confirmed dementia-free at final assessment	Negative	407	580	464	478
Loss to follow-up: alive or vital status unknown, not assessed at final wave	Excluded	1,212	1,228	2,191	3,578
Death: died before final wave, any dementia history	Excluded	2,869	3,868	2,825	3,797
Died, no dementia observed	Excluded	2,467	3,167	2,398	3,112
Died, dementia observed	Excluded	402	701	427	685
Analytic sample size	Positive or negative	13,255	12,240	14,227	11,868