

Q1B Brainstorming

Jonah Cooke SID: 111059487

January 1, 2026

What is known?

The EOM:

$$\ddot{z} = \frac{a(1 - \dot{z}^2)}{az + 1} \quad (1)$$

A solution:

$$z(t) = \frac{\sqrt{a^2 t^2 + 1} - 1}{a} \quad (2)$$

The Lagrangian:

$$L = -m(z)\sqrt{1 - \dot{z}^2}, \quad L = -\frac{m_0}{az + 1}\sqrt{1 - \dot{z}^2} \quad (3)$$

The action:

$$S[z] = -\int_{z_i}^{z_f} \frac{m_0}{az + 1} \sqrt{1 - \dot{z}^2} dz \quad (4)$$

Questions

Question: Can there be multiple solutions to the EOM?

Thought: yes so long as the solution shares the same endpoints and extremizes the action of the system, namely so long as it is identical to the current solution within the region being extremized by the current solution, it is also a solution.

Thought: Or if there are multiple paths that extremize the action between the two endpoints.

Study: I need to research uniqueness theorems for EOMs derived from lagrangians.

Question: What are the ways in which this can be exploited to find new solutions, or how does it simplify the problem?

Thought: If there is only one extrema to the action between the two endpoints why would we care about multiple solutions as it only makes sense to care about the function within the region being analyzed,

Thought: If it is shown later that a particular form of the solution is easier to work with, it would then be better to have said solution.