

Environmental Reporting BC Enhancements – Methods for SO₂ and NO₂

METRIC VALUES

SO₂

- (i) The 3-year average of the annual 99th percentile of the SO₂ daily maximum 1-hour average concentrations
- (ii) The arithmetic average over a single calendar year of all 1-hour average SO₂ concentrations

NO₂

- (i) The 3-year average of the annual 98th percentile of the NO₂ daily maximum 1-hour average concentrations
- (ii) The arithmetic average over a single calendar year of all 1-hour average NO₂ concentrations

METRIC CALCULATIONS

1. 3-year average of the annual 98th/99th percentile of the daily maximum 1-hour average concentrations

1.1 Calculating the Daily 1-hour Maximum Concentration

The daily 1-hour maximum concentration refers to the maximum 1-hour value reported over a 24-hour period (midnight to midnight, local standard time). The value is to be reported to the nearest 0.1 ppb.

1.2 Calculating the Annual 98th or 99th Percentile Value

The 98th percentile is the daily value out of a year of monitoring data for which 98 percent of all values are less than or equal to this level, and 2% are over. The 99th percentile is the daily value out of a year of monitoring data for which 99 percent of all values are less than or equal to this level, and 1% are over. Annual 98th and 99th percentiles are to be reported to the nearest 0.1 ppb.

The annual 98th percentile (98P) is determined and defined as follows:

- Sort all the daily 1-hour maximum concentration values for the given year into an array of numbers ordered from highest to lowest
- Repeat equal values as many times as they occur.
- Calculate the number "id" defined as,
 - id = 0.98*n (the product of 0.98 and n), where
 - i = the integer part of the number
 - d = the decimal part of the number
 - n = total number of the daily 24-hour concentration values
- The annual 98P is then defined to be the (n-i)th largest value.
- Data are reported to the nearest 0.1 ppb.

A similar procedure is used for the calculation of the 99th percentile, substituting id=0.99*n.

The table below describes the rank of the 98th and 99th percentile values for a given number of days.

Rank equivalent to 98th and 99th percentile values over range of valid days of reporting (data sorted in descending order).

No. Valid Days	Rank Equivalent to Annual 98 th Percentile (NO ₂)	Rank Equivalent to Annual 99 th Percentile (SO ₂)
0-50	1	1
51-100	2	1
101-150	3	2
151-200	4	2
201-250	5	3
251-300	6	3
301-350	7	4
351-366	8	4

**** Note:** A similar procedure has been developed for PM_{2.5} in the `rcaqs` package (`pm_98_percentile` and `quantile2` functions) and should be a good starting point.

1.3 Calculating the 3-year Average Annual 98th or 99th Percentile Value

The Annual 98th or 99th Percentile Value is repeated for each year of a 3-year period and averaged (e.g. $99P_1 + 99P_2 + 99P_3/3$). The 3-year average annual 98th or 99th percentile values are reported to the nearest integer.

2. Annual average over a single calendar year of all 1-hour average concentrations

An annual average value reflects the average of all hourly values. The annual average concentrations should be reported to the nearest 0.1 ppb.

Intermediate calculations should retain all available digits and decimal places.

DATA COMPLETENESS CRITERIA

1. Data completeness criteria for the 3-year average of the annual 98th/99th percentile of the daily maximum 1-hour average concentrations

1.1. For the purpose of determining a daily 1-hour maximum concentration, a valid daily value is calculated where at least 18 hourly measurements are available in a day.

1.2. For the purpose of determining an annual 98th/99th percentile an annual data set should be considered complete if there are available at least 75% of all daily-maximum 1-hour measurements AND at least 60% of all daily-maximum 1-hour measurements in each calendar quarter (Q1 Jan-Mar, Q2 Apr-Jun, Q3 Jul-Sept, Q4 Oct-Dec).

1.3 For the purpose of determining a 3-year average of the annual 98th/99th percentile of the daily maximum 1-hour average concentrations, 2 of the 3 years must be available.

**** Note:** some of the above functionality is already implemented in `rcaqs` for PM_{2.5}, SO₂, and NO₂, and should be a good starting point for the above.

2. Data completeness criteria for the annual average over a single calendar year of all 1-hour average concentrations

2.1. An annual data set should be considered complete if there are available at least 75% of all hourly measurements in a year AND at least 60% of all hourly measurements in each calendar quarter (Q1 Jan-Mar, Q2 Apr-Jun, Q3 Jul-Sept, Q4 Oct-Dec).