

# Next generation platform for implementing fast gates in ion trap quantum computation

D. Webb S. Saner O. Bazavan M. Minder C.J. Ballance

Ion Trap Quantum Computing Group, Department of Physics, University of Oxford

Scalable trapped-ion quantum computation relies on the development of high-fidelity fast entangling gates in a many ion crystal. Conventional geometric phase gates either suffer from scattering errors or off-resonant carrier excitations. A potential route to achieve fast entanglement is creating a standing wave which can suppress the unwanted carrier coupling.

## Non-Adiabatic Mølmer Sørensen Gates

- Two qubit gates are implemented by coupling spin with shared motion of the ions. XXX Figure of geometric phase gate. XXX
- Fast entangling gates enable deeper quantum computational circuit depths for given level of incoherent error.
- But going fast excites multiple motional modes ("spectator modes") which can introduces errors. XXX Figure of multiple motions. XXX

- Mølmer Sørensen (MS) interaction is common two-qubit gate which requires a bichromatic field incident on the ions. Using travelling waves gives the Hamiltonian

$$\hat{H}_{MS-TW} = \hbar\Omega\hat{S}_{\phi-\pi/2}\cos(\delta t) + \hbar\Omega\eta\hat{S}_{\phi}\cos(\delta t)(\hat{a}e^{-i\omega_z t} + \hat{a}^\dagger e^{i\omega_z t})$$



with the first term being the carrier whilst the second is the desired coupling. XXX Figure of TW MS.

XXX

- As these terms do not commute, in the ??interaction picture?? this Hamiltonian may be expressed as [Xref Canzz?X]:

$$\hat{H}_{MS-TW} = \hbar\eta\Omega(J_0(2\Omega/\delta) + J_2(2\Omega/\delta)) \cdot \cos(\delta t)\hat{S}_{\phi}(\hat{a}e^{-i\omega_z t} + \hat{a}^\dagger e^{i\omega_z t})$$

- Carrier term:** causes the spin dependent force coupling to be modulated by (J0+J2).
- Spectator excitation:** Amplitude shaped pulses [ref vera] effectively remove "spectator" error by closing phase loops of all excited modes.




## Standing Wave Single and MS Gates

Bichromatic standing wave Hamiltonian where ions are separated by  $n\lambda/2$ :

$$\hat{H}_{MS-SW} = \hbar 2\Omega\hat{S}_{\phi}\cos(\delta t)\sin(\Delta\phi/2) + \hbar 2\Omega\eta\hat{S}_{\phi}\cos(\delta t)(\hat{a}e^{-i\omega_z t} + \hat{a}^\dagger e^{i\omega_z t})\cos(\Delta\phi/2)$$


XXX Figure of SW MS (1b). XXX

- Setting  $\Delta\phi = 0$  (ions sitting at antinodes) we suppress the carrier term and maximise sideband coupling.
- Enables fast gates by preventing the saturation effect seen in the travelling MS.
- Using optical lattice gives complete control over phase visible to ions.



Results from Cnulling:

## Phase Stabilization



- Optical lattice requires phase stabilisation to perform coherent interactions.
- Phase stabilised passively using enclosure and actively by two-step process:
  - 1) Fast drifts removed utilising interference of light from the two branches on a photodiode (PD).
  - 2) As PD lockpoint is 30 cm away from ions, a second feedback loop using the ion as a sensor is required. We do this by performing a  $\pi/2$ -pulse using arm 1 ( $\pi/2$ , b1) followed immediately by a  $\pi/2$ -pulse using arm 2 ( $\pi/2$ , b2). This is equivalent to a zero-delay Ramsey sequence, which gives a signal sensitive on the difference in phase between the two pulses, hence the relative phase between the branches.
- No detriment using lattice: Standard Randomized Benchmarking gave quality of our single qubit rotations per gate to be  $\epsilon = 0.173(3)\%$  for a travelling wave and  $\epsilon = 0.144(3)\%$  for the standing wave.

## New Platform

- New ion trap experiment in development for exploring fast gate regime.
- Two high NA lenses allow optical access on both faces of the trap to create an array of singly addressing standing waves.
- 3D segmented trap design from NPL facilitates low heating rates whilst enabling ion shuttling and crystal rotations.
- Calcium 40
- quadrupole transition used.
- MuMetal shield and permanent magnets.
- fig: solidworks of new experiment

