

Question				Marking details	Marks available				Maths	Prac
					AO1	AO2	AO3	Total		
9	(a)	(i)		Use of intensity = $\frac{P}{4\pi R^2}$ or correct substitution (1) Intensity = $1\,415\text{ W m}^{-2}$ (1)	1	1		2	2	
		(ii)		[Using $1\,400\text{ W m}^{-2}$]: $20\% = 280\text{ W m}^{-2}$ $15\% = 42\text{ W m}^{-2}$ (1) for both efficiency calculations Area = $\frac{8 \times 10^9}{42}$ (ecf) = $1.9 \times 10^8\text{ m}^2$ (1) (unit) Accept 190 km^2 or equivalent		2		2	2	
		(iii)		Greater intensity of radiation in Portugal (accept increased/longer sunshine hours throughout year) Do not accept more efficient			1	1		
	(b)	(i)		$\lambda_{\text{max}}T = \text{constant}$	1			1		
		(ii)		$\lambda_{\text{max}} = \frac{0.0029}{290}$ (1) $\lambda_{\text{max}} = 10 \times 10^{-6}\text{ [m]}$ so in IR (1)		2		2	2	
		(iii)	I	Surface emits IR in region $5\text{ }\mu\text{m}$ to $20\text{ }\mu\text{m}$ (1) [Partially] absorbed by water vapour (and CO_2) (1) Correct reference to numerical data from graphs e.g. either water: greatest absorption between $5\text{ }\mu\text{m}$ to $8\text{ }\mu\text{m}$ or $12\text{ }\mu\text{m}$ to $20\text{ }\mu\text{m}$ or CO_2 : (greatest absorption between $10\text{ }\mu\text{m}$ to $20\text{ }\mu\text{m}$) (1) [IR] re-emitted in all directions (1)	1 1		1 1	4		
			II	Higher temperatures will lead to increased water vapour in atmosphere (1) Leading to further absorption and increase in temperature (1) Alternative: As the ice cap melts leaving darker surface (1) which absorbs more visible radiation and increase in temperature (1) OR Melting permafrost (1) means more methane in atmosphere (1)			2	2		