

Question				Marking details	Marks available					
					AO1	AO2	AO3	Total	Maths	Prac
	(c)			Possible answers: $2 \times 3.2 \text{ [m/s}^2\text{]} = 6.4 \text{ [m/s}^2\text{]}$ (1) $\neq 5.6 \text{ [m/s}^2\text{]}$ so suggestion not correct (1) $16 \text{ [N]} / 5.6 \text{ [m/s}^2\text{]} = 2.9 \text{ [kg]}$ (1) $\neq 2.5 \text{ [kg]}$ ecf from (b)(ii), so suggestion not correct $16 \text{ [N]} / 2.5 \text{ [kg]}$ (ecf from (b)(ii)) = $6.4 \text{ [m/s}^2\text{]}$ (1) $\neq 5.6 \text{ [m/s}^2\text{]}$ so suggestion not correct Alternative: Adding accelerations from forces adding up to 16 N, e.g. $4.8 + 0.8 + 0.8 = 6.4$ (1) $\neq 5.6 \text{ [m/s}^2\text{]}$ so suggestion not correct (1) NB. Use of (4.0 N, 2.6 m/s^2) not penalised in this part, e.g. from 12 N + 4 N $4.8 \text{ [m/s}^2\text{]} + 2.6 \text{ [m/s}^2\text{]} = 7.4 \text{ [m/s}^2\text{]}$ (1) $\neq 5.6 \text{ [m/s}^2\text{]}$ so suggestion not correct. (1) Alternative: Showing clearly that a different force, i.e. 14 N ecf, gives an acceleration of 5.6 m/s^2 (1) $\neq 16 \text{ [N]}$ so suggestion not correct (1)			2	2		
				Question 4 total	1	6	2	9	7	8