

- (b) **Figure 1** shows the number and rate of reported cases of food poisoning in Wales between 1992 and 2016.

Figure 1

Year	Total number of reported cases	Rate (number per 100 000 population)
1992	3 590	124.5
1998	5 946	205.0
2000	4 716	162.2
2006	4 301	145.4
2010	4 980	165.6
2014	4 516	146.1
2016	4 504	145.7

The percentage of reported cases of food poisoning during each quarter of a year is shown in **Figure 2**.

Figure 2

Quarter	% of total for year		
	2012	2014	2016
Jan-Mar	18	17	19
Apr-Jun	26	26	25
Jul-Sep	32	32	33
Oct-Dec	24	25	23

Figure 3 shows the rate of food poisoning amongst females by age group in 2016.



Figure 3

Figure 4 shows the rate of food poisoning amongst males by age group in 2016.

Figure 4



Use the data in **Figures 1 to 4** to answer the following questions.

- (i) Since 1998, egg-laying hens have been vaccinated against a species of *Salmonella*. Explain how this has affected the number of reported cases of food poisoning. [2]

.....

.....

.....

- (ii) Explain the variation in the percentage of reported cases of food poisoning in each of the quarters in a year. [3]

.....

.....

.....

.....

.....

- (iii) It is suggested that the rate of food poisoning does not depend on gender or age. Explain whether this suggestion is confirmed by the data. [2]

.....

.....

.....