

6.




Following flooding in parts of Wales in 2015, environmental scientists investigated nitrate pollution in three different streams. They counted the numbers of eight different indicator species in equal lengths of the three streams, on the same day. The indicator species they counted are shown in the chart below.

GROUP	INDICATOR SPECIES		POLLUTION LEVEL
A	 Mayfly nymph	 Stonefly nymph	If you find these animals in your stream, then there is LITTLE OR NO POLLUTION
B	 Caddis fly larva	 Freshwater shrimp	If you find these animals, but none from Group A, then there may be SLIGHT POLLUTION
C	 Hoglouse	 Bloodworm	If you find these animals, but none from Groups A or B, then there is probably MEDIUM POLLUTION
D	 Worm (<i>Tubifex</i>)	 Rat-tailed maggot	If you find these animals, but none from Groups A, B or C, then there is HIGH POLLUTION
E	No live animals found		If you find no animals at all, then there is VERY HIGH POLLUTION



Examiner
only

Graph showing the number of indicator species
found in each of the streams 1, 2 and 3



(a) (i) Using the graph above and the chart on the opposite page state the pollution level in each of the three streams. [2]

stream 1

stream 2

stream 3

(ii) Suggest **two** possible **sources** of nitrate pollution in streams. [2]

.....

.....

(b) Name **one** type of indicator species which could be used to assess levels of air pollution. [1]

.....

5

