

| Question |     |       |  | Marking details                                                                                                                                                                                                                                                                                                                            | Marks Available |     |     |       |       |      |
|----------|-----|-------|--|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------|-----|-----|-------|-------|------|
|          |     |       |  |                                                                                                                                                                                                                                                                                                                                            | AO1             | AO2 | AO3 | Total | Maths | Prac |
| 3        | (a) | (i)   |  | A. HCl converts pepsinogen into pepsin (1)<br>(ignore activates / reacts with)<br>B. Prevents pepsin being activated inside cells (1)<br>C. Pepsin would {hydrolyse / break down} <u>proteins</u> within the<br>(stomach wall / the chief) cells /<br>Pepsin would cause autolysis of cells (1)<br>(ignore digests)                        |                 | 3   |     | 3     |       |      |
|          |     | (ii)  |  | {Produce / secrete} mucus (1)                                                                                                                                                                                                                                                                                                              | 1               |     |     | 1     |       |      |
|          |     | (iii) |  | <b>Any two (x1) from:</b><br>A. Endopeptidases are secreted {before / owtte} exopeptidases (1)<br>B. Endopeptidases {hydrolyse / break} {peptide bonds in the middle of the {protein / polypeptide chain} / non terminal peptide bonds} (1)<br>C. This provides <u>more</u> {terminal / end} (peptide) bonds for exopeptidases / owtte (1) |                 | 2   |     | 2     |       |      |
|          | (b) |       |  | X: epithelium<br>NOT villus / epidermis / endothelium<br>Y: muscle<br>2 correct = 1 mark<br>1/0 correct = 0 marks                                                                                                                                                                                                                          |                 | 1   |     | 1     |       |      |